

Analysis of first permanent molar condition on Kindergarten student aged 5-6 years old in Eastern Suburban Area of Makassar City

by Admin Admin

Submission date: 20-Apr-2020 06:36PM (UTC+0700)

Submission ID: 1092503655

File name: aty_Yunus,_Hendrastuty_Handayani,_Sumintarti,_Sri_Ramadhany.docx (33.64K)

Word count: 2064

Character count: 11295

Analysis of first permanent molar condition on Kindergarten student aged 5-6 years old in Eastern Suburban Area of Makassar City

Marhamah¹, Rafikah Hasyim², Dwi P. Wulansari³, Muliaty Yunus³, Hendrastuty Handayani¹, Sumintarti⁴, Sri Ramadhany⁵

¹Department of Pediatric Dentistry, Faculty of Dentistry, Hasanuddin University, Makassar

²Department of Oral Biology, Faculty of Dentistry, Hasanuddin University, Makassar

³Department of Dental Radiology, Faculty of Dentistry, Hasanuddin University, Makassar

⁴Department of Oral Medicine Department, Faculty of Dentistry, Hasanuddin University, Makassar

⁵Department of Public Health Department, Faculty of Medicine, Hasanuddin University, Makassar

Abstract

Objective: The aim of this study was to analyze the first permanent molar condition on kindergarten student aged 5-6 years old in eastern suburban area of Makassar city.

Material and method: This study was a descriptive observational study on 100 kindergarten students from four schools in Tamalanrea District, Makassar City. Data was collected with clinical examination and interview. Data was gathered in terms of DMFT status of first permanent molar and OHI-S. Data was analyzed with descriptive statistics and t-test. In this descriptive cross-sectional study. Statistical analysis was performed by descriptive statistics and t-test.

Results: Mean DMFT of first permanent molars was 1.00 ± 1.10 , with a higher value seen among boys. Mean decayed teeth component was 0.95, followed by filled (0.10) and missing (0.0) components.

Conclusion: The observed oral hygiene status among the 5-6 year-olds necessitates implementing pre-ventive as well as restorative measures to improve the oral health status of the young population.

Keywords: First permanent molar, Suburban area, 5-6 years old

13

Corresponding author:

Email: rafikahasyim@gmail.com

p-ISSN: 25030817; e- ISSN: 25030825

DOI: 10.15562/jdmfs.v5i1.1011

Article history

Received 30 September 2019

Revised 5 January 2020

Accepted 28 March 2020

Available online 1 April 2020

Introduction

First permanent molar is the earliest tooth of secondary dentition. First permanent molar erupt when deciduous teeth still exist in the children mouth, thereby included in mixed dentition period. First permanent molar is considered important due to its vital role in stomatognathic system, masticatory efficiency, serve as a guide for the eruption of the remaining molar teeth, and the key in Angle occlusion.¹ The first permanent molar is the most important oral masticatory unit playing a fundamental role in favorable occlusion evolution, and due to its wide occlusal surface, it is more effective in chewing food than any other teeth. Because first permanent molar is erupted during mixed dentition period, parents are often unaware of the

presence of this teeth, thus they have high risk of exposed to caries risk factor thereby often result in teeth destruction or premature loss of first permanent molar.²

Decay of first permanent molar constitutes the biggest component of decayed missing filled tooth index (DMFT) among children and adolescents. Severe caries on first permanent molar often cause pain and infection thus diminished dietary intake and malocclusion.³ Dental caries is one of the most researched oral diseases and results from a chronic process that occurs after a certain amount of exposure to a cariogenic diet and to tooth-susceptible microorganisms. Caries is considered a multifactorial condition that is transmitted locally and is one of the most prevalent pathologies in childhood.⁴

Dental caries is a multifactorial disease that requires a host, nutrient and bacteria which produce acid. The bacterial plaque in the mouth feeds from substrates and produces acid causing tooth decay.

An inappropriate diet of the child and unbalanced consumption of chocolate and sweets, in addition the inability to observe the oral health properly, cause tooth decay. Since children are not aware of the importance of oral hygiene and are not able to brush their teeth properly, it is their parent's responsibility to take care of their children oral and dental health.⁵

Kindergarten students aged 5-6 years old have at least one erupted first permanent molar should received a better attention from their parents in terms of dental and oral health care, because these kids are susceptible to first permanent molar caries due to their habits in chocolate and sweets consumption as well as improper tooth brushing method.

A study from China demonstrated that 75% of children consumed snacks in between meals showed a statistically significant correlation between snacks and high caries prevalence (35%). In Spain, children who brushed their teeth only once a day had a higher frequency (55%) of caries compared with children who brushed twice a day (20%).⁶

Considering the variation in the available information, this study aims to analyze the first permanent molar condition on kindergarten student aged 5-6 years old in eastern suburban area of Makassar city along with the associated factors.

1 Materials and Methods

This study was a descriptive observational study on 100 kindergarten students from four schools in Tamalanrea District, Makassar City. Data was collected with clinical examination. Data was gathered in terms of DMFT status of first permanent molar and OHI-S. Data was analyzed with descriptive statistics and t-test. In this descriptive cross-sectional study, 100 5-6-year-old students were selected randomly. Data was gathered with clinical examination and a questionnaire, collecting data of DMFT of first permanent molars, OHI-S, oral hygiene and nutrition habits. Statistical analysis was performed by descriptive statistics and t-test.

All study samples have at least one first permanent molar erupted. Kindergarten students whom first permanent molar not erupted yet, were excluded from this study. Study samples were examined clinically with a dental mirror and an explorer under artificial light. The students were given oral hygiene instructions and introduced to a clinic to receive dental treatments, if necessary. The evaluated variables included gender, DMFT, OHI-S, frequency of tooth brushing, and nutrition habits.

The simplified oral hygiene index (OHI-S) is a common index, in which six dental surfaces are scored, four from posterior and two from anterior teeth. The scoring can be done fairly rapidly and consistently. OHI-S scores 0-1.2 were considered to indicated good, 1.3-3.0 moderate, and 3.1-6.0 poor oral health status. While the DMFT index is one of the simplest and most commonly used indices in epidemiologic surveys of dental caries. It quantifies dental health status based on the number of decay, missing and filled teeth.

Questionnaire was filled by the examiner by interviewing the study sample in regards of nutrition habits. Data were analyzed with descriptive statistics and t-test using computer software. In this study, P value < 0.05 was considered statistically significant.

Results

A total of 100 kindergarten students aged 5-6 years old were included in this study, but only 60 students were included due to the emergence of their first permanent molar. DMFT score for first permanent molar are presented in table 1.

Table 1 showed there were no missing first permanent molar in study sample, but the decay score were slightly higher in boy compared to the girl.

Table 1. DMFT score for first permanent molars on 5-6 years old kindergarten students

	DMFT		Decayed		Missing		Filling	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Boy	1.21	1.40	1.10	1.01	0.0	0.30	0.10	0.31
Girl	1.01	1.30	0.85	0.96	0.0	0.28	0.10	0.30
Total	1.00	1.10	0.95	1.01	0.0	0.29	0.10	0.29

Table 2. Number (%) of decayed, missing, and filled of DMFT score according to tooth number in 5-6 years old kindergarten students

Teeth	Right upper	Left upper	p value*	Left lower	Right lower	p value*	p value**
	molar (16)	molar (26)		molar (36)	molar (46)		
decayed	8(11.4)	8(11.4)	p=0.100	10(14.29)	14(20)	p=0.08	0.045
filled	2(2.86)	2(2.86)	p=0.100	4(5.71)	3(4.29)	p=0.100	0.051
missing	0 (0)	0 (0)		0 (0)	0 (0)		

* comparison of left and right sides; t-test

** comparison of four first permanent molar; t-test

Table 2 showed number of decayed, missing, and filled teeth of DMFT score in relationship to tooth number in 5-6 years old kindergarten students. Table 2 showed that lower teeth have higher number of decayed as well as filled teeth .

Table 3. Mean DMFT of first permanent molar in relationship with OHI-S

OHI-S	Total	Mean DMFT	SD
Good	22 (31.43%)	1.01	1.00
Moderate	38 (54.29%)	1.98	1.50
Poor	10 (14.29%)	3.10	1.65
Total	70	1.65	1.25

OHI-S score 0-1.2 indicating good, 1.3-3.0 moderate, 3.1-6.0 poor oral health status

Table 3 showed that out of 70 samples, there were 38 have moderate oral health status, 22 have good oral health status, and 10 have poor oral health status, while oral health status in girls (mean OHI-S score, 1.051 ± 0.871) was better than boys (mean OHI-S score, 1.475 ± 1.085).

Discussion

This study was aimed to analyze the condition of first permanent molar on 5-6 years old kindergarten students in Tamalanrea district, one of district in eastern suburban area of Makassar city, the capital of South Sulawesi province.

In this present study, mean DMFT of boys (1.21±1.40) were higher compared to the girls (1.01±1.30). The result of this study was in line with the study by Hiremath⁷ where the mean dmft in girls were 2.88 and boys were 3.15. In explaining the consistent trend of caries rates being higher in females than males, all contributing factors must be considered, such as genetic contribution, saliva, substrate/diet, psychosocial and economic factors, time, as well as systemic correlation.⁸

The result of this study also showed that lower teeth have higher number of decayed as well as filled teeth. The higher amount of caries in the lower first permanent molars could be due to the depth and the form of occlusal pits and fissures, plaque retention as a result of gravity, and a broader masticatory area.⁹

The present study showed that none of the participants had their first permanent molar that were restored with fissure sealants, indicate the underutilization of fissure sealants. And the increased awareness of oral hygiene among individuals over time as well as differences in diet and oral hygiene habits may be reasons for the lower amount of tooth loss in the evaluated samples of this study.

Out of 70 samples, there were 38 have moderate oral health status, 22 have good oral health status, and 10 have poor oral health status, while oral health status in girls (mean OHI-S score, 1.051 ± 0.871) was better than boys (mean OHI-S score, 1.475 ± 1.085).

The observed oral hygiene status among the 5-6-year-olds necessitates implementing preventive as well as restorative measures to improve the oral health status of the young population. The underlying problem should be addressed by emphasizing on prevention of dental caries, which could save lot of resources from a public health perspective. Community based preventive oral health programs on practices of adequate oral hygiene should be implemented. School program should be promoted through integration into the school curriculum and oral health care preventive services to target the growing problem of dental caries among school children.¹⁰

Oral health education should be raised in the community through oral health campaigns and adequate access to public preventive and restorative dental programs and services should be ensured for all the children.

Acknowledgment

None.

18

Conflict of Interest

The authors report no conflict of interest.

References

1. Pontigo-Loyola AP, Márquez-Corona ML, Minaya-Sánchez M, et al. Correlation between the caries status of the first permanent molars and the overall DMFT Index: a cross-sectional study. *Med (Baltimore)* 2020;99: e19061.
2. Espinoza-Espinoza G, Corsini G, Rojas R, et al. The cost-utility of school-based first permanent molar sealants programs: a Markov model. *BMC Oral Health* 2019;19: 293.
3. Borges, Silvana T, Schwanke, et al. Factors associated with caries: a survey of students from southern Brazil. *Rev Paul Pediatr* 2016;34: 489-494.
4. Jetpurwala M. Parental perception of the importance of the permanent first molar in their children. *J Dentist Child* 2020;87: 26-30.

5. Heydari, Alireza. Parental knowledge and awareness of the first permanent molar. *Int J Clin Pediatric Dentist* 2018;11: 382.
6. Maciulskiene, Vita. Estimation of caries treatment needs in first permanent molars of Lithuanian 5-6-year-old children, based on caries lesion activity assessment. *Medicina* 2020;56: 105.
7. Hiremath A, Murugaboopathy V, Ankola AV, et al. Prevalence of dental caries among Primary School Children of India: a cross-sectional study. *J Clin Diagn Res* 2016;10: ZC47–ZC50.
8. Ferraro M, Vieira AR. Explaining gender differences in caries: a multifactorial approach to a multifactorial disease. *Int J Dent* 2010;2010: 649643.
9. Soumya SG, Shashibhushan KK, Pradeep MC, et al. Evaluation of oral health status among 5-15-Year-old School Children in Shimoga City, Karnataka, India: a cross-sectional study. *J Clin Diagn Res* 2017;11: ZC42–ZC47.
10. Nazir MA, Al-Ansari A, Al-Khalifa KS, et al. First permanent molar caries and its association with carious lesions in other permanent teeth. *J Clin Diagnostic Res* 2019;13: ZC36-ZC39.

Analysis of first permanent molar condition on Kindergarten student aged 5-6 years old in Eastern Suburban Area of Makassar City

ORIGINALITY REPORT

23%

SIMILARITY INDEX

18%

INTERNET SOURCES

18%

PUBLICATIONS

12%

STUDENT PAPERS

PRIMARY SOURCES

1

dentistry.tbzmed.ac.ir

Internet Source

1%

2

www.ncbi.nlm.nih.gov

Internet Source

1%

3

worldwidescience.org

Internet Source

1%

4

www.jcdr.net

Internet Source

4%

5

www.elsevier.es

Internet Source

3%

6

academicjournals.org

Internet Source

2%

7

www.jourlib.org

Internet Source

1%

8

Feifei Li, Peipei Jiang, Fanyuan Yu, Chunjie Li, Si Wu, Jing Zou, Xin Xu, Ling Ye, Xuedong

1%

Zhou, Liwei Zheng. "Comparison between Fissure Sealant and Fluoride Varnish on Caries Prevention for First Permanent Molars: a Systematic Review and Meta-analysis", Scientific Reports, 2020

Publication

9

"Free Communication Sessions 01-20 and Poster Sessions 01-16", International Dental Journal, 2015.

Publication

1%

10

downloads.hindawi.com

Internet Source

1%

11

Oystein Fardal. "Tooth loss during maintenance following periodontal treatment in a periodontal practice in Norway", Journal Of Clinical Periodontology, 7/2004

Publication

1%

12

Submitted to University of Sunderland

Student Paper

1%

13

Muhammad Reza Setiawan, Nurdiana Dewi, Ika Kustiyah Oktaviyanti. "Ekstrak ikan haruan (Channa striata) meningkatkan jumlah neokapiler pada penyembuhan luka (Extract of haruan (Channa striata) increases neocapillaries count in wound healing process)", Journal of Dentomaxillofacial Science, 2015

Publication

1%

14 Arghavan Behbahani Rad, Hassan Joulaei, Mehrdad Vossoughi, Ali Golkari. "Assessing Oral Health Status and Behaviors in 6-Year-Old School Children in Rural and Urban Areas of Shiraz, Southern Iran", International Journal of School Health, 2015
Publication 1%

15 América Patricia Pontigo-Loyola, María de Lourdes Márquez-Corona, Mirna Minaya-Sánchez, Salvador Eduardo Lucas-Rincón et al. "Correlation between the caries status of the first permanent molars and the overall DMFT Index", Medicine, 2020
Publication 1%

16 journals.lww.com
Internet Source 1%

17 Submitted to Universitas Brawijaya
Student Paper 1%

18 journals.sagepub.com
Internet Source 1%

19 Submitted to Eiffel Corporation
Student Paper <1%

20 emedicine.medscape.com
Internet Source <1%

21 Yusrini Selviani, Nurul Waqiah Mas'ud, Arsmin Nur Idul Fitri, Atikah Balqis Ferry, Rini F Lestari, <1%

Rasmidar Samad. "Inorganic component of saliva during fasting and after fast break",
Journal of Dentomaxillofacial Science, 2016

Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On