

Article_OSJO_2015_50_2_-
S18_Nurjannah_Nurdin__LAST.
pdf
by

Submission date: 09-Mar-2021 12:52AM (UTC-0800)

Submission ID: 1528264899

File name: Article_OSJO_2015_50_2_-S18_Nurjannah_Nurdin__LAST.pdf (2.67M)

Word count: 3915

Character count: 19490

Multisensor and Multitemporal Data from Landsat Images to Detect Damage to Coral Reefs, Small Islands in the Spermonde Archipelago, Indonesia

Nurjannah Nurdin^{1,3*}, Teruhisa Komatsu², Agus³, M. Akbar AS³, Abdul Rasyid Djalil¹, and Khairul Amri¹

¹Marine Science Department, Hasanuddin University, Makassar 95245, Indonesia

²Atmosphere and Ocean Research Institute, The University of Tokyo, Kashiwa 277-8564, Japan

³Center for Regional Development & Spatial Information, Hasanuddin University, Makassar 95245, Indonesia

Received 26 August 2014; Revised 20 January 2015; Accepted 25 May 2015
© KSO, KIOST and Springer 2015

Abstract—Coral reefs are important because of their high biodiversity and their key role in the tropical marine biosphere. Furthermore, coral reefs are very valuable as a socioeconomic resource as they make important contributions to the gross domestic product of many countries. Thus, it is very important to monitor dynamic spatial distributions of coral reefs and related habitats dominated by coral rubble, dead coral, and bleached corals. Despite these natural and socio-economic advantages, many factors are threatening coral reefs. The study site was selected in Spermonde archipelago, South Sulawesi, Indonesia because this area is included in the Coral Triangle, recognized as the epicenter of coral diversity and a priority for conservation. Images of Landsat MSS, Landsat TM, Landsat ETM, Landsat ETM+, and Landsat 8 data were used to examine changes in the coral reefs of Suranti Island in the Spermonde Archipelago during forty one years from 1972 to 2013. The image processing includes gap fills, atmospheric corrections, geometric corrections, image composites, water column corrections, unsupervised classifications, and reclassifications. Fill Gap processing was done on Landsat ETM+ SLC-off. Subsequently, a multi-component change detection procedure was applied to define changes. Shallow water bottom types classification was divided into live coral, rubble and sand habitats, dead coral with algae, rubble, and sand. Preliminary results showed significant changes during the period 1972-2013 as well as changes in coral reefs, likely explained partly by destructive fishing practices.

Key words – coral reef, landsat, destructive fishing

1. Introduction

The world's most species rich coastal ecosystems occur within the Coral Triangle (Veron et al. 2009), a region which

includes all or part of six Indo-Pacific countries; Indonesia, the Philippines, Malaysia (Sabah), Timor-Leste, Papua New Guinea and the Solomon Islands. Coral reefs are one of the most important ecosystems in marine and coastal areas. Coral reef ecosystems are also important for the fish community and various marine biotas as feeding, nursery, and spawning grounds. Ecologically, a coral reef protects other components of the marine and coastal ecosystem from waves and storms. Compared to other ecosystems, coral reefs can be easily destroyed. The coral reefs in the Coral Triangle have been directly threatened by human activities and natural threats. Coral reef destruction has occurred in the small islands of the Spermonde Archipelago as a result of human activity and rising SST (COREMAP II 2010). Suranti Island is one of these small islands in the Pangkep District. Suranti Island contributes a great deal to the local society in the sense that many local people make a living the shallow water resources. However, the condition of the coral reefs will worsen if the illegal fishing and SST in Suranti Island continues to increase every year. Therefore, spatial dynamic mapping and spatial prediction models of coral reefs are needed to create good spatial planning for coastal areas. Satellite or airborne remote sensing has increasingly been employed to map coral reef communities worldwide (Green et al. 1996; Andrefouet and Guzman 2005). While a range of these studies have recently used high-resolution data, e.g. IKONOS (Knudby et al. 2013; Andrefouet et al. 2003; Riegl and Purkis 2005; Elvidge et al. 2004), Quickbird (Knudby et al. 2013; Botha et al. 2013; Mishra et al. 2006), Worldview (Botha et al. 2013), and CASI (Botha et al. 2013; Mishra et al. 2006), there are

*Corresponding author. E-mail: nurj_din@yahoo.com

number of studies that have used medium spatial resolution data, e.g. Landsat (Barnes et al. 2014; Torres-Pulliza et al. 2013; Palandro et al. 2008). Thematic mapping can be used as a reference for spatial dynamics and habitat distribution of coral reefs. The objective of this research was to analyze the dynamics of coral reefs by using the multisensor (Landsat MSS, TM, ETM, ETM+, and OLI TIRS) and multitemporal data accumulated over 41 years (1972–2013).

2. Data and Method

Study area

Suranti is one of the islands in the Spermonde Archipelago, Pangkep District, Indonesia. Suranti has a land area of 0.044 km². Geographically, Suranti Island is located at longitude 119°8'15.483"E and latitude 04°39'10.135"S.

Data

The Landsat MSS data have four spectral bands, with a spatial resolution of 60 m. Landsat TM data have seven spectral bands with a spatial resolution of 30 m for bands 1–5 and 7. The Landsat ETM+ data consisted of eight spectral bands with a spatial resolution of 30 m for bands 1–7. The Landsat 8 data have nine spectral bands with a spatial resolution of 30 m for bands 1–7 and 9. The following table shows the types of data (Table 1).

Classification bottom types consisting of three classes (live coral, rubble and sand, and dead coral with algae) were examined by Landsat MSS which have a 60 meter resolution and Landsat TM, ETM, ETM+, and OLI that have a 30 meter spatial resolution examined four classes (live coral, rubble, sand, and dead coral with algae).




Fig. 1. Study area in Suranti Island, one of the small islands in Spermonde Archipelago, Indonesia

Table 1. Types of data

N	Satellite	Sensor	Resolution (m)	Acquisition	Path/Row
1	Landsat_1	MSS	60	1972-10-28	122/063
2	Landsat_2	MSS	60	1981-10-26	122/063
3	Landsat_4	TM	30	1990-12-16	114/063
4	Landsat_5	TM	30	1996-04-28	114/063
5	Landsat_7	ETM	30	2002-06-24	114/063
6	Landsat_7	ETM	30	2008-08-19	114/063
7	Landsat_8	OLI_TIRS	30	2013-10-04	114/063

Image processing

The image processing was composed of seven main steps including: (1) gap filling, (2) atmospheric corrections, (3) geometric correction, (4) image composite, (5) water column correction, (6) unsupervised classification, and (7) reclassification.

Filling Gaps

The Landsat 7 ETM+ scan-line corrector is a mechanism designed to correct the under sampling of the primary scan mirror, and it failed in May 2003. This research used the Gap method with Frame and Fill Software developed by Richard Irish at NASA Goddard Space Flight Center. The working principle of this software is to fill gap of Landsat image with an other Landsat image acquired at a different time, as they will have different gaps.

Atmospheric Corrections

This research used simple Dark Object Subtraction to method remove the effects of scattering from the image data. It requires only the information contained in the digital image data. It derives the corrected DN (Digital Number) values solely from the digital data with no outside information (Chavez 1998). Haze DN value is directly selected from the DN frequency Histogram of a digital Image.

Geometric Corrections

The geometric corrections used the 2013 image as references. Landsat image of 1972 and 1981 was corrected using the Landsat 8 image as a reference (Landsat 8 corrected). The corrected image was considered acceptable if RMSE (Root Mean Square Error) is one-half pixel wide (RMSE = 0.5). Overall, RMSE errors of less than 0.5 pixel were achieved for each transformation.

Image Composite (True Color)

The combination band of 321 is used in this interpretation

of Landsat 1, 2, 4, 5, 7 images and the combination band of 432 for Landsat 8. This band combination was used because, in interpretation, the appearance of shallow water bottom types could be seen clearly.

Water Column Correction

Water column correction performed the Lyzenga algorithm for reducing the attenuation effect (Lyzenga 1978):

$$Y_{ij} = \ln(b_i) + \frac{k_i}{k_j} \ln(b_j) \quad (1)$$

$$\frac{K_i}{K_j} = a + \sqrt{(a^2 + 1)} \quad (2)$$

$$a = \frac{\sigma^2 b_i - b^2 b_j}{2\sigma(b_i, b_j)} \quad (3)$$

where Y_{ij} is the result of information extraction channel bottom. Channel bottom are band i and band j that have high penetration into water column, b_i is reflectance on band i , b_j is reflectance on band j , $\frac{K_i}{K_j}$ is the ratio of attenuation coefficient between band i and band j , $\sigma^2 b_i$ is variance of band i , $\sigma^2 b_j$ is variance of band j , $\sigma(b_i, b_j)$ is covariance of band ij . To obtain the attenuation coefficient between the two bands, the selection of 30 sample points of live coral, dead coral with algae rubble, and sand was carried out to calculate the variance value of band i and band j and the covariance to band ij .

Unsupervised Classification

This research used an Unsupervised Classification ISOCCLASS algorithm with 20 unlabelled classes and 100 times of iterations.

Ground truth and reclassification

Ground truth was conducted to determine the actual shallow water bottom types on the ground. The result of the ground truth was compared with the result of classification image. The number of sampling points in the field survey is 83. There

Table 2. Class definitions

Classes	Class Definitions
Live coral	Habitat dominated by a mix of live scleractinian corals
Dead coral with algae	Substrate predominantly made of dead corals covered by algae (turf and fleshy algae)
Rubble	Pieces of broken corals, generally found in reef flats
Sand	Sediment from carbonate origin (dead corals and skeleton for calcifying organisms)
Rubble and sand	Sedimentary mix of rubble and sand

are five dominant classes of shallow water bottom types. They are live coral, rubble and sand, dead coral with algae, rubble, and sand (Table 2).

Post classification


A post-classification change matrix function was applied between 1972 to 1981, 1990 to 1996, 1996 to 2002, 2002 to

Table 3. Band combination and attenuation coefficient

	Satellite/Sensor and acquisition	Band Combination	ki/kj
1	Landsat_4 TM, 1990	1 and 2	0.847919
2	Landsat_5 TM, 1996	1 and 2	0.935253
3	Landsat_7 ETM, 2002	1 and 2	0.893087
4	Landsat_7 ETM, 2008	1 and 2	0.882156
5	Landsat_8 OLI_TIRS, 2013	2 and 3	0.720855

Table 4. Areal estimates of major bottom types in 1972, 1981, 1990, 1996, 2002, 2008, and 2013

Classes	Areal of Suranti Island (Ha)						
	1972	1981	1990	1996	2002	2008	2013
Live Coral	527.66	505.51	325.65	258.61	210.97	143.74	119.46
Rubble	0	0	184.46	248.03	289.15	329.45	336.91
Sand	0	0	60.10	56.10	55.87	55.33	68.39
Dead coral with algae	8.49	14.76	50.74	58.42	65.47	93.23	95.78
Rubble and Sand	59.41	78.60	0	0	0	0	0

**Fig. 2.** Coral reef classification maps of 1972 and 1981

2008, and 2008 to 2013 classification results using the ArcGIS 2008, and 2008 to 2013 classification results using the ArcGIS 10.2.1 software.

Accuracy assessment

Accuracy was tested using confusion matrices of the different classifications. Overall Accuracy (OA), Producer's Accuracy (PA), and User's Accuracy (UA) were computed from the confusion matrices (Foody et al. 1992).

3. Results

In this study, a geometric correction process was performed using 7 point GCP. The result of the geometric correction shows

that the average RMSE obtained from Landsat images was 0.02.

Based on the calculation of digital values that have been extracted from band i and j at the research location. A coefficient value of k_i/k_j was obtained (Table 3).

Annual estimate of changes

Our results show that for this shallow water, coral reef change can be seen by comparing the coral reef maps at different time series. The results of changes in the coral reef areas during the seven acquisition years are represented in Table 4.

Statistically, the change detection obtained from the classified scenes of 1972 to 2013 reveal a significant decrease in the live coral, while dead coral with algae and rubble are increasing every year.




Fig. 3. Coral reef classification maps of 1990, 1996, 2002, 2008, and 2013

Table 5. Areal changes of bottom types in 1972 to 1981, 1990 to 1996, 1996 to 2002, 2002 to 2008, and 2008 to 2013

Bottom types	Areal (Ha)				
	Live coral	Rubble	Dead coral with algae	Sand	Rubble and sand
1972 to 1981					
Live coral	505.51	0.00	6.27	0.00	15.87
Dead coral with algae	0.00	0.00	8.49	0.00	0.00
Rubble and Sand	0.00	0.00	0.00	0.00	59.41
1990 to 1996					
Live coral	258.61	63.08	3.96	0.00	0.00
Rubble	0.00	155.45	25.22	3.78	0.00
Dead coral with algae	0.00	21.50	29.24	0.00	0.00
Sand	0.00	8.00	0.00	52.04	0.00
1996 to 2002					
Live Coral	210.97	45.94	1.70	0.00	0.00
Rubble	0.00	212.01	28.14	7.88	0.00
Dead coral with algae	0.00	22.80	35.63	0.00	0.00
Sand	0.00	8.29	0.00	47.80	0.00
2002 to 2008					
Live Coral	143.74	56.11	11.13	0.00	0.00
Rubble	0.00	240.59	39.79	8.65	0.00
Dead coral with algae	0.00	23.16	42.31	0.00	0.00
Sand	0.00	9.60	0.00	46.26	0.00
2008 to 2013					
Live coral	119.46	23.91	0.36	0.00	0.00
Rubble	0.00	272.58	39.41	17.47	0.00
Dead coral with algae	0.00	37.22	56.01	0.00	0.00
Sand	0.00	3.20	0.00	50.93	0.00

The change matrix (Table 5) shows that 6.27 ha of live coral changed between 1972 to 1981 to dead coral with algae, and 15.87 ha had changed to rubble and sand. The largest category of live coral change was loss to rubble (63.08 ha) from 1990 to 1996. Similar patterns of change were observed from 1996 to 2002, from 2002 to 2008, and from 2008 to 2013. The values are presented in Table 5.

Bottom types of shallow water show the changes in coral reef areas during the period between 1972 and 2013. It can be observed that the changes from 1972 to 2013 show that live coral decreased approximately by 408.19 ha (from 527.66 ha to 119.46 ha), while dead coral with algae increased approximately by 87.29 ha (from 8.49 ha to 95.78 ha), and rubble increased approximately by 345.89 ha (from 59.41 ha to 405.30 ha%). Image classification in 1972 - 2008 is validated by interviews with fisherman who settled around Suranti Island 20 to 40 years ago.

Accuracy assessment

The results of coral reef classification maps were checked

with an accuracy assessment method in order to obtain the level of map accuracy. As previously mentioned, standard map accuracy was more than 70%. Based on the confusion matrix on Table 6, it was found that the OA of the classification image for 2013 in Suranti is was 76.14%. Generally, differences between the classification results and reality in the field are caused by water clarity and spatial resolution of satellite derived data. This accuracy assessment is valid only for the present time (2013).

4. Discussion

Although shallow water may change from year to year due to different human activities and oceanographic parameters, destructive fishing is a significant factor in the substantial decrease in the area of live coral and the increase in dead coral, rubble and sand. The phenomenon of coral bleaching that occurred at the end of 2009 and continued until the middle of 2010 was caused by the La Nina phenomena and increased the SST as a result of the westward movement of




Fig. 4. Post classification of coral reef of Suranti Island: From 1972 to 1981, 1990 to 1996, 1996 to 2002, 2002 to 2008, and 2008 to 2013

the warmpool from the Central Pacific to the Indonesian Seas (Yusuf and Jompa 2012).

Destructive fishing methods such as blast and chemical fishing are still widely practiced in many areas around small islands in the Spermonde Archipelago, Indonesia. Blast fishing

is considered to be ²⁹one of the most destructive anthropogenic threats to coral reefs because of its pernicious effects. Despite national and local government policy to control these illegal methods, destructive fishing is still widely practised by local fishermen. There are several reasons why these practices are

Table 6. Confusion matrix for coral reef classification

Data Classify	Reference Data				Row Total	Commisi	UA
	Live Coral	DCA	Rubble	Sand			
Live Coral	25	3	2	1	31	6	96.15
DCA	0	24	2	0	26	2	80.00
Rubble	0	2	13	0	15	2	72.22
Sand	1	1	1	8	11	3	88.89
Coloum Total	26	30	18	9	83	13	
Ommisi	1	6	5	1	0	13	
PA	80.65	92.31	86.67	72.73			

difficult to eradicate: 1) rapid population growth, particularly in small islands: 2) over-fishing in coastal areas due to limited boat capacity and the traditional fishing gears used by fishermen: 3) poor knowledge of the ecosystem's carrying capacity: 4) weak government control and monitoring. All these issues needed to be seriously addressed and managed by the government to conserve the coastal ecosystems of small islands and marine resources. In order to stop blast and chemical fishing practices, the government should assist local fishermen in seeking alternative livelihoods that provide other means of income. This a serious problem and while many blast fishermen are aware of the destruction their practices entail, they argue that they have no real alternatives.

5. Conclusion

The results indicate that decreases in live coral areas resulted in increased areas of dead coral with algae and rubble during the forty one year period from 1972 to 2013. Live coral from 1972 to 2013 was decreased, while dead coral with algae and rubble were increased. Field observations on the shallow water of Suranti revealed that coral reef habitat destruction was largely caused by human activities. This could be concluded because a lot of loose fragments of rubble that typically result from destructive activities such as explosive fishing (bombs) were found. Another possible cause of the decrease in the area of live corals in the study sites was probably due to the coral bleaching phenomenon which badly affected many Indonesian coral reefs in 2010.

Acknowledgments

We would like to thank Jamaluddin Jompa, Marine Biology Laboratory, Hasanuddin University, Indonesia, for his assistance in preparing the manuscript. We gratefully acknowledge the

sponsors of this study, which was conducted under the network of the Asian CORE Program of the Japan Society for the promotion of Science, "Establishment of research and education network on coastal marine science in Southeast Asia", and the Ocean Remote Sensing Project for Coastal Habitat Mapping (WESTPAC-ORSP: PAMPEC III) of the Intergovernmental Oceanographic Commission for the Western Pacific supported by Japanese Funds-in-Trust provided by the Ministry of Education, Culture, Sports, Science and Technology in Japan.

References

- Andréfouët S, Guzman HM (2005) Coral reef distribution, status and geomorphology-biodiversity relationship in Kuna Yala (San Blas) archipelago, Caribbean Panama. *Coral Reefs* **24**:31–42
- Andréfouët S, Kramer P, Torres-Pulliza D, Joyce KE, Hochberg EJ, Garza-Pérez R, Mumby PJ, Rieglh B, Yamanoi H, Whitej WH, Zubiak M, Brock JC, Phinnd SR, Naseer A, Hatcherl BG, Muller-Karger FE (2003) Multi-site evaluation of IKONOS data for classification of tropical coral reef environments. *Remote Sens Environ* **88**:128–143
- Barnes BB, Hu C, Holkamp KL, Blonski S, Spiering BA, Palandro D, Lapointe B (2014) Use of Landsat data to track historical water quality changes in Florida Keys marine environments. *Remote Sens Environ* **140**: 485–496
- Botha EJ, Brando VE, Anstee JM, Dekker AG, Sagar S (2013) Increased spectral resolution enhances coral detection under varying water conditions. *Remote Sens Environ* **131**: 247–261
- Chavez PS (1988) An improved dark-object subtraction technique for atmospheric scattering correction of multispectral data. *Remote Sens Environ* **24**:459–479
- Contreras-Silva AI, Lopez-Caloca AA, Tapia-Silva FO, Cerdeira-Estrada S (2012) Satellite remote sensing of coral reef habitats mapping in shallow waters at banco chinchorro reefs, México: a classification approach. In: Escalante-Ramirez B (ed) *Remote Sensing - Applications*. Intech, Mexico, pp 331–354
- Coral Reef Rehabilitation and Management Project (2010) Final

- Report: data base of Coral Reef Status in South Sulawesi. Makassar (ID) Coremap II dan Dinas Kelautan dan Perikanan Provinsi Sulawesi Selatan.
- Elvidge CD, Dietz JB, Berkelmans R, Andréfouët S, Skirving W, Strong AE, Tuttle BT (2004) Satellite observation of Keppel Islands (Great Barrier Reef) 2002 coral bleaching using IKONOS data. *Coral Reefs* **23**:123-132
- English S, Wilkinson C, Baker V (1997) Survey manual for tropical marine resources, 2nd edn. Australian Institute of Marine Science, Townsville, 390 p
- Foody GM, Campbell NA, Trodd NM, Wood TF (1992) Derivation and applications of probabilistic measures of class membership from the maximum-likelihood classification. *Photogramm Eng Rem S* **58**(9):1335-1341
- Gomez ED, Yap HT (1988) Monitoring reef condition. In: Kenchington RA, Hudson BET (eds) Coral reef management hand book. UNESCO Regional Office for Science and Technology for South East Asia, Jakarta, pp 171-178
- Green EP, Mumby PJ, Edwards AJ, Clark CD (1996) A review of remote sensing for the assessment and management of tropical coastal resources. *Coast Manage* **24**:1-40
- Knudby A, Jupiter S, Roelfsema C, Lyons M, Phinn S (2013) Mapping coral reef resilience indicators using field and remotely sensed data. *Remote Sens* **5**(3):1311-1334
- Kotta J, Kutser T, Teeveer K, Vahtmäe E, Pärmoja M (2013) Predicting species cover of marine macrophyte and invertebrate species combining hyperspectral remote sensing, machine learning and regression techniques. *PLoS ONE* **8**(6):e63946
- Lyzenga DR (1978) Passive remote sensing techniques for mapping water depth and bottom features. *Appl Optics* **17**:379-383
- Mishra DR, Narumalani S, Rundquist D, Lawson M (2006) Benthic habitat mapping in tropical marine environments using Quickbird multispectral data. *Photogramm Eng Rem S* **72**:1037-1048
- Palandro D, Andréfouët S, Hu C, Hallock P, Müller-Karger FE, Dustan P, Callahan MK, Kranenburg C, Beaver CR (2008) Quantification of two decades of coral reef habitat decline in the Florida Keys National Marine Sanctuary using Landsat data (1984-2002). *Remote Sens Environ* **112**:3388-3399
- Riegl BM, Purkis SJ (2005) Detection of shallow subtidal corals from IKONOS satellite and QTC View (50, 200 kHz) single-beam sonar data (Arabian Gulf; Dubai, UAE). *Remote Sens Environ* **95**:96-114
- Torres-Pulliza D, Wilson JR, Darmawan A, Campbell SJ, Andréfouët S (2013) Ecoregional scale seagrass mapping: a tool to support resilient MPA network design in the coral triangle. *Ocean Coast Manage* **80**:55-64
- Yusuf S, Jompa J (2012) Indonesia First Quantitative Assessment of Coral Bleaching on Indonesian Reefs. In: Proceedings of the 12th International Coral Reef Symposium. Cairns, Australia 9-13 July 2012
- Veron JEN, De Vantier LM, Turak E, Green AL, Kininmonth S, Stafford-Smith M, Petersen N (2009) Delineating the Coral Triangle. *Galaxea J Coral Reef Stud* **11**:91-100

ORIGINALITY REPORT

18%

SIMILARITY INDEX

11%

INTERNET SOURCES

16%

PUBLICATIONS

9%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Chen, Chi-Farn, Nguyen-Thanh Son, Ni-Bin Chang, Cheng-Ru Chen, Li-Yu Chang, Miguel Valdez, Gustavo Centeno, Carlos Thompson, and Jorge Aceituno. "Multi-Decadal Mangrove Forest Change Detection and Prediction in Honduras, Central America, with Landsat Imagery and a Markov Chain Model", *Remote Sensing*, 2013.
Publication 2%
- 2 citeseerx.ist.psu.edu
Internet Source 2%
- 3 Sawayama, Shuhei, Teruhisa Komatsu, Nurjannah Nurdin, Naoto Ebuchi, Delu Pan, and Toshiro Saino. "Coral reef habitats mapping of Spermonde Archipelago using remote sensing compared with in situ survey of fish abundance", *Remote Sensing of the Marine Environment II*, 2012.
Publication 1%
- 4 Torres-Pulliza, Damaris, Joanne R. Wilson, Arief 1%

Darmawan, Stuart J. Campbell, and Serge Andréfouët. "Ecoregional scale seagrass mapping: A tool to support resilient MPA network design in the Coral Triangle", Ocean & Coastal Management, 2013.

Publication

5

researchrepository.murdoch.edu.au

Internet Source

1%

6

Mohammad Shawkat Hossain, Aidy M. Muslim, Muhammad Izuan Nadzri, Abdul Wafi Sabri et al. "Coral habitat mapping: a comparison between maximum likelihood, Bayesian and Dempster–Shafer classifiers", Geocarto International, 2019

Publication

1%

7

jurnal.big.go.id

Internet Source

1%

8

Submitted to University of Plymouth

Student Paper

1%

9

Baoying Ye, Ling Liu. "Remote sensing monitoring land use change in Donglutian coal mine, Shuozhou City", IOP Conference Series: Earth and Environmental Science, 2017

Publication

1%

10

Minaei, Masoud, and Wolfgang Kainz. "Watershed Land Cover/Land Use Mapping

1%

Using Remote Sensing and Data Mining in Gorganrood, Iran", ISPRS International Journal of Geo-Information, 2016.

Publication

11

repository.ipb.ac.id:8080

Internet Source

1%

12

Attachai Kantachumpoo, Shinya Uwai, Thidarat Noiraksar, Teruhisa Komatsu. "Systematics of marine brown alga Sargassum from Thailand: A preliminary study based on morphological data and nuclear ribosomal internal transcribed spacer 2 (ITS2) sequences", Ocean Science Journal, 2015

Publication

<1%

13

M S Thalib, A Faizal, Y A La Nafie. "Remote Sensing Analysis of Seagrass Beds in Bontosua Island, Spermonde Archipelago", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1%

14

Submitted to University of Leicester

Student Paper

<1%

15

Chu Van Cuong, Michael Russell, Sharon Brown, Peter Dart. "Using Shoreline Video Assessment for coastal planning and restoration in the context of climate change in Kien Giang, Vietnam", Ocean Science Journal, 2015

<1%

16 eprints.nottingham.ac.uk <1 %
Internet Source

17 ojs.uajy.ac.id <1 %
Internet Source

18 www.clim-past.net <1 %
Internet Source

19 Chen, Chi-Farn, Va-Khin Lau, Ni-Bin Chang, Nguyen-Thanh Son, Phuoc-Hoang-Son Tong, and Shou-Hao Chiang. "Multi-temporal change detection of seagrass beds using integrated Landsat TM/ETM+/OLI imageries in Cam Ranh Bay, Vietnam", Ecological Informatics, 2016.
Publication

20 José Lucas Safanelli. "Leveraging the application of Earth observation data for mapping and monitoring cropland soils", Universidade de Sao Paulo, Agencia USP de Gestao da Informacao Academica (AGUIA), 2020
Publication

21 Aramita Livia Ardis, Mega Laksmini Syamsudin, Herman Hamdani, Lantun Paradhita Dewanti. "Coral Reef Zone Analysis in Development of Segmentation Ecotourism in Karimunjawa National Park", Asian Journal of Fisheries and

Aquatic Research, 2019

Publication

-
- | | | |
|----|--|-----|
| 22 | epdf.tips
Internet Source | <1% |
|----|--|-----|
-
- | | | |
|----|---|-----|
| 23 | www.sfjo-lamer.org
Internet Source | <1% |
|----|---|-----|
-
- | | | |
|----|--|-----|
| 24 | "Full Issue Article Volume 32, Issue 4",
Fisheries, 2007
Publication | <1% |
|----|--|-----|
-
- | | | |
|----|---|-----|
| 25 | Elizabeth McLeod, Russell Moffitt, Axel Timmermann, Rodney Salm et al. "Warming Seas in the Coral Triangle: Coral Reef Vulnerability and Management Implications",
Coastal Management, 2010
Publication | <1% |
|----|---|-----|
-
- | | | |
|----|--|-----|
| 26 | Salcedo-Sanz, S., L. Cuadra, and M.J.A. Vermeij. "A review of Computational Intelligence techniques in coral reef-related applications",
Ecological Informatics, 2016.
Publication | <1% |
|----|--|-----|
-
- | | | |
|----|---|-----|
| 27 | Xiuling Zuo, Fenzhen Su, Huanting Zhao, Junjue Zhang, Qi Wang, Di Wu. "Regional hard coral distribution within geomorphic and reef flat ecological zones determined by satellite imagery of the Xisha Islands, South China Sea",
Chinese Journal of Oceanology and Limnology, 2016 | <1% |
|----|---|-----|

28

journals.plos.org

Internet Source

<1%

29

nsuworks.nova.edu

Internet Source

<1%

30

www.frontiersin.org

Internet Source

<1%

31

M. H. Ahmed. "Application of remote sensing to site characterisation and environmental change analysis of North African coastal lagoons", *Hydrobiologia*, 04/2009

Publication

<1%

32

Pataporn Kuanui, Suchana Chavanich, Voranop Viyakarn, Makoto Omori, Chiahsin Lin. "Effects of temperature and salinity on survival rate of cultured corals and photosynthetic efficiency of zooxanthellae in coral tissues", *Ocean Science Journal*, 2015

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off