

LEPTIN,_KORTISOL,_SM_GUR U-11-16.pdf

by

FILE	LEPTIN,_KORTISOL,_SM_GURU-11-16.PDF (202.88K)		
TIME SUBMITTED	23-SEP-2020 06:48AM (UTC+0700)	WORD COUNT	3277
SUBMISSION ID	1394338935	CHARACTER COUNT	16342

Leptin and Cortisol: Relationships with Metabolic Syndrome in Male and Female Teachers

Nurzakiah Hasan^{1,2}, Veni Hadju³, Nurhaedar Jafar³, Ridwan Mochtar Thaha⁴

¹ Doctor Program, Faculty of Public Health, Hasanuddin University, Makassar-South Sulawesi, Indonesia, ²Health College Baramuli Pinrang, South Sulawesi, Indonesia, ³Department of Nutrition, Faculty of Public Health, ⁴Health Promotion Department, Faculty of Public Health, Hasanuddin University, Makassar-South Sulawesi, Indonesia

ABSTRACT

Background : Increasing prevalence of metabolic syndrome causes the need for prevention of risk factors and markers, some of them are the role of leptin and cortisol. The aim of this study was to investigate the relationship between leptin and cortisol levels as risk factors of metabolic syndrome among men and women in the teacher group.

Method: A cross section study was performed with 86 teachers (16 men and 70 women). Characteristic sample, Anthropometry, Lipid profile, fasting blood glucose, blood pressure, cortisol and leptin were measured for all samples.

Results: Leptin levels are higher in women than in men (30.64 ± 15.50 vs 7.87 ± 6.02 ; $p=0.005$). While cortisol levels are higher in men than in women (12.09 ± 4.94 vs 8.64 ± 4.15 ; $p<0.001$). Age, stress levels, leptin and cortisol showed a significant association with metabolic syndrome. Leptin correlates significantly with High Density Lipoprotein/ HDL levels ($r=0.391$) for all samples. In men, leptin is significantly correlated with triglycerides/ TG ($r=0.529$) and systolic blood pressure ($r=0.510$), whereas in men, leptin correlates with abdominal circumference ($r=0.479$). Cortisol was significantly correlated with Fasting Blood Glucose/ FBG ($r=0.30$) in all samples. In men cortisol was significantly correlated with Body mass index/ BMI ($r=0.612$) while in women it was significantly correlated with FBG ($r=0.328$).

Conclusions: Leptin levels are higher in women than in men, but cortisol is higher in men than in women. In men, triglyceride levels and systolic blood pressure correlate with an increase in leptin, whereas in women is the abdominal circumference. In men, BMI correlates with cortisol and in women fasting blood glucose levels.

Keyword: markers, metabolic syndrome, leptin, cortisol

INTRODUCTION

Non-communicable diseases (NCD) cause the death of 40 million people each year, equivalent to 70% of deaths globally. The highest causes of death were vascular disease, chronic lung cancer, diabetes, and

other NCD (44,25%; 22%; 9,75%; 4%; and 20%)¹. Of this amount, 85% are in developing countries, one of which is Indonesia.

One of the main risk factors for NCD is metabolic syndrome². The main parameters are blood glucose level, abdominal circumference, blood pressure, HDL levels, and triglyceride levels³. Several markers were then developed to detect an increased risk of metabolic syndrome including cortisol and leptin^{4,5}.

The hormone cortisol is a hormone that is associated with stress, not only in negative conditions, but also

Corresponding author:

Nurzakiah Hasan,
Health College Baramuli, Pinrang, South Sulawesi,
Indonesia, e-mail, nurzakiah15p@student.unhas.ac.id,
081342226001 yumiatinurung7@gmail.com

17 a comfortable and happy condition⁶. Chronic stress is associated with hypercortisolemia and long-term sympathetic nervous system (SNS) activation that results in fat accumulation, especially in the abdomen⁷. Excess fat in the abdomen is one of the parameters of the metabolic syndrome. Identifying risk factors for cortisol is important for stress management as an effort to prevent metabolic syndrome.

Leptin is commonly known as the obese gene. People who are obese have high leptin levels. Leptin is identified as a regulator in regulating body weight. Errors in transportation can cause leptin resistance and cause obesity⁸. A literature review shows that of several markers available, leptin is an appropriate biomarker to identify metabolic syndrome⁹.

Research on metabolic syndrome in Indonesia is still very limited. The results of the analysis of the Riskesdas data 2013 conducted by Nurhaedar Jafar showed that the prevalence of metabolic syndrome was 5.2% which increased along with the increasing prevalence of obesity¹⁰. Research on leptin and cortisol as a marker of metabolic syndrome has never been done before in Indonesia. This study aims to determine the relationship between differences in levels of leptin and cortisol in men and women, the influence of risk factors on metabolic syndrome and risk factors that correlate with leptin and cortisol in men and women in the teacher group.

Method

1 The study with a cross sectional study design was conducted on a group of teachers in Makassar City. This study involved 12 selected schools and was part of a cohort study, educating teachers as an effort to prevent metabolic syndrome.

The number of samples that can be analyzed for this study is 86 people (16 men and 70 women). Demographic characteristics (age and sex), stress levels were measured using a questionnaire through interviews with respondents. Interview and measurement of anthropometry (weight, height, waist circumference/WC) was carried out by trained personnel taken from undergraduate nutrition students public health faculty of Hasanuddin University. Blood collection is carried out by the prodia laboratory.

24 Metabolic syndrome

Metabolic syndrome is defined using criteria from the results of harmonization of several groups in the world. The following are the limits for determining the risk of metabolic syndrome parameters.

12 HDL levels, risk if <40 mg / dl male and <50 mg / dl female

Triglyceride levels, risk if \geq 150 mg / dl

21 Glucose blood sugar levels, risk if fasting blood glucose levels \geq 100 mg / dl

Blood pressure, risk if \geq 130/85 mmHg

6 Abdominal circumference, risk if > 90 cm for men and > 80 cm for women

Blood samples were taken after fasting respondents for 12-14 hours were taken by medical personnel from the Prodia laboratory. HDL examination is carried out by Homogenous Enzymatic Colorimetric Assay method, examination of triglycerides by using enzymatic colorimetric method, whereas fasting blood glucose examination using the Hexokinase method. Blood pressure was measured in the condition of the respondent being seated, and being relaxed using Mercurial Sphygmomanometer.

Leptin and cortisol

Leptin and cortisol were measured using the enzyme immunoassay test method. Reagents used in the Diagnostic Biochem Canada Inc brand, where cortisol uses reagents with Ref can-C-270 and leptin using reagents with Ref: can-L-4260. Leptin and cortisol examinations were carried out at the Hasanuddin University Hospital Laboratory.

RESULT

29 Characteristics of samples based on sex can be seen in Table 1. The average age of male samples is higher than women (50,81 vs 48,89) but the stress level in women is higher than in men (29,32 vs 27,56). There are differences in anthropometry in men and women ($p < 0,001$). 23 There are differences in levels of leptin and 28 cortisol in men and women ($p < 0,001$ and $p = 0,005$). Fasting blood glucose and triglyceride levels do not show the difference between men and women. However, there are significant differences in HDL levels, systolic blood pressure, diastolic and abdominal girth, where women are better than men.

The relationship between how many risk factors for MetS can be seen in table 2. The risk factors for age and stress are higher in respondents who experience MetS than those who are only at risk (consecutive p=0,010 and p=0,026). All MetS parameters show a meaningful relationship with MetS (p<0,05) as well as levels of the hormone leptin and cortisol. Respondents who have lower levels of leptin are at risk of developing metabolic syndrome (p=0,016) and respondents who have higher cortisol levels are at risk of developing metabolic syndrome (p=0,014). The relationship of MetS risk

factors stratified based on sex can be seen in table 3. There were significant differences in age, systolic blood pressure, fasting blood glucose levels, triglyceride levels, HDL levels, between men and women in respondents who experienced metabolic syndrome.

The relationship between leptin and cortisol in several MetS risk factors and MetS parameters can be seen in Table 4. Leptin hormone is significantly associated with HDL levels and abdominal circumference (p <0,05) while cortisol hormone is significantly associated with fasting blood glucose levels (p < 0,001).

Table 1. Characteristics of Samples Based on Sex

	Variable	Men (n=16)	Women (n=70)	P value*
	Age (Mean±SD)	50.81± 3.89	48.89 ± 5.79	0.210
	Stress level (Mean±SD)	27.56±4.77	29.00±8.28	0.506
Anthropometry	weight (Mean±SD)	73.37±6.48	59.69±5.87	<0.001
	height (Mean±SD)	166.20±5.61	153.05±5.51	<0.001
	BMI (Mean±SD)	26.57±1.99	25.49±2.23	0.081
Hormone	Leptin (Mean±SD)	7.87±6.02	30.64±15.50	<0.001
	Cortisol (Mean±SD)	12.09±4.94	8.64±4.15	0.005
MetS parameter	FBG (Mean±SD)	99.81±25.41	94.46±22.95	0.411
	TG (Mean±SD)	176.81±94.95	135.14±67.70	0.113
	HDL (Mean±SD)	42.06±7.51	58.16±10.90	<0.001
	Systole (Mean±SD)	130.00±12.65	119.71±12.74	0.005
	Diastole (Mean±SD)	85.63±6.29	81.00±7.45	0.024
	WC (Mean±SD)	93.42±3.30	86.66±5.06	<0.001

*Sex difference are using T test

Table 2. MetS Risk Factors

	Mets (n=24)	Risk Mets (n=62)	P value
Age	51.21±3.39	48.48±5.99	0.010*
Stress	31.71±12.01	27.58±4.94	0.026*
BMI	26.29±2.21	25.46±2.19	0.122
WC	89.99±5.57	87.11±5.23	0.027*
Systole	130.42±15.17	118,23±10.79	<0.001**
Diastole	84.58±8,84	80.81±6.60	0.034*
FBG	113.58±34.06	88.44±11.99	<0.001**
TG	198.46±84.08	121.39±58.50	<0.001**
HDL	46.08±10.26	58.68±10.90	<0.001**
Leptin	19.46±12.49	29.09±17.50	0.016*
Cortisol	11.17±5.04	8.54±4.07	0.014*

* P < 0,05

** P < 0,001

Table 3. MetS Risk Factor by Sex

	Mets		Risk Mets	
	Male (n=10)	Female (n=14)	Male (n=6)	Female (n=56)
Age	50.40±4.11	51.79±2.77*	51.50±3.73	48.16±6.13
BMI	27.01±1.97	25.77±2.31	25.83±1.99	25.42±2.23
WC	93.62±3.74	87.41±5.29	93.08±2.69	86.47±5.04
Systole	136±11.74*	126.43±16.46*	120.00±6.33	118.04±11.19
Diastole	86.00±6.99	83.57±10.08	85.00±5.48	80.36±6.59
FBG	101.50±31.37	122.21±34.34*	97.00±12.23	87.52±11.69
TG	212.10±103.80*	188.71±69.29*	118.00±31.88	121.75±60.85
HDL	38.80±6.32*	51.29±9.43*	47.50±6.35	59.88±10.64
Leptin	10.09±6.59	26.16±11.42	4.18±2.06	31.76±16.26
Cortisol	12.69±5.52	10.08±4.55	11.08±4.04	8.28±4.01

*P < 0.05

**P < 0,001

Table 4. Correlation Table Between Leptin and Cortisol With MetS Risk Factors and MetS Parameters

	Leptin			Cortisol		
	Men (n=16)	Women (n=70)	Total (n=86)	Men (n=16)	Women (n=70)	Total (n=86)
Stress	-0.263	-0.083	-0.038	0.130	-0.055	-0.051
Spiritual	-0.069	-0.178	-0.198	-0.293	-0.057	-0.063
Age	-0.226	-0.021	-0.099	0.124	-0.059	0.012
FBG	0.172	-0.150	-0.147	0.153	0.328**	0.302*
TG	0.529*	-0.032	-0.095	-0.352	0.027	-0.003
HDL	-0.212	0.180	0.391**	0.074	0.041	-0.120
Sistole	0.510*	-0.008	-0.136	0.007	0.006	0.097
Diastole	-0.078	0.107	-0.053	0.321	-0.089	0.059
WC	-0.074	0.479**	0.074	-0.483	0.014	0.099
BMI	0.420	0.050	0.067	-0.612*	-0.079	-0.083

*P < 0.05

**P < 0,001

DISCUSSION

This study shows the relationship between levels of leptin and cortisol with metabolic syndrome in teachers who are distinguished by sex. Leptin levels are higher in women than in men. In the group that experienced the MetS and risk of MetS, female respondents had higher levels of leptin than men. Leptin hormone levels are associated with obesity. Research conducted in Korea

shows that serum leptin is associated with metabolic syndrome, especially in the body mass index⁵. A meta-analysis conducted by Zeng, et al showed that there was a relationship between leptin and an increased risk of heart and stroke¹³.

Leptin is a hormone associated with regulating food intake and energy balance¹⁴. Leptin is closely related to the level of obesity, where obese people also have higher hormone levels than those who do not obese¹⁵. This

study showed that the average abdominal circumference and BMI of men were higher than women, however, based on the results of the analysis it was seen a positive correlation with the increase in levels of leptin hormone with abdominal circumference in women. The higher the abdominal circumference, the higher the level of leptin hormone. This study is in line with research conducted in Saudi Arabia, where leptin levels are higher in women and are positively correlated with BMI and abdominal circumference¹⁶.

The hormone cortisol shows a significant relationship with the metabolic syndrome, where respondents who experience metabolic syndrome have higher cortisol levels than those at risk. Hormone cortisol is higher in men than in women, as well as in respondents who experience mets and are at risk of MetS, men have higher cortisol levels than women. This is the same as research conducted by Esteghamati, et al in Tehran, which shows high levels of serum cortisol in men compared to women after being justified by age, BMI, and abdominal circumference¹⁷. High cortisol levels are strongly associated with a person's stress level⁷.

The hormone cortisol can be a marker of metabolic syndrome. One mechanism that shows the relationship between metabolic syndrome and cortisol is hypothalamic-pituitary-adrenal (HPA) active in respondents who experience Mets. One of the active activities of HPA is due to sustained levels of stress⁴. One of the factors associated with stress is work¹⁸.

In this study, stress showed a significant relationship with the metabolic syndrome, but did not show a significant relationship with cortisol levels. A meta-analysis was conducted on 29 cross sectional studies by Pan, et al. Which showed that respondents who experienced higher stress had a higher prevalence of metabolic syndrome than those who experienced less stress¹⁹. Some mechanisms that can show this relationship are obesity²⁰, the occurrence of inflammation²¹ and an increase in oxidative stress in respondents who are obese²².

CONCLUSION

Leptin levels are higher in women than in men, but cortisol levels are higher in men than in women. Increased parameters of the metabolic syndrome also increase levels of leptin and cortisol, but there are different parameters that increased in men and women.

This study strengthens that the hormone leptin and the hormone cortisol are markers for the determination of the metabolic syndrome.

Conflict of Interest: There is no any conflict of interest within this study and publication

Ethical Clearance: Taken from Hasanuddin University Ethics Committee with number: 969/H4.8.4.5.31 /PP36-KOMETIK / 2017.

Source of Funding : Ministry of Research, Technology and Higher Education, Indonesia.

REFERENCES

1. WHO. WHO Fact Sheet. 2017; <http://www.who.int/mediacentre/factsheets/fs355/en/>. Accessed 16 April, 2017.
2. Kaur J. A comprehensive review on metabolic syndrome. *Cardiology research and practice*. 2014;2014.
3. Alberti K, Zimmet P, Shaw J. The metabolic syndrome—a new worldwide definition. *Lancet*. 2005;366.
4. Anagnostis P, Athyros VG, Tziomalos K, Karagiannis A, Mikhailidis DP. The Pathogenetic Role of Cortisol in the Metabolic Syndrome: A Hypothesis. *The Journal of Clinical Endocrinology & Metabolism*. 2009;94(8):2692-2701.
5. Uzcátegui E, Valery L, Uzcátegui L, Gomez Perez R, Marquina D, Baptista T. Prevalence of the metabolic syndrome, insulin resistance index, leptin and thyroid hormone levels in the general population of Mérida (Venezuela). *Investigación Clínica*. 2015;56(2).
6. Shier D, Butler J, Lewis R. *Human anatomy and physiology*. McGraw-Hill Boston, MA, USA; 2001.
7. Kyrou I, Tsigos C. Stress hormones: physiological stress and regulation of metabolism. *Current opinion in pharmacology*. 2009;9(6):787-793.
8. Nappo A, Gonzalez-Gil E, Ahrens W, et al. Analysis of the association of leptin and adiponectin concentrations with metabolic syndrome in children: Results from the IDEFICS study. *Nutrition, Metabolism and Cardiovascular Diseases*. 2017;27(6):543-551.
9. Falahi E, Rad AHK, Roosta S. What is the best biomarker for metabolic syndrome diagnosis?

- Diabetes & Metabolic Syndrome: Clinical Research & Reviews. 2015;9(4):366-372.
10. Jafar N. Sindroma metabolik dan epidemiologi. *Media Gizi Masyarakat Indonesia*. 2012;1(2).
 11. Bertoni K, Eckel RH, Grundy SM, et al. Harmonizing the metabolic syndrome a joint interim statement of the international diabetes federation task force on epidemiology and prevention; national heart, lung, and blood institute; American heart association; world heart federation; international atherosclerosis society; and international association for the study of obesity. *Circulation*. 2009;120(16):1640-1645.
 12. Yun JE, Kimm H, Jo J, Jee SH. Serum leptin is associated with metabolic syndrome in obese and nonobese Korean populations. *Metabolism*. 2010;59(3):424-429.
 13. Zeng R, Xu C-H, Xu Y-N, Wang Y-l, Wang M. Association of leptin levels with pathogenetic risk of coronary heart disease and stroke: a meta-analysis. *Arquivos Brasileiros de Endocrinologia & Metabologia*. 2014;58(8):817-823.
 14. Zhang F, Chen Y, Heiman M, DiMarchi R. Leptin: structure, function and biology. *Vitamins & Hormones*. 2005;71:345-372.
 15. Al-Sultan AI, Al-Elq AH. Leptin levels in normal weight and obese Saudi adults. *Journal of family & community medicine*. 2006;13(3):97.
 16. Al-Amodi HS, Abdelbasit NA, Fatani SH, Babakr AT, Mukhtar MM. The effect of obesity and components of metabolic syndrome on leptin levels in Saudi women. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2018;12(3):357-364.
 17. Esteghamati A, Novin L, Nakhjavani M. Association of serum cortisol levels with parameters of metabolic syndrome in men and women. *Clinical and Investigative Medicine (Online)*. 2011;34(3):E131.
 18. Bergmann, Natasha C, Gyntelberg, Finn F, Jens. Chronic stress and the development of the metabolic syndrome: a systematic review of prospective cohort studies. *Endocrine connections*. 2014;EC-14-0031.
 19. Pan A, Keum N, Okereke OI, et al. Bidirectional association between depression and metabolic syndrome: a systematic review and meta-analysis of epidemiological studies. *Diabetes care*. 2012;35(5):1171-1180.
 20. Xu Q, Anderson D, Lurie-Beck J. The relationship between abdominal obesity and depression in the general population: A systematic review and meta-analysis. *Obesity research & clinical practice*. 2011;5(4):e267-e278.
 21. Howren MB, Lamkin DM, Suls J. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosomatic medicine*. 2009;71(2):171-186.
 22. Furukawa S, Fujita T, Shimabukuro M, et al. Increased oxidative stress in obesity and its impact on metabolic syndrome. *The Journal of clinical investigation*. 2004;114(12):1752-1761.

ORIGINALITY REPORT

%20
SIMILARITY INDEX

%13
INTERNET SOURCES

%17
PUBLICATIONS

%2
STUDENT PAPERS

PRIMARY SOURCES

1 worldwidescience.org **%2**
Internet Source

2 f1000research.com **%2**
Internet Source

3 "1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020 **%1**
Publication

4 Ebrahim Falahi, Amir Hossein Khalkhali Rad, Sajjad Roosta. "What is the best biomarker for metabolic syndrome diagnosis?", Diabetes & Metabolic Syndrome: Clinical Research & Reviews, 2015 **%1**
Publication

5 Ji Eun Yun, Heejin Kimm, Jaeseong Jo, Sun Ha Jee. "Serum leptin is associated with metabolic syndrome in obese and nonobese Korean populations", Metabolism, 2010 **%1**
Publication

6 link.springer.com **%1**
Internet Source

7

hal.inrae.fr

Internet Source

% 1

8

Robberecht, Harry, and Nina Hermans. "Biomarkers of Metabolic Syndrome: Biochemical Background and Clinical Significance", *Metabolic Syndrome and Related Disorders*, 2016.

Publication

% 1

9

cardiab.biomedcentral.com

Internet Source

% 1

10

Yıldız Dinçer, Tülay Akçay, Nilgün Çelebi, İlhami Uslu, Özlem Özmen, Hüsrev Hatemi. "Glutathione S-Transferase and O - Methylguanine DNA Methyl Transferase Activities in Patients with Thyroid Papillary Carcinoma ", *Cancer Investigation*, 2002

Publication

% 1

11

Submitted to University of West London

Student Paper

% 1

12

Ghandehari, H.. "Prevalence and extent of dyslipidemia and recommended lipid levels in US adults with and without cardiovascular comorbidities: The National Health and Nutrition Examination Survey 2003-2004", *American Heart Journal*, 200807

Publication

% 1

13

"Track 3 - Disease and Treatment", International Journal of Obesity, 05/2007

Publication

<% 1

14

www.frontiersin.org

Internet Source

<% 1

15

Helen Truby, Sue Baic, Anne deLooy, Kenneth R Fox et al. "Randomised controlled trial of four commercial weight loss programmes in the UK: initial findings from the BBC "diet trials"", BMJ, 2006

Publication

<% 1

16

www.tandfonline.com

Internet Source

<% 1

17

physrev.physiology.org

Internet Source

<% 1

18

cimonline.ca

Internet Source

<% 1

19

iptek.its.ac.id

Internet Source

<% 1

20

Jamaluddin M. Sakung, Saifuddin Sirajuddin, Andi Zulkifli, Stang Abdul Rahm, A. Arsunan Ar, Masni ., Nurhaedar Jafar, Bohari .. "Influence of Nutrition Education and Provision of Instant Chayote to Change in Blood Pressure", Pakistan Journal of Nutrition, 2018

Publication

<% 1

21

www.medicalnewstoday.com

Internet Source

<% 1

22

citeseerx.ist.psu.edu

Internet Source

<% 1

23

Y H Khang. "Trends in socioeconomic health inequalities in Korea: use of mortality and morbidity measures", *Journal of Epidemiology & Community Health*, 2004

Publication

<% 1

24

circ.ahajournals.org

Internet Source

<% 1

25

www.researchsquare.com

Internet Source

<% 1

26

www.ncbi.nlm.nih.gov

Internet Source

<% 1

27

Evangelista, L.S.. "Gender differences in health perceptions and meaning in persons living with heart failure", *Heart & Lung - The Journal of Acute and Critical Care*, 200106

Publication

<% 1

28

Emily Wolfe Phillips, Deepa P. Rao, Leonard A. Kaminsky, Grant R. Tomkinson, Robert Ross, Justin J. Lang. "Criterion-referenced mCAFT cut-points to identify metabolically healthy cardiorespiratory fitness among adults aged 18–

<% 1

69 years: an analysis of the Canadian Health Measures Survey", Applied Physiology, Nutrition, and Metabolism, 2020

Publication

29

eresearch.qmu.ac.uk

Internet Source

<% 1

30

Tohru Funahashi, Yuji Matsuzawa. "Metabolic syndrome: Clinical concept and molecular basis", Annals of Medicine, 2009

Publication

<% 1

31

Yoon, J.H.. "The ratio of serum leptin to adiponectin provides adjunctive information to the risk of metabolic syndrome beyond the homeostasis model assessment insulin resistance: The Korean Genomic Rural Cohort Study", Clinica Chimica Acta, 20111120

Publication

<% 1

32

Sara García-Jiménez, German Bernal Fernández, Maria Fernanda Martínez Salazar, Antonio Monroy Noyola et al. "Serum Leptin is Associated With Metabolic Syndrome in Obese Mexican Subjects", Journal of Clinical Laboratory Analysis, 2015

Publication

<% 1

33

Shivam Champaneri, Gary S. Wand, Saurabh S. Malhotra, Sarah S. Casagrande, Sherita Hill Golden. "Biological Basis of Depression in

<% 1

Adults with Diabetes", Current Diabetes Reports, 2010

Publication

34

vuir.vu.edu.au

Internet Source

<% 1

35

Maria I. Pikilidou, Anastasios N. Lasaridis, Pantelis A. Sarafidis, Ioannis M. Tziolas et al. "Blood Pressure and Serum Potassium Levels in Hypertensive Patients Receiving or Not Receiving Antihypertensive Treatment", Clinical and Experimental Hypertension, 2009

Publication

<% 1

36

H A Elshinnawy, S M Shawky, M A Abdelgawad, M H M Ebrahim. "Assessment of 25-hydroxy vitamen d in patients on regular hemodialysis in ain shams university hospital", QJM: An International Journal of Medicine, 2020

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES

< 5 WORDS