Workflow jurnal Padjung et al. 2021 di Biodiversitas Journal Q3 SJR 0.26

Proses submit



Proses Pre review





Proses Review



Proses review ke 2



Accepted Process



Proses Pembayaran



Proses produksi



Proses copyediting



Published



BIODIVERSITAS
Volume 22, Number 9, September 2021
Pages: 4028-4035

ISSN: 1412-033X
E-ISSN: 2085-4722
DOI: 10.13057/biodiv/d220951

Drought-adapted maize line based on morphophysiological selection index

RUSNADI PADJUNG[1,♥], MUH FARID[1,♥♥], YUNUS MUSA[1], MUHAMMAD FUAD ANSHORI[1], AMIN NUR[2], ANDI MASNENONG[3]

[1]Department of Agronomy, Faculty of Agriculture, Universitas Hasanuddin. Jl. Perintis Kemerdekaan Km 10, Makassar 90245, South Sulawesi, Indonesia. Tel. +62 81355041712, ♥email: rusnadi2015@gmail.com, ♥♥email: farid_deni@yahoo.co.id
[2]Assessment Institute for Agriculture Technology of Gorontalo. Jl. Mohamad Van Gobel No. 270, Ilohelumа, Bone Bolango 96583, Gorontalo, Indonesia
[3]Program of Agrotechnology, Faculty of Agricultural, Universitas Hasanuddin. Jl. Perintis Kemerdekaan Km 10, Makassar 90245, South Sulawesi, Indonesia

Manuscript received: 21 May 2021. Revision accepted: 29 August 2021.

Abstract. *Padjung R, Farid M, Musa Y, Anshori MF, Nur A, Masnenong A. 2021. Drought-adapted maize line based on morphophysiological selection index. Biodiversitas 22: 4028-4035.* Synthetic line formation is an effort to increase maize productivity in drought-stressed areas. This process requires systematic selection in determining adaptability levels involving important secondary