



[Skip to Main Content](#)

Hydrogen peroxide concentration and DNA fragmentation of buffalo oocytes matured in sericin-supplemented maturation medium

by H Hasbi

FILE	MATURED_IN_SERICIN-SUPPLEMENTED_MATURATION_MEDIUM_SAJAS_2019.PDF (244.57K)	WORD COUNT	5141
TIME SUBMITTED	06-JUL-2019 11:19AM (UTC+0700)	CHARACTER COUNT	27357
SUBMISSION ID	1149571585		

Hydrogen peroxide concentration and DNA fragmentation of buffalo oocytes matured in sericin-supplemented maturation medium

S. Gustina¹, N.W.K. Karja^{3*}, H. Hasbi², M.A. Setiadi³ & I. Supriatna³

¹ Division of Animal Science, Faculty of Animal Science and Fisheries, Sulawesi Barat University, Majene 91412, Sulawesi Barat, Indonesia

² Department of Animal Production, Faculty of Animal Science, Hasanuddin University, Makassar 90245, Sulawesi Selatan, Indonesia

³ Division of Reproduction and Obstetrics, Department of Veterinary Clinic, Reproduction, and Pathology, Bogor Agricultural University, Bogor 16680, Jawa Barat, Indonesia

(Received 10 September 2018; Accepted 8 January 2019; First published online 3 April 2019)

Copyright resides with the authors in terms of the Creative Commons Attribution 4.0 South African Licence.

See: <http://creativecommons.org/licenses/by/4.0/za>

Condition of use: The user may copy, distribute, transmit and adapt the work, but must recognise the authors and the South African Journal of Animal Science.

Abstract 49

This study was conducted to investigate the effects of sericin on meiotic maturation, hydrogen peroxide (H₂O₂) concentration and deoxyribonucleic acid (DNA) fragmentation in buffalo oocytes. Oocytes were matured *in vitro* in tissue culture medium (TCM-199) with *in vitro* maturation (IVM) groups under several conditions, namely without bovine serum albumin (-BSA), (+BSA), and 0.025%, 0.05%, 0.1% and 0.25% (*w/v*) sericin. The results showed that supplementation of the maturation medium with 0.05% sericin significantly increased the rate of oocytes that reach metaphase II compared with other groups, except for the 0.025% sericin-treated group. Intracellular H₂O₂ concentrations in oocytes of all groups were measured using 2',7'-dichlorodihydrofluorescein diacetate (DCHFDA). The concentration of H₂O₂ in matured oocytes treated with 0.05% sericin was lower than in other groups. DNA fragmentation and the nuclear status of oocytes were examined using the terminal deoxyribonucleotidyl transferase-mediated dUTP digoxigenin nick end-labelling (TUNEL) method. The total proportion of TUNEL-positive oocytes at the MII stage was lower in the 0.05% sericin group. The results indicate that addition of 0.05% sericin to the maturation medium may improve nuclear maturation and may inhibit DNA fragmentation in oocytes by decreasing H₂O₂ concentrations.

Keywords: antioxidant, apoptosis, *in vitro* maturation, reactive oxygen species

* Corresponding author: karjanwk13@gmail.com

Introduction 57

In buffalo, the efficiency of *in vitro* embryo production (IVEP) is still low (Neglia *et al.*, 2003; Drost, 2007). Approximately only 20% of buffalo oocytes that are fertilized *in vitro* form blastocysts (Gasparrini, 2002; Nandi *et al.*, 2002; Manjunatha *et al.*, 2008). This inefficiency in buffalo IVEP could be associated with the quality of the oocytes at the beginning of the maturation process, although oocytes were selected based on the compaction of cumulus cells and the homogeneity of ooplasm (Chauhan *et al.*, 1998).

One of the critical factors that could affect the quality of oocytes matured *in vitro* is the oxygen tension under which oocytes are cultured. *In vitro* culture environments differ from *in vivo* conditions. The oxygen tension in *in vitro* culture systems (\approx 20%) is often higher than in the female reproductive tract (3% to 9%) (Rocha-Frigoni *et al.*, 2016). This condition is believed to possibly induce the generation of reactive oxygen species (ROS), such as superoxide anions, hydroxyl radicals and hydrogen peroxide (H₂O₂) (Morado *et al.*, 2009). Reactive oxygen species can modify cell function and endanger the survival of cells that cause fragmentation of DNA, protein oxidation and lipid peroxidation (Yang *et al.*, 1998). High oxygen tension during *in vitro* maturation and culture might contribute to decreased oocyte quality (Hashimoto *et al.*, 2000) and alter the development of mammalian embryos (Karja *et al.*, 2004). Several enzymatic and non-enzymatic

antioxidant mechanisms have been developed that protect oocytes and embryos against the harmful effects of ROS (Johnson & Nasr-Esfahani, 1994). Antioxidants have the roles that they can i) prevent the formation of ROS, ii) inactivate oxidants and iii) possibly limit the deleterious effects of oxidants or repair oxidative damage (Agarwal *et al.*, 2004).

Sericin is a family of proteins that covers fibroin, and as the glue in silkworm cocoons (Takahashi *et al.*, 2003). Sericin is known to have a skin moisturizing and antiwrinkle action owing to its high serine content (Engel *et al.*, 1987). Currently, sericin is used for various purposes, including functional biomaterials (Altman *et al.*, 2003; Cao & Zhang, 2016), pharmacological purposes (Prommuak *et al.*, 2008), medical uses (Prasong, 2011; Kunz *et al.*, 2016), and food supplements (Zhang, 2002). Sericin is a high molecular weight protein with adhesive properties (Aramwit *et al.*, 2010) and is derived from silkworm cocoons (Dash *et al.*, 2008). Sericin has ROS-scavenging, anti-tyrosinase, anti-aging (Chlapanidas *et al.*, 2013) and antibacterial properties (Sarovart *et al.*, 2003). It can prevent cell death and supports the growth of cells (Agarwal *et al.*, 2004), increasing the growth of fibroblast cells (Aramwit *et al.*, 2009), and stimulates mammalian cell proliferation (Terada *et al.*, 2002). Sato *et al.* (2011) explained that sericin promotes growth in a variety of cells and affects mitosis in mammalian cells through various signalling pathways. Furthermore, Aramwit *et al.* (2009) reported that with its high methionine content, sericin can promote the fastest growth rate of fibroblast cells. Methionine increases the accumulation of extracellular matrix components such as collagen type I and activates matrix metalloproteinases-9 as an active part in extracellular matrix modelling. Kato *et al.* (1998) demonstrated the antioxidant action of sericin by showing it suppresses *in vitro* lipid peroxidation and inhibits tyrosinase activity in rat brains. Previous studies by Chlapanidas *et al.* (2013) on *in vitro* stimulated peripheral blood mononuclear cells showed that sericin can serve as a ROS-scavenger. Following its addition to the culture medium for embryo production *in vitro*, sericin increased the percentage of maturation and *in vitro* fertilization of sheep oocytes in maturation medium without BSA (Yasmin *et al.*, 2015), and increased the quality and pre-implantation development of bovine embryos that were cultured individually (Isobe *et al.*, 2012). However, the role of sericin in DNA fragmentation or the concentration of H₂O₂ in oocytes matured *in vitro* has not been studied. Therefore, the objective of this study was to evaluate DNA fragmentation and the concentration of H₂O₂ of buffalo oocytes matured in maturation medium supplemented with sericin.

Materials and Methods

Buffalo ovaries were brought from the abattoir to the laboratory in 0.9% NaCl supplemented with 100 IU/mL penicillin and 100 µg/mL streptomycin sulphate. Each ovary was sliced with a scalpel blade to release the cumulus-oocyte complexes in phosphate buffered saline (PBS) supplemented with 0.3% bovine serum albumin (BSA) (Sigma-Aldrich) and antibiotics. Oocytes that were surrounded by compact cumulus layers and uniform cytoplasm (cumulus-oocyte complexes) were used for this study. The basic maturation medium comprised tissue culture medium-199 (TCM-199) (Sigma, USA) supplemented with 10 IU/mL pregnant mare serum gonadotrophin (Intergonan) (Intervet Deutschland GmbH), 10 IU/mL human chorionic gonadotrophin (Chorulo) (International Intervet BV Boxmeer-Holland), and 50 µg/mL gentamycin (Sigma, USA). Selected oocytes were washed three times and divided into groups, namely those matured in maturation medium without BSA (-BSA group), those matured with BSA only (+BSA group), and without BSA but with sericin (Wako Pure Chem. Industries, Ltd. Osaka, Japan) in varying concentrations [0.025% (w/v) (0.025 Ser group), 0.05% (0.05 Ser group), 0.1% (0.1 Ser) or 0.25% (0.25 Ser)]. Oocytes were matured at a temperature of 38.5 °C with 5% CO₂ for 24 hours.

After maturation, oocytes were denuded from the cumulus cells by repeated pipetting in PBS, which contained 0.25% hyaluronidase enzyme (Sigma, USA). After denudation, oocytes were fixed in methanol (1:3 v/v), for 48 to 72 hours and then stained with aceto-orcein (2% orcein and 45% acetic acid). Nuclear maturation was observed under a phase-contrast microscope and classified based on the stage of maturation as germinal vesicle (GV), germinal vesicle breakdown (GVBD), metaphase I (MI), anaphase I (ATI), metaphase II (MII) and degenerated.

The concentrations of H₂O₂ of matured oocytes were measured using DCHFDA (Sigma) as described by Karja *et al.* (2006) with minor modifications. Oocytes in each group were incubated for 15 min in maturation medium containing 10 µM DCHFDA, and then washed in fresh medium before being placed on glass slides and covered with cover slips. Fluorescence emissions were recorded with a digital camera (Zeiss AxioCam HRc, Germany) attached to a fluorescence microscope (Zeiss Axio Imager A2) after excitation at 480 nm and 510 nm emission. The fluorescent image was converted to TIFF files using Adobe Photoshop CS3 (Adobe Systems Inc., San Jose, CA), then analysed with ImageJ software 1.47 (Sun Microsystems, Inc., California, USA). The fluorescent image was measured by counting the number of pixels after colour inversion. The fluorescence intensity represents the intracellular H₂O₂ concentration.

Fragmentation of DNA and nuclear status of matured oocytes in each group were analysed using the combined technique for nuclear staining and TUNEL (*in situ* cell death detection system) (Roche Diagnostics Corporation, Indianapolis, IN, USA) (procedure modified after those described by Wongsrikeao *et al.* (2004)). Oocytes were fixed overnight at 4 °C in 3.7% (wt/vol) paraformaldehyde diluted in PBS. After overnight fixation, the oocytes were washed four times in PBS, which contained 3% (w/v) polyvinyl alcohol (PVA) and permeabilized with 0.5% (v/v) Triton-X100 for 1 hour, and then incubated in a blocking solution (PBS + 10 mg/ml BSA) overnight at 4 °C. After being washed in PBS-PVA, positive controls and all treated oocytes were incubated in fluorescein conjugated dUTP and TdT (TUNEL reagent) at 38.5 °C for 1 hour in the dark. As positive controls, one to two oocytes per TUNEL analysis were incubated in 1000 IU/mL deoxyribonuclease I (DNase I) (Sigma) for 20 min. Negative controls were incubated in fluorescein dUTP without TdT. After TUNEL, oocytes were washed three times in PBS-PVA, and later stained with 50 µg/mL propidium iodide (PI) for 20 min to label all nuclei. Oocytes were washed extensively in blocking solution and placed on a glass slide and covered with a glass cover. The oocytes were examined under a fluorescence microscope (Zeiss Axio Imager A2) using excitation at a wavelength of 488 nm and 568 nm to detect the reaction of TUNEL and PI. The images were captured with a digital camera (Zeiss AxioCam HRc, Germany) (Figure 1). To assess the relationship between nuclear status and DNA fragmentation, chromatin staining of oocytes was used, classify them based on configuration from GV to MII. Those oocytes with diffusely stained cytoplasm characteristics of non-viable cells and those in which chromatin was not visible were excluded from DNA fragmentation analyses.




Figure 1 DNA fragmentation in buffalo oocytes stained by terminal deoxyribonucleotidyl transferase-mediated dUTP-digoxigenin nick end-labelling (TUNEL) (left) and propidium iodide (PI) (right). MII-oocyte TUNEL-negative (red nucleus) with the polar body TUNEL-positive (A,a); MII-oocyte TUNEL-positive (bright nucleus) (B,b); and MII-oocyte TUNEL-negative (C,c)
Scale bars represent 40 µm

The results of the meiotic maturation rate, concentration of H₂O₂, and total proportion of TUNEL-positive oocytes are presented as the mean ± SEM. Data were analysed by one-way analysis of variance (ANOVA). Statistical analysis was performed using SPSS for Windows version 21. *P* ≤ 0.05 was the criterion for statistical significance.

Results

The highest proportion of oocytes reaching MII was found when oocytes were cultured in 0.05% sericin (89.2%) ($P \leq 0.05$) (Table 1). Although the MII rate of oocytes that were cultured in 0.025% sericin was significantly higher than that of oocytes cultured in (-BSA) ($P \leq 0.05$), the rate was not significantly different from oocytes in the (+BSA) group. The proportion of oocytes reaching MII decreased when the concentration of sericin was increased ($P < 0.05$) (79.3%, 77.0%, respectively, for 0.1 and 0.25% groups).

Table 1 Meiotic maturation of buffalo oocytes with supplementation of sericin in maturation medium

Groups of IVM	No. oocytes examined	GV (No)	GVBD (No)	MI (No)	AI/TI (No)	MI (No)	Degenerated (No)
(-BSA)	89	2.4 ± 1.8 (2)	12.8 ^a ± 5.0 (10)	10.2 ± 3.1 (10)	0.0 ± 0.0 (0)	71.9 ^a ± 3.2 (65)	2.6 ± 1.8 (2)
(+BSA)	85	1.1 ± 1.1 (1)	3.1 ^b ± 2.2 (3)	17.6 ± 4.5 (13)	0.0 ± 0.0 (0)	73.8 ^{ab} ± 3.0 (64)	1.9 ± 1.3 (2)
0.025 Ser	84	0.8 ± 0.8 (1)	3.3 ^b ± 1.6 (3)	8.3 ± 2.9 (6)	2.8 ± 1.4 (3)	83.0 ^{bc} ± 2.4 (69)	1.9 ± 1.3 (2)
0.05 Ser	95	0.0 ± 0.0 (0)	2.5 ^b ± 1.3 (3)	7.7 ± 2.2 (8)	0.0 ± 0.0 (0)	89.2 ^c ± 2.6 (83)	0.6 ± 0.6 (1)
0.1 Ser	97	0.6 ± 0.6 (1)	4.4 ^{ab} ± 2.5 (5)	10.8 ± 4.3 (11)	1.0 ± 1.0 (1)	79.3 ^{ab} ± 3.6 (76)	3.9 ± 2.1 (3)
0.25 Ser	87	1.0 ± 1.0 (1)	7.7 ^{ab} ± 4.4 (7)	12.2 ± 4.1 (10)	0.0 ± 0.0 (0)	77.0 ^{ab} ± 4.3 (67)	2.1 ± 1.5 (2)

Results show the mean of percentage ± SE

Means with different superscripts (^{a, b, c}) in the same column differ significantly ($P \leq 0.05$)

Ser: sericin; BSA: bovine serum albumin; AI/TI: anaphase I and telophase I; GV: germinal vesicle; GVBD: germinal vesicle breakdown; MI: metaphase I; MII: metaphase II; No: number of oocytes

Fluorescent photomicrographs of buffalo oocytes stained with DCHFDA are shown in Figure 2. The concentration of intracellular H_2O_2 in oocytes is shown in Figure 3. The concentration of H_2O_2 in oocytes treated with 0.05% sericin (33.6 FIU/oocyte) was significantly lower ($P \leq 0.05$) than in the (-BSA) group (39.9 FIU/oocyte), but not significantly different ($P > 0.05$) from other groups of (+BSA), namely 0.025%, 0.1% and 0.25% (34.4%, 34.7%, 35.1% and 37.9%, respectively).




Figure 2 Fluorescent photomicrographs of buffalo oocytes stained with 2',7'-dichlorodihydrofluorescein diacetate sericin
Scale bars represent 140 μ m




Figure 3 Hydrogen peroxide concentration in buffalo oocytes matured with and without sericin
FIU: values represent fluorescence intensity units
Within each end point, bars with different letters (a, b) are significantly different ($P \leq 0.05$)
The numbers on the bars represent the number of oocytes used for the assay

TUNEL results (Table 2) revealed no significant differences in the proportion of DNA fragmentation in oocytes at all of the nuclear stages after maturation among all groups ($P > 0.05$). However, the 0.05% sericin group has the lowest percentage mean of DNA fragmentation (7.1) in oocytes that remained at the MII stage, whereas the (-BSA), (+BSA), 0.025, 0.1 and 0.25 groups had mean DNA fragmentation rates in oocytes that remained at MII of 15.8, 9.4, 9.1, 9.2 and 10.1, respectively ($P > 0.05$).

Table 2 Nuclear DNA fragmentation of buffalo oocytes matured with sericin

Groups of IVM	No. oocytes examined	Percentage of TUNEL-positive oocytes				
		GV	GVBD	MI	ATI	MII
(-BSA)	58	0.0 ± 0.0 (0/0)	1.6 ± 1.6 (1/1)	10.1 ± 3.5 (6/14)	1.6 ± 1.6 (1/1)	15.8 ± 3.1 (9/42)
(+BSA)	64	0.0 ± 0.0 (0/0)	0.0 ± 0.0 (0/0)	7.8 ± 3.1 (4/13)	1.4 ± 1.4 (1/2)	9.4 ± 3.0 (6/49)
0.025 Ser	65	0.0 ± 0.0 (0/0)	0.0 ± 0.0 (0/0)	6.9 ± 2.9 (5/20)	1.8 ± 1.8 (1/1)	9.1 ± 4.5 (6/44)
0.05 Ser	65	0.0 ± 0.0 (0/0)	1.6 ± 1.6 (1/1)	4.9 ± 2.4 (3/11)	0.0 ± 0.0 (0/0)	7.1 ± 2.9 (5/53)
0.1 Ser	61	0.0 ± 0.0 (0/0)	0.0 ± 0.0 (0/0)	6.1 ± 2.3 (4/18)	1.3 ± 1.2 (1/1)	9.2 ± 3.2 (6/42)
0.25 Ser	63	1.4 ± 1.4 (1/1)	1.8 ± 1.8 (1/1)	3.0 ± 2.0 (2/11)	0.0 ± 0.0 (0/0)	10.1 ± 4.5 (6/50)

Results show the mean of percentage ± SEM, with the number of terminal deoxyribonucleotidyl transferase-mediated dUTP-digoxigenin nick end-labelling (TUNEL)-positive oocytes/total number of examined oocytes
AI/TI: anaphase I and telophase I; GV: germinal vesicle; GVBD: germinal vesicle breakdown
MI: metaphase I; MII: metaphase II

Discussion

During maturation and culturing, various factors can affect the quality and development of oocytes and embryos, especially environmental conditions. Oxygen tension during the culture process is influential on embryo development. Culture conditions under high oxygen tension have a negative effect that can induce the generation of ROS (Rocha-Frigoni et al., 2016). ROS can alter most types of cellular molecules, and block the development of pre-implantation embryos (Guerin et al., 2001).

In this study, the addition of 0.05% sericin significantly increased the percentage of meiotic maturation of buffalo oocytes compared with the (-BSA) and (+BSA) groups. These data are consistent with the findings of Isobe *et al.* (2012) that the addition of 0.5% sericin can improve the quality and preimplantation development of bovine embryos. In sheep, maturation and fertilization rates can be improved with the addition of 0.1% sericin (Yasmin *et al.*, 2015). On the other hand, the authors found that the level of nuclear maturation in the high concentration sericin group (0.25%) showed a significant decrease, possibly owing to a toxic effect of sericin. According to Aramwit *et al.* (2010), sericin can increase collagen production, depending on the concentration, but high concentrations can cause cellular toxicity.

Furthermore, in this study, 0.05% sericin supplementation reduced the concentrations of intracellular H_2O_2 compared with the (-BSA) group. Dash *et al.* (2008) reported that sericin reduced levels of catalase, lactate dehydrogenase and the activity of malondialdehyde (markers of oxidative stress) in fibroblast exposed H_2O_2 . Isobe *et al.* (2012) reported that sericin may have a unique antioxidant potential. The mechanism of the antioxidant effect of sericin is chelation of trace elements, because of its high content of hydroxyl amino acids (serine and threonine) (Kato *et al.*, 1998).

Hydrogen peroxide penetrates biological membranes (Silva & Coutinho, 2010). The mechanism of damage by H_2O_2 in cultured oocytes involves ROS derivatization. Reactive oxygen species can spread and pass through cell membranes and alter cellular molecules, such as proteins, lipids and nucleic acids, resulting in alterations in the mitochondria, embryo cell blockage, depletion of adenosine triphosphate (ATP) and DNA fragmentation (Guerin *et al.*, 2001). DNA fragmentation indicates apoptosis, in which the generation of ROS can result in DNA strand breaks that arrest the cell cycle or even cause cell death (Fahrudin *et al.*, 2002). Therefore, the authors also observed the index of DNA fragmentation in oocyte nuclei matured in sericin. Apoptosis is a normal process during preimplantation embryo development which may be due to environmental stress and chromosome abnormalities (Matwee *et al.*, 2000). Apoptosis is characterized by loss of phospholipid balance in the plasma membrane, chromatin condensation, DNA fragmentation and blebbing of the plasma membrane. In fragmentation, the cell becomes condensed and divides into several fragments, causing cytoplasmic condensation, condensed nuclei (Wyllie *et al.*, 1980), and DNA breakdown, resulting in DNA fragments of 180 - 200 base pairs (Elmore, 2007).

This study demonstrated that although sericin can enhance the number of oocytes that reach MII, there were no significant differences in H_2O_2 concentration or the total of oocytes undergoing DNA fragmentation in all treatment groups. Although the reasons for this finding are currently unclear, the authors presume that it may be due to the effect of glutathione (GSH), which is a natural intracellular antioxidant. Levels of intracellular GSH increase during the oocyte maturation process in the ovary, and peak when oocytes reach MII (Deleuze & Goudet, 2010). Similar to other species, buffalo oocytes can synthesize GSH *de novo* during IVM (Gasparini *et al.*, 2003). Nevertheless, the results of this study show that the addition of sericin to maturation media may serve as a replacement for BSA to increase the maturation rate, and reduce the concentration of H_2O_2 and the proportion of DNA fragmentation. Bovine serum albumin is the main soluble protein in the circulatory system and has many physiological functions. BSA serves as a depot protein and as a transport protein for a variety of compounds (Papadopoulou *et al.*, 2005). It is generally used as a supplement in culture medium. Although BSA can protect cells and has many advantages, there are sanitary risks such as disease transmission, and it may serve as a contamination vector (Thibier, 2006). BSA is not only expensive in the long term, but it leads to enhanced risk of many zoonotic diseases (Cao & Zhang, 2016). This finding is in agreement with other studies, which were reported by Yasmin *et al.* (2015) in sheep oocytes, and Terada *et al.* (2002) in several mammalian cell lines in which sericin was used to accelerate cell proliferation. Sericin may serve as an alternative supplement in serum-free media for cell culture (Tunma *et al.*, 2013). Yanagihara *et al.* (2006) reported that sericin is a powerful alternative to the production of the adenoviral vector in human cells.

Conclusion

In conclusion, the current research findings suggest that supplementation of 0.05% sericin in the maturation medium enhances meiotic maturation of buffalo oocytes. Additionally, sericin can replace BSA as a supplement in serum-free medium, which at 0.05% may decrease the concentration of intracellular H_2O_2 in buffalo oocytes. Further research is warranted to evaluate the effects of sericin supplementation in the quality and development of buffalo embryos.

Acknowledgments

This work was supported by the Ministry of Research, Technology and Higher Education of the Republic of Indonesia (Grant No. 2604/E4.4/2012).

Author's Contributions

SG contributed to the conception and design of the study, data collection, statistical analysis, interpretation of data and drafting the manuscript. HH contributed data collection, statistical analysis and interpretation of data. NWKK, MAS, and IS were involved in conception and design of study, and critical revision of manuscript.

Conflict of Interest Declaration

The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

References

- Agarwal, A., Nallella, K.P., Allamaneni, S.S. & Said, T.M., 2004. Role of antioxidants in treatment of male infertility: An overview of the literature. *Reprod. Biomed. Online*, 8, 616-627.
- Altman, G.H., Diaz, F., Jakuba, C., Calabro, T., Horan, R.L., Chen, J., Lu, H., Richmond, J. & Kaplan, D.L., 2003. Silk-based biomaterials. *Biomaterials* 24, 401-416.
- Aramwit, P., Kanokpanont, S., De-Eknamkul, W., Kamei, K. & Srichana, T., 2009. The effect of sericin with variable amino-acid content from different silk strains on the production of collagen and nitric oxide. *J. Biomater. Sci.* 20, 1295-1306.
- Aramwit, P., Kanokpanont, S., Nakpheng, T. & Srichana, T., 2010. The effect of sericin from various extraction methods on cell viability and collagen production. *Int. J. Mol. Sci.* 11, 2200-2211.
- Cao, T.-T. & Zhang, Y.-Q., 2016. Processing and characterization of silk sericin from *Bombyx mori* and its application in biomaterials and biomedicines. *Mater. Sci. Eng. C* 61, 940-952.
- Chauhan, M.S., Singla, S.K., Palta, P., Manik, R.S. & Tomer, O.S., 1998. Development of in vitro produced buffalo (*Bubalus bubalis*) embryos in relation to time. *Asian-Australas. J. Anim. Sci.* 11, 398-403.
- Chlapanidas, T., Farago, S., Lucconi, G., Perteghella, S., Galuzzi, M., Mantelli, M., Avanzini, M.A., Tosca, M.C., Marazzi, M., Vigo, D., Torre, M.L. & Faustini, M., 2013. Sericins exhibit ROS-scavenging, anti-tyrosinase, anti-elastase, and in vitro immunomodulatory activities. *Int. J. Biol. Macromol.* 58, 47-56.
- Dash, R., Acharya, C., Bindu, P.C. & Kundu, S.C., 2008. Antioxidant potential of silk protein sericin against hydrogen peroxide-induced oxidative stress in skin fibroblasts. *BMB Reports* 41, 236-241.
- Deleuze, S. & Goudet, G., 2010. Cysteamine supplementation of in vitro maturation media: A review. *Reprod. Domest. Anim.* 45, e476-e482.
- Drost, M., 2007. Advanced reproductive technology in the water buffalo. *Theriogenol.* 68, 450-453.
- Elmore, S., 2007. Apoptosis: A review of programmed cell death. *Toxicol. Pathol.* 35, 495-516.
- Engel, W., Hoppe, U., Pape, W. & Sauermaun, G., 1987. Eigenschaften eines sericinhaltingen, puffernden Shampoos. *Arztl. Kosmetol.* 17, 91-110.
- Fahrudin, M., Otoi, T., Karja, N.W.K., Mori, M., Murakami, M. & Suzuki, T., 2002. Analysis of DNA fragmentation in bovine somatic nuclear transfer embryos using TUNEL. *Reproduction* 124, 813-819.
- Gasparini, B., 2002. In vitro embryo production in buffalo species: State of the art. *Theriogenol.* 57, 237-256.
- Gasparini, B., Sayoud, H., Neglia, G., de Matos, D.G., Donnay, I. & Zicarelli, L., 2003. Glutathione synthesis during in vitro maturation of buffalo (*Bubalus bubalis*) oocytes: effects of cysteamine on embryo development. *Theriogenol.* 60, 943-952.
- Guerin, P., El Mouatassim, S. & Menezo, Y., 2001. Oxidative stress and protection against reactive oxygen species in the pre-implantation embryo and its surroundings. *Hum. Reprod. Update.* 7, 175-189.
- Hashimoto, S., Minami, N., Takakura, R., Yamada, M., Imai, H. & Kashima, N., 2000. Low oxygen tension during in vitro maturation is beneficial for supporting the subsequent development of bovine cumulus-oocyte complexes. *Mol. Reprod. Dev.* 57, 353-360.
- Isobe, T., Ikebata, Y., Onitsuka, T., Wittayarat, M., Sato, Y., Taniguchi, M. & Otoi, T., 2012. Effect of sericin on preimplantation development of bovine embryos cultured individually. *Theriogenol.* 78, 747-752.
- Johnson, M.H. & Nasr-Esfahani, M.H., 1994. Radical solutions and cultural problems: could free oxygen radicals be responsible for the impaired development of preimplantation mammalian embryos in vitro? *Bioessays* 16, 31-38.
- Karja, N.W.K., Wongsrikeao, P., Murakami, M., Agung, B., Fahrudin, M., Nagai, T. & Otoi, T., 2004. Effects of oxygen tension on the development and quality of porcine in vitro fertilized embryos. *Theriogenol.* 62, 1585-1595.
- Karja, N.W.K., Kikuchi, K., Fahrudin, M., Ozawa, M., Somjai, T., Ohnuma, K., Noguchi, J., Kaneko, H. & Nagai, T., 2006. Development to the blastocyst stage, the oxidative state, and the quality of early developmental stage of porcine embryos cultured in alteration of glucose concentrations in vitro under different oxygen tensions. *Reprod. Biol. Endocrinol.* 4, 1-12.
- Kato, N., Sato, S., Yamanaka, A., Yamada, H., Fuwa, N. & Nomura, M., 1998. Silk protein, sericin, inhibits lipid peroxidation and tyrosinase activity. *Biosci. Biotechnol. Biochem.* 62, 145-147.
- Kunz, R.I., Brancalhão, R.M.C., Ribeiro, L.d.F.C. & Natali, M.R.M., 2016. Silkworm sericin: Properties and biomedical applications. *Biomed. Res. Int.* 1-19.
- Manjunatha, B., Gupta, P., Ravindra, J., Devaraj, M. & Nandi, S., 2008. In vitro embryo development and blastocyst hatching rates following vitrification of river buffalo embryos produced from oocytes recovered from slaughterhouse ovaries or live animals by ovum pick-up. *Anim. Reprod. Sci.* 104, 419-426.
- Matwee, C., Betts, D.H. & King, W.A., 2000. Apoptosis in the early bovine embryo. *Zygote* 8, 57-68.
- Morado, S.A., Cetica, P.D., Beconi, M.T. & Dalvit, G.C., 2009. Reactive oxygen species in bovine oocyte maturation in vitro. *Reprod. Fertil. Dev.* 21, 608-614.

- Nandi, S., Raghu, H., Ravindranatha, B. & Chauhan, M., 2002. Production of buffalo (*Bubalus bubalis*) embryos in vitro: Premises and promises. *Reprod. Dom. Anim.* 37, 65-74.
- Neglia, G., Gasparini, B., di Brienza, V.C., Di Palo, R., Campanile, G., Presicce, G.A. & Zicarelli, L., 2003. Bovine and buffalo in vitro embryo production using oocytes derived from abattoir ovaries or collected by transvaginal follicle aspiration. *Theriogenol.* 59, 1123-1130.
- Papadopoulou, A., Green, R.J. & Frazier, R.A., 2005. Interaction of flavonoids with bovine serum albumin: A fluorescence quenching study. *J. Agric. Food Chem.* 53, 158-163.
- Prasong, S., 2011. Screening of antioxidant activity of some *Samia ricini* (Eri) silks: Comparison with *Bombyx mori*. *J. Biol. Sci.* 11, 336-339.
- Prommuak, C., De-Eknankul, W. & Shotipruk, A., 2008. Extraction of flavonoids and carotenoids from Thai silk waste and antioxidant activity of extracts. *Sep. Purif. Technol.* 62, 444-448.
- Rocha-Frigoni, N.A.S., Leão, B.C.S., Dall'Acqua, P.C. & Mingoti, G.Z., 2016. Improving the cytoplasmic maturation of bovine oocytes matured in vitro with intracellular and/or extracellular antioxidants is not associated with increased rates of embryo development. *Theriogenol.* 86, 1897-1905.
- Sarovart, S., Sudatis, B., Meesilpa, P., Grady, B.P. & Magaraphan, R., 2003. The use of sericin as an antioxidant and antimicrobial for polluted air treatment. *Rev. Adv. Mater. Sci.* 5, 193-198.
- Sato, W., Fukumoto, K., Yanagihara, K., Sasaki, M., Kunitomi, Y. & Terada, S., 2011. Mitogenic effect of sericin on mammalian cells. *BMC Proc. BioMed Central* P121.
- Silva, J.P., Coutinho, O.P., 2010. Free radicals in the regulation of damage and cell death — basic mechanisms and prevention. *Drug Discov. Ther.* 4, 144-167.
- Takahashi, M., Tsujimoto, K., Yamada, H., Takagi, H. & Nakamori, S., 2003. The silk protein, sericin, protects against cell death caused by acute serum deprivation in insect cell culture. *Biotechnol. Lett.* 25, 1805-1809.
- Terada, S., Nishimura, T., Sasaki, M., Yamada, H. & Miki, M., 2002. Sericin, a protein derived from silkworms, accelerates the proliferation of several mammalian cell lines including a hybridoma. *Cytotechnology* 40, 3-12.
- Thibier, M., 2006. Biosecurity and the various types of embryos transferred. *Reprod. Domest. Anim.* 41, 260-267.
- Tunma, S., Song, D-H., Kim, S-E., Kim, K-N., Han, J-G. & Boonyawan, D., 2013. Immobilization of sericin molecules via amorphous carbon plasma modified-polystyrene dish for serum-free culture. *Appl. Surf. Sci.* 283, 930-940.
- Wongsrikeao, P., Otoi, T., Murakami, M., Karja, N.W.K., Budiyanoto, A., Murakami, M., Nii, M. & Suzuki, T., 2004. Relationship between DNA fragmentation and nuclear status of in vitro-matured porcine oocytes: Role of cumulus cells. *Reprod. Fertil. Dev.* 16, 773-780.
- Wyllie, A.H., Kerr, J.R. & Currie, A., 1980. Cell death: the significance of apoptosis. *Int. Rev. Cytol.* 68, 251-306.
- Yanagihara, K., Terada, S., Miki, M., Sasaki, M. & Yamada, H., 2006. Effect of the silk protein sericin on the production of adenovirus-based gene-therapy vectors. *Biotechnol. Appl. Biochem.* 45, 59-64.
- Yang, H.W., Hwang, K.J., Kwon, H.C., Kim, H.S., Choi, K.W. & Oh, K.S., 1998. Detection of reactive oxygen species (ROS) and apoptosis in human fragmented embryos. *Hum. Reprod.* 13, 998-1002.
- Yasmin, C., Otoi, T., Setiadi, M.A. & Karja, N.W.K., 2015. Maturation and fertilisation of sheep oocytes cultured in serum-free medium containing silk protein sericin. *Acta. Vet. Hung.* 63, 110-117.
- Zhang, Y-Q., 2002. Applications of natural silk protein sericin in biomaterials. *Biotechnol. Adv.* 20, 91-100.

Hydrogen peroxide concentration and DNA fragmentation of buffalo oocytes matured in sericin-supplemented maturation medium

ORIGINALITY REPORT

% **21**
SIMILARITY INDEX

% **10**
INTERNET SOURCES

% **17**
PUBLICATIONS

% **6**
STUDENT PAPERS

PRIMARY SOURCES

- 1** Masashi Nagano, Seiji Katagiri, Yoshiyuki Takahashi. "ATP content and maturational/developmental ability of bovine oocytes with various cytoplasmic morphologies", *Zygote*, 2006
Publication % **1**
- 2** Manabu Ozawa. "Addition of glutathione or thioredoxin to culture medium reduces intracellular redox status of porcine IVM/IVF embryos, resulting in improved development to the blastocyst stage", *Molecular Reproduction and Development*, 08/2006
Publication % **1**
- 3** krishikosh.egranth.ac.in
Internet Source % **1**
- 4** Cao, Ting-Ting, and Yu-Qing Zhang. "Processing and characterization of silk sericin from *Bombyx mori* and its application in biomaterials and biomedicines", *Materials* % **1**

Science and Engineering C, 2016.

Publication

-
- | | | |
|-----------|---|-------------|
| 5 | NI WAYAN KURNIANI KARJA, MOKHAMAD FAHRUDIN, KAZUHIRO KIKUCHI. "Inhibitory Effect of Iodoacetate on Developmental Competence of Porcine Early Stage Embryos In Vitro", HAYATI Journal of Biosciences, 2009
Publication | % 1 |
| <hr/> | | |
| 6 | Aramwit, Pornanong, Sorada Kanokpanont, Wanchai De-Eknamkul, Kaeko Kamei, and Teerapol Srichana. "The Effect of Sericin with Variable Amino-Acid Content from Different Silk Strains on the Production of Collagen and Nitric Oxide", Journal of Biomaterials Science Polymer Edition, 2009.
Publication | % 1 |
| <hr/> | | |
| 7 | Xiao, J.. "CdTe quantum dots (QDs) improve the affinities of baicalein and genistein for human serum albumin in vitro", Journal of Inorganic Biochemistry, 201011
Publication | % 1 |
| <hr/> | | |
| 8 | Submitted to Universiti Putra Malaysia
Student Paper | % 1 |
| <hr/> | | |
| 9 | pubs.sciepub.com
Internet Source | <% 1 |
| <hr/> | | |
| 10 | Systems Biology of Free Radicals and | <% 1 |

Antioxidants, 2014.

Publication

-
- | | | |
|----|---|------|
| 11 | molehr.oxfordjournals.org
Internet Source | <% 1 |
|----|---|------|
-
- | | | |
|----|---|------|
| 12 | Dash, Rupesh, Chitragada Acharya, P.C. Bindu, and S.C. Kundu. "Antioxidant potential of silk protein sericin against hydrogen peroxide-induced oxidative stress in skin fibroblasts", BMB Reports, 2008.
Publication | <% 1 |
|----|---|------|
-
- | | | |
|----|---|------|
| 13 | www.cambridge.org
Internet Source | <% 1 |
|----|---|------|
-
- | | | |
|----|--|------|
| 14 | T Tsujioka. "Effect of Post-IVF Developmental Kinetics on In Vitro Survival of Vitrified-warmed Domestic Cat Blastocysts", Reproduction in Domestic Animals, 6/2008
Publication | <% 1 |
|----|--|------|
-
- | | | |
|----|---|------|
| 15 | ibic.lib.ku.ac.th
Internet Source | <% 1 |
|----|---|------|
-
- | | | |
|----|--|------|
| 16 | Submitted to Heriot-Watt University
Student Paper | <% 1 |
|----|--|------|
-
- | | | |
|----|---|------|
| 17 | b4brain.blogspot.com
Internet Source | <% 1 |
|----|---|------|
-
- | | | |
|----|---|------|
| 18 | Submitted to University of Babylon
Student Paper | <% 1 |
|----|---|------|
-

19 Soottawat Benjakul, Supatra Karnjanapratum, Wonnop Visessanguan. " Hydrolysed collagen from skin: its acute toxicity and impact on cell proliferation and collagen production of fibroblasts ", International Journal of Food Science & Technology, 2018
Publication

<% 1

20 Faranak Aghaz, Mozafar Khazaei, Asad Vaisi-Raygani, Mitra Bakhtiyari. "Cryoprotective effect of sericin supplementation in freezing and thawing media on the outcome of cryopreservation in human sperm", The Aging Male, 2018
Publication

<% 1

21 Submitted to University of Florida
Student Paper

<% 1

22 SASAKI, Masahiro, Hideyuki YAMADA, and Norihisa KATO. "A Resistant Protein, Sericin Improves Atropine-Induced Constipation in Rats.", Food Science and Technology Research, 2000.
Publication

<% 1

23 eprints.ucm.es
Internet Source

<% 1

24 Submitted to Institute of Chemical Technnology, Mumbai
Student Paper

<% 1

25 Submitted to Saint Edwards University <% 1
Student Paper

26 Matteo Bonoli, Vito Verardo, Emanuele Marconi, Maria Fiorenza Caboni. " Antioxidant Phenols in Barley (L.) Flour: Comparative Spectrophotometric Study among Extraction Methods of Free and Bound Phenolic Compounds ", Journal of Agricultural and Food Chemistry, 2004 <% 1
Publication

27 Atef Ali, Marc-André Sirard. "The effects of 17 β -estradiol and protein supplement on the response to purified and recombinant follicle stimulating hormone in bovine oocytes", Zygote, 2002 <% 1
Publication

28 idus.us.es <% 1
Internet Source

29 sutir.sut.ac.th:8080 <% 1
Internet Source

30 R. Verheijen. "Molecular cloning of a major CENP-B epitope and its use for the detection of anticentromere autoantibodies", Molecular Biology Reports, 02/1992 <% 1
Publication

31

Internet Source

<% 1

32

Submitted to Higher Education Commission
Pakistan

Student Paper

<% 1

33

Ting-Ting Cao, Yu-Qing Zhang. "Viability and proliferation of L929, tumour and hybridoma cells in the culture media containing sericin protein as a supplement or serum substitute", *Applied Microbiology and Biotechnology*, 2015

Publication

<% 1

34

refubium.fu-berlin.de

Internet Source

<% 1

35

scialert.net

Internet Source

<% 1

36

Tola, Esra Nur, Muhittin Tamer Mungan, Abdülhadi Cihangir Uğuz, and Mustafa Naziroğlu. "Intracellular Ca²⁺ and antioxidant values induced positive effect on fertilisation ratio and oocyte quality of granulosa cells in patients undergoing in vitro fertilisation", *Reproduction Fertility and Development*, 2013.

Publication

<% 1

37

ijas.pagepress.org

Internet Source

<% 1

epdf.tips

38

Internet Source

<% 1

39

real.mtak.hu

Internet Source

<% 1

40

Ping Qian. "Evaluation of DNA damage and antioxidant capacity of sericin by a DNA electrochemical biosensor based on dendrimer-encapsulated Au-Pd/chitosan composite", *Microchimica Acta*, 01/22/2010

Publication

<% 1

41

espace.curtin.edu.au

Internet Source

<% 1

42

Submitted to Università di Bologna

Student Paper

<% 1

43

B. Roy, S. Rajput, S. Raghav, P. Kumar, A. Verma, A. Jain, T. Jain, D. Singh, S. De, S.L. Goswami, T.K. Datta. " Characterization of oocyte-expressed gene in buffalo and mapping of its TSS and putative regulatory elements ", *Zygote*, 2012

Publication

<% 1

44

www.ncbi.nlm.nih.gov

Internet Source

<% 1

45

etd.lib.metu.edu.tr

Internet Source

<% 1

46

K. Jena, J.P. Pandey, Ruchi Kumari, A.K. Sinha, V.P. Gupta, G.P. Singh. "Tasar silk fiber waste sericin: New source for anti-elastase, anti-tyrosinase and anti-oxidant compounds", International Journal of Biological Macromolecules, 2018

Publication

<% 1

47

link.springer.com

Internet Source

<% 1

48

Agung, B.. "Effects of oxygen tension and follicle cells on maturation and fertilization of porcine oocytes during in vitro culture in follicular fluid", Theriogenology, 20100415

Publication

<% 1

49

academic.oup.com

Internet Source

<% 1

50

C. Matwee. "The effects of antibodies to heat shock protein 70 in fertilization and embryo development", Molecular Human Reproduction, 09/01/2001

Publication

<% 1

51

Jo, Yoon Nam, Do Gyu Bae, and In Chul Um. "The Effect of Extraction Conditions and Film Side on the Molecular Conformation of Silk Sericin Film", International Journal of Industrial Entomology, 2013.

Publication

<% 1

52

K.M. Banwell, M. Lane, D.L. Russell, K.L. Kind, J.G. Thompson. "Oxygen concentration during mouse oocyte in vitro maturation affects embryo and fetal development", Human Reproduction, 2007

Publication

<% 1

53

Catherine Guyader-Joly. "Precursors of taurine in female genital tract: Effects on developmental capacity of bovine embryo produced in vitro", Amino Acids, 03/1998

Publication

<% 1

54

MA Gil. "Advances in Swine In Vitro Embryo Production Technologies : Advances in Swine In Vitro Embryo Production", Reproduction in Domestic Animals, 05/17/2010

Publication

<% 1

55

F. Marco-Jiménez, J.S. Vicente, M.P. Viudes-de-Castro. " Effect of lanosterol on the maturation in semi-defined culture system of prepubertal ewe oocytes ", Zygote, 2011

Publication

<% 1

56

A.M. Brad. "Glutathione and adenosine triphosphate content of in vivo and in vitro matured porcine oocytes", Molecular Reproduction and Development, 04/2003

Publication

<% 1

Satoshi Sugimura, Manabu Kawahara, Takuya

57

Wakai, Ken-ichi Yamanaka, Hiroshi Sasada, Eimei Sato. "Effect of cytochalasins B and D on the developmental competence of somatic cell nuclear transfer embryos in miniature pigs", *Zygote*, 2008

Publication

<% 1

58

Mattioli, Samantha, Sabata Martino, Francesco D'Angelo, Carla Emiliani, Josè Maria Kenny, and Ilaria Armentano. "Nanostructured polystyrene films engineered by plasma processes: Surface characterization and stem cell interaction", *Journal of Applied Polymer Science*, 2014.

Publication

<% 1

EXCLUDE QUOTES ON
EXCLUDE ON
BIBLIOGRAPHY

EXCLUDE MATCHES < 5
WORDS