

B-2

Angle in the space of p - summable sequences

by

Submission date: 26-Nov-2021 12:41PM (UTC+0700)

Submission ID: 1712964579

File name: 10.3934_math.2022155.pdf (227.25K)

Word count: 3591

Character count: 14557



Research article

Angle in the space of p -summable sequences

Muh Nur^{1,*}, Moch Idris² and Firman¹

¹ Department of Mathematics, Hasanuddin University, 90245, Indonesia

² Department of Mathematics, Universitas Lambung Mangkurat, 70714, Indonesia

* Correspondence: Email: muhammadnur@unhas.ac.id.

Abstract: The aim of this paper is to investigate completeness of A that equipped with usual norm on p -summable sequences space where A is subspace in p -summable sequences space and $1 \leq p < \infty$. We also introduce a new inner product on A and prove completeness of A using a new norm that corresponds this new inner product. Moreover, we discuss the angle between two vectors and two subspaces in A . In particular, we discuss the angle between 1-dimensional subspace and $(s - 1)$ -dimensional subspace where $s \geq 2$ of A .

Keywords: completeness; vectors; subspaces; ℓ^p spaces; angles

Mathematics Subject Classification: 15A03, 46B20, 46A45, 51N15

1. Introduction

In an inner product space, we can calculate angles between two subspaces. Let $(X, \langle \cdot, \cdot \rangle)$ be a real inner product space. If $U = \text{span}\{u\}$ is a 1-dimensional subspace and $V = \text{span}\{v_1, \dots, v_r\}$ is a r -dimensional subspace of X , then the angle between subspaces U and V is defined by θ with $0 \leq \theta \leq \frac{\pi}{2}$ and $\cos^2 \theta = \frac{\langle u, u_V \rangle^2}{\|u\|^2 \|u_V\|^2}$. In formula, u_V denotes the (orthogonal) projection of u on V and $\|\cdot\| = \langle \cdot, \cdot \rangle^{1/2}$. Gunawan et al. [3] show that the value of $\cos \theta$ is equal to the ratio between the length of the projection of u on V and the length of u ($\cos^2 \theta = \frac{\|u_V\|^2}{\|u\|^2}$). Likewise, if $U = \text{span}\{u, w_2, \dots, w_s\}$ and $V = \text{span}\{v, w_2, \dots, w_s\}$ are s -dimensional subspaces of X that intersects on $(s - 1)$ -dimensional subspace $W = \text{span}\{w_2, \dots, w_s\}$ with $s \geq 2$ then the angle between U and V is defined by θ with $0 \leq \theta \leq \frac{\pi}{2}$ and $\cos^2 \theta = \frac{\langle u_W^\perp, v_W^\perp \rangle^2}{\|u_W^\perp\|^2 \|v_W^\perp\|^2}$ with u_W^\perp, v_W^\perp are the orthogonal complement of u and v , respectively, on W . Gunawan et al. [3] show that the value of $\cos \theta$ is equal to the ratio between the volume of the s -dimensional parallelepiped spanned by the projection of u, w_2, \dots, w_s on V and the volume of the s -dimensional parallelepiped spanned by u, w_2, \dots, w_s .

Now suppose $(X, \|\cdot\|)$ is a normed space. As it is known, not all normed spaces are inner product

spaces. For instance, the space ℓ^p for $1 \leq p < \infty$ with norm $\|x\|_p = \left[\sum_{n=1}^{\infty} |x_n|^p \right]^{\frac{1}{p}}$ is not an inner product space except for $p = 2$ (see [6]). Konca et al. [5] define a (weighted) inner product $\langle \cdot, \cdot \rangle_v$ on ℓ^p where $2 < p < \infty$, but $(\ell^p, \langle \cdot, \cdot \rangle_v)$ is not Banach space. They also find that the inner product is actually defined on a larger space namely $\ell_{2,v}$ that contains ℓ^p and $(\ell_{2,v}, \langle \cdot, \cdot \rangle_v)$ is Banach space. More recent works may be found in [4, 8, 12]

Now suppose $(A, \|\cdot\|)$ is a normed space where A is subspace of X . Two questions arise: What are A complete? Can we define a norm on A which satisfies the parallelogram law? The reason why we are interested in the parallelogram law is because we eventually wish to define an inner product, so that we can define angle between two subspaces and many other notions on this space. The concept of the angle between two vectors and two subspaces in normed space has been studied intensively, see [1, 2, 7, 9–11].

Let $(\ell^p, \|\cdot\|_p)$ be a normed space and A is subspace in ℓ^p for $1 \leq p < \infty$. In this paper, we discuss completeness of A that equipped with usual norm on ℓ^p . We also introduce a new inner product $\langle \cdot, \cdot \rangle_{b,2}$ on A such that $(A, \|\cdot\|_{b,2})$ is complete where $\|x\|_{b,2} = (\langle x, x \rangle_{b,2})^{\frac{1}{2}}$ denotes the induced norm in A . Motivated by this fact, we shall discuss the angle between two subspaces in A .

2. Main results

2.1. The completeness of subspaces on ℓ^p

Let $\{x_1, \dots, x_n\}$ be linearly independent set on ℓ^p for $1 \leq p < \infty$. Define $A = \text{span}\{x_1, \dots, x_n\}$. Then we observe that $(A, \|\cdot\|_p)$ is a subspace of $(\ell^p, \|\cdot\|_p)$. [6], we know that ℓ^p is Banach space. Here, A will be proved as Banach space. Before this result, we have the following lemma.

Lemma 2.1. Let $C = \sum_{i=1}^n a_i$ where $a_1, \dots, a_n \in \mathbb{R}$.

(a) If $\text{sign}(a_j) = \text{sign}(C)$ for $j = 1, \dots, n$ then the equality

$$|ta_j| + \left| (1-t)a_j + \sum_{\substack{i=1 \\ i \neq j}}^n a_i \right| = |C|$$

holds for some $t \in (0, 1)$.

(b) If $\text{sign}(a_j) = -\text{sign}(C)$ for $j = 1, \dots, n$ then the equality

$$|ta_j| + \left| (t+1)a_j + \sum_{\substack{i=1 \\ i \neq j}}^n a_i \right| = |C|$$

holds for some $t \in (0, 1)$.

Proof. (a) Without loss of generality, writing $j = 1$. Suppose that $\text{sign}(a_1) = \text{sign}(C)$. Choose $t \in (0, 1)$ such that

$$D = (1-t)a_1 + \sum_{i=2}^n a_i.$$

and $\text{sign}(D) = \text{sign}(C)$. We can write $C = ta_1 + D$. Hence, $|C| = |ta_1| + |D|$.

(b) Without loss of generality, writing $j = 1$ and $\text{sign}(C) = 1$. Suppose that $\text{sign}(a_1) = -\text{sign}(C)$. Choose $t \in (0, 1)$ such that

$$\begin{aligned} C &= -ta_1 + (1+t)a_1 + \sum_{i=2}^n a_i \\ &= -ta_1 + E \end{aligned}$$

and $\text{sign}(E) = \text{sign}(C)$. We observe that $C, -ta_1, E$ are positive real. Hence, we can write $|C| = |ta_1| + |E|$. \square

Using Lemma 2.1, We shall now show that $(A, \|\cdot\|_p)$ is Banach space.

Theorem 2.2. Let $\{x_1, \dots, x_n\}$ be linearly independent set on ℓ^p where $1 \leq p < \infty$. If $A = \text{span}\{x_1, \dots, x_n\}$ then $(A, \|\cdot\|_p)$ is Banach space.

Proof. We consider any Cauchy sequence (y_k) in A , writing $y_k = \alpha_{k1}x_1 + \alpha_{k2}x_2 + \dots + \alpha_{kn}x_n$. Since (y_k) is Cauchy sequence, we have for every $\epsilon > 0$ there exist N_ϵ such that

$$\begin{aligned} \|y_k - y_l\|_p &= \left\| \sum_{i=1}^n (\alpha_{ki} - \alpha_{li})x_i \right\|_p \\ &= \left(\sum_{j=1}^{\infty} \left| \sum_{i=1}^n (\alpha_{ki} - \alpha_{li}) \zeta_{ij} \right|^p \right)^{\frac{1}{p}} < \epsilon \end{aligned}$$

for all $k, l > N_\epsilon$. It follows that for every $j = 1, 2, \dots$ we have

$$\left| \sum_{i=1}^n (\alpha_{ki} - \alpha_{li}) \zeta_{ij} \right| < \epsilon_1.$$

Case 1. For some $i = 1, \dots, n$, $\text{sign} \left(\sum_{i=1}^n (\alpha_{ki} - \alpha_{li}) \zeta_{ij} \right) = \text{sign}(\alpha_{ki} - \alpha_{li}) \zeta_{ij}$ holds. By Lemma 1, we can find $t \in (0, 1)$ such that $|t(\alpha_{ki} - \alpha_{li}) \zeta_{ij}| < \epsilon_1$.

Case 2. For some $i = 1, \dots, n$, $\text{sign} \left(\sum_{i=1}^n (\alpha_{ki} - \alpha_{li}) \zeta_{ij} \right) = \text{sign}(\alpha_{ki} - \alpha_{li}) \zeta_{ij}$ holds. By Lemma 1, we can find $t \in (0, 1)$ such that $|t(\alpha_{ki} - \alpha_{li}) \zeta_{ij}| < \epsilon_1$.

So, we obtain $|t\zeta_{ij}| |(\alpha_{ki} - \alpha_{li})| < \epsilon_1$ for all $k, l \geq N_\epsilon$. Thus, for each fixed $i \in \{1, \dots, n\}$, (α_{ki}) is a Cauchy sequence of real numbers. Hence, it is convergent, say $\alpha_{ki} \rightarrow \alpha_i$ as $k \rightarrow \infty$. Now, we can view that $\alpha = (\alpha_1, \dots, \alpha_n)$ and we define $y := \alpha_1 x_1 + \dots + \alpha_n x_n$. Its obvious that $y \in A$. Because α_k converge to α then y_k converge y . Hence, $(A, \|\cdot\|_p)$ is Banach Space. \square

Next, we discuss a new inner product in ℓ^p and prove that A with the new inner product is Hilbert space. Write $b = \sum_{i=1}^n |x_i|$. We define a following mapping

$$\langle y, z \rangle_{b,2} := \sum_{k=1}^{\infty} b_k^{p-2} y_k z_k \quad (2.1)$$

for every $y, z \in \ell^p$. Using Hölder's inequality, we have

$$\sum_{k=1}^{\infty} b_k^{p-2} y_k z_k \leq \left[\sum_{k=1}^{\infty} b_k^p \right]^{\frac{p-2}{p}} \left[\sum_{k=1}^{\infty} y_k^p \right]^{\frac{1}{p}} \left[\sum_{k=1}^{\infty} z_k^p \right]^{\frac{1}{p}}.$$

Hence, the mapping $\langle \cdot, \cdot \rangle_{b,2}$ is defined on ℓ^p . Next, we have the following proposition.

Proposition 2.3. *The mapping $\langle \cdot, \cdot \rangle_{b,2}$ in (2.1) defines an inner product on ℓ^p .*

Proof. For every $y, z, w \in \ell^p$, we will verify that $\langle \cdot, \cdot \rangle_{b,2}$ satisfies the four properties of an inner product.

1) Since $\sum_{k=1}^{\infty} b_k^{p-2} y_k^2 \geq 0$, we have $\langle y, y \rangle_{b,2} \geq 0$.

Next, we show that $\langle y, y \rangle_{b,2} = 0$ if only if $y = 0$. Suppose that $y = 0$, then

$$\langle 0, 0 \rangle_{b,2} = \sum_{k=1}^{\infty} b_k^{p-2} 0 = 0.$$

Conversely, if $\langle y, y \rangle_{b,2} = 0$, then $\sum_{k=1}^{\infty} b_k^{p-2} y_k^2 = 0$. Since $b_k^{p-2} \neq 0$, we obtain $y = 0$.

2) Observe that

$$\langle y, z \rangle_{b,2} = \sum_{k=1}^{\infty} b_k^{p-2} y_k z_k = \sum_{k=1}^{\infty} b_k^{p-2} z_k y_k = \langle y, z \rangle_{b,2}.$$

3) Observe that

$$\langle \alpha y, z \rangle_{b,2} = \sum_{k=1}^{\infty} b_k^{p-2} \alpha y_k z_k = \alpha \langle y, z \rangle_{b,2}.$$

4) Observe that

$$\begin{aligned} \langle y, z + w \rangle_{b,2} &= \sum_{k=1}^{\infty} b_k^{p-2} y_k (z_k + w_k) \\ &= \sum_{k=1}^{\infty} b_k^{p-2} y_k z_k + \sum_{k=1}^{\infty} b_k^{p-2} y_k w_k \\ &= \langle y, z \rangle_{b,2} + \langle y, w \rangle_{b,2}. \end{aligned}$$

Therefore $\langle \cdot, \cdot \rangle_{b,2}$ defines an inner product on ℓ^p . □

Corollary 2.4. *The following function*

$$\|y\|_{b,2} = \left[\sum_{k=1}^{\infty} b_k^{p-2} y_k^2 \right]^{\frac{1}{2}} \quad (2.2)$$

defines a norm that corresponds to the inner product $\langle \cdot, \cdot \rangle_{b,2}$ on ℓ^p .

Using a norm that corresponds to the inner product $\langle \cdot, \cdot \rangle_{b,2}$, we have the following result.

Theorem 2.5. The space $(A, \|\cdot\|_{b,2})$ is complete. Accordingly, $(A, \langle \cdot, \cdot \rangle_{b,2})$ is a Hilbert space.

Proof. We consider any Cauchy sequence (y_k) in A , writing $y_k = \alpha_{k1}x_1 + \dots + \alpha_{kn}x_n$ and $\alpha_k = (\alpha_{k1}, \dots, \alpha_{kn})$. Since y_k is Cauchy, for every $\epsilon > 0$ there is an N_ϵ such that for all $k, l > N_\epsilon$,

$$\begin{aligned} \|y_k - y_l\|_{b,2} &= \left\| \sum_{i=1}^n (\alpha_{ki} - \alpha_{li}) x_i \right\|_{b,2} \\ &= \left[\sum_{j=1}^{\infty} \sum_{i=1}^n b_j^{p-2} ((\alpha_{ki} - \alpha_{li}) \zeta_{ij})^2 \right]^{\frac{1}{2}} < \epsilon. \end{aligned}$$

It follows that for every $j = 1, 2, \dots$ we have

$$b_j^{p-2} \sum_{i=1}^n (\zeta_{ij} \alpha_{ki} - \zeta_{ij} \alpha_{li})^2 < \epsilon^2.$$

So, we obtain $|\zeta_{ij}| b_j^{\frac{p-2}{2}} |\alpha_{ki} - \alpha_{li}| < \epsilon$ for all $k, l \geq N_\epsilon$. Thus, for each fixed $i \in \{1, \dots, n\}$, (α_{ki}) is a Cauchy sequence of real numbers. Hence, it is convergent, say $\alpha_{ki} \rightarrow \alpha_i$ as $k \rightarrow \infty$. Now, we can view that $\alpha = (\alpha_1, \dots, \alpha_n)$ and we define $y := \alpha_1 x_1 + \dots + \alpha_n x_n$. It's obvious that $y \in A$. Because α_k converge to α then y_k converge y . Hence, $(A, \|\cdot\|_{b,2})$ is complete. \square

2.2. Angle between two subspaces on A

We know that $(A, \langle \cdot, \cdot \rangle_{b,2})$ is a Hilbert space with $A = \text{span}\{x_1, \dots, x_n\}$ and, as before, $\{x_1, \dots, x_n\}$ is a linearly independent set on ℓ^p . Using the Gram-Schmidt process, we have an orthonormal set $\{y_1, \dots, y_n\}$. As a consequence, $\text{span}\{x_1, \dots, x_n\} = \text{span}\{y_1, \dots, y_n\}$. For every $u, v \in A$, we can write $u = \sum_{i=1}^n c_i y_i$ and $v = \sum_{i=1}^n d_i y_i$ where $c_i, d_i \in \mathbb{R}$ for every $i = 1, \dots, n$. Moreover,

$$\begin{aligned} \langle u, v \rangle_{b,2} &= \left\langle \sum_{i=1}^n c_i y_i, \sum_{i=1}^n d_i y_i \right\rangle_{b,2} \\ &= \sum_{i=1}^n \sum_{j=1}^n c_i d_j \langle y_i, y_j \rangle_{b,2} \end{aligned}$$

Because $\langle y_i, y_j \rangle_{b,2} = 0$ for $i \neq j$ and $\langle y_i, y_i \rangle_{b,2} = \|y_i\|_{b,2}^2$ then

$$\begin{aligned} \langle u, v \rangle_{b,2} &= \sum_{i=1}^n c_i d_i \|y_i\|_{b,2}^2 \\ &= \sum_{i=1}^n c_i d_i = \langle c, d \rangle, \end{aligned}$$

with $c = (c_1, \dots, c_n)$ and $d = (d_1, \dots, d_n)$. If $u = v$, then $\|u\|_{b,2} = \left(\sum_{i=1}^n c_i^2 \right)^{\frac{1}{2}}$.

According to the above form, we conclude that an inner product on A can be viewed as the inner product of the Euclidean space \mathbb{R}^n . Hence, an explicit formula of the angle between u and v of A , denoted θ , is given by

$$\cos \theta = \frac{\sum_{i=1}^n c_i d_i}{\left(\sum_{i=1}^n c_i^2 \sum_{i=1}^n d_i^2 \right)^{\frac{1}{2}}}.$$

Note that the angle between two vectors of A is also the angle between two lines of A . Moreover, the angle between two vectors of A coincides with the angle between two vectors of \mathbb{R}^n .

Example 2.6. Let $A = \text{span} \{x_1, \dots, x_5\}$ with $\{x_1, \dots, x_5\}$ be the orthonormal set on ℓ^p . If $u = x_1 + 2x_2 + 3x_4 + 4x_5$ and $v = x_2 + 3x_3 + 5x_4 + 2x_5$ then the angle between u and v of A is

$$\cos \theta = \frac{25}{(30.39)^{\frac{1}{2}}} = \frac{25}{34.2}.$$

Next, we can discuss angle between 1-dimensional subspace and $(n-1)$ -dimensional subspace where $n \geq 2$ of A . The result is shown as follows.

Proposition 2.7. If $U = \text{span}\{x_k\}$ where $k = 1, \dots, n$ and $V = \text{span}\{x_{i_2(k)}, \dots, x_{i_n(k)}\}$ where $\{i_2(k), \dots, i_n(k)\} = \{1, 2, \dots, n\} - \{k\}$ of A , then the angle between subspaces U and V is θ ($0 \leq \theta \leq \frac{\pi}{2}$) with

$$\cos^2 \theta = \frac{1}{\|x_k\|_{b,2}^2} \sum_{j \in \{1, \dots, n\} - \{k\}} \langle x_k, y_j \rangle_{b,2}^2.$$

Proof. Without loss of generality, write $k = 1$. Using the Gram-Schmidt process, we have an orthonormal set $\{y_2, \dots, y_n\}$. As a consequence, $\text{span} \{x_2, \dots, x_n\} = \text{span} \{y_2, \dots, y_n\}$. The projection of x_1 on V is given by

$$x_1^V = \sum_{j=2}^n \langle x_1, y_j \rangle_{b,2} y_j.$$

Then, we have

$$\begin{aligned} \langle x_1^V, x_1^V \rangle_{b,2} &= \left\langle \sum_{j=2}^n \langle x_1, y_j \rangle_{b,2} y_j, \sum_{j=2}^n \langle x_1, y_j \rangle_{b,2} y_j \right\rangle_{b,2} \\ &= \sum_{j=2}^n \sum_{i=2}^n \langle x_1, y_j \rangle_{b,2} \langle x_1, y_i \rangle_{b,2} \langle y_j, y_i \rangle_{b,2} \\ &= \sum_{j=2}^n \langle x_1, y_j \rangle_{b,2}^2. \end{aligned}$$

Hence, we obtain

$$\cos^2 \theta = \frac{\|x_1^V\|_{b,2}^2}{\|x_1\|_{b,2}^2} = \frac{1}{\|x_1\|_{b,2}^2} \sum_{j=2}^n \langle x_1, y_j \rangle_{b,2}^2.$$

□

Example 2.8. Let $A = \text{span}\{x_1, x_2, x_3\}$ with $\{x_1, x_2, x_3\}$ be the orthonormal set in ℓ^1 . Take $x_1 = (1, 0, 0, \dots)$, $x_2 = (0, 1, 0, \dots)$ and $x_3 = (0, 0, 1, \dots)$, so that we have $b = (1, 1, 1, \dots)$. Clearly $\|x_1\|_{b,2}^2 = 1$, $\langle x_1, x_2 \rangle_{b,2}^2 = 0$ and $\langle x_1, x_3 \rangle_{b,2}^2 = 0$. If $U = \text{span}\{x_1\}$ and $V = \text{span}\{x_2, x_3\}$ of A then angle between U and V is θ with

$$\begin{aligned}\cos^2 \theta &= \frac{1}{\|x_1\|_{b,2}^2} [\langle x_1, x_2 \rangle_{b,2}^2 + \langle x_1, x_3 \rangle_{b,2}^2] \\ &= 0.\end{aligned}$$

Hence $\theta = \frac{\pi}{2}$.

Before we can discuss an explicit formula for the cosine of the angle between two subspaces of $A = \text{span}\{x_1, \dots, x_n\}$, we recall definition of angle two subspace in inner product space as follows.

Definition 2.9. [3] Let $(X, \langle \cdot, \cdot \rangle)$ be a real inner product space. If $U = \text{span}\{u_1, \dots, u_t\}$ is a t -dimensional subspace and $V = \text{span}\{v_1, \dots, v_s\}$ is a s -dimensional subspace of X with $t \leq s$, then the angle between subspaces U and V is defined by θ ($0 \leq \theta \leq \frac{\pi}{2}$) and

$$\cos^2 \theta = \frac{\|u_1^V, \dots, u_t^V\|^2}{\|u_1, \dots, u_t\|^2},$$

where u_i^V denote the projection of u_i on V for each $i = 1, \dots, t$.

Using Definition 2.9, we have the following theorem.

Theorem 2.10. If $U = \text{span}\{x_1, \dots, x_{n_1}\}$ is a n_1 -dimensional subspace and $V = \text{span}\{x_{n_1+1}, \dots, x_n\}$ is a $(n - n_1)$ -dimensional subspace of A with $n_1 \leq \frac{n}{2}$, then the angle between subspaces U and V is defined by θ and

$$\cos^2 \theta = \frac{\det([\langle x_i, y_l \rangle_{b,2}] [\langle x_i, y_l \rangle_{b,2}]^T)}{\det([\langle x_i, x_j \rangle_{b,2}])}$$

where $[\langle x_i, y_l \rangle_{b,2}]$ is a $(n_1 \times (n - n_1))$ matrix and $[\langle x_i, y_l \rangle_{b,2}]^T$ is its transpose for $i, j = 1, \dots, n_1$.

Proof. Suppose that $\{x_{n_1+1}, \dots, x_n\}$ is linearly independent. Using the Gram-Schmidt process, we obtain the orthonormal set $\{y_{n_1+1}, \dots, y_n\}$. Here $\text{span}\{x_{n_1+1}, \dots, x_n\} = \text{span}\{y_{n_1+1}, \dots, y_n\}$. For each $i = 1, \dots, n_1$, the projection of x_i on V is given by

$$x_i^V = \sum_{l=n_1+1}^n \langle x_i, y_l \rangle_{b,2} y_l.$$

So, for $i, j = 1, \dots, n_1$, we have

$$\begin{aligned}\langle x_i^V, x_j^V \rangle_{b,2} &= \langle x_i, x_j^V \rangle_{b,2} = \left\langle x_i, \sum_{l=n_1+1}^n \langle x_j, y_l \rangle_{b,2} y_l \right\rangle_{b,2} \\ &= \sum_{l=n_1+1}^n \langle x_i, y_l \rangle_{b,2} \langle x_j, y_l \rangle_{b,2}.\end{aligned}$$

Next, using formula angle in [3], we obtain

$$\|x_1^V, \dots, x_{n_1}^V\|^2 = \det \left[\sum_{i=1}^{n_1} \langle x_i, y_i \rangle_{b,2} \langle x_j, y_i \rangle_{b,2} \right] = \det \left([\langle x_i, y_i \rangle_{b,2}] [\langle x_i, y_i \rangle_{b,2}]^T \right),$$

where $[\langle x_i, y_i \rangle_{b,2}]$ is a $(n_1 \times (n - n_1))$ matrix and $[\langle x_i, y_i \rangle_{b,2}]^T$ is its transpose. Therefore, cosine of the angle between U and V is

$$\cos^2 \theta = \frac{\det \left([\langle x_i, y_i \rangle_{b,2}] [\langle x_i, y_i \rangle_{b,2}]^T \right)}{\det \left([\langle x_i, x_j \rangle_{b,2}] \right)},$$

where $[\langle x_i, x_j \rangle_{b,2}]$ is a $(n_1 \times n_1)$ matrix. □

Next, we discuss angle between two subspaces that intersects on a subspace of A . Write $A = \text{span} \{x_1, \dots, x_{n_1}, x_{n_1+1}, \dots, x_{n_1+n_2}, x_{n_1+n_2+1}, \dots, x_{n_1+n_2+n_3}\}$ with $n = n_1 + n_2 + n_3$. Suppose now that $U = \text{span} \{x_1, \dots, x_{n_1}, x_{n_1+1}, \dots, x_{n_1+n_2}\}$ and $V = \text{span} \{x_1, \dots, x_{n_1}, x_{n_1+n_2+1}, \dots, x_{n_1+n_2+n_3}\}$. We observe that U and V are subspace on $A \subseteq \ell^p$. Moreover, using Definition 2.9, we have angle between U and V as follows.

Theorem 2.31. If $U = \text{span} \{x_1, \dots, x_{n_1}, x_{n_1+1}, \dots, x_{n_1+n_2}\}$ is a $(n_1 + n_2)$ -dimensional subspace and $V = \text{span} \{x_1, \dots, x_{n_1}, x_{n_1+n_2+1}, \dots, x_{n_1+n_2+n_3}\}$ is a $(n_1 + n_3)$ -dimensional subspace of A with $n_2 \leq n_3$ that intersects on n_1 -dimensional subspace $W = \text{span} \{x_1, \dots, x_{n_1}\}$ then the angle between subspaces U and V is defined by θ with

$$\cos^2 \theta = \frac{\det \left[\left\langle (x_i^V)_{W^\perp}^\perp, (x_j^V)_{W^\perp}^\perp \right\rangle_{b,2} \right]}{\det \left[\left\langle (x_i)_{W^\perp}^\perp, (x_j)_{W^\perp}^\perp \right\rangle_{b,2} \right]},$$

where $(x_i^V)_{W^\perp}^\perp$, $(x_j^V)_{W^\perp}^\perp$, $(x_i)_{W^\perp}^\perp$ and $(x_j)_{W^\perp}^\perp$ are the orthogonal complement of u_i^V , u_j^V , u_i and u_j , respectively, on W for $i, j = n_1 + 1, \dots, n_1 + n_2$.

Proof. The projection of x_i on V is x_i^V . Next, we may write $x_i^V = (x_i^V)_W + (x_i^V)_{W^\perp}^\perp$ where $(x_i^V)_W$ is the projection of x_i^V on W and $(x_i^V)_{W^\perp}^\perp$ is the orthogonal complement of x_i^V on W . In line with this, we may write $x_i = (x_i)_W + (x_i)_{W^\perp}^\perp$ where $(x_i)_W$ is the projection of x_i on W and $(x_i)_{W^\perp}^\perp$ is the orthogonal complement of x_i on W for $i = n_1 + 1, \dots, n_1 + n_2$. Using the standard $(n_1 + n_2)$ -norm and properties of determinants, we obtain

$$\begin{aligned} \cos^2 \theta &= \frac{\|x_1, \dots, x_{n_1}, x_{n_1+1}^V, \dots, x_{n_1+n_2}^V\|^2}{\|x_1, \dots, x_{n_1}, x_{n_1+1}, \dots, x_{n_1+n_2}\|^2} \\ &= \frac{\|x_1, \dots, x_{n_1}, (x_{n_1+1}^V)_W + (x_{n_1+1}^V)_{W^\perp}^\perp, \dots, (x_{n_1+n_2}^V)_W + (x_{n_1+n_2}^V)_{W^\perp}^\perp\|^2}{\|x_1, \dots, x_{n_1}, (x_{n_1+1})_W + (x_{n_1+1})_{W^\perp}^\perp, \dots, (x_{n_1+n_2})_W + (x_{n_1+n_2})_{W^\perp}^\perp\|^2} \\ &= \frac{\|x_1, \dots, x_{n_1}\|^2 \|(x_{n_1+1}^V)_{W^\perp}^\perp, \dots, (x_{n_1+n_2}^V)_{W^\perp}^\perp\|^2}{\|x_1, \dots, x_{n_1}\|_{n_1}^2 \|(x_{n_1+1})_{W^\perp}^\perp, \dots, (x_{n_1+n_2})_{W^\perp}^\perp\|^2} \end{aligned}$$

$$\begin{aligned}
&= \frac{\| (x_{n_1+1}^V)_{\bar{W}}^{\perp}, \dots, (x_{n_1+n_2}^V)_{\bar{W}}^{\perp} \|^2}{\| (x_{n_1+1})_{\bar{W}}^{\perp}, \dots, (x_{n_1+n_2})_{\bar{W}}^{\perp} \|^2} \\
&= \frac{\det \left[\langle (x_i^V)_{\bar{W}}^{\perp}, (x_j^V)_{\bar{W}}^{\perp} \rangle_{b,2} \right]}{\det \left[\langle (x_i)_{\bar{W}}^{\perp}, (x_j)_{\bar{W}}^{\perp} \rangle_{b,2} \right]}.
\end{aligned}$$

with $[\langle (x_i^V)_{\bar{W}}^{\perp}, (x_j^V)_{\bar{W}}^{\perp} \rangle_{b,2}]$ and $[\langle (x_i)_{\bar{W}}^{\perp}, (x_j)_{\bar{W}}^{\perp} \rangle_{b,2}]$ are $(n_2 \times n_2)$ matrix. \square

3. Conclusions

Based result has been given on the Sections 2, we have known that A that equipped with usual norm on ℓ^p is complete. We have introduced $\langle \cdot, \cdot \rangle_{b,2}$ on A and have shown that $(A, \|\cdot\|_{b,2})$ is complete. Next, we have got angle between two vectors and between 1-dimensional subspace and $(s-1)$ -dimensional subspace where $s \geq 2$ of A . Moreover, we have got angle between two subspaces that intersects on a subspace of A .

Acknowledgments

The first author is supported by PDUPT Program 2021 No. 752/UN4.22/PT.01.03/2021.

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. V. Balestro, A. G. Horváth, H. Martini, R. Teixeira, Angles in normed spaces, *Aequationes Math.*, **91** (2017), 201–236. doi: 10.1007/s00010-016-0445-8.
2. H. Gunawan, J. Lindiarni, O. Neswan, P -, I -, g -, and D -angles in normed spaces, *J. Math. Fund. Sci.*, **40** (2008), 24–32. doi: 10.5614/itbj.sci.2008.40.1.3.
3. H. Gunawan, O. Neswan, W. Setya-Budhi, A formula for angles between two subspaces of inner product spaces, *Beitr. Algebra Geom.*, **46** (2005), 311–320.
4. S. Konca, Some notes on the space of p -summable sequences, *Kuwait J. Sci.*, **45** (2018), 6–13.
5. S. Konca, M. Idris, H. Gunawan, p -summable sequence spaces with inner products, *Beu J. Sci. Techn.*, **5** (2015), 37–41. doi: 10.18532/caam.47117.
6. E. Kreyszig, *Introductory functional analysis with applications*, John Wiley & Sons, Inc., New York, 1978. doi: 10.1137/1021075.
7. P. M. Miličić, On the B -angle and g -angle in normed spaces, *J. Inequal. Pure Appl. Math.*, **8** (2007), 1–9.
8. M. Nur, H. Gunawan, Three equivalent n -norms on the space of p -summable sequences, *Fundam. J. Math. Appl.*, **2** (2019), 123–129. doi: 10.33401/fujma.635754.

9. M. Nur, H. Gunawan, A new orthogonality and angle in a normed space, *Aequationes Math.*, **93** (2019), 547–555. doi: 10.1007/s00010-018-0582-3.
10. M. Nur, H. Gunawan, A note on the g -angle between subspaces of a normed space, *Aequationes Math.*, **95** (2021), 309–318. doi: 10.1007/s00010-020-00742-1.
11. M. Nur, H. Gunawan, O. Neswan, A formula for the g -angle between two subspaces of a normed space, *Beitr. Algebra Geom.*, **59** (2018), 133–143. doi: 10.1007/s13366-017-0355-5.
12. J. Srivastava, P. Singh, On some Cauchy sequences defined in ℓ^p -summable sequences, *J. Rajasthan Acad. Phys. Sci.*, **15** (2016), 107–116.
13. C. Zhi-Zhi, L. Wei, L. Lü-Lin, J. You-qing, Projections, Birkhoff orthogonality and angles in normed spaces, *Comm. Math. Res.*, **27** (2011), 378–384. doi: 10.1007/s11401-011-0630-y.



AIMS Press

© 2022 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)

Angle in the space of p-summable sequences

ORIGINALITY REPORT

16%

SIMILARITY INDEX

11%

INTERNET SOURCES

12%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

1	personal.fmipa.itb.ac.id Internet Source	2%
2	dergipark.ulakbim.gov.tr Internet Source	2%
3	John Douglas. Journal of Earthquake Engineering, 2003 Publication	1%
4	L. P. Lebedev, I. I. Vorovich, G. M. L. Gladwell. "Functional Analysis", Springer Science and Business Media LLC, 1996 Publication	1%
5	journals.itb.ac.id Internet Source	1%
6	bioinfo.cipic.ucdavis.edu Internet Source	1%
7	Progress in Mathematical Physics, 2015. Publication	1%
8	F. Dadipour, F. Sadeghi, A. Salemi. "An orthogonality in normed linear spaces based	1%

on angular distance inequality", Aequationes
mathematicae, 2015

Publication

9	caam.fatih.edu.tr Internet Source	1 %
10	www.87994.com Internet Source	<1 %
11	Leonid Khinkis. "On a global measure of nonlinearity and its application in parameter estimation in nonlinear regression", Mathematics for Application, 2020 Publication	<1 %
12	Xiangrong Wang, Moeness Amin, Xianbin Cao. "Analysis and Design of Optimum Sparse Array Configurations for Adaptive Beamforming", IEEE Transactions on Signal Processing, 2018 Publication	<1 %
13	Anirban Kundu, Tarapada Bag, Sk. Nazmul. "A new notion of orthogonality involving area and length", Mathematica Slovaca, 2021 Publication	<1 %
14	Giorgio Gubbiotti, Andrew P Kels. "Algebraic entropy for face-centered quad equations", Journal of Physics A: Mathematical and Theoretical, 2021 Publication	<1 %

15 Alexey A. Kryukov. "Linear algebra and differential geometry on abstract Hilbert space", International Journal of Mathematics and Mathematical Sciences, 2005
Publication <1 %

16 Chao Ding, Sebastian Bock, Klaus Gürlebeck. "Special bases for solutions of a generalized Maxwell system in 3-dimensional space", Mathematical Methods in the Applied Sciences, 2018
Publication <1 %

17 Nicolás García Trillos. "Gromov–Hausdorff limit of Wasserstein spaces on point clouds", Calculus of Variations and Partial Differential Equations, 2020
Publication <1 %

18 P. Sannuti. "Theory of LTR for nonminimum phase systems, recoverable target loops, recovery in a subspace. I. Analysis", 29th IEEE Conference on Decision and Control, 1990
Publication <1 %

19 Swartz, . "Function Spaces", Measure Integration And Function Spaces, 1994.
Publication <1 %

20 theses.ucalgary.ca
Internet Source <1 %

21 www.coursehero.com

Internet Source

<1 %

22

www.emis.de

Internet Source

<1 %

23

www.eudoxuspress.com

Internet Source

<1 %

24

Boo Rim Choe, Young Joo Lee, Kyunguk Na.
"Toeplitz operators on harmonic Bergman
spaces", Nagoya Mathematical Journal, 2016

Publication

<1 %

25

Reijer Leijssen, Wyger Brink, Xin An, Andrew
Webb, Rob F. Remis. "TRANSVERSE-EPT: A
LOCAL FIRST ORDER ELECTRICAL PROPERTIES
TOMOGRAPHY APPROACH NOT REQUIRING
ESTIMATION OF THE INCIDENT FIELDS",
Progress In Electromagnetics Research M,
2021

Publication

<1 %

26

purehost.bath.ac.uk

Internet Source

<1 %

27

doi.org

Internet Source

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On