

B-16

The Existence of Global Attractor for a Strongly Continuous Semigroup in Metric Space

by

Submission date: 28-Nov-2021 07:53PM (UTC+0700)

Submission ID: 1714098574

File name: ng_ICMSCSME_Makassar_2015_The_Existence_of_Global_Attractor.pdf (234.58K)

Word count: 2075

Character count: 9498



5 The Existence of Global Attractor for a Strongly Continuous Semigroup in Metric Space

Naimah Aris¹, A. Muh. Amil Siddik², Muh. Nur³

¹Department of Mathematics, Faculty of Mathematics and Natural Sciences,
Hasanuddin University,
email: newima@unhas.science.ac.id

²Department of Mathematics, Faculty of Mathematics and Natural Sciences,
Hasanuddin University,

³Department of Mathematics, Faculty of Mathematics and Natural Sciences,
Hasanuddin University,

Jl. Perintis Kemerdekaan Km.10 Tamalanrea, Makassar, Indonesia Postal Address

ABSTRACT

The behavior of dynamic systems in long term dynamics can be described by the global attractor. Let V be a metric space, the global attractor is a nonempty, compact, and invariant set. A subset V which attracts every bounded subset of V . This paper proves three necessary and sufficient conditions that guarantee the existence of the global attractor for a strongly continuous semigroup in metric space: the semigroup is point dissipative, asymptotically smooth, and keeps bit of bounded set bounded. The existence of global attractor is proved by using ω -limit compactness of the semigroup.

Key Words: global attractor, strongly continuous semigroup.

1. Introduction

Many mathematical physics problems can be put into the perspective of infinite dimensional systems and that system can be described by a strongly continuous semigroup in proper function spaces. One important object to describe the long time dynamics of the infinite dimensional system is the global attractor. In the past forty years, some mathematicians have successfully applied the theory in dynamical systems, real analysis, and functional analysis to show the existence of the global attractor. In lecture notes [1] and [2], a dynamical system is defined as a family of operator semigroups $\{T_t\}$ which map a metric space V to itself and are strongly continuous. That semigroup can describe asymptotic behavior of one dynamical system for a long time.

The global attractor is a basic concept and tool to study asymptotic behaviors of solutions of nonlinear evolution equations. A global attractor is an invariant compact set which absorbs all of the bounded sets as time goes to infinity. This theory has been investigated by several authors (cf. Cholewa and Dlotko [1], Ladyzhenskaya [3], Lan dan Shi [4]) and some of techniques seems to be useful for our problem.



The object of this paper is to show the existence of the global attractor for the strongly continuous semigroup in metric space. In [5], some techniques introduced to show the existence of the global attractor in abstract parabolic problem. Such a measure of noncompactness of the semigroup was used by Ma, Wang and Zhong [6]. But here, we show the existence of the global attractor by using ω -limit compactness of the semigroup.

2. Theory

Based on [5], [6], [7], some definitions in real analysis, functional analysis and dynamical systems are defined as follows.

Definition 2.1. Let V be a metric space. The evolution of dynamical system in V is described by a family of operators $\{T(t)\}$ of maps V into itself. The family is called a C^0 semigroup if it satisfies

- (i) T_0 is the identity map on V .
- (ii) $T_{(t+s)} = T_t \circ T_s$ for every $t, s \geq 0$,
- (iii) The function $T : [0, \infty) \times V \rightarrow V$
 $\ni (t, x) \rightarrow T_t(x) \in V$

is continuous at each point $(t, x) \in [0, \infty) \times V$.

Definition 2.2. Let set K in metric space V . The set K is said to be compact if for every open K open bounded set subset in K .

Definition 2.3. Let V, Y be metric, map $T: V \rightarrow Y$ is continuous at $v_0 \in V$ if given any $\varepsilon > 0$ there exist $\delta > 0$ such that

$$d'(T(v), T(v_0)) < \varepsilon \text{ with } d(v, v_0) < \delta.$$

By geometrical views, let $B_Y(T(v_0), \varepsilon)$ open ball in $T(v_0)$ with radius ε , and $B_V(v_0, \delta)$ open ball at v_0 with radius δ , then

$$T(v) \in B_Y(T(v_0), \varepsilon) \text{ with } v \in B_V(v_0, \delta)$$

or

$$T(B_V(v_0, \delta)) \subseteq B_Y(T(v_0), \varepsilon).$$

Theorem 2.1 Let A, B be metric spaces and $f: A \rightarrow B$. If $\{S_i\}$ subset A and $i, n \in \mathbb{N}$, then

$$f\left(\bigcup_{i=1}^n S_i\right) = \bigcup_{i=1}^n f(S_i).$$

Definition 2.4. Let V be a metric space. Given sets $W_1, W_2 \subset V$, W_2 $\{T_t\}$ absorbed by W_1 if there exist $t_0 \geq 0$ such that

$$T_t W_2 \subset W_1 \text{ for every } t \geq t_0.$$

Definition 2.5. Let V is a metric space. A Global attractor is a non empty, compact and $\{T_t\}$ -invariant set $\mathcal{A} \subset V$ which attract every bounded set.

Definition 2.6. The semigroup $\{T_t\}$ is called point dissipative if and only if there exist a non empty, bounded set $W_0 \subset V$ which attract every point in V . In other words, if $W_0 \subset V$ attracts every point in V , then $\forall v \in (V, d)$ there exist an open neighborhood N_{W_0} of W_0 which absorb v , $\forall v \in (V, d) \exists t_0 \geq 0 \forall t \geq t_0 T_t(v) \subset N_{W_0}$.



Definition 2.7. The semigroup $\{T_t\}$ is called asymptotically smooth if and only if there exist a non empty, closed, bounded, positively invariant set $W_0 \subset V$ contains a non empty, compact subset C which attracts W_0 .

Theorem 2.2. Let $\{T_t\}$ be C^0 semigroup in metric space V . If $\{T_t\}$ is asymptotically smooth, B a non empty subset V , and for some $t_B \geq 0$, set $\bigcup_{s \geq t_B} T_s(B)$ bounded, then $\omega(B)$ is a non empty, compact, and invariant set, moreover $\omega(B)$ attract B .

3. Result and Discussion

In this section we give the existence of the global attractor for strongly continuous semigroup in metric space V describe in Theorem 3.1. below.

Theorem 3.1. Let $\{T_t\}$ be a C^0 semigroup in metric space V . If $\{T_t\}$ is point dissipative, asymptotically smooth and keep orbit of bounded set bounded then $\{T_t\}$ has a global attractor in V .

Proof. The proof proceeds in two steps.

Step 1.

We shall first show the existence of a bounded set $O \subset V$ such that for every bounded closed set $C \subset V$ there is a neighborhood \mathcal{N}_C which is absorbed by O .

Since semigroup $\{T_t\}$ is point dissipative, from Definition 2.6 there exist a non empty and bounded set $W_0 \subset V$ which attracts every point in V , therefore $\forall v \in V$ every open neighborhood \mathcal{N}_{W_0} from W_0 absorbs v . We can say that \mathcal{N}_{W_0} is an absorbing point of V . From Definition 2.6 \mathcal{N}_{W_0} absorbs every point v if $\forall v \in V \exists t_0 \geq 0 \forall t \geq t_0$ such that $T_t(v) \subset \mathcal{N}_{W_0}$. (3.1)

Since $\{T_t\}$ is a C^0 semigroup in metric space V , from Definition 2.3 $\{T_t\}$ continuous in every point in V , hence $\forall v \in V \forall \epsilon > 0 \exists \delta > 0$ such that $T_t(B_V(v, \delta)) \subset B_V(T_t(v), \epsilon)$.

Because \mathcal{N}_{W_0} is an absorbing point in V , then $\forall v \in (V, d) \exists t_0 \geq 0 \forall t \geq t_0 T_t(v) \subset \mathcal{N}_{W_0}$, so for $B_V(T_t(v), \epsilon) \subset \mathcal{N}_{W_0}$, we can choose $B_V(v, \delta)$ such that $T_t(B_V(v, \delta)) \subset \mathcal{N}_{W_0}$, and we have $\forall v \in V \exists B_{V(v, \delta)} \exists t_0 \geq 0$ such that

$$T_t(B_V(v, \delta)) \subset \mathcal{N}_{W_0}. \quad (3.2)$$

Let us choose $t_{\mathcal{N}_{W_0}} \geq 0$ such that

$$O := \bigcup_{t \geq t_{\mathcal{N}_{W_0}}} T_t(\mathcal{N}_{W_0})$$

bounded.

Since $\{T_t\}$ is a strongly continuous semigroup in V , then O invariant positive, we obtain,

$$T_{t'}(O) \subset O, t' \geq 0.$$

Furthermore,

$$T_{t'}(O) = T_{t'}\left(\bigcup_{t \geq t_{\mathcal{N}_{W_0}}} T_t(\mathcal{N}_{W_0})\right), t' \geq 0.$$

From Theorem 2.1. there follows



$$T_{t'}(\mathcal{O}) = T_{t'}\left(\bigcup_{t \geq t_{N_{W_0}}} T_t(N_{W_0})\right) = \bigcup_{t \geq t_{N_{W_0}}} T_{t'}(T_t(N_{W_0})). \quad (3.3)$$

According assumptions that $\{T_{t'}\}$ and $\{T_t\}$ are strongly continuous semigroup and from Definition 2.1 Part (ii) we have

$$\bigcup_{t \geq t_{N_{W_0}}} T_{t'}(T_t(N_{W_0})) = \bigcup_{t \geq t_{N_{W_0}}} T_{t'+t}(N_{W_0}).$$

Next from

$$\bigcup_{t \geq t_{N_{W_0}}} T_{t'+t}(N_{W_0}) = T_{t'+t_{N_{W_0}}}(N_{W_0}) \cup T_{t'+t_{N_{W_0}+1}}(N_{W_0}) \cup \dots \quad (3.4)$$

And because $t' \geq 0$

$$\begin{aligned} \bigcup_{t \geq t_{N_{W_0}}} T_t(N_{W_0}) &= T_{t_{N_{W_0}}}(N_{W_0}) \cup T_{t_{N_{W_0}+1}}(N_{W_0}) \cup \dots \cup T_{t'+t_{N_{W_0}}}(N_{W_0}) \cup \\ &T_{t'+t_{N_{W_0}+1}}(N_{W_0}) \cup \dots \\ &= T_{t_{N_{W_0}}}(N_{W_0}) \cup T_{t_{N_{W_0}+1}}(N_{W_0}) \cup \dots \cup \bigcup_{t \geq t_{N_{W_0}}} T_{t'+t}(N_{W_0}) \cup \\ &\dots \quad (3.5) \end{aligned}$$

Therefore from Equations (3.3), (3.4) and (3.5) we obtain

$$T_{t'}(\mathcal{O}) = \bigcup_{t \geq t_{N_{W_0}}} T_{t'+t}(N_{W_0}) \subset \bigcup_{t \geq t_{N_{W_0}}} T_t(N_{W_0}) = \mathcal{O}.$$

So, $T_{t'}(\mathcal{O}) \subset \mathcal{O}$, or \mathcal{O} positive invariant.

Next, because N_{W_0} absorbs every point in V ,

$$\forall v \in V \exists t_0 \geq 0 \text{ such that } T_{t_0}(v) \subset N_{W_0}, \quad t \geq t_0$$

Then there is $t_0 \geq 0$ such that

$$T_{t_{N_{W_0}}}(T_{t_0}(v)) \subset T_{t_{N_{W_0}}}(N_{W_0}) \subset \bigcup_{t \geq t_{N_{W_0}}} T_t(N_{W_0}) = \mathcal{O}, \quad t_{N_{W_0}} \geq t_0.$$

Hence \mathcal{O} absorbs every point in V .

According to Definition 2.1 Part (ii), $\{T_t\}$ is strongly continuous, hence

$$T_{t_{N_{W_0}}}(T_{t_0}(v)) = T_{t_{N_{W_0}}+t_0}(v).$$

With the aid of Equation (3.2) and Definition 2.1 part (ii) also with the fact that \mathcal{O} is the absorbing point in V , $\forall v \in V \exists t_v = t_{N_{W_0}} + t_0 \geq 0 \exists B_{V(v, \delta)}$ such that

$$T_t(B_{V(v, \delta)}) \subset \mathcal{O}, \quad t \geq t_v. \quad (3.6)$$

Consider any compact set $C \subset V$, certainly from Definition 2.2,

$$\begin{aligned} C &\subset \bigcup_{i=1}^n B_{V(v_i, \delta_i)} \\ C &\subset B_{V(v_1, \delta_1)} \cup B_{V(v_2, \delta_2)} \cup \dots \cup B_{V(v_n, \delta_n)} =: \mathcal{N}_C \quad (3.7) \end{aligned}$$

For some $n \in \mathbb{N}$, $v_1, v_2, \dots, v_n \in C$ and because T_t continuous, then

$$T_t(C) \subset T_t(\mathcal{N}_C) = T_t \bigcup_{i=1}^n B_{V(v_i, \delta_i)} = \bigcup_{i=1}^n T_t(B_{V(v_i, \delta_i)}).$$

According to Equation (3.6) we obtain

$$T_t(B_{V(v_1, \delta_1)}) \subset \mathcal{O}, \text{ for } t \geq t_{v_1}$$

$$T_t(B_{V(v_2, \delta_2)}) \subset \mathcal{O}, \text{ for } t \geq t_{v_2}$$



⋮

$$T_t(B_V(v_n, \delta_n)) \subset \mathcal{O}, \text{ for } t \geq t_{v_n}.$$

We have for $t \geq \max\{t_{v_1}, t_{v_2}, \dots, t_{v_n}\} \geq 0$ that $T_t(\mathcal{N}_C) \subset \mathcal{O}$.

So we can show that there exists a set $\mathcal{O} \subset V$ such that every compact set $\mathcal{C} \subset V$ has a neighborhood \mathcal{N}_C which is absorbed by set \mathcal{O} .

Step 2

We show that there is a compact, $\{T_t\}$ invariant set $\mathcal{A} \subset V$ which attract every bounded set in V .

Let $B \subset V$ be a non empty bounded set. Since

- (i) $\{T_t\}$ is asymptotically smooth and
- (ii) From property of orbit of bounded set keep bounded we have

$$\gamma(B) = \bigcup_{s \geq 0} T_s(B)$$

is bounded, therefore for some $t_B \geq 0$, set

$$\bigcup_{s \geq t_B} T_s(B)$$

is bounded too, then from Theorem 2.1 we can conclude according to Theorem 2.2 that $\omega(B)$ compact and attract B , i.e.

$$\forall \mathcal{N}_{\omega(B)} \exists t_B \geq 0 \quad T_{t_1} B \subset \mathcal{N}_{\omega(B)} \text{ for } t_1 \geq t_B \quad (3.8)$$

As shown in step I, there is some neighborhood of compact set $(\omega(B)); (\mathcal{N}_{\omega(B)})$; which absorbed by \mathcal{O} , i.e.

$$\exists t_0 T_{t_3}(\mathcal{N}_{\omega(B)}) \subset \mathcal{O} \text{ for } t_3 \geq t_0$$

Take $t_2 = t_3 + t_1$,

$$\begin{aligned} T_{t_2}(B) &= T_{t_3+t_1}(B) \\ &= T_{t_3}(T_{t_1}(B)) \subset T_{t_3}(\mathcal{N}_{\omega(B)}) \subset \mathcal{O}. \end{aligned}$$

We obtain, there is $\tau_B = t_B + t_0$ such that

$$\forall B \subset V \exists \tau_B \geq 0 \quad T_{t_2}(B) \subset \mathcal{O} \text{ for } t_2 \geq \tau_B. \quad (3.9)$$

Let, $\mathcal{A} := \omega(\mathcal{O})$. Because

- (i) $\{T_t\}$ is asymptotically smooth and
- (ii) Set \mathcal{O} bounded and positive invariant such that $T_{t_2}(\mathcal{O}) \subset \mathcal{O}$, hence

$$\exists \tau_B \geq 0 \bigcup_{t_2 \geq \tau_B} T_{t_2}(\mathcal{O}) \text{ is bounded.}$$

From Theorem 2.2, $\mathcal{A} := \omega(\mathcal{O})$ is a non empty, compact set, moreover, \mathcal{A} attracts \mathcal{O} .

Now we will show that \mathcal{A} attracts every bounded set in V .

\mathcal{A} attracts \mathcal{O} means $\forall \mathcal{N}_A \exists t_0 \geq 0 T_t(\mathcal{O}) \subset \mathcal{N}_A$ for every $t \geq t_0$.

From Equation (3.9), let $t_4 = t + t_2$

$$T_{t_4}(B) = T_{t+t_2}(B) = T_t(T_{t_2}(B)) \subset T_t(\mathcal{O}) \subset \mathcal{N}_A$$

there is $\tau_2 = t_0 + \tau_B$ such that



$$\forall B \subset V \exists \tau_2 = t_0 + \tau_B \geq 0 \quad T_{t_4}(B) \subset \mathcal{N}_{\mathcal{A}} \text{ for every } t_4 \geq \tau_2.$$

Hence \mathcal{A} attract every bounded set in V . The proof is complete. ■.

5. Conclusion

Let $\{T_t\}$ be a C^0 semigroup in metric space V . There exist a global attractor in V if and only if the semigroup $\{T_t\}$ is point dissipative, asymptotically smooth and keep orbit of bounded set bounded.

REFERENCES

- [1] J. W. Cholewa, dan T. Dlotko (2000). *Global Attractor in Abstract Parabolic Problem*. Cambridge University Press.
- [2] Temam, Roger (1998). *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*. Springer.
- [3] O. Ladyzhenskaya (1991). *Attractor for Semigroup and Evolution Equation*. Cambridge University Press.
- [4]. Lan dan Y. Shi (2013). *Notes on the Global Attractors for Semigroup*. Soochow University.
- [5] M Nakao, N. Aris (2007). On Global Attractor For Nonlinear Parabolic Equation of m -Laplacian Type. *J. Math. Anal. Appl.* **331**: 793-809.
- [6] Q. Ma, S. Wang and C. Zhong (2002). Necessary and Sufficient Conditions for Existence of Global Attractors for Semigroups and Applications. *Indiana University Mathematics Journal.* **51**(6): 1541-1559.

The Existence of Global Attractor for a Strongly Continuous Semigroup in Metric Space

ORIGINALITY REPORT

10%

SIMILARITY INDEX

4%

INTERNET SOURCES

10%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Yunhua Zhou, Xingyou Zhang. " Stability of semigroups which possess global attractors ", Dynamical Systems, 2010
Publication 1%
- 2** Universitext, 2015.
Publication 1%
- 3** www.jstage.jst.go.jp
Internet Source 1%
- 4** "Global attractor for a quasilinear parabolic equation of mean curvature type", International Journal of Dynamical Systems and Differential Equations, 2007
Publication 1%
- 5** "Nonlinear Parabolic-Hyperbolic Coupled Systems and Their Attractors", Springer Science and Business Media LLC, 2008
Publication 1%
- 6** Yantao Guo, Pengyan Ding. " Global attractor of the fractional damping wave equation on \mathbb{R}^n ", Applicable Analysis, 2020 1%

-
- | | | |
|-----------|---|------|
| 7 | Chowdhury, M.S.R.. "Minimax inequalities on G-convex spaces with applications to generalized games", Nonlinear Analysis, 200101
Publication | 1 % |
| 8 | Lee, Mi Jin. "GLOBAL ATTRACTOR FOR SOME BEAM EQUATION WITH NONLINEAR SOURCE AND DAMPING TERMS", East Asian Mathematical Journal, 2016.
Publication | 1 % |
| 9 | Song, H.. "Attractors of non-autonomous reaction-diffusion equations in L^p ", Nonlinear Analysis, 20080401
Publication | 1 % |
| 10 | Cholewa, J.W.. "Extremal equilibria for monotone semigroups in ordered spaces with application to evolutionary equations", Journal of Differential Equations, 20100801
Publication | <1 % |
| 11 | aimsciences.org
Internet Source | <1 % |
| 12 | eprints.ucm.es
Internet Source | <1 % |
| 13 | Weisheng Niu, Chengkui Zhong. "Global attractors for the p-Laplacian equations with | <1 % |

nonregular data", Journal of Mathematical
Analysis and Applications, 2012

Publication

14

Hongtao Li, Shan Ma. "Asymptotic Behavior of
a Class of Degenerate Parabolic Equations",
Abstract and Applied Analysis, 2012

Publication

<1%

15

Weisheng Niu, Hongtao Li. "Asymptotic
Behavior of Approximated Solutions to
Parabolic Equations with Irregular Data",
Abstract and Applied Analysis, 2012

Publication

<1%

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On