

PROSIDING



ANNUAL SCIENTIFIC MEETING

PERHIMPUNAN AHLI ANATOMI INDONESIA (PAAI)
"FROM BASIC TO CLINICAL APPLICATION"


MAKASSAR, 9 - 12 AGUSTUS 2017

Editor:

dr. M. Iqbal Basri, Sp.S

dr. Rahmawati Mihnjajat, Ph.D, Sp.PD

dr. Asty Amalia



PERHIMPUNAN AHLI ANATOMI INDONESIA
PROSIDING
PERTEMUAN ILMIAH TAHUNAN
PERHIMPUNAN AHLI ANATOMI INDONESIA
2017

Tema:
From Basic to Clinical Application

ISBN 978-602-99925-7-1

Editor:
dr. M. Iqbal Basri, Sp.S
dr. Rahmawati Mihnajat, Ph.D, Sp.PD
dr. Asty Amalia

Penerbit dan Redaksi:
Fakultas Kedokteran Universitas Hasanuddin
Jl. Perintis Kemerdekaan Km. 10
Kampus UNHAS Tamalanrea Makassar 90245
Telp//Fax: (0411) 586010
Email: fkunhas@med.unhas.ac.id
Website: <http://med.unhas.ac.id>

Cetakan Pertama, Agustus 2017

Hak cipta dilindungi undang-undang
Dilarang memperbanyak, mencetak dan menerbitkan sebagian atau seluruh isi karya tulis ini
dalam bentuk dan dengan cara apapun tanpa ijin tertulis dari penulis dan penerbit.

**DAFTAR ISI ANTAR
PERTEMUAN ILMIAH NASIONAL
PERHIMPUNAN AHLI ANATOMI INDONESIA
2017**

Sampul	i
Kata Pengantar Dekan Fakultas Kedokteran Universitas Hasanuddin	iii
Sambutan Ketua Pengurus Besar Perhimpunan Ahli Anatomi Indonesia	iv
Kata Pengantar Ketua Panitia Pertemuan Ilmiah Nasional Perhimpunan Ahli Anatomi Indonesia	vi
Susunan Acara Pertemuan Ilmiah Nasional Perhimpunan Ahli Anatomi Indonesia	vii
Daftar Isi	viii
Kumpulan Materi Plenary Session	
Research Trends in Histology..... <i>Ahmad Aulia Jusuf</i>	2
Challenges And Future Directions Of Anatomy: Advanced Study Of Brain Microcircuitry Connections <i>Taruna Ikrar</i>	4
Inovasi Pembelajaran Anatomi..... <i>Sigit Moerjono</i>	7
Basic Anatomy: Teaching Methodology and Innovation..... <i>Fundhy Sinar Ikrar Prihatanto</i>	9
The Drivers for Evolution..... <i>Eiichi Hondo</i>	12
The Role of Skull Anatomic Landmarks in Indonesia Suicide Bombings & Criminal Cases Identification..... <i>Peter Sahelangi</i>	14
Vascular Anatomy and Its Clinical Application in Treatment Of Disease. Case Reports..... <i>Mohd Rizal Roslan</i>	15
Kumpulan Materi Simposium Paralel	
Sistem Pembelajaran Berbasis Teman Sebaya Sebagai Penunjang Pembelajaran Anatomi di Sistem Modul Mahasiswa FK Undip..... <i>Surya Pratama Brilliantika</i>	18
Variation of Hepatic Artery: A Clinical Review..... <i>Nikmatia Latief</i>	18
Human Endothelial Progenitor Cell: Biological Characteristics and Clinical Application.....	18

Rahmawati Minhajat

Evaluasi Variasi Arteri Renalis pada Pasien yang Dilakukan Pemeriksaan MDCT-Scan Angiografi Abdomen..... 19

Rosdianah, Bachtiar Murtala, Nikmatia Latief

Aspek Anatomi Femur pada Penurunan Densitas Mineral Tulang..... 25

Sitti Rafiah

Bone Growth 26

Shelly Salmah

Rehabilitasi Fraktur Ekstremitas Bawah..... 27

Yose Waluyo

Retina 29

Batari Todja

Perubahan Anatomi dan Fisiologi pada Kasus Ablatio Retina 31

A.M.Ichsan, Andi Pratiwi

Prefrontal Cortex: Basic to Clinic 40

Taufik Pasiak

The Role of Mirror Neurons System in Mental Imagery Processing 45

Muhammad Iqbal Basri

Clinical Neurophysiology in Evaluation of Cognitive Function..... 51

Manfaluthy Hakim

Kumpulan Materi Presentasi Oral

Hubungan Lompatan Jumping Smash dan Bentuk Arcus Pedis Dengan Kejadian Sprain Pergelangan Kaki pada Atlet Bulutangkis..... 54

Rochman Basuki, Sigit Moerjono, Muhammad Arif Nurohman

Variasi Rugae Palatina pada Suku Jawa dan Batak..... 56

Fidya, Iswara Rizki Arum Lestari, Diwya Nugrahini

A Study of Adult Human Cerebellum from The Anatomy Cadavers..... 61

Kalanjati VP., Dewi AK., Santoso MWA

Identification of Sutural Bones in Indonesian Skulls 62

Safrina D Ratnaningrum

Fleksibilitas Lumbal dan Hubungannya dengan Indeks Massa Tubuh pada Remaja Putra 64

Tri Suciati

Gambaran Garis Lipat Tangan pada Anak Normal dan Anak Retardasi Mental di SLB Jakarta Pusat 64

Etty Widayanti, Titiek Djannatun, Endang Purwaningsih, Restu Samsul Hadi, Mirfat

Peran Nitrit Oksida Intraluminal Pada Taut Gastroesofagus / Gastroesophagus junction (GEJ) 68

Tena Djuartina, Ahmad Aulia, Ari Fahrial Syam

Pancreatic Beta Cells Protective Effect of Okra Extract in Alloxan-Induced Type II Diabetic Rats..... 70

Karim Susanto, Anita Anjani

Kadar Hormon FSH Basal dan LH Basal Perempuan Sindrom Ovarium Polikistik (SOPK) Yang Mengikuti Program In Vitro Fertilization..... 74

Roselina Panghiyangan, Dwi Anita Suryandari, Budi Wiweko, Mala Kurniati

Gambaran Histopatologi Hepar pada Tikus yang Diinfeksi *Plasmodium berghei* Pasca Pemberian Oksigen Hiperbarik 75

Herin Setianingsih, Riami, Prawesty Diah Utami

Hepatocyte Expression of TNF- α in High Fat Diet Fed Wistar Rat with *Andrographis paniculata* Extract Therapy 80

Hariadi, T.A.S., Jong, F.X.H.H

Theurapetic Effects of *Andrographis Paniculata* Ethanol Extract in The Liver of Wistar Rat Treated with A High Fat Diet..... 88

FX Himawan Haryanto Jong, Tania Ardiani Saleh Hariadi, Ari Gunawan, Mochamad Wirono Aman Santoso, Ni Wajan Tirthaningsih, Susilowati Andajani, Achmad Basori

Museum Anatomi Sebagai Salah Satu Sarana Pembelajaran Pendidikan Dokter..... 92

Robilrawan dan Liliana Sugiharto

E-Forum in Anatomy Learning: Challenge and Advantages..... 95

Lucky Prasetiowati, Sakina, Fundhy Sinar Ikrar Prihatanto

Process in Building E-Quiz Histology as An Innovative Assessment Model in Block Curriculum Faculty of Medicine Universitas Airlangga 95

Rimbun, Tri Hartini Yuliawati, Dewi Ratna Sari, Kusuma Eko Purwantari, Anggun Raksyantoro

Student assistant in anatomy: A form of student engagement in the educational process .. 96

Sakina, Fundhy Sinar Ikrar Prihatanto, Lucky Prasetiowati

Belajar Anatomi Traktus Substansia Alba Otak Menggunakan Media *Magnetic Resonancy Imaging Fiber Tractography*..... 96

Herlina Uinarni

The Exposure Effect of Al-Qur'an Recitation to Spatial Memory on *Rattus novergicus* 99

Abdurrahman Afa Haridhi, Nanang Wiyono, Yunia Hastami, Muthmainah

Penerapan Multivariate Adaptive Regresi Spline untuk Mengidentifikasi Faktor yang Berpengaruh pada Penggemukan Sapi di Desa Samaran Kabupaten Bojonegoro 99

Benjamin Christoffel Tehupuring dan Soeharsono

Inhibisi Sel Punca CD90⁺ Sumsum Tulang dan Sel Punca CD34⁺ Darah Tali Pusat Terhadap Aktivitas Fibrosis Sel Stelata Hepatik 100

Radiana D Antarianto, Ahmad Aulia Jusuf, Atikah C Barasila, Wahyunia L Septiana, Ervina Julien Sitanggung

Signaling pathways involved in peripheral neuronal generation..... 101

Ria Margiana, Jeanne Adiwinata Pawitan, Ahmad Aulia Jusuf

Environmental Enrichment (EE) Mengurangi Perilaku Depresif Akibat Paparan Stres Kronis pada Tikus Putih (*Rattus norvegicus*) 105

Maia Thalia Giani, Muthmainah, Dyah Ratna Budiani, Nanang Wiyono

Concentration of Reactive Oxygen Species (ROS) and Telomeric Repeat Binding Factor2 (TERF2) On Individuals By Age and Gender..... 108

Endang Purwaningsih, Tripanjiasih Susmiarsih, Yenni Zulhamidah, Achmad Sofwan

Mekanisme Kerja Obat Hipoglikemik Oral dan Hipoglikemik Herbal..... 115

Yuliana

Evaluation Antacid Contain Magnesium Hidroxide Treatment Based on Parameter of Reactive Gastropathy 117

Arni Kusuma Dewi, Chairul Anwar

Valproic Acid Reduces Pancreatic Beta Cells in Rat Offspring 117

Ade Komariah, Bambang Kiranadi, Adi Winanto, Wasmen Manalu, Ekowati Handharyani

Evaluasi gambaran histologis penyembuhan luka bakar full thickness dengan pewarnaan Masson trichrome 118

Dewi Sukmawati, Astheria Eryani, Elvin Clara Angmalisang, Lia Damayanti

Pengaruh Tipe Bentuk Jari Kaki, Model dan Ukuran *Flat Shoes* terhadap Kenyamanan Kaki pada Mahasiswi Fakultas Kedokteran Universitas Katolik Indonesia Atma Jaya 118

Tan Fei Fan, Paula Chintya, Robi Irawan, Nawanto Agung Prastowo

Understanding The Anatomy of Chromosome and Its Abnormality: Role of An Anatomist .. 119

Ni Wajan Tirthaningsih, Tri Hartini Yuliawati, Sakina, Lucky Prasetiowati

Kajian Pustaka: PPAR γ agonis dan Sekresi Dopamin pada Kontrol *Food Intake*..... 120

Fifi Veronica, Ronny Lesmana, Hanna G, Leonardo Lubis

Secang Wood Ethanol Extracts (*Caesalpinia sappan*) with potential effect on the quality of spermatozoa parameter of Wistar Strain Male Rats 122

Andri Rezano, Nadiyah, Raden Ghita Sariwidyantry, Sunarjati Sugdoadi

Pengaruh Frekuensi Penggantian Balutan Kompres Betaine Polyhexanide 0,1% Terhadap Pembentukan Jaringan Epitel pada Luka Bakar Derajat II A Model Tikus Putih galur Wistar 129

Rizky Oktavia Primasari, Heri Kristianto, Danik Agustin Purwantiningrum

Efek Kuratif Sari Kukusan Brokoli (<i>Brassica oleracea L. var italica</i>) Terhadap Kadar Nf-Kb Pada Mencit Model Kolitis	133
<i>Sijani Prahastuti, Jeanny Ervie Ladi, Teresa Liliana Wargasetia</i>	
Model MARS Pola Pertumbuhan Proventriculus, Ventriculus dan Intestinum tenue Ayam Pedaging Umur 1 hingga 4 minggu	135
<i>Siti F. Masruroh, B. C. Tehupuring, Surjo Kuncorojakti, Hana Eliyani, Suharsono</i>	
Pengaruh Pemberian Ekstrak Daun Teh Hijau Terhadap Konsentrasi dan Kecepatan Spermatozoa Tikus (<i>Rattus norvegicus</i>) Setelah Paparan Asap Rokok	138
<i>Tri Panjiasih Susmiarsih, Kenconoviyati, Kuslestari</i>	
Effect of Physical Stress on Cortisol Level and Diameter of White Pulp in Rats	143
<i>Tri Hartini Yulawati, Rimbun, Viskasari P Kalanjati</i>	
Kumpulan Materi Presentasi Poster	
Transisi Epitel-Mesenkim dan Progres Kanker	150
<i>Oeij Anindita Adhika</i>	
Aktivitas Ekstrak Air Daun Kersen (<i>Muntingia calabura</i>) Terhadap Gambaran Histologis Sel Hepar <i>Mus Musculus Balb/C</i> yang Diinduksi D-Galaktosa	153
<i>Cempaka Jaga Paramudita, Dwi Nur Ahsani, Ika Fidiansih, Evy Sulistyoningrum</i>	
Kajian literatur: Perbandingan Pengaruh Paparan Musik dan Bising Selama Periode Prenatal Terhadap Neurogenesis dan Sinaptogenesis Otak.....	154
<i>Fitri, Isabella Kurnia Liem</i>	
Effect of BCCAO Duration and Animal Models Sex on Brain Ischemic Volume After 24 Hours Reperfusion	165
<i>Ety S. Handayani, Titis Nurmasitoh, Saefudin Ali Akhmad, Afifah Nur Fauziah, Rizky Rizani, Rika Yulita Rahmawati, Angga Afriandi</i>	
Perbedaan Distribusi Tebal Lipatan Kulit antara Remaja Perempuan Madura di Sumenep dengan Surabaya	165
<i>Anung Putri Illahika, Abdurahman, Myrtati Dyah Artaria</i>	
Pengaruh Pemberian Propolis Terhadap Gambaran Histopatologi Gaster Tikus (<i>Sprague dawley</i>) yang Diberi Perlakuan Stres Isolasi Sosial	168
<i>Wisika Habiburohman Efendi, Kuswati, Zainuri Sabta Nugraha</i>	
Kajian Mikroskopis Ovarium dan Kompetensi Maturasi Oosit Domba dari Ovarium yang Disimpan Pada Suhu 4° C	168
<i>Masturi M</i>	
Comparison of The Crushing Point on Rat's Sciatic and Tibial Nerve for Peripheral Nerve Regeneration Study Model	169
<i>May Valzon, Ayu Permata</i>	

Pola Perkembangan Vertebra Cervicalis pada Mamalia <i>Msy. Rulan Adnindya, Isabella Kurnia Liem</i>	170
Effect of Ethanolic Extract of <i>Moringa oleifera</i> Leaves on Histopathology Image of Kidney <i>Subhan Amal Romis, Zainuri Sabta Nugraha, Kuswati</i>	170
The Effect of Caffeine Per Oral on The Neuron Cell Number and the Granuler Layer thickness of Dentate Gyrus, Formatio Hippocampalis..... <i>Kusuma Eko Purwantari, Tri Hartini Yuliawati, Dewi Ratna Sari, Rimbun</i>	171
Korelasi Panjang Lengan Atas dan Panjang Sternum Perkutaneus dengan Tinggi Badan Mahasiswa Suku Banjar Fakultas Kedokteran Universitas Lambung Mangkurat..... <i>Iwan Aflanie, Lena Rosida, Cynthia Pratiwi, Ramadhan Maulana Hikmat</i>	171
Anthropometric Indicators of Insulin Resistance on Obese Adults in Surabaya..... <i>Dewi Ratna Sari, Rimbun, Tri Hartini Yuliawati, Kusuma Eko Purwantari</i>	175
Korelasi Nilai Ujian Praktikum Anatomi Dengan Indeks Prestasi Kumulatif Mahasiswa Fakultas Kedokteran Univesitas YARSI Angkatan 2013 <i>Yenni Zulhamidah, Kenconoviyati, Achmad Sofwan, Kuslestari, Edward Syam, Aryenti, Zakiyah</i>	176
Profil Pasien Demam Berdarah dan Kadar Hemoglobin, Trombosit, dan Hematokrit pada Pasien Anak di RSUD Ratu Zalecha Martapura <i>Ida Yuliana</i>	178
Hubungan antara Penggunaan Tas Sekolah dan Keluhan Muskuloskeletal pada Siswa Sekolah Dasar di Kecamatan Ilir Barat I Kota Palembang..... <i>Legiran, Tri Suciati, Meirisa Rahma Pratiwi</i>	181
Pengaruh Suplementasi Madu terhadap Migrasi dan Diferensiasi Sel Human Dermal Fibroblast (HDF) Sebagai Model Uji Luka In Vitro <i>Restu Syamsul Hadi, Nadira, Yoan Rahmah Aprilia</i>	188
Disfungsi Kandung Kemih Non-Neurogenik pada Anak: Diagnosis dan Tata Laksana <i>Bernadetha Nadeak</i>	192
Hubungan Paparan Gas Formaldehida di Ruang Praktikum Anatomi dengan Efek Iritasi Mata <i>Sagala, Antonius Junedi, Margiana, Ria</i>	201
Air Pollution as A Cause Of Malignancy of Respiratory Organs in Riau Province..... <i>Laode Burhanuddin, Erizon, Delta Kurnia</i>	202
Factors that influence the academic performance of Malaysian Students in Faculty of Medicine Hasanuddin University <i>Mohd. Azzam, Asty Amalia, St. Rafiah, M. Iqbal Basri</i>	205
Efek Kuratif Sari Kukusan Brokoli (<i>Brassica oleracea L. var italica</i>) Terhadap Gambaran Histopatologis Kolon Proksimal Dan Medial Pada Mencit Model Kolitis <i>Mohd. Azzam, Asty Amalia, St. Rafiah, M. Iqbal Basri</i>	209

Jeanny Ervie Ladi, Sijani Prahastuti, Teresa Liliانا Wargasetia

Gambaran Anatomi dan Histologi Saluran Pencernaan Ikan Bungo (<i>Glossogobius Cf. aureus</i>) dari Danau Tempe Sulawesi Selatan.....	209
<i>Satrya Adi Pratama, Wahyuni, I Ketut Mudite Adnyare, Andi Tamsil, Dwi Kesuma Sari</i>	
Hubungan Kadar Profil Lipid dengan Kejadian Penyakit Jantung Koroner di RSUP Dr. Wahidin Sudirohusodo Makassar	215
<i>Najdah Hidayah, A. Armyn Nurdin</i>	
Hubungan Status Menopause dengan Tingkat Depresi pada Wanita Usia 45-60 Tahun di Puskesmas Jumpang Baru Periode Juni 2016.....	215
<i>Reza Kumiawan Arta, Andi Alfian Zainuddin</i>	

Kumpulan Materi Workshop

Immunohistochemistry: Principles and Methods	218
<i>Rahmawati Minhajat</i>	
Preparation Of Histological Slide for Light Microscopy	218
<i>Triani Hastuti Hatta</i>	
Item Development dan Item Review Soal MCQ Berbasis Kompetensi	219
<i>Sri Asriyani</i>	
Cognitive Assessment in Anatomy	220
<i>Irwin Aras</i>	

HUMAN ENDOTHELIAL PROGENITOR CELL: BIOLOGICAL CHARACTERISTICS AND CLINICAL APPLICATION

Rahmawati Minhajat**

[‡]Internal Medicine Department, ^{*}Histology Department
Medical Faculty Hasanuddin University, Makassar

Abstract

Endothelial progenitor cells (EPCs) are cells derived from bone marrow and circulate into the peripheral circulation, they are a population of adult stem cells. These cells have characteristics similar to stem cells, but their ability to proliferate and differentiate is more limited. EPC discovery has changed the old paradigm in the field of vascular biology and it brings huge implications in medicine as EPCs can mediate the processes of vasculogenesis and maintain the vascular integrity.

In 1997, Asahara et al, has successfully isolated EPC from peripheral blood for the first time. The cells have positive surface antigen for mucosialin (CD34) and vascular endothelial growth factor receptor-2 (VEGFR-2/FLK-1). In vitro, the cells have a potency to develop into mature endothelial cells and in vivo they have roles in neoangiogenesis. Shi et al, have demonstrated that mononuclear cells which contain bone-marrow-derived-CD34⁺ surface antigen can be mobilized to peripheral circulation and can differentiate into mature endothelial cells. Since the discovery, mononuclear cells of peripheral blood which have CD34⁺ surface antigen are believed to be derived from the bone marrow and the cells are originated from the same precursor of hematopoietic stem cells (hemangioblast) since both of them show the same surface antigen, i.e. which is positive to CD34, prominin-1 (CD133) and FLK-1.

The EPC discovery has changed the old paradigm in the field of vascular biology, which believes that the process of vasculogenesis exclusively occurs only during embryogenesis. Some evidences indicate that postnatal neovascularization is not only derived from proliferation, migration and remodeling of endothelial cells on vascular wall (angiogenesis), but it also involve EPC recruitment from bone marrow, a process that has been known as vasculogenesis. The discovery has brought huge implications in medicine as EPCs can mediate the processes of vasculogenesis and maintain the vascular integrity.

Endothelial progenitor cells brings huge implications in the medical field as an EPC can mediate processes vasculogenesis and maintain the integrity of blood vessels. Thus, EPC could potentially be used for treatment of disease caused by endothelial dysfunction and other ischemic disease. Increasing the amount of EPC in the circulation is important because it positively correlates with reendothelialisasi and neovascularization.

I. Introduction

Endothelial progenitor cells (EPCs) are cells derived from bone marrow and circulate into the peripheral circulation, they are a population of adult stem cells. These cells have characteristics similar to stem cells, but their ability to proliferate and differentiate is more limited. EPC discovery has changed the old paradigm in the field of vascular biology and it brings huge implications in medicine as EPCs can mediate the processes of vasculogenesis and maintain the vascular integrity.¹

In 1997, Asahara et al² has successfully isolated EPC from peripheral blood for the first time. The cells have positive surface antigen for mucosialin (CD34) and vascular endothelial growth factor receptor-2 (VEGFR-2/FLK-1). In vitro, the cells have a potency to develop into mature endothelial cells and in vivo they have roles in neoangiogenesis. Shi et al³ have demonstrated that mononuclear cells which contain bone-marrow-derived-CD34+ surface antigen can be mobilized to peripheral circulation and can differentiate into mature endothelial cells. Since the discovery, mononuclear cells of peripheral blood which have CD34+ surface antigen are believed to be derived from the bone marrow and the cells are originated from the same precursor of hematopoietic stem cells (hemangioblast) since both of them show the same surface antigen, i.e. which is positive to CD34, prominin-1 (CD133) and FLK-1.

The EPC discovery has changed the old paradigm in the field of vascular biology, which believes that the process of vasculogenesis exclusively occurs only during embryogenesis. Some evidences indicate that postnatal neovascularization is not only derived from proliferation, migration and remodeling of endothelial cells on vascular wall (angiogenesis), but it also involve EPC recruitment from bone marrow, a process that has been known as vasculogenesis.⁴⁻⁶ The discovery has brought huge implications in medicine as EPCs can mediate the processes of vasculogenesis and maintain the vascular integrity.²

Endothelial progenitor cells (EPCs) are promising for cancer therapy because they specifically target tumors. They have the capacity to home to, invade, migrate within and incorporate into tumor structures. They are easily expanded and can be armed with therapeutic payloads protected within the progenitor cells. Once in the tumor, armed EPCs can be triggered to induce cell death in surrounding tumor cells while being transiently protected from premature demise. In preclinical studies, therapeutic EPCs attenuated tumor growth and increased survival. Enhancing homing, self-protection and collateral tumor cell damage will increase the efficacy of EPCs for cancer gene therapy.⁷

II. Source, Mobilization, Differentiation and Homing of EPCs

The main source of EPCs is bone marrow, but the cells can also be isolated from peripheral and umbilical blood. Investigators have isolated or generated EPCs from many sources: mouse embryos, mouse or human embryonic stem cells, fetal liver, human umbilical cord blood, postnatal bone marrow and peripheral blood.⁷ The advantage of embryonic EPCs is their unlimited proliferative capacity and the ease of genetic manipulation. These cells have the potential for systemic cancer gene therapy, as shown in a proof-of-principle study.⁸ However, mature embryonic stem cell derivatives are immunogenic and ethical considerations may limit their generation. Thus, additional sources of EPCs have been explored.

In normal condition, the number of EPCs from those various sources is very limited; while the mobilization from bone marrow and the number in the circulation are extremely affected by endogenous, exogenous factors as well as physiological and pathological conditions.⁹⁻¹¹ The release of EPCs from bone marrow is affected by various growth factors, enzymes, ligands and surface receptors. The natural response of body tissues when there is hypoxia is increasing the production and secretion of factors that stimulate neovascularization in order to reduce hypoxia. In hypoxia state, the hypoxia-inducible transcription factor-1 α (HIF-1 α) induces transcription of various proangiogenic proteins such as VEGF, stromal cell-derived factor-1 (SDF-1) and monocytes chemoattractant protein-1 (MCP-1), which will actively recruit EPCs from bone marrow to the circulation and then guide them to hypoxic site. Moreover, the local condition of bone marrow has important role in EPC mobilization. The cytokines of Granulocyte colony-stimulating factor (G-CSF), matrix metalloproteinases-9 (MMP-9), VEGF, SDF-1, endothelial nitric oxide synthases (eNOS) and Nitric oxide (NO) induce mobilization by interfering the interaction between EPC and the stroma cells of bone marrow, which allow EPCs to be released from bone marrow through endothelial sinusoid and entering the blood circulation. The process is the initial phase of EPCs mobilization from the bone marrow.^{12,13}

Morphologically, EPCs cannot be identified, but there are some specific markers that can be used. In the early phase, the cells showed positive response to CD133, CD34 and VEGFR-2. Cells that show positive response to those three markers are mainly still in the bone marrow; while the cells which are already present in peripheral circulation show diminishing expression of CD133; however, the expressions of CD34 and VEGFR-2 are still present. The next development is the advanced phase of EPC (the mature endothelial cells), in which the expression of CD34 has diminished, but the VEGFR-2 is still positive and the expressions of vascular endothelial cadherin (VE-cad), platelet endothelial cell adhesion molecule-1 (PECAM-1/ CD31) and von Willebrand factor (vWF) is initiated. Markers of EPCs mobilization include SDF-1, MMP-9, G-CSF and eNOS. Marker which shows the quantity of EPCs is the expression of total CD34; while the marker for EPCs quality are viable CD34, VEGFR-2 and EPC culture cells.¹³⁻¹⁵

III. EPCs in Physiological Vasculogenesis and Angiogenesis

During development, the vasculature is formed by vasculogenesis. In this process, endothelial progenitors differentiate to endothelial cells (ECs) that form a primary capillary network. Developmental vasculogenesis depends on endothelial progenitor cells (EPCs) that are derived from a common precursor of both the hematopoietic system and the vascular system called the hemangioblast. Vascular endothelial growth factor (VEGF) and the receptor VEGFR2 are pivotal for embryonic vasculogenesis as shown by the phenotype of both VEGFR2 and VEGF knockout mice, who die in utero owing to lack of endothelial and blood cells.⁷ In the adult, new vessels are built solely in healing wounds, the cycling endometrium and in growing tumors. Postnatal vessel formation was thought to proceed exclusively by sprouting from existing vessels (angiogenesis). However, it was shown that EPCs from the bone marrow participate in normal and pathological vessel formation in the adult (vasculogenesis).^{4,5} Adult EPCs proliferate rapidly and share other characteristics of embryonic angioblasts. They are recruited from the bone marrow by stimuli emanating from angiogenic sites.

At these sites, the EPCs differentiate into mature ECs that to some degree incorporate into vessels. Thus, vasculogenesis appears not to be restricted to embryogenesis but to be operative in adults also. Endothelial progenitor cells from the bone marrow participate in normal and pathological vessel formation in the adult (vasculogenesis). Angiogenesis and vasculogenesis may be complementary mechanisms for vessel formation in the adult.¹⁶

IV. ENDOTHELIAL DYSFUNCTION AND THE ROLE OF EPC

Endothelial cell is a layer of cell covering the inner side of vascular wall. The cell responds to each physical or chemical stimulation by releasing appropriate substances to maintain vasomotor balance and vascular hemostasis. A condition, in which the endothelial cell loses its ability to maintain the balance and when the endothelial cells have lost its physiological capacity to promote vasodilation, fibrinolysis and antiaggregation effect, is called as endothelial dysfunction. The pathophysiology of endothelial dysfunction is very complex and involves various mechanism through oxidative stress, i.e. a condition with increased reactive oxygen species (ROS) production and reduced antioxidant level. Increased ROS causes deprivation in NO availability, i.e. a substance produced by endothelial cells through eNOS activation.¹⁶ Nitric oxide not only has a role in EPCs mobilization, but it also has a role in EPCs migration and proliferation. Moreover, increased oxidative stress will initiate NF- κ B that may lead to inflammatory process as a beginning of atherosclerosis process. There is a dynamic correlation between inflammation, oxidative stress and EPCs mobilization. Oxidative stress stimulates inflammatory process, in which proinflammatory cytokines will stimulate the production of growth factors such as VEGF, that will further stimulate EPCs mobilization to the circulation. Inflammation may have good and bad impacts on EPCs. The limited and temporary inflammatory response can stimulate EPCs mobilization; while excessive and continuous response will cause reduced EPCs in the circulation. An in vitro study has demonstrated that inflammation and oxidative stress may affect EPC mobilization. The presence of continuous factors causes damages or continuous endothelial dysfunction which may ultimately cause reduction or exhaustion in EPCs supply. The number of EPCs in the circulation has negative correlation with the degree of endothelial dysfunction in patients with various cardiovascular risk factors. There is a correlation between the number of EPCs and endothelial dysfunction, i.e. reduced number of EPCs can predict the incidence of cardiovascular disease in the future. The role of EPCs in vascular repair is through normalization of endothelial function and improvement of blood flow in injured vascular area. Since endothelial cells are mature cells which has poor capacity to proliferate, i.e. only about 0.01%,²⁰ therefore, EPCs play an important role in maintaining endothelial layer through reendothelialization and neovascularization. It also indicates that EPCs can be used as a potential therapy for treating endothelial dysfunction.

V. CONDITION AND FACTORS AFFECTING THE NUMBER AND FUNCTION OF EPCS

The number of EPCs in the circulation of healthy individual is very limited, i.e. less than 1% of total bone marrow cells and less than 0.01% of total mononuclear cells in peripheral blood; however various factors and conditions can affect its number and function.⁶ Evidences of some studies show that patients with cardiovascular risk factors such as age, sex, smoking habits, hypertension, diabetes mellitus (DM) and dyslipidemia have reduced number and function of EPCs. In contrast, some cytokines, hormones, medicines and physical activity can improve its

Pertemuan Ilmiah Tahunan Pertemuan Ahli Anatomi Indonesia (PAAI) 2017

number and function. A study has demonstrated that the number of EPCs is increased significantly for a short of time after acute myocardium infarction (AMI) attack and it reaches peak level on the seventh day after the attack and subsequently, the level decreases. Another study that evaluated the number of CD34+ cells in patients with heart failure shows that there is increasing biphasic response, i.e. increased CD34+ cells in patients with Class I New York Heart Association (NYHA), but it is reduced in patients with Class IV NYHA and the number is even lower when compared with control. There are physiological conditions affect the number and function of EPCs, which include different age, sex and physical activity. Estrogen has been known as a rapid stimulator for endothelial NO production and eNOS activation. In addition, estrogen can also lower the level of endogenous asymmetric dimethylarginin (ADMA), which is the inhibitor of eNOS. Fandini et al²⁵ have reported that the number of EPC is greater in fertile female than male subjects; however, the difference has not been found in postmenopausal women compared to men of the same age. The difference can represent the risk of cardiovascular risk in men and postmenopausal women, in which it is correlated to the low number of EPCs. In aging process, there is also an imbalance between production of free radicals and the availability of antioxidant. There is a correlation between ROS, inflammation and age, i.e. increased ROS level will potentially stimulate chronic inflammation that ultimately will cause impaired EPCs mobilization. Thus, it has also been reported that there is reduced response of EPC migration to VEGF in elderly age as well as reduced clonogenic capacity of EPC that has begun starting from the middle age. Heiss et al²⁷ have reported that the survival, migration and proliferation capacity of the EPCs are lower in those with older age. The findings show that there is a negative effect of age on EPC differentiation and proliferation, which is correlated to cardiovascular risk. Physical exercise causes increased NO production. In a study that evaluate the effect of physical exercise on EPC in patients with stable coronary artery disease (CAD), it found that the number of EPC in the circulation increases significantly and the apoptosis is reduced after physical exercise for 28 days compared to the first day of exercise. Increased EPC number is correlated to higher NO availability after physical exercise. The study demonstrates that physical exercise increases the number of EPC in bone marrow and peripheral blood, in which the upregulation of EPC in physical exercise depends on NO and VEGF.²⁸ Smoking is one of risk factors causing endothelial dysfunction. The high level of oxidative stress in smoker can potentially affect EPC mobilization and survival in vivo, in which there is 50% reduction of EPC number in smokers compared to the control. Hypertension is also characterized by endothelial dysfunction and reduced NO availability. Endothelial dysfunction in patients with early stage of hypertension can induce the development of EPCs mobilization factor as the mechanism of compensation in the body. In the advanced stage of hypertension, which has continuous process of oxidative stress and inflammation, it will give contrary effect, i.e. reduced quality and quantity of EPCs.^{2,13} In patients with hypertension there is accelerated aging of EPCs and reduced activity of telomerase that affect vascular remodeling. Dysfunction of EPCs in hypertension causes endothelial repair and neoangiogenesis cannot take place and it will worsen the microvascular abnormality and atherosclerosis, which is the beginning of target organ damage in hypertension. In a study of subjects with refractory hypertension, the number of EPCs was reduced to 76.7% compared to the control. Angiotensin II accelerates the onset of EPCs aging through increased oxidative stress. A clinical trial in CAD patients has demonstrated that ramipril can increase the number of EPCs 2-5 folds as well as its function. Treatment using angiotensin II receptor antagonist also increases

the number of EPCs significantly, in which olmesartan treatment can cause increased EPCs from 231±24 to 465±71; while irbesartan treatment increases the number from 196±15 to 300±23 after 4-week treatment.³⁵ In patients with type 1 and type 2 DM, there is reduced number and function of EPC in circulation. Tapper et al³⁶ have demonstrated that isolated EPCs from peripheral blood of patients with type 2 DM have reduced proliferation capacity in the culture up to 48% compared to normal individuals. Loomans et al³⁷ also have reported similar results in patients with type 1 DM, i.e. there is reduced EPCs number of 44% compared to the non-diabetics control group. There is a significant increase of EPCs number in DM patients who were treated with insulin or oral diabetic agents. Insulin stimulates NO production; while in insulin resistance, there is an imbalance between endothelial damage and the ability of repair. This condition is correlated to reduced availability of NO, increased ROS production and PI3K/Akt down-regulation. It is potentially disrupt the process of vascular repair, which is the role of EPCs, including impaired EPC mobilization from the bone marrow. Increased plasma cholesterol level has significant correlation with endothelial damage and dysfunction. It has been proven that the number of EPCs is reduced significantly in patients with hypercholesterolemia and the number of EPCs has negative correlation with the level of total cholesterol and low density lipoprotein (LDL) cholesterol. Anti-inflammatory agents and antioxidants such as HMG-CoA reductase inhibitors (statin) have been proven to have positive effects on EPC. Statin increases mobilization and function of EPC through upregulation of PI3K/ Akt. The effect of statin in mobilizing EPC to the circulation is correlated with increased reendothelialization and reduced neointima resulting in reduced restenosis. EPCs in tumor neovascularization and target for tumor therapy

The formation of new vessels within a tumor have two possible sources of endothelial cells are; migration and co-option of pre-existing vascular walls endothelial cells or recruitment of EPCs from the bone marrow. Once EPCs arrive at the site of tumor mass, they can take part in neovascularization in three ways: (1) EPCs are directly incorporated in new vessels, (2) EPCs differentiate in mature ECs, (3) EPCs produce and secrete proangiogenic factor and cytokines with paracrine effects. In cancer patients, the number of circulating EPCs is increased.^{17-19.}

The need of endothelial progenitors for tumor vasculogenesis is the fundamental reason why these cells are being considered for therapeutic targeting of the tumor vasculature. EPC specifically target tumors because they have the capacity to home to, invade, migrate within and incorporate into tumor structures. EPCs can be easily expanded and manipulated without showing problems of immunological intolerance and they are easily expanded and can be armed with therapeutic payloads protected within the progenitor cell. Armed EPCs can be triggered to induce cell death in surrounding tumor cells while being transiently protected from premature demise. Therapeutic EPCs attenuated tumor growth and increased survival. Enhancing homing, self-protection and collateral tumor cell damage will increase the efficacy of EPCs for cancer gene therapy.⁷

Strategies by targeting tumour-associated EPCs:¹⁹

- Blocking EPC mobilisation inhibits vasculogenesis and impairs the formation of macro-metastasis in vivo
- Blocking genes involved in the homing of EPCs to tumour vasculature may carry the potential for improving antiangiogenic and antitumor effects. Another anti-tumour strategy could be to arrest the mobilisation of EPCs from the bone marrow by inhibiting some of the factors involved in their recruitment.
- Transplantation of genetically modified bone marrow progenitors may represent a vehicle for the transport of cytotoxic genes
- EPCs can be genetically engineered ex vivo by transduction with retrovirus and lentivirus vectors, which allow long-term transgene expression

VI. Summary

The EPC discovery has changed the old paradigm in the field of vascular biology, which believes that the process of vasculogenesis exclusively occurs only during embryogenesis. Some evidences indicate that postnatal neovascularization is not only derived from proliferation, migration and remodeling of endothelial cells on vascular wall (angiogenesis), but it also involve EPC recruitment from bone marrow, a process that has been known as vasculogenesis. The discovery has brought huge implications in medicine as EPCs can mediate the processes of vasculogenesis and maintain the vascular integrity.

Endothelial progenitor cells brings huge implications in the medical field as an EPC can mediate processes vaskulogenesis and maintain the integrity of blood vessels. Thus, EPC could potentially be used for treatment of desease caused by endothelial dysfunction and other ischemic disease. Increasing the amount of EPC in the circulation is important because it positively correlates with reendotelialisasi and neovascularization.

EPCs from the bone marrow participate in normal and pathological vessel formation in the adult (vasculogenesis). The need of endothelial progenitors for tumor vasculogenesis is the fundamental reason why these cells are being considered for therapeutic targeting of the tumor vasculature

References

1. Minhajat R, Nilasari D, Bakri S. The Role of Endothelial Progenitor Cell in Cardiovascular Disease Risk Factors. *Acta Med Indones*. 2015;47(4):340-7
2. Asahara T, Murohara T, Sullivan A, et al. Isolation of putative progenitor endothelial cells for angiogenesis. *Science*. 1997;275:964-7.
3. Shi Q, Rafii S, Wu MH, et al. Evidence for circulating bone marrow-derived endothelial cells. *Blood*. 1998;92:362-7.
4. Smadja DM, Cornet A, Emmerich J, et al. Endothelial progenitor cells: characterization, in vitro expansion, and prospects for autologous cell therapy. *Cell Biol Toxicol*. 2007;23:223–39.
5. Zampetaki A, Kirton PK, Xu Q. Vascular repair by endothelial progenitor cells. *Cardiovasc Res*. 2008;78:413-21.
6. Eguchi M, Masuda H, Asahara T. Endothelial progenitor cells for postnatal vasculogenesis. *Clin Exp Nephrol*. 2007;11:18-25.
7. Debatin K-M, Wei J and Beltinger C. Endothelial progenitor cells for cancer gene therapy. *Gene Ther* 2008;15:780–786
8. Wei J, Blum S, Unger M, Jarmy G, Lamparter M, Geishauser A et al. Embryonic endothelial progenitor cells armed with a suicide gene target hypoxic lung metastases after intravenous delivery. *Cancer Cell* 2004; 5: 477–488.
9. Smadja DM, Cornet A, Emmerich J, et al. Endothelial progenitor cells: characterization, in vitro expansion, and prospects for autologous cell therapy. *Cell Biol Toxicol*. 2007;23:223–39.
10. Gross P, Herbrig K. Role of endothelial progenitor cells in cardiovascular. *Pathology Roczn Akad Med Białymst*. 2004;49:174-7.
11. Hristov M, Weber PC. Endothelial progenitor cells: characterization, pathophysiology, and possible clinical relevance. *J Cell Mol Med*. 2004;8:498- 508.
12. Krenning G, van Luyn MJ, Harmsen MC. Endothelial progenitor cell-based neovascularization: implications for therapy. *Trends Mol Med*. 2009;15(4):180-
13. Tousoulis D, Andreou I, Antoniadou C, et al. Role of inflammation and oxidative stress in endothelial progenitor cells function and mobilization: therapeutic implications for cardiovascular disease. *Atherosclerosis*. 2008;201:236-47.
14. Esper RJ, Nordaby RA, Vilarino LO, et al. Endothelial dysfunction: a comprehensive appraisal. *Cardiovasc Diabetol*. 2006;23(5):1-18.
15. Endemann DH, Schiffrin EL. Endothelial dysfunction. *J Am Soc Nephrol*. 2004;15:83-92.
16. Marçola M, Rodrigues CE. Endothelial Progenitor Cells in Tumor Angiogenesis: Another Brick in the Wall. *Stem Cells Int*. 2015; 2015: 832649.
17. Botelho MC, Helena Alves H. Endothelial Progenitor Cells in Breast Cancer. *Int J Immunother Cancer Res*. 2016; 2: 1–2
18. Yajuan Su, Lei Zheng, Qian Wang, et al. Quantity and clinical relevance of circulating endothelial progenitor cells in human ovarian cancer. *J Exp Clin Cancer Res*. 2010; 29(1): 27.
19. Flamini V, Jiang WG, Lane J, et al. Significance and therapeutic implications of endothelial progenitor cells in angiogenic-mediated tumour metastasis. *Crit Rev Oncol Hematol*. 2016 Apr;100:177-89



Certificate

THIS IS TO CERTIFY THAT

dr. Rahmawati Minhajat, Ph.D, Sp.PD-KHOM

has participated as **SPEAKER**
in the

**SYMPOSIUM OF ANNUAL SCIENTIFIC MEETING
PERHIMPUNAN AHLI ANATOMI INDONESIA (PAAI)
THEME "FROM BASIC TO CLINICAL APPLICATION"**

Held on August 11-12, 2017 at Clarion Hotel Makassar

dr. H. Ahmad Aulif Jusuf, AHK, PhD
Chairman of PAAI (Perhimpunan Ahli Anatomi Indonesia)

Speaker 8 SKP, Participant 6 SKP, Moderator 2 SKP, Committee 2 SKP
No.: 032/IDI-WIL/SS/VII/2017/CPD

dr. Muhammad Iqbal Basri, Sp.S, M.Kes
Chairman of the Committee