

Quinone Oxidoreductase 2 Serum Level that affecting the Psychotic Disorders of Methamphetamine Abusers

by Andi Jayalangkara

Submission date: 16-Mar-2022 03:57AM (UTC+0700)

Submission ID: 1785122605

File name: affecting_the_Psychotic_Disorders_of_Methamphetamine_Abusers.pdf (2.58M)

Word count: 4214

Character count: 25443

ISSN-0971-720X (Print) • ISSN-0973-1283 (Electronic)

29

Volume 20, Number 3 July-September 2020



Medico-Legal Update

An International Journal



www.medicolegalupdate.org

Medico-Legal Update

Editor-in Chief

6

Prof. (Dr) R K Sharma

Former Head, Department of Forensic Medicine & Toxicology
All-India Institute of Medical Sciences, New Delhi-110029
E-mail: medicolegalupdate@gmail.com

ASSOCIATE EDITOR

1. **S.K. Dhattarwal** (Professor)
Forensic Medicine, PGIMS, Rohtak, Haryana
2. **Dr. Adarsh Kumar** (Additional Professor)
Forensic Medicine, AIIMS, New Delhi
3. **Dr. Vijaynath V** (Associate Professor)
Forensic Medicine, Vinayaka Mission Medical college, Tamil Nadu
4. **Ms. Roma Khan** (5)
Forensic Sciences, INSAAF Mumbai
5. **Dr. Imran Sabri** (Assistant Professor)
Department of Bio-Medical Sciences, College of Medicine, King Faisal University, Saudi Arabia

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **B. N. Yadav** (Professor)
Forensic Medicine, BP Koirala Institute of Medical Sciences, Nepal
2. **Dr. Vasudeva Murthy Challakere Ramaswam** (Senior Lecturer)
Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur, Malaysia
3. **Babak Mostafazadeh** (Associate Professor)
Department of Forensic Medicine & Toxicology, Shahid Beheshti University of Medical Sciences, Tehran-Iran
4. **Dr. Sarathchandra Kodikara** (Lecturer)
Forensic Medicine Department of Forensic Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka

NATIONAL EDITORIAL ADVISORY BOARD

1. **Prof. N.K. Agarwal** (Professor) Forensic Medicine, UCMS, Delhi
2. **P.K. Chattopadhyay**, (Professor)
Forensic Sciences, Amity University, Noida
3. **Dalbir Singh** (Professor) Forensic Medicine, PGIMER, Chandigarh
4. **Dr. Harish Pathak**, Mumbai
5. **J. Gargi** (Professor) GGS Medical College, Faridkot
6. **P.C. Dikshit** (Professor)
Forensic Medicine, Jamia Hamdard Medical College, New Delhi
7. **Anil Mittal** (Professor)
Forensic Medicine, Vardhman Mahavir Medical college, New Delhi
8. **Balbir Kaur** (Professor)
Forensic Medicine, MM institute of Medical Sciences, Ambala
9. **Mukesh Yadav** (Professor) Forensic Medicine, School of Medical Sciences and research, Greater Noida
10. **T.K.K. Naidu** (Professor) Forensic Medicine, Prathima Institute of Medical Sciences Andhra Pradesh
11. **S. Das** (Professor) Forensic Medicine, Himalayan Institute of Medical Sciences Dehradun
12. **Col Ravi Rautji**, Forensic Medicine, Armed Forces Medical College, Pune
13. **Dr. Manish Nigam** (Professor and Head)
Department of Forensic Medicine & Toxicology Sri Aurobindo Institute of Medical Sciences, INDORE (M.P.)
14. **Dr. Shailesh Kudva** (Principal)
Rajasthan Dental College and Hospital Jaipur-302026
15. **Usmanganishah Makandar** (Associate Professor)
Anatomy, AIMS, Bhatinda
16. **Dr. Pratik Patel** (Professor and Head) Forensic Medicine, Smt NHL Municipal Medical College Ahmedabad
17. **Basappa S. Hugar** (Associate Professor)
Forensic Medicine, Ramaiah Medical College, Bangalore

NATIONAL EDITORIAL ADVISORY BOARD

18. **Dr. Vandana Mudda** (Awati) (Associate Prof)
Dept of FMT, M.R. Medical College, Gulbarga, Karnataka, India
19. **Dr. Harish Kumar. N.** (Associate Professor)
Dept. of Forensic Medicine, Sri Siddhartha Medical College, Tumkur
20. **Dr. Gowri Shankar** (Associate Professor)
Forensic Medicine, SNMC, Bagalkot
21. **Dr. Manjunath Badni** (Reader) Dept of Oral pathology Maharana Pratap college of Dentistry and Research Centre, Gwalior
22. **Dr. L. Ananda Kumar** (Associate Professor) Forensic Medicine, Rajiv Gandhi Institute of Medical Sciences, (RIMS), Kadapa
23. **Dr. Ramesh Nanaji Wasnik** (Associate Professor and Head)
Forensic Medicine Late B.R.K.M. Govt. Medical college, Jagdalpur
24. **Dr. Sachin Sinha** (Reader), Dept. of Oral Pathology & Microbiology
Daswani Dental College & Research Centre, Rajasthan
25. **Dr. Sasi Kanth**, Asst. Professor, A.C.S.R Government Medical College,
Nellore, Andhra Pradesh.

Medico Legal Update is a scientific journal which brings latest knowledge regarding changing medico legal scenario to its readers. The journal caters to specialties of Forensic Medicine, Forensic Science, DNA fingerprinting, Toxicology, Environmental hazards, Sexual Medicine etc. The journal has been assigned international standard serial number (ISSN) 0971-720X. The journal is registered with Registrar of Newspaper for India vide registration numbers 63757/96 under Press and Registration of Books act, 1867. The journal is also covered by EMBASE (Excerpta Medica Database) from 1997 and by INDEX COPERNICUS, POLAND. Medico legal update is a half yearly peer reviewed journal. The journal has also been assigned E-ISSN 0973-1283 (Electronic version). The first issue of the journal was published in 1996.

6 Website: www.medicolegalupdate.org

© All Rights reserved The views and opinions expressed are of the authors and not of the Medico Legal Update. The Medico Legal Update does not guarantee directly or indirectly the quality or efficacy of any products or service featured in the advertisement in the journal, which are purely commercial.

Editor

Dr. R.K. Sharma
Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Printed, published and owned by

Dr. R.K. Sharma
Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)

Published at

Institute of Medico-legal Publications
Logix Office Tower, Unit No. 1704, Logix City Centre Mall,
Sector- 32, Noida - 201 301 (Uttar Pradesh)



Medico-Legal Update

CONTENTS

Volume 20, Number 3

July-September 2020

1. Carbamate Poisoning – An Autopsy Study 1
Jinu.M, Thankamma.P.George, N.A.Balaram, Sujisha.S.S
2. Profile of Burn Deaths: A Study based on Postmortem Examination of Burn Cases at R.N.T. Medical College, Udaipur 8
Akhilesh Sharma, Manish Kumar Sharma, Ashish Kumar Garg
3. Psychological Autopsy: the Psychological Assessment of the Dead Individual's Role in the Death 13
Anand Kumar Vasudevan, Hanumantha, Lynne Eccleston
4. Legality In Hospitals- Medico-Legal Case Management 19
Ashwini Pandit, Dhvani Patel, Arushi Chawla
5. To Assess Association of Bullous Pemphigoid and Neurological Disorders- A Clinical Study 24
Balkrishna Pralhadrao Nikam, Dhanraj D Chavan, Nischhal Shrivastava
6. To Assess Clinical Profile of Patients with Dermatophytosis Infection 28
Balkrishna Pralhadrao Nikam, Dhanraj D Chavan, Nischhal Shrivastava, Namita Narkhede
7. The Study of Trends and Out Come of Acute Poisoning in a Tertiary Care Hospital Khammam, Telangana; India 32
Bharath Kumar Guntheti, Kondru Laxman
8. Correlation between various Measurements of Sternum with Stature - An Autopsy based Study 40
Bhumesht Darade, Pawan Sabale, Shailesh Mohite, Swapnil Akhade, Chaitanya Kulkarni
9. Assessment of Depression among the Housewives of Alcohol Consuming Husbands in Selected Community Areas, Kancheepuram District, Tamilnadu 46
Bhuvana.A, Ajaykumar.P, Yaazhini.T, Dineshkumar.S, Devaki. K
10. A Study to Assess the Knowledge on Oral Health and Self Esteem among Young Adults in Selected College, in Kancheepuram District, Tamilnadu, India 51
Gini T.R , Ashika Parveen, Sharmilavathi.A, Avinash.U, Anto Benson, Shanthi
11. Knowledge and Attitude of Final Year Medical Students Towards Dentistry and Dental Consultation: A Questionnaire-based Study 55
Girish Suragimath, Mrunal B Mohite, Ashwinirani SR, Sameer Zope, Siddhartha Varma
12. Consensual Sexual Intercourse among children Vs Current Legal Provisions under POCSO: A Scientific Review with Prospective Quantitative Analytical Study 60
H. S. Tatiya, A. A. Taware, H. V. Vaidya, V. T. Jadhav, A. L. Bandgar, S. B. Punpale

X

128. Proximate Characteristics and Nutritional Value of White Anchovy Flour 744
Hendrayati, Theresia Dewi KB, Laras Budyghifari, Adriyani Adam
129. Distribution of RNA M. Leprae Household Contact in Endemic Area Indonesia 750
A. Rizki Amelia
130. Analysis of Environmental Risk Factors and Treatment Efforts Malaria Import Patients in Puskesmas in the Work Area of District Health Center of Segeri, Pangkep Regency 2019 757
Mugfira Mayangsari Putri, Hasanuddin Ishak, Indar, Anwar Mallongi
131. Formulation and Effectiveness Test of Matoa Stem (*Pometia Pinnata*) Skin Extract Cream Combination of Honey on *Staphylacoccus Aureus* Bacteria..... 763
Norma
132. Relationship between Level of Knowledge of Childbearing Age Women About Cervical Cancer and Behavior of Preventing Cervical Cancer in Mayamuk District, Sorong Regency 770
Sunaeni, S.ST, M.Keb
133. ¹⁹ The Affectivity of Knee and Hip Muscle Strengthening Exercise to Improve Muscle Strength, Balance, Pain, and Functional Activity of People with Osteoarthritis 775
Djohan Aras, Fatimatu Zahrah Syarif, Natalia Toding Padang, Nurhikmawaty Hasbiah, Herdin Rusli, Ita Rini
134. ⁸ Quinone Oxidoreductase 2 Serum Level that affecting the Psychotic Disorders of Methamphetamine Abusers 782
Dany H. Ludong, A. Jayalangkara Tanra, Mochammad Hatta, Sonny T. Lisal, Ismail Fikri, Reza Nasution, Tri Supartini
135. Application Pis (Indonesia Wellness Progsam) PK Model with Decision and Matriarchal Patriarchal Family of Compliance on the Family of the Suffering of Disease TB Lung Health in the Work Remu City Sorong 788
Maria Loihala
136. Decompression Sickness Indicators in Traditional Divers in Bajo, Boalemo District 794
Syamsiar S. Russeng, Lalu Muhammad Saleh, Muhammad Syafar Muliadi Rais
137. ³³ The Coparation between Exclusive Breastfeeding and Infant's Development at the Age of 6 Months in Rskia Pertiwi Makassar South Sulawesi Indonesia 800
Sundari, Andi masnilawati, Daswati, Lina Fitriani, Sitti Hadriyanti, Sumiaty, Nurlina Akbar
138. ²⁰ ¹⁶ Effects of Classical Music Therapy on Anxiety Level of Caesarean Section Mother 806
Sumarni Marwang, Rosita Passe, Rosdianah Aswan, Nurhidayat Triananinsi, Nahira Iskandar, Fadjriah Ohorella, Sumarni Sikki
139. ¹⁸ Effectiveness of Baby Spa and Music Therapy on Growth and Development of Baby 811
Sri Handayani, Anwar Mallongi
140. The Role of Kamoro Traditional Leaders in Communities Empowerment and Health in Mimika District 817
Vince Tebay, Anwar Mallongi

Quinone Oxidoreductase 2 Serum Level that affecting the Psychotic Disorders of Methamphetamine Abusers

Dany H. Ludong¹, A. Jayalangkara Tanra², Mochammad Hatta³, Sonny T. Lisal⁴, Ismail Fikri⁵, Reza Nasution⁶, Tri Supartini⁷

¹Doctor of Rehabilitation Center of Baddoka National Narcotics Agency, Makassar; ²Professor of Department of Psychiatry, Faculty of Medicine, Hasanuddin University, Indonesia; ³Professor of Department of Molecular Biology and Immunology, Faculty of Medicine, Hasanuddin University; ⁴ Professor of Department of Psychiatry, Faculty of Medicine, Hasanuddin University; ⁵Postgraduate School of Medicine Doctoral Study Program, Hasanuddin University; ⁶Postgraduate School of Medicine Doctoral Study Program, Hasanuddin University; ⁷Jaffray Bible College, Makassar, Indonesia

Abstract

Background: Some evidence suggests that genetic factors contribute to susceptibility to drug abuse such as methamphetamine and that dopamine-quinone produced by dopamine might be involved in the mechanism of symptoms associated with psychotic methamphetamine (MAP). Quinone detoxification is catalyzed by a group of proteins, namely quinone oxidoreductase (NQO). In this study we analyzed serum NQO2 levels by examining ELISA Sandwich in clients who used methamphetamine. **Methods:** This research is a Case-Control to determine the contribution of NQO2 to the emergence of psychotic disorders due to methamphetamine abuse. The case group is residents who consume methamphetamine with psychotic characteristics, while the control group is clients who consume methamphetamine without psychotic characteristics. **Results:** The mean value of NQO2 gene protein in the Elisa Sandwich examination for the case group was lower than the value of the control group (1,246 < 3,657). The t test showed the value of sig (Equal variances not assumed) = 0.00 that value is less than 0.05 (<0.05), this shows the difference in the value of content protein NQO2 between the case group and the control group. Then the cut-off point of the Receiver Operating Characteristic (ROC) curve in this study is recommended statistically between 1,960 and 2,165, meaning that if the expression value of protein content > 2.165 then the potential for psychotic suffering is great, on the other hand if < 1.960, the potential is small for suffering from psychotics. **Conclusion:** The protein content of the NQO2 gene in the Elisa Sandwich examination between the case and control groups is different, the case group is lower than the control. While the CURVE ROC (Receiver Operating Characteristic) Cut-Off value is recommended statistically between 1,960 and 2,165.

Keywords: Methamphetamine, Dopamine, NQO2 serum level, Methamphetamine Psychotic.

Introduction

Recently, methamphetamine abuse has become a major problem and developed into an international problem²⁰. According to the *United Nations Office on Drugs and Crime* (UNODC), methamphetamine is a dangerous substances and the 2nd ranks after hashish/cannabis in the 1st rank of the 20 types of substances that are often misused¹⁹. The prevalence of narcotics abuse in the world from 2006 to 2013 has increased¹⁹. In the last five years, the trend of amphetamine type has declined by around 15% in various countries, while the use of ecstasy

has been reported to be stable (12). However, there has been a drastic increase (158%) in the last five years namely the consumption of methamphetamine^{14,19,20}. Methamphetamine abusers can also cause psychotic symptoms similar to schizophrenia (Schizophrenia-like psychosis) but there are also methamphetamine abusers who do not experience psychotic symptoms^{24,25}. As a reference from several previous studies that contributed to our research, there are two sources of reference that we can point out, namely:

a. Functional polymorphisms of the NQO2 gene is associated with methamphetamine psychosis. (Ohgake, S. at all,^{14,17}.

b. Cognitive impairment is related to oxidative stress and chemokine levels in first psychotic episodes. (Martinez-Cengotitabengoa M at all, Schizophr Res 2012, 137:66–72).

7 **Material and Method**

Study Design and Subject

This research is a case-control to determine the effect of NQO2 gene on the onset of psychotic symptoms due to methamphetamine abuse^{1,22}. The Selection Criteria for clients who only consume methamphetamine with a diagnosis of Behavioral Mental Disorders due to the use of methamphetamine addictive substances with psychotic characteristics for the case group, while those who do not have psychotic characteristics as a control group and are currently undergoing inpatient rehabilitation programs at the BNN Baddoka Rehabilitation Center. Inclusion Criteria: Methamphetamine abusers, aged 20 - 50 years old, qualify as methamphetamine abuse. Exclusion Criteria: Clients suffering from Mental Disorders and Behavior due to addictive substances other

than methamphetamine. There is a history of hereditary disorders in the Psychotic group, suffering from Severe Physical Disease and consumption of polysubstance during the past year^{9,13}.

Population of Research: a. Case group: Methamphetamine users have Psychotic disorders, after being examined with the criteria of the Brief Psychiatric Rating Scale (BPRS) and the Diagnostic And Statistical Manual of Mental Disorder (DSM V).

b. Control group: Methamphetamine type who do not experience Psychotic disorders (examination instruments also with BPRS and DSM V).

2.3 Statistical Analysis

Analysis of the data in this study (Elisa’s protein content) using statistical application assistance, IBM SPSS version 24. To see the difference between the two samples, namely the case group and the control group, an independent t test was performed. Then to determine the cut-off point between psychotic and non-psychotic (Specificity and Sensitivity), by Receiver Operational Characteristics Curve (ROC) test and then performed AUC test (Area Under Curve) so that cut-off from data obtained in both groups.

Results

Table 1. Characteristic of subject based on age and gender

Distribution of case subjects (n = 30)			Distribution of control subjects (n = 30)		
Ages	n	%	Ages	n	%
20-35 yrs	27	90	20-35 yrs	27	90
36-50 yrs	3	10	36-50 yrs	3	10
Gender			Gender		
Men	29	96,7	Men	30	100
Women	1	3	Women	0	0

32 Table 1 shows that the age of the subjects in the case group ranged from 20-35 years (90%), most of the men were 29 (96.7%), and the rest were women.

T-Test

Independent t test is a test to see the difference between the two sample groups, namely cases and controls.

The mean value of Elisa protein levels in the case group (1,246) is smaller than the control group (3,657).

3


Table 2. Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
Content Sandwich Elisa	Equal variances assumed	10,668	,002	12,144	58	,000	2,41100	,19853	2,01359	2,80841
	Equal variances not assumed			12,144	49,684	,000	2,41100	,19853	2,01217	2,80983

15

The result of *t-test* showed the sig. value of 0.000 was less than 0.05 (< 0.05). It shows that H_0 rejected and H_a accepted, meaning that there were significant differences between case and control groups.

ROC (Receiver Operating Characteristic) KURVE



17

Figure 1. ROC Curves in Case Control Groups

The ROC (Receiver Operating Characteristic) curve shows that the value is far from the 50% line and close to 100%.

27

Table 3. Area Under the Curve

Test Result Variable(s): PROTEIN LEVELS SANDWICH ELISA

10 Area	Std. Error ^a	Asymptotic Sig. ^b	Asymptotic 95% Confidence Interval	
			Lower Bound	Upper Bound
,986	,012	,000	,962	1,000

4

The test result variable(s): PROTEIN LEVELS SANDWICH ELISA has at least one tie between the positive actual state group and the negative actual state group. Statistics may be biased.

a. Under the nonparametric assumption

b. Null hypothesis: true area = 0.5

The AUC (Area under the Curve) value is 98.6% (IK95%; 96.2% - 100%), meaning that if the ELISA Sandwich protein content / content score is used to diagnose the presence or absence of psychotic symptoms due to methamphetamine abuse in 100 clients, then the right conclusion will be obtained at 98.6 clients.

11

Table 4. Coordinates of the Curve

Test Result Variable(s): PROTEIN LEVELS SANDWICH ELISA

No.	Positive if Greater Than or Equal To ^a	Sensitivity	1 - Specificity
25	1,9600	1,000	,100
26	2,1650	1,000	,067

The test result variable(s): PROTEIN LEVELS SANDWICH ELISA has at least one tie between the positive actual state group and the negative actual state group.

The smallest cut-off value is the minimum observed test value minus 1, and the largest cutoff value is the maximum observed test value plus 1. All the other cut-off values are the averages of two consecutive ordered observed test values.

Discussion

As is known that methamphetamine is one type of narcotics that are stimulant, if consumed will affect all

parts of the brain, especially in the limbic system, namely the Nucleus Acumbens and Ventral Tegmental Area to activate dopamine (DA), then dopamine is released¹⁶. from storage. divescular sites to the cytoplasm so that the amount of DA increases in the brain. Furthermore, DA can carry out automatic oxidation to produce DA-Quinones. DA-Kuinon is a very reactive molecule and easily produces semi-quinones that are not stable by the reduction process. Semi-quinones enter the redox cycle together with oxygen molecules to form reactive oxygen species (ROS).

Then, ROS and semi-quinone cause oxidative stress. Therefore, it can be said that this oxidative

stress plays an important role in the mechanism of MAP, where this is evidenced by the involvement of DA-Kuinon in the mechanism of MAP which induces neurotoxicity and psychotic symptoms in humans^{14, 15}. DA-quinone is usually detoxified by genes catalyzed by the Quinoneoxidoreductase (NQOs) protein group and also this gene protects brain cell damage from oxidative stress. In humans, genetic evidence indicates that Quinoneoxidoreductase (NQOs) has ²³ gene loci, both of which have been identified as **NAD (P) H-quinone oxidoreductase 1 (NQO1)** (Ross et al.) and **NRH quinoneoxiductase 2 (NQOs) NQO2**^{9,11}.

In a research article such as that conducted in Japan by Ohgake, S., et.al with the title “NQO2 gene polymorphism associated with methamphetamine psychosis”, in his research found a percentage (11.7%) D / D of NQO2 genes in patients with prolonged psychosis had double the control (4.8%). On the other hand, this study shows that the NQO1 polymorphism (Pro187Ser), in which this gene ⁵ is not associated with MAP abuse. The NQO1 gene **is proven to be a risk factor for breast, lung and leukemia**^{3, 14, 18}. It has been explained that methamphetamine is a type of narcotic that is stimulant, because it affects the ²²ase of excessive dopamine neurotransmitters from **Nucleus Accumbens (NA) and Ventral Tegmental Area (VTA) in the Limbic System** that causes psychotic disorders. Meanwhile, the NQO2 gene serves to detoxify the effects of excessive dopamine, as well as protect brain organs from the effects of excessive dopamine poisoning which can cause permanent brain damage¹⁴.

This NQO2 gene is “Helper”, so it is normatively assumed that the lack of regulation of the NQO2 gene causes expression of the protein content of the low NQO2 gene, this can cause psychotic disorders in methamphetamine users^{14,15}, due to insufficiency. detoxification function and protection of the NQO2 gene. Further studies such as those conducted by Ohgake et al. (regarding the NQO2 gene polymorphism), we then tried to conduct another study, namely how the influence or contribution of protein levels from the NQO2 gene to psychotic disorders in methamphetamine users^{6, 11}. Analysis of NQO2 gene protein content through Sandwich Elisa examination for occupants / perpetrators of methamphetamine abuse in the control group (Mental Disorders and Causative Behavior of Amphetamine

Type Non Psychotic Stimulant Type) and case groups (Mental Disorders and Behavior of Addictive Substances of Amphetamine Type of Stimulant with Addictive Substance ²⁵ Psychotic Type)) gives different results, where **the average value of the protein content in the control group is higher (3,657) than in the case (1,246)**. Whereas the recommended KURVE ROC (Receiver Operating Characteristic) cut-off is statistically recommended between 1,960 and 2,165, with a sensitivity value of 100% and specificity $\geq 90\%$, it means that a cut-off value of 1,960 to 2,165 which can be used as a reference for the presence or absence potential psychotic disorders in methamphetamine abuse

Conclusion

a. The mean value of the NQO2 gene protein level in the Sandwich Elisa examination for the case group ie MAP users = 1,246, lower than the value of the control ¹² group ie non Psychotic MA users = 3,657. b. The t-test results show the value of sig. (2-tailed) = 0,000 (smaller than alpha = 0.05 (<0.05)), so it can be assumed that **there** are differences between the samples in the case and control groups. c. Cut-Off CURVE ROC (Receiver Operating Characteristic) recommended statistically in both groups is between 1.960 and 2.165, meaning that if the expression value of protein content > 2.165 then the potential for psychotic suffering is great, on the other hand if < 1.960 , the potential is small for suffering from psychotics^{2,21}.

Source of Funding : This study was not sponsored by government or pharmaceutical company fund.

⁵ **Ethical Clearance :** Taken from University Ethical committee

Conflict of Interest: Nil

References

1. APA. The Diagnostic and Statistical Manual of Mental Disorders (5th ed.). Washington DC: American Psychiatric Association Publishing. 2013.
2. Barr, A. . et al. The need for speed: an update on methamphetamine addiction. *J Psychiatry Neurosci*, 2006; 31(5), 301–313.
3. Bevens, K. M. G. & T. D. L. S. M. W. & M. L. & S. F. S. H. E. G. & G. C. & R. A. Methamphetamine-

- Associated Psychosis. 2012; New York: Neuroimmune Pharmacol.
4. Cruickshank, CC.; Dyer, K. A review of the clinical pharmacology of methamphetamine. *Addiction*, 2009; 104(7), 1085–1099. Retrieved from <http://onlinelibrary.wiley.com/doi/10.1111/j.1360-0443.2009.02564.x/full>
 5. Daniel F Hermens, Dan I Lubman, Philip B Ward, Sharon L Naismith, I. B. H. Amphetamine psychosis: a model for studying. *MJA*, 2009; 190.
 6. Darke, S.; Kaye, S.; McKetin, R.; Duffou, J. Major physical and psychological harms of methamphetamine use. *Drug Alcohol Rev*, 2008; 27(3), 253–262.
 7. Gabbard, G. *Psychodynamic Psychiatry in Clinical Practice* (5th ed.). Virginia: American Psychiatric Association Publishing, 2014.
 8. Hatta, M. *Pelatihan Aplikasi Teknik Biologi Molekular dan Imunologi dalam Penelitian Bidang Kesehatan*. Makassar: Laboratorium Biologi Molekular dan Imunologi, 2018; FK UNHAS.
 9. Joewana, S. *Gangguan Mental dan Perilaku Akibat Penggunaan Zat Psikoaktif: Penyalahgunaan Napza/Narkoba* (2nd ed.). 2004; Jakarta: ECG.
 10. John. *PCP Direct ELISA*. Catalog Number MBS580067. 2017
 11. Long DJ II, J. A. NRHQuinone oxidoreductase2 (NQO2). *ChemBiol Interact*, 2000; 129, 99–112.
 12. Neil A Campbell, Jane B. Reece, dan L. G. M. *Biologi*. 2002; Jakarta: Erlangga.
 13. NIDA. *Genetics of addiction: A research update*. Retrieve dfrom<http://www.drugabuse.gov/tib/genetics/html>, 2008.
 14. Ohgake, S. K. H. E. S. Functional polymorphism of the NQO2 gene is associated with methamphetamine psychosis. *Addiction Biology*, 2005; 10, 145–148. <https://doi.org/10.1080/13556210500123423>
 15. Okubo T, Harada S, Higuchi S, M. S. Association analyses between polymorphisms of the phase II detoxification enzymes (GTSM1, NQO1, NQO2) and alcohol withdrawal symptoms. *Alcohol Clin Exp Res*, 2003; 27(8), 68–71.
 16. Oyler JM, Cone EJ, Joseph RE, Moolchan ET, H. M. Duration of detectable methamphetamine and amphetamine excretion in urine after controlled oral administration of methamphetamine to humans. Retrieved August 23, 2018, from <http://www.clinchem.org/cgi>
 17. Ross D, Kepa JK, Winski SL, Beall HD, Anwar A, S. D. (2000). Nad(p)h:Quinone oxidoreductase 1 (nqo1): Chemoprotection, bioactivation, gene regulation and genetic polymorphisms. *Chem Biol Interact*, 2002; 129, 77–79.
 18. Suryo. *Genetika*. Jakarta: Departemen P dan K Direktorat Jendral Pendidikan Tinggi. 1996
 19. UNODC, L. and S. S. *The Challenge of Synthetic Drugs in East and South-East Asia and Oceania : Trends and Patterns of Amphetamine-type Stimulants and New Psychoactive Substances*. 2015; Vienna.
 20. UU No.35, *Tentang Narkotika*. Indonesia. 2009
 21. Anwar Mallongi, Stang, Syamsuar, Hasnawati Amqam, Aminuddin Syam, Syahrul Basri, Muh. Saleh. 2019., *Target Cancer Risks Due To The Exposure From Silica Among The Communities Living Surround Cement Tonasa Industry Pangkep, Indonesia*. *Interciencia Journal*. 2019 44(11)
 22. Lequin, RM “Enzyme Immunoassay (EIA)/ Enzyme-Linked Immunosorbent Assay (ELISA)”. *Clinical Chemistry*. 2005; 51 (12): 2415–2418.
 23. Koes Irianto. *Biologi Molekuler*, Penerbit ALFABETA, Bandung, 2017; 546-547.
 24. Hatta, M, Eko E. Surachmanto, Andi Asadul Islam, Syarifuddin Wahid. Expression of mRNA IL-17F and sIL-17F in atopic asthma patients. *BMC Research Notes*. 2017; 10:202. DOI: 10.1186/s13104-017-2517-9. Published: 12 June (2017)
 25. Martínez-Cengotitabengoa M, Mac-Dowell KS, Leza JC, Micó JA, Fernandez M, Echevarría E, Sanjuan J, Elorza J, González-Pinto A: Cognitive impairment is related to oxidative stress and chemokine levels in first psychotic episodes. *Schizophr Res* 2012; 137:66–72

Quinone Oxidoreductase 2 Serum Level that affecting the Psychotic Disorders of Methamphetamine Abusers

ORIGINALITY REPORT

24%

SIMILARITY INDEX

20%

INTERNET SOURCES

12%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to University of Mosul Student Paper	2%
2	Submitted to Queen Margaret University College, Edinburgh Student Paper	2%
3	www.researchgate.net Internet Source	1%
4	Submitted to University of Strathclyde Student Paper	1%
5	repo.unand.ac.id Internet Source	1%
6	Submitted to Western Governors University Student Paper	1%
7	ir.ymlib.yonsei.ac.kr Internet Source	1%
8	repository.poltekkes-mks.ac.id Internet Source	1%

9	Internet Source	1 %
10	learnmem.cshlp.org Internet Source	1 %
11	Submitted to Panjab University Student Paper	1 %
12	journal.unesa.ac.id Internet Source	1 %
13	www.incliva.es Internet Source	1 %
14	Purnamanita Syawal, Budu Budu, Mochammad Hatta, Muhammad Nasrum Massi, Andi Muhammad Ichsan, Rahmawati Minhajat. "Comparison between the triamcinolone and bevacizumab subconjunctivals and changes in Interleukin-1 mRNA expression in pterygium", Journal of Taibah University Medical Sciences, 2022 Publication	1 %
15	eprints.eudl.eu Internet Source	1 %
16	Submitted to Forum Perpustakaan Perguruan Tinggi Indonesia Jawa Timur Student Paper	1 %
17	Tufan Çınar, Mert Ilker Hayıroğlu, Veysel Ozan Tanık, Emre Aruğaslan et al. "The predictive	1 %

value of the CHA2DS2-VASc score in patients with mechanical mitral valve thrombosis", *Journal of Thrombosis and Thrombolysis*, 2018

Publication

18

Submitted to Adtalem Global Education

Student Paper

<1 %

19

unhas.ac.id

Internet Source

<1 %

20

Ayça Solt Kırca, Tuba Kızılkaya. "Effects of music medicine on premenstrual symptoms levels and quality of life: A randomized controlled trial", *Complementary Therapies in Clinical Practice*, 2022

Publication

<1 %

21

Mariko Murakami, Yoshio Nagai, Ayumi Tenjin, Yasushi Tanaka. "Proposed cut-off value of CA19-9 for detecting pancreatic cancer in patients with diabetes: a case-control study", *Endocrine Journal*, 2018

Publication

<1 %

22

Submitted to Royal Agricultural College

Student Paper

<1 %

23

Xing Gong, Labanyamoy Kole, Karim Iskander, Anil K. Jaiswal. "NRH:Quinone Oxidoreductase 2 and NAD(P)H:Quinone Oxidoreductase 1 Protect Tumor Suppressor p53 against 20S

<1 %

Proteasomal Degradation Leading to Stabilization and Activation of p53", Cancer Research, 2007

Publication

24	www.burmalibrary.org Internet Source	<1 %
25	bestjournal.untad.ac.id Internet Source	<1 %
26	bmcp psychiatry.biomedcentral.com Internet Source	<1 %
27	ccforum.biomedcentral.com Internet Source	<1 %
28	ijmmu.com Internet Source	<1 %
29	repository.umi.ac.id Internet Source	<1 %
30	Andrea Toccaceli, Andrea Giampaoletti, Lucia Dignani, Carla Lucertini, Cristina Petrucci, Loreto Lancia. "The role of shock index as a predictor of multiple-trauma patients' pathways", Nursing in Critical Care, 2016 Publication	<1 %
31	digitalcommons.unl.edu Internet Source	<1 %
32	documents.mx Internet Source	<1 %

33 ejournal.undip.ac.id
Internet Source

<1 %

34 Marieke M. ter Wee, Birgit I. Lissenberg-Witte.
"A Quick Guide on How to Conduct Medical
Research", Springer Nature, 2019
Publication

<1 %

35 RUSSELL EISENMAN, M. L. DANTZKER, LEE
ELLIS. "Self Ratings of Dependency/Addiction
Regarding Drugs, Sex, Love, and Food: Male
and Female College Students", Sexual
Addiction & Compulsivity, 2004
Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On