

JPNR_-_S02_-_2023_-_121.pdf

by Arni Irawaty

Submission date: 28-Feb-2023 12:05PM (UTC+0700)

Submission ID: 2025018380

File name: JPNR_-_S02_-_2023_-_121.pdf (274.37K)

Word count: 5842

Character count: 33365

The Utilization Of Milk Fish Scales Chitosan For Bone And Soft Tissue Regeneration After Tooth Extraction : A Literature Review

Arni Irawaty Djais¹, Surijana Mappangara¹, Asdar Gani¹, Shinta Rahma Mansyur², Febrianty², Nurfitra Abd Fatah²

¹Department of Periodontics Faculty of Dentistry, Hasanuddin University, Makassar, Indonesia

²Periodontology Specialist Educational Program, Faculty of Dentistry, Hasanuddin University, Makassar, Indonesia

*Corresponding Author : irawatyarni1975@gmail.com

DOI: 10.47750/pr.2023.14.502.121

Abstract

Introduction : Chitosan from milkfish scale chitin is a polymer material that is biocompatible, biodegradable, and non-toxic. Chitosan has been used as an antimicrobial, anti-inflammatory, bioactivity with dental materials, hemostasis, wound healing, and bone repair. Wound healing after tooth extraction is a pathophysiological process involves proliferation, cell migration, and tissue remodeling. The ability of chitosan to regenerate tissue is use as a soft tissue regeneration material and scaffold for bone regeneration. This literature review aims to discuss the ability of chitosan contained in milkfish scales to regenerate bone tissue and soft tissue after tooth extraction.

Methods: Data were collected through article searches on PubMed and Cochrane published from 2016 to 2022. Data search was conducted using keywords: ((Tooth Extraction) OR (Bone Regeneration)) OR (Soft Tissue Healing) AND (Chitosan).

Result: An initial search of Pubmed and Cochrane found 392 articles. Furthermore, 357 articles were excluded because they did not meet the inclusion criteria desired by the author in the title and abstract, or there were duplications, so that 35 articles were obtained. Then from these 35 articles, 23 articles were excluded, and the final results were 12 articles that would be reviewed and included in the synthesis table.

Conclusion: Milkfish scale chitosan has the potential as an anti-inflammatory agent, stimulates the activity of fibroblasts, osteoblasts, and mesenchymal stem cell differentiation, and osteoconductive bone regeneration scaffold. However, the utilization will be better when combined with polymers/biomaterials, and/or other bioactive molecules to improve mechanical properties, protein absorption, and biomineralization of chitosan.

Keywords: Bone Regeneration, Chitosan, Tooth extraction, Wound healing

INTRODUCTION

The development of the use of natural materials in the field of dentistry is currently growing rapidly, one of which is chitosan. Chitosan is an acetylated, non-toxic, and biocompatible chitin-derived polymer material. Chitosan can be derived from crustacean shells, cell walls of fungi and algae, insect exoskeletons, and radula molluscs.¹ In addition to these sources, several studies have stated that milkfish scales can be used as the basic material for chitosan because they contain chitin.²⁻⁵ Milkfish (*Chanos chanos*) is a brackish water fishery product that is abundant in Indonesia and is widely consumed by the public and is known as a fish that has many bones. During this time, milkfish processing mostly only uses the meat, so fish scales are only considered as waste.^{6,7} In dentistry, chitosan has been widely used as antimicrobial, anti-inflammatory, bioactivity with dental materials, hemostasis, wound healing, and bone repair.⁸

Tooth extraction is an action that can cause minimal trauma by leaving a wound in the tooth socket. The process of wound healing after tooth extraction has the same principle as wound healing in general. Physiologically, the socket healing process can be divided into four successive overlapping phases, namely the hemostasis phase, the inflammatory phase, the proliferative phase, and the maturation/remodeling phase.^{9,10} The inflammatory phase will occur immediately after tooth extraction, where the wound will be dominated

by neutrophil cells, then a phagocytosis process occurs followed by macrophage cells. Macrophages will increase osteoclast activity and release proinflammatory cytokines, anti-inflammatory and growth factors, which will degrade the extracellular matrix, stimulate cell differentiation and proliferation for the restoration of damaged tissue, and continue the process of bone formation.^{11,12} Gupta et al reported that chitosan was effective in accelerating the process of wound healing and osteogenesis in tooth extraction sockets by enhancing the function of inflammatory cells, such as PMN leukocytes, macrophages, fibroblasts, and osteoclasts.¹³ Research on milkfish scale chitosan on bone and bone tissue regeneration has not been done much. Thus, this literature review will discuss the potential of milkfish scale waste chitosan for soft and hard tissue regeneration after tooth extraction through an overview of health research on chitosan.

MATERIAL AND METHOD

Data Source

Data were collected through article searches on PubMed and Cochrane published from 2016 to 2022. Data search was conducted using keywords: ((Tooth Extraction) OR (Bone Regeneration)) OR (Soft Tissue Healing)) AND (Chitosan).

Research criteria

A. Inclusion Criteria

- Published articles from 2016-2022
- Articles in English and Indonesian
- Articles assessing soft and hard tissue regeneration in tooth socket or bone defects or on cell activity

B. Exclusion Criteria

- Articles in the form of systematic reviews, literature reviews, books, meta-analysis, and case reports
- Full text is not available for free

Data collection

The data used in this review literature are secondary. The data is obtained from articles which are then reviewed based on the criteria made by the author.




Figure 1. Review journal search flowchart

RESULT

An initial search of Pubmed and Cochrane found 392 articles. Furthermore, 357 articles were excluded because they did not meet the inclusion criteria desired by the author in the title and abstract, or there were duplications, so that 35 articles were obtained. Then from these 35 articles, 23 articles were excluded, and the final results were 12 articles that would be reviewed and included in the synthesis table.

Table 1. The effect of the combination of chitosan with polymers/biomaterials and/or other bioactive molecules on soft and hard tissue regeneration

No	Reference	Preparation	Polymer/ Biomaterial Combination	Combination of Bioactive Molecules	Method	Results
1	Hendrijanti et al ¹⁴ (2018) The effect of combination spirulina- chitosan on angiogenesis, osteoclast, and osteoblast cells in socket models of hyperglycemic Rattus norvegicus	Gel	Combination of 20% chitosan and 12% spirulina		A laboratory study involved 36 Rattus norvegicus, divided into three groups (non diabetic Mellitus (DM), uncontrolled DM, and controlled DM) and further divided into six subgroups. The control group (K1, K2, and K3) was induced with 3% carboxymethyl cellulose Na, while the treatment group was induced with 12% spirulina and 20% chitosan. On day 14, the lower jaws of the rats were removed. Capillary lumen, osteoblasts, and osteoclasts were calculated by examining the hypothalamus-pituitary-adrenal examination and the results were analyzed using Shapiro-Wilk, Levene's, one-way ANOVA, and Tukey's post hoc difference test.	The combination of 12% spirulina and 20% chitosan increases angiogenesis and the number of osteoblast cells and decreases the number of osteoclasts in diabetic conditions. This combination can inhibit the accumulation of reactive oxygen species (ROS) so that it also inhibits the expression of pro-inflammatory cytokines (IL-1 β , IL-6, and TNF- α) which play a role in bone resorption. A decrease of IL-6 will increase the amount of osteoprotegerin that inhibits RANK and RANKL interactions. This condition causes a decrease in the number of osteoclasts. The osteoinductive properties and osteointegrability of chitosan stimulate bone regeneration.
2	Maryani et al ⁵ (2018) Effect of Gel Combination Platelet Rich Plasma and Chitosan to Increase Osteoblasts Bone Regeneration in	Gel		Platelet Rich Plasma	The true experimental post- test only control group design was carried out on 28 Wistar rats. The samples were divided into 4	There was a significant increase in the number of osteoblasts in the combination PRP and chitosan gel groups compared to the PRP, chitosan gel, and

<p>Wound Healing Post Extraction Teeth in Mice Wistar</p>				<p>groups, namely PRP gel, combination PRP and chitosan gel, chitosan gel, and Povidone iodine control group. The mandibular right incisor was extracted and treated according to the group. The number of osteoblasts in the post-extraction socket was observed microscopically after 14 days using IHC staining. Data were analyzed using the One-Way ANOVA parametric test followed by the Post Hoc LSD test.</p>	<p>povidone-iodine groups 14 days after tooth extraction.</p>
<p>3 Xu et al¹⁶ (2019) Influence of in vitro differentiation status on the in vivo bone regeneration of cell/chitosan microspheres using a rat cranial defect model</p>	<p>Microsphere</p>		<p>Bone-marrow-derived mesenchymal stromal cells (BMSCs)</p>	<p>Stromal mesenchymal cells (BMSCs) derived from mouse bone marrow were inserted into the apatite-coated chitosan microsphere. The constructs were osteogenically differentiated for 0, 7, 14, and 21 days followed by implantation of the in vivo calvarial defect for up to 8 weeks.</p>	<p>There was the closure of the bone defect in the chitosan scaffold group cultured with osteogenic media for 14 days, after eight weeks. The new bone is thick and almost covers the entire defect.</p>
<p>4 Gahne et al¹⁷ (2018) Comparative Investigation of porous nano-hydroxyapatite/chitosan, nano-zirconia/chitosan and novel nano-calcium</p>	<p>Composite</p>	<p>Nano hydroxyapatite/Nano ZnO₂ /Nano CaZrO₃</p>		<p>The porous composite scaffold was developed using the freeze-drying technique. Cell culture studies were carried out using mouse</p>	<p>The addition of bio-ceramicnanopowder to the chitosan scaffold increased mechanical strength, cell proliferation, and cell</p>

<p>5</p> <p>Saravanan et al¹⁸ (2018)</p> <p>Chitosan-based thermoresponsive hydrogel containing graphene oxide for bone tissue repair</p>	<p>Hydrogel</p>	<p>Glycerophosphate + graphene oxide</p>	<p>pre-osteoblast cells (OB-6). The sterilized samples were incubated with culture media for 5 hours at 37°C and 5% CO₂. The scaffold was then removed and 200 µl of cell suspension containing 50,000 to 105 cells was added to the top of the scaffold slowly so that the cells could spread over the entire surface, then incubated at 37°C and 5% CO₂ for 3 hours. The proliferation of pre-osteoblasts along the surface and into the scaffold was observed using a confocal laser scanning microscope (CLSM) (Leica, USA) after staining cells with calcein AM</p>	<p>The hydrogel is biocompatible with mesenchymal stem cells and is metabolically active. Hydrogels increase osteogenic differentiation of mouse mesenchymal stem cells by</p>
<p>zirconate/chitosan composite scaffolds for their potential applications in bone regeneration</p>			<p>spread on the scaffold.</p>	

				<p>rate, and bionumerization. In in vitro biological studies, mesenchymal stem cells derived from mouse bone marrow (BMSCs) were isolated. BMSCs were used for in vitro biocompatibility analysis and mouse mesenchymal stem cells (C3H10T1/2) were used for differentiation studies. BMSCs were homogeneously suspended in HEC solution (2 x 10⁵ cells / ml; 20 mg / ml HEC dissolved in DMEM), and mixed with CS / GP gel solution. To qualitatively assess the viability of the encapsulated cells, FDA and DAPI exogenous stains were performed to assess their distribution in the hydrogel. After 4 days, cell hydrogels were washed and fixed with 2.5% glutaraldehyde for 1 hour, washed with ethanol series (gradient elution).</p>	<p>upregulating Runx-related transcription factor 2 (Runx2), Alkaline phosphatase (ALP), collagen type I (COL-1), and osteocalcin (OC) in osteogenic conditions.</p>
--	--	--	--	---	--

				dried, gold-sputtered, and analyzed by SEM to assess cell distribution in the hydrogel.	
6	Keller et al ¹⁹ (2019) Preclinical safety study of a combined therapeutic bone wound dressing for osteoarticular regeneration	Composite	Poly-ε-caprolactone	BMP-2	<p>A 1.5 mm osteochondral round defect was induced with a short drill in the femur to subchondral bone bleeding (approx. 2 mm). A NanoMI - BMP2 membrane was placed at the bottom of the break, which was then filled with the hMSCs / hydrogel mixture and gelled by adding 102 mM calcium chloride (SigmaAldrich), for</p> <p>5 minutes. Rats in experimental group 1 (ARTCAR; n = 20 rats; 10 males and 10 females; 3.5 µl hydrogel containing 105,000 ± 10% cells) and group 2 (n = 20 mice; 10 males and 10 females; 3.5 µl control hydrogel) underwent the same procedure with different ingredients. After gelation, the articulation capsule is closed, the muscles and skin are sutured and the</p> <p>Bone and cartilage regeneration was seen in both experimental animals without cytotoxicity. BMP-2 is released gradually, thereby reducing the side effects of inflammation.</p>

				<p>wound is disinfected. Rats were observed daily for wound healing, leg mobility, morbidity, mortality, and signs of toxicity, and twice a week for each weight loss for up to 90 days. Hematocrit, hemoglobin concentration, erythrocyte count, leukocyte count, mean cell volume, and platelet count is determined in a blood sample. Histopathological analysis was performed on animals used for blood tests.</p>	
7	<p>Ansari-zadeh et al²⁰ (2019) Fabrication, modeling and optimization of lyophilized advanced platelet-rich fibrin in combination with collagen-chitosan as a guided bone regeneration membrane</p>	Membrane	Collagen	<p>Advanced platelet-rich fibrin (A-PRF)</p> <p>Response surface methodology (RSM) was used to design experimental conditions and to correlate effect parameters, including the weight ratio of chitosan/collagen (chi/coll) and A-PRF concentration to Young's modulus, the viability of mesenchymal stem cells (MSCs), and degree of membrane degradation.</p>	<p>Analysis of alkaline phosphatase (ALP) showed an increase in osteoblast differentiation due to the addition of A-PRF</p>

8	<p>Elango et al.²¹ (2019) Chitosan-Collagen 3D Matrix Mimics Trabecular Bone and Regulates RANKL-Mediated Paracrine Cues of Differentiated Osteoblast and Mesenchymal Stem Cells for Bone Marrow Macrophage-Derived Osteoclastogenesis</p>	Composite	Chondroitin sulfate + hydroxyapatite	<p>Collagen is cross-linked with chitosan, hydroxyapatite (H), and chondroitin sulfate (Cs), to produce a natural, bone-like 3D structure and to evaluate its effect on bone homeostasis using bone marrow mesenchymal stem cells, osteoblasts, and bone marrow macrophages. XRD and micro-CT data confirm the arrangement of H crystals in the three-dimensional (3D) chitosan-collagen-H-Cs (CCHCs) matrix and the three-dimensional structure of the matrix. The stimulatory osteoblastogenic and osteoclastogenic activities of the 3D matrix were identified using differentiated osteoblasts and osteoclasts.</p>	<p>There is increased osteoblast differentiation through increased cellular ALP and bone mineral. The matrix regulates RANKL which can promote osteoclastogenesis so that bone resorption is limited and bone regeneration is enhanced.</p>
9	<p>Sularsih et al.²² (2016) Influenced of Using Chitosan with Weight Molecule Different to Tumour</p>	Gel		<p>Rattus norvegicus strain Wistar was divided into three treatment groups.</p>	<p>Chitosan gel with high molecular weight showed an increase in the amount of TNF-α</p>

<p>10</p> <p>Gupta et al¹ (2019) Efficacy of Chitosan in promoting wound healing in extraction socket: A prospective study</p>	<p>Wound dressing</p>	<p>Expression of Necrosis Factor Alpha (TNF-α) in Wound Healing Post Extraction Teth Rattus Norvegicus</p>	<p>namely group I with chitosan gel treatment which has a high molecular weight, group II with chitosan gel treatment which has a low molecular weight, and group III without chitosan gel. Chitosan is applied to the tooth socket. The mandibular jaw was decapitated 3 and 4 days after treatment, then an anatomical histopathological examination was carried out to observe the angiogenesis process</p>	<p>expression that was greater than that of the treatment group with chitosan which had low molecular weight both at 3 and 4 days of observation. Chitosan with high molecular weight contains more N-acetyl so that it will stimulate macrophage cells to release more TNF-α cytokines. Also, the polymer chains are long so that the molecular bonds are getting stronger. The more N-acetyl monomer, the higher the effect of accelerating wound healing.</p> <p>5.2</p> <p>Chitosan is effective in accelerating wound healing and early osteogenesis of the post-extraction socket.</p>
---	-----------------------	--	--	--

					observing the lamina dura and extraction socket density.	
11	Gani et al. ²¹ (2022) Effectiveness of Combination of Chitosan Gel and Hydroxyapatite from Crabs Shells (<i>Portunus pelagicus</i>) Waste as Bone graft on Periodontal Tissue Regeneration through IL-1 and BMP-2 Analysis	Gel	Hydroxyapatite	Experimental laboratory research and clinical trials with posttest only control group design. Twenty-seven Wistar rats were divided into three groups. The femoral bone defect was made; the positive control group was given placebo gel, the positive control group was given BATAN hydroxyapatite, and the test group was given a combination of chitosan gel and hydroxyapatite crab shells. Wistar rats were sacrificed on days 7, 14, and 21, and the femur bone was then taken for immuno histochemical analysis to determine the levels of IL-1 and BMP-2. Kolmogorov mirnov test, Levene test, and one-way ANOVA analyzed the data.	On days 7, 14, and 21, the expression levels of IL-1 and BMP2 were significantly different between the three groups. The group added with chitosan gel and crab shell HA showed a faster decrease in IL-1 expression than the control group. BMP-2 expression increased in the test group compared to the control group.	
12	Djans et al. ²⁴ (2022) South Sulawesi Milkfish (<i>Chanos Chanos</i>) Scale Waste as a New Anti-inflammatory Material in Socket Preservation	Gel	Hydroxyapatite	This is a post-test-only control group design study. Thirty-two Cavia cobaya were divided into four groups: (1) Socket preservation using milkfish scales, chitosan, (2) milkfish scales + chitosan + bovine	On days 3, 7, 14, and 28, groups with chitosan added showed lower levels of TNF- α and a faster decrease in IL-6 expressions compared to those without chitosan.	

						<p>xenograft, (3) bovine xenograft as a positive control, and (4) placebo as a negative control, then were sacrificed on 3rd, 7th, 14th, and 28th days. The mandible jaw specimen was taken for immunohistochemical analysis to determine the levels of TNF-α and IL-6. The data were analyzed using the Kolmogorov-Smirnov test, Levene's test, and one-way analysis of variance.</p>	
--	--	--	--	--	--	---	--

LITERATURE REVIEW

Wound healing is a biological process of repairing or replacing functional tissue damaged by injury. This process involves inflammation and bone healing which sequentially goes through several stages, characterized by the presence of blood clots, migration and proliferation of mesenchymal stem cells, formation of granulation tissue, inflammatory infiltrates, angiogenesis, fibroblast proliferation and collagen synthesis which ends with bone remodeling.^{9,25}

1. Milkfish Scales Chitosan

Chitosan is a natural polysaccharide derived from N-deacetylated chitin and a semicrystalline polysaccharide consisting of glucosamine and N-acetyl glucosamine units in -(1-4)-2-acetamide-2-deoxy-D-glucopyranol bonds. The main source of chitosan is chitin which can come from the cell walls of fungi, crustacean exoskeletons, and insects.²⁶ The process of obtaining chitosan is carried out by chemical extraction methods including deproteinization, demineralization, and deacetylation of chitin.⁴ Chitosan has a composition of carbon (40.30%), nitrogen (6.35%), and hydrogen (5.83%), and is non-toxic, biodegradable, biocompatible, antibacterial, bioactivity and can be used in various forms of solutions, gels, pastes, mixtures, sponges, membranes, tablets, and micro-granules depending on the application.^{20,27} Fish scales are waste that has not been used optimally. Fish scales contain chitin, calcium, proximate, alkaloids, steroids, saponins, phenol hydroquinone, molisch, benedict, biuret, and ninhydrin. Fish scales can be used as raw material for chitin extraction and further modified into chitosan. Extraction and modification of chitin into chitosan begins with the stages of preparation, demineralization, deproteinization, and deacetylation. Fish scales with a content similar to bone contain potential antimicrobial materials that can be used as dental materials.²⁸ Chitosan from milkfish scales (*Chanos chanos*) has antimicrobial activity against *Candida albicans*. The higher the chitosan concentration given, the larger the diameter of the formed inhibition zone. Also, milk scales chitosan gel can also inhibit the growth of *Aggregatibacteria actinomycetemcomitans* and *Porphyromonas gingivalis* which are the bacteria that cause periodontitis, and the higher the concentration of milk scales chitosan gel, the higher the inhibition zone produced.^{2,29}

2. Effect of Chitosan on Hard Tissue Regeneration

One of the treatments for bone regeneration includes the development of a scaffold that resembles the composition and structure of the extracellular matrix of bone. Structures with high porosity and topography and suitable surface chemical properties can facilitate cell adhesion, cell growth, proliferation, diffusion of oxygen and nutrients, as well as facilitate the disposal of metabolic wastes that occur during the regeneration process.²⁴ Chitosan can be used as a drug, as a conductor of growth factors, and as a scaffold material for bone regeneration.^{26,30,31} Currently, chitosan has been combined with polymer materials or other natural materials, biomaterials, or bioactive molecules with the aim of increasing mechanical resistance, protein absorption, and protein biomineralization.^{1,17,21,32,33} The combination of bioceramic nanopowder and chitosan can increase the mechanical strength, cell proliferation, and cell dispersion on the chitosan scaffold.¹⁷ In addition, the formation of bone tissue due to the increased osteoconductive properties of the combination of hydroxyapatite and chitosan materials.³⁴ Recent research using natural chitosan sources from crab shells with a combination of hydroxyapatite showed an increase in the expression of BMP-2 which plays a major role in the differentiation of osteoblasts for bone tissue regeneration.²³

3. Effect of Chitosan on Soft Tissue Regeneration

Chitosan has been shown to promote tissue healing, stimulate platelet growth factor production, and exhibit antibacterial activity that can reduce postoperative discomfort.³⁵ Chitosan has an antimicrobial effect because it can cause destabilization of the outer membrane of gram-negative bacteria and permeabilization of the microbial plasma membrane. Chitosan can also increase granulation tissue and angiogenesis, and accelerate wound healing by enhancing the function of inflammatory cells, such as macrophages, polymorphonuclear leukocytes (PMN), and fibroblasts or osteoclasts.¹³ Chitosan is metabolized by lysozyme so that it can be broken down and can act as a tissue engineering scaffold because it has a similar structure to glycosaminoglycans and is hydrophilic.³⁶

The production of interleukin-8 (IL-8) due to fibroblast stimulation plays an important role in the process of chemotaxis and angiogenesis. Complement activation occurs by increasing the production of C5a which triggers the migration of neutrophils and monocytes to the vessel wall. Chitosan can activate macrophages, cytokine production, giant cell migration, and stimulate type IV collagen synthesis. In addition, chitosan can also promote the formation of adequate granulation tissue accompanied by angiogenesis and collagen fiber deposition.¹

DISCUSSION

Milkfish scale waste is very abundant in Indonesia, but its utilization is not optimal. Fish scales contain chitin, calcium, proximate, alkaloids, steroids, saponins, phenol hydroquinone, molisch, benedict, biuret, and ninhydrin. The content of chitosan in milkfish comes from chitin which can reach 37.4% after dehydration.³⁷ Chitosan consists of -1,4-linked N-acetyl-D and glucosamine-D units, which are natural cationic polysaccharides. N-acetyl glucosamine functions as an anti-inflammatory which is synthesized in the human body from glucose. Chitosan exerts an anti-inflammatory effect by inhibiting the expression of prostaglandin E2 and cyclooxygenase-2 proteins and attenuating the pro-

inflammatory cytokines tumor necrosis factor- α and interleukin-1 β , as well as increasing the expression of the anti-inflammatory cytokine interleukin-10.³⁶

Sularsih et al (2016) reported that chitosan gel can stimulate macrophage cells to release TNF- cytokines on days 3 and 4 so that it has the effect of accelerating the wound healing process.²² Similar results were found in the study of Gupta et al (2019) who reported that chitosan was effective in accelerating wound healing and early osteogenesis in post-extraction sockets.¹³ Chitosan has potential as a scaffold material because it is biocompatible, degraded by tissue formation, does not cause inflammatory and allergic reactions, adequate porosity, and low degradation products.³⁸⁻⁴⁰ The hydrophilic nature of chitosan can support cell adhesion and proliferation. Several in vitro studies have proven that chitosan can increase the adhesion and proliferation of osteogenic cells and mesenchymal stem cells.^{41,42} Osteogenic cell culture in chitosan produces extracellular matrix which is mineralized into bone tissue. Chitosan also enhances the osteogenic differentiation of mesenchymal stem cells.⁴³ This is shown in the study of Maryani et al¹⁵, Sularsih et al²², and Gupta et al¹³ which stated that the use of chitosan in dental sockets showed a significant increase in osteoblasts and collagen, as well as a decrease in the number of osteoclasts so that they could support bone formation in the sockets.

Chitosan has several disadvantages, such as low water solubility at neutral or high pH and low mechanical properties.^{39,42} Therefore, currently many studies are conducting surface modification of chitosan or combining chitosan with bioactive inorganic materials (hydroxyapatite, tricalcium phosphate) or synthetic polymers (poly (vinyl alcohol) and poly (ethylene glycol)) and natural polymers (collagen) with the aim of improve mechanical resistance, protein absorption, and biomineralization.^{32,42} Xu et al²² bioceramic nanopowder added to chitosan increased mechanical strength, cell proliferation, and cell spread resulting in a new bone that was thick and almost completely covered the defect 14 days after defect creation. Danilchenko et al³⁴ and Asdar Gani et al²³ also showed the formation of bone tissue due to the increased osteoconductive properties of the combination of hydroxyapatite and chitosan. Combinations with bioactive molecules, for example, PRP, BMP-2, BMSCs, and A-PRF carried out by each Maryaniet al¹⁵, Keller et al¹⁹, Xu et al¹⁶ showed there was a significant increase in activation and osteogenic differentiation compared to only using chitosan. In gel form, chitosan protects the wound area and has a cooling effect that can reduce pain. Several studies have shown that chitosan gel with a degree of acetylation of 80% -84% and a molecular weight of 150-252 kDa is the most optimal in increasing bone regeneration.²⁶

CONCLUSION

Chitosan from milkfish scales can be used as an anti-inflammatory agent and stimulates the activity of fibroblasts, osteoblasts, and mesenchymal stem cell differentiation. In addition, chitosan can also be used as a scaffold for osteoconductive bone regeneration, but optimization of its utilization can be done by combining it with polymers/biomaterials and/or bioactive molecules to improve mechanical properties, protein absorption, and biomineralization of chitosan.

ACKNOWLEDGEMENT

We are very thankful to all the staff members of the Department of Periodontics, Hasanuddin University for their valuable suggestions and encouragement during making this journal

REFERENCES

1. Matica MA, Achmann FL, Tøndervik A, Sletta H, Ostafe V. Chitosan as a wound dressing starting material: Antimicrobial properties and mode of action. *Int J Mol Sci*. 2019;20(23):1–34.
2. Achmad H, Djais AI, Jannah M, Carmelita AB, Uinami H, Arifin EM, et al. Antibacterial chitosan of milkfish scales (*Chanos chanos*) on bacteria *prophyromonas gingivalis* & *agregatibacter actinomycetemcomitans*. *Syst Rev Pharm*. 2020;11(6):836–41.
3. Khaira Ummah Z, Sari N, Fortuna J, Boy E. Comparison of the Effectiveness of Milkfish Scales Chitosan with Gentamicin on the Development of *Escherichia Coli*. *Yars Med J*. 2017;25(2):108.
4. Djais A, Mappangara S, Gani A, Achmad H, Endang S, Tjokro J, et al. The effectiveness of Milkfish (*Chanos Chanos*) scales Chitosan on soft and hard tissue regeneration intooth extraction socket: A literature review. *Ann Rom Soc Cell Biol*. 2021;25(3):8729–52.
5. Ilmiyah Z, Rohmah J. Characterization of Chitosan Nanoparticles from Milkfish Scales as an Alternative Preservatives of Fresh Pangas Catfish (*Pangasius hypophthalmus*). *Medicra (Journal Med Lab Sci*. 2020;3(1):12–20.
6. Banggalino H, Akbar AMI. Utilization of Milkfish Scales as Raw Material for Chitosan. *SNP2M*. 2017;3(2):105–8.
7. Nur RM, Asy'ari A. The Utilitation of Fish Scale Waste as A Chitosan. *Agrikan J Agribisnis Perikan*. 2020;13(2):269–73.
8. Kmiec M, Pighinelli L, Tedesco M, MM Silva, Reis V. Chitosan-Properties and Applications in Dentistry. *Adv Tissue Eng Regen Med Open Access*. 2017;2(4):205–11.
9. Vieira AE, Repeke CE, De Barros Ferreira S, Colavite PM, Bigueti CC, Oliveira RC, et al. Intramembranous bone healing process subsequent to tooth extraction in mice: Micro-computed tomography, histomorphometric and molecular characterization. *PLoS One*. 2015;10(5):1–22.
10. Gomes P de S, Daugeia P, Poskevicius L, Mariano L, Femandes MH. Molecular and Cellular Aspects of Socket Healing in the Absence and Presence of Graft Materials and Autologous Platelet Concentrates: a Focused Review. *J Oral Maxillofac Res [Internet]*. 2019 Sep 5 [cited 2021 Mar 17];10(3):3–5. Available from: /pmc/articles/PMC6788423/
11. Ismardianita E, Elianora D, Rosalina W, Nofrike L, Khairani VY. The effectiveness methanol extract *clausena excavate* on number of fibroblast and density of collagen fibers after tooth extraction. *J Dentomaxillofacial Sci*. 2019;4(3):170–5.

12. Lunardi LC, Kresnadi U, Agustono B. The effect of a combination of propolis extract and bovine bone graft on the quantity of fibroblasts, osteoblasts and osteoclasts in tooth extraction sockets. *Dent J (Majalah Kedokt Gigi)*. 2019;52(3):126.
13. Gupta A, Rattan V, Rai S. Efficacy of Chitosan in promoting wound healing in extraction socket: A prospective study. *J Oral Biol Craniofacial Res*. 2019 Jan;9(1):91–5.
14. Nike Hendrijantini, Rostiny, Mefina Kuntjoro, Kent Sidharta, Dea Syarifina Putri Wiyono, Alocitta Anindyanari SS. **The Effect of Combination Spirulina-Chitosan on Angiogenesis, Osteoclast, and Osteoblast Cells in Socket Models of Hyperglycemic Rattus Norvegicus**. *Contemp Clin Dent*. 2019;9(4):582–6.
15. Maryani I, Rochmah YS, Parmana AD. Analysis of Platelet Rich Plasma and Chitosan Combination Gel on Increasing Number of Osteoblasts as Bone Regeneration in Wistar Rats Post Tooth Extraction Wounds. *ODONTO Dent J*. 2018;5(2):89.
16. Xu F, Wu Y, Zhang Y, Yin P, Fang C, Wang J. Influence of in vitro differentiation status on the in vivo bone regeneration of cell/chitosan microspheres using a rat cranial defect model. *J Biomater Sci Polym Ed [Internet]*. 2019;30(12):1008–25. Available from: <https://doi.org/10.1080/09205063.2019.1619959>
17. Gaihe B, Jayasunya AC. Comparative investigation of porous nano-hydroxyapatite/chitosan, nano-zirconia/chitosan and novel nano-calcium zirconate/chitosan composite scaffolds for their potential applications in bone regeneration. *Mater Sci Eng C [Internet]*. 2018;91(2017):330–9. Available from: <https://doi.org/10.1016/j.msec.2018.05.060>
18. Saravanan S, Vimalraj S, Anuradha D. **Chitosan based thermoresponsive hydrogel containing graphene oxide for bone tissue repair**. *Biomed Pharmacother [Internet]*. 2018;107(July):908–17. Available from: <https://doi.org/10.1016/j.biopha.2018.08.072>
19. Keller L, Pijnenburg L, Idoux-Gillet Y, Bornert F, Benameur L, Tabrizian M, et al. Preclinical safety study of a combined therapeutic bone wound dressing for osteoarticular regeneration. *Nat Commun [Internet]*. 2019;10(1):1–10. Available from: <http://dx.doi.org/10.1038/s41467-019-10165-5>
20. Ansarizadeh M, Mashayekhan S, Saadatmand M. Fabrication, modeling and optimization of lyophilized advanced platelet rich fibrin in combination with collagen-chitosan as a guided bone regeneration membrane. *Int J Biol Macromol [Internet]*. 2019;125:383–91. Available from: <https://doi.org/10.1016/j.ijbiomac.2018.12.078>
21. Elango J, Saravanakumar K, Rahman SU, Henrotin Y, Regensten JM, Wu W, et al. **Chitosan-collagen 3d matrix mimics trabecular bone and regulates rankl-mediated paracrine cues of differentiated osteoblast and mesenchymal stem cells for bone marrow macrophage-derived osteoclastogenesis**. *Biomolecules*. 2019;9(5).
22. Sularsih, Soeprijanto. Comparison of the number of osteoblast cells in wound healing between the use of 1% and 2% chitosan gel. *J Mater Kedokt Gigi* 2012;1(2):163. Available from: jurnal.pdgi.or.id/index.php/jmk/article/view/181/164
23. Gani A, Yulianty R, Supriaty S, Rusly M, Dwipa Asri G, Eka Satya D, et al. Effectiveness of Combination of Chitosan Gel and Hydroxyapatite from Crabs Shells (*Portunus pelagicus*) Waste as Bonegraft on Periodontal Network Regeneration through IL-1 and BMP-2 Analysis. *Int J Biomater*. 2022;2022.
24. Djais AI, Mappangara S, Gani A, Achmad H, Endang S, Tjokro J, et al. South Sulawesi Milkfish (*Chanos Chanos*) Scale Waste as a New Anti-inflammatory Material in Socket Preservation. *Open Access Maced J Med Sci*. 2022;10(D):221–8.
25. Sjuhada Oki A, Amalia N, Tantiana. Wound healing acceleration in inflammation phase of post-tooth extraction after aerobic and anaerobic exercise. *Sci Sport [Internet]*. 2020;35(3):168 e1–168 e6. Available from: <https://doi.org/10.1016/j.scispo.2019.06.001>
26. Jennings JA BJ. *Chitosan Based Biomaterials Vol 1 : Fundamental*. Duxford: Elsevier; 2017.
27. Gayen K, Pabale S, Shirotkar S, Sarkar S, Roychowdhury S. Chitosan biomaterials : Natural resources for dentistry. 2022;(February).
28. Rumengan IFM, Suptijah P, Salindeho N, Wullur S LA. Nanochitosan From Fish Scales. 2018. 117 p.
29. Dewi R, Nur RM, Nebore IDY. Antimicrobial Activity of Chitosan from Milkfish Scales (*Chanos chanos*) on the Oral Pathogen *Candida Albicans*. *Int J Nurs Heal Sci [Internet]*. 2019;6(4):54–8. Available from: <http://www.openscienceonline.com/journal/ijnhs>
30. Hartono BT, Firdaus FG. Utilization of Chitosan Biomaterials in the Field of Oral Surgery. *B-Dent J Kedokt Gigi Univ Baiturrahmah*. 2019;6(1):62–70.
31. Husain S, Al-Samadani KH, Najeeb S, Zafar MS, Khurshid Z, Zohaib S, et al. materials Chitosan Biomaterials for Current and Potential Dental Applications. 2017; Available from: www.mdpi.com/journal/materials
32. Georgopoulou A, Papadogiannis F, Batsali A, Marakis J, Alpentaki K, Eliopoulos AG, et al. Chitosan/gelatin scaffolds support bone regeneration. *J Mater Sci Mater Med*. 2018 May;29(5).
33. Khoshkhalgh P, Rabiee SM, Kiaee G, Heidari P, Miri AK, Moradi R, et al. Development and characterization of a bioglass/chitosan composite as an injectable bone substitute. *Carbohydr Polym [Internet]*. 2017;157:1261–71. Available from: <http://dx.doi.org/10.1016/j.carbpol.2016.11.003>
34. Danilchenko SN, Kalinkevich O V., Pogorelov M V., Kalinkevich AN, Sklyar AM, Kalinichenko TG, et al. Characterization and in vivo evaluation of chitosan-hydroxyapatite bone scaffolds made by one step coprecipitation method. *J Biomed Mater Res - Part A*. 2011;96 A(4):639–47.
35. Pippi R, Santoro M, Cafolla A. The Use of a Chitosan-Derived Hemostatic Agent for Postextraction Bleeding Control in Patients on Antiplatelet Treatment. *J Oral Maxillofac Surg [Internet]*. 2017;75(6):1118–23. Available from: <http://dx.doi.org/10.1016/j.joms.2017.01.005>
36. Ahmed S, Ikram S. Chitosan Based Scaffolds and Their Applications in Wound Healing. *Achiev Life Sci [Internet]*. 2016;10(1):27–37. Available from: <http://dx.doi.org/10.1016/j.als.2016.04.001>
37. Cadano JR, Jose M, Lubi AG, Maling JN, Moraga JS, Shi QY, et al. A comparative study on the raw chitin and chitosan yields of common bio-waste from Philippine seafood. *Environ Sci Pollut Res [Internet]*. 2020 [cited 2021 Feb 28]; Available from: <https://pubmed.ncbi.nlm.nih.gov/32198682/>
38. Levensgood SKL, Zhang M. Chitosan-based scaffolds for bone tissue engineering. *J Mater Chem B*. 2014;2(21):3161–84.
39. Dash M, Chiellini F, Ottenbrite RM, Chiellini E. Chitosan - A versatile semi-synthetic polymer in biomedical applications. *Prog Polym Sci [Internet]*. 2011;36(8):981–1014. Available from: <http://dx.doi.org/10.1016/j.progpolymsci.2011.02.001>
40. Keller L, Regiel-Futyr A, Gimeno M, Eap S, Mendoza G, Andreu V, et al. Chitosan-based nanocomposites for the repair of bone defects. *Nanomedicine Nanotechnology, Biol Med*. 2017;13(7):2231–40.
41. Aguilar A, Zein, Harmouch, Hafdi, Bornert, Offner, et al. Application of Chitosan in Bone and Dental Engineering. Belinha J, Natal Jorge RM, Reis Campos JC, Vaz MAP, Manuel J, Tavares RS, editors. *Molecules [Internet]*. 2019 Aug 19;24(16):3009. Available from: <https://www.taylorfrancis.com/books/9780429555848>
42. Ezoddini-Ardakani F, Navab Azam A, Yassaei S, Fatehi F, Rouhi G. Effects of chitosan on dental bone repair. *Health (Irvine Calif)*. 2011;3(3(4)):200–5.
43. Jain KG, Mohanty S, Ray AR, Malhotra R, Atran B. Culture & differentiation of mesenchymal stem cell into osteoblast on degradable biomedical composite scaffold: In vitro study. *Indian J Med Res*. 2015 Dec;142(6):747–58.

ORIGINALITY REPORT

22%

SIMILARITY INDEX

14%

INTERNET SOURCES

14%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1 m.scirp.org
Internet Source

1
%

2 www.em-consulte.com
Internet Source

1
%

3 academictree.org
Internet Source

1
%

4 www.nanobe.org
Internet Source

1
%

5 Jian Dong, Hiroko Kojima, Toshimasa Uemura, Masanori Kikuchi, Tetsuya Tateishi, Junzo Tanaka. "In vivo evaluation of a novel porous hydroxyapatite to sustain osteogenesis of transplanted bone marrow-derived osteoblastic cells", Journal of Biomedical Materials Research, 2001

1
%

Publication

-
- 6 Shahram Ghanaati, Patrick Booms, Anna Orłowska, Alica Kubesch et al. "Advanced Platelet-Rich Fibrin: A New Concept for Cell-Based Tissue Engineering by Means of Inflammatory Cells", *Journal of Oral Implantology*, 2014
Publication **1** %

-
- 7 Submitted to University of Western Australia
Student Paper **1** %

-
- 8 Ahmad Sukari, Chin Keong. "Chapter 12 Biomedical-Grade Chitosan in Wound Management and Its Biocompatibility In Vitro", *IntechOpen*, 2010
Publication **1** %

-
- 9 Claudia Cristina Biguetti, Franco Cavalla, Elicia M. Silveira, Angélica Cristina Fonseca et al. "Oral implant osseointegration model in C57Bl/6 mice: microtomographic, histological, histomorphometric and molecular characterization", *Journal of Applied Oral Science*, 2018
Publication **1** %

-
- 10 Hasanuddin Thahir, Arni Irawaty Djais, Mansjur Nasir, Ayu Rahayu Febalina, Afdalia Annisa, Nir Etriyani, Harun Achmad. "Virgin Coconut Oil as a New Concept for Periodontal Tissue **1** %

Regeneration via Expressions of TNF- α and TGF- β 1",
International Journal of Biomaterials, 2022
Publication

11 www.stipwunaraha.ac.id
Internet Source 1 %

12 Feng Rao, Yanhua Wang, Diaying Zhang, Changfeng Lu et al.
"Aligned chitosan nanofiber hydrogel grafted with peptides
mimicking bioactive brain-derived neurotrophic factor and
vascular endothelial growth factor repair long-distance sciatic
nerve defects in rats", Theranostics, 2020
Publication 1 %

13 Fateme Vasei, Farahnaz Sharafeddin. "Effect Of Chitosan
Treatment On Shear Bond Strength Of Composite To Deep
Dentin Using Self-Etch And Total-Etch Adhesive Systems",
Brazilian Dental Science, 2021
Publication 1 %

14 www.sci.muni.cz
Internet Source 1 %

15 acris.aalto.fi
Internet Source 1 %

16 Submitted to Johns Hopkins University
Student Paper

<1 %

17 Morteza Asadollahzadeh, Sayed Mahmood Rabiee, Hamed Salimi - Kenari. "In vitro apatite formation of calcium phosphate composite synthesized from fish bone", International Journal of Applied Ceramic Technology, 2019
Publication

<1 %

18 biologicalproceduresonline.biomedcentral.com
Internet Source

<1 %

19 Submitted to Consorcio CIXUG
Student Paper

<1 %

20 Sourav Sarkar, N.T. Prashanth, E.S. Shobha, Vinod Rangan, G. Nikhila. "Efficacy of Platelet Rich Fibrin versus chitosan as a hemostatic agent following dental extraction in patients on antiplatelet therapy", Journal of Oral Biology and Craniofacial Research, 2019
Publication

<1 %

21 Roberto Pippi. "Effectiveness of local hemostatic measures after tooth extraction in patients on antithrombotic therapy", Clinical Oral Investigations, 2019

<1 %

22 Submitted to University of Portsmouth
Student Paper

<1 %

23 repository.akfarsurabaya.ac.id
Internet Source

<1 %

24 www.science.gov
Internet Source

<1 %

25 M. Nabizadeh, M.R. Naimi-Jamal, M. Rohani, P. Azerang, A. Tahghighi. "Hydrazone analogues with promising antibacterial profiles: Synthesis, morphology, in vitro and in silico approaches", Letters in Applied Microbiology, 2022
Publication

<1 %

26 biocoreopen.org
Internet Source

<1 %

27 medicopublication.com
Internet Source

<1 %

28 Janela Rose Agpangan Limocon, Lyka Marie Cawaling Madalag, Paulyn Jade Balungcas Reliquias, Janina Veana Soriano Tionko et al. "Small but Terrible: Utilizing Chitosan-Based Nanoparticles as

<1 %

Drug Carriers to Treat Tuberculosis in the Philippines", *Frontiers in Pharmacology*, 2021
Publication

29 docplayer.net
Internet Source <1 %

30 ijb.whoice.com
Internet Source <1 %

31 unsworks.unsw.edu.au
Internet Source <1 %

32 www.electrospinning.co.uk
Internet Source <1 %

33 jbiomedsci.biomedcentral.com
Internet Source <1 %

34 bmcbiotechnol.biomedcentral.com
Internet Source <1 %

35 Andreia Espindola Vieira, Carlos Eduardo Repeke, Samuel de Barros Ferreira Junior, Priscila Maria Colavite et al.
"Intramembranous Bone Healing Process Subsequent to Tooth Extraction in Mice: Micro-Computed Tomography, <1 %

Histomorphometric and Molecular Characterization", PLOS ONE, 2015
Publication

36 www.unboundmedicine.com
Internet Source

<1 %

37 Shu-Jyun Cai, Ching-Wen Li, Daphne Weihs, Gou-Jen Wang.
"Control of cell proliferation by a porous chitosan scaffold with multiple releasing capabilities", Science and Technology of Advanced Materials, 2017
Publication

<1 %

38 Submitted to The Maldives National University
Student Paper

<1 %

39 www.ac1.hhu.de
Internet Source

<1 %

40 Xionglin Chen, Xiaoming Cao, He Jiang, Xiangxin Che, Xiaoyuan Xu, Baicheng Ma, Jie Zhang, Tao Huang. "SIKVAV-Modified Chitosan Hydrogel as a Skin Substitutes for Wound Closure in Mice", Molecules, 2018
Publication

<1 %

41 Georgia Vagropoulou, Maria Trentsiou, Anthie Georgopoulou, Eleni Papachristou et al. "Hybrid

chitosan/gelatin/nanohydroxyapatite scaffolds promote odontogenic differentiation of dental pulp stem cells and in vitro biom mineralization", Dental Materials, 2020
Publication

<1 %

42 Khoshakhlagh, Parastoo, Sayed Mahmood Rabiee, Gita Kiaee, Pedram Heidari, Amir K. Miri, Roshanak Moradi, Fathollah

Moztarzadeh, and Roya Ravarian. "Development and characterization of a bioglass/chitosan composite as an injectable bone substitute", Carbohydrate Polymers, 2016.
Publication

<1 %

43 Sumintarti Sugiharto, Sri Ramadany, Hendrastuti Handayani, Harun Achmad et al. "Assessment of the Anti-inflammatory

Activities of the Moringa Leaf Extract in Periodontitis Cases through IL-6 Cytokine Analysis in Wistar (Rattus norvegicus)", Open Access Macedonian Journal of Medical Sciences, 2022
Publication

<1 %

44 mail.aiims.edu
Internet Source

<1 %

45 repository.derby.ac.uk
Internet Source

<1 %

46 Ahmad Taha Khalaf, Yuanyuan Wei, Jun Wan, Jiang Zhu et al.
"Bone Tissue Engineering through 3D Bioprinting of Bioceramic Scaffolds: A Review and Update", Life, 2022
Publication

<1 %

47 Andi Zhang, Ya Liu, Di Qin, Mengjie Sun, Ting Wang, Xiguang Chen. "Research status of self-healing hydrogel for wound management: A review", International Journal of Biological Macromolecules, 2020
Publication

<1 %

48 Mary L Bouxsein, Stephen K Boyd, Blaine A Christiansen, Robert E Guldberg, Karl J Jepsen, Ralph Müller. "Guidelines for assessment of bone microstructure in rodents using micro-computed tomography", Journal of Bone and Mineral Research, 2010
Publication

<1 %

49 f1000research.com
Internet Source

<1 %

50 he01.tci-thaijo.org

<1 %

Internet Source

<1 %

51 **researchmap.jp**
Internet Source

<1 %

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

Off

10

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

