

**BUKTI  
SUBMIT  
MANUSKRIP**



Andi Dian Permana &lt;andi.dian.permana@farmasi.unhas.ac.id&gt;

## Thank you for your submission to Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

1 message

**Spectrochimica Acta Part A** <em@editorialmanager.com>  
Reply-To: Spectrochimica Acta Part A <support@elsevier.com>  
To: Andi Dian Permana <andi.dian.permana@farmasi.unhas.ac.id>

Thu, Aug 4, 2022 at 6:33 AM

Dear Dr. Permana,

Thank you for sending your manuscript Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: Application to in-vitro release and ex-vivo dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria. for consideration to Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy. Please accept this message as confirmation of your submission.

When should I expect to receive the Editor's decision?

We publicly share the average editorial times for Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy to give you an indication of when you can expect to receive the Editor's decision. These can viewed here: [http://journalinsights.elsevier.com/journals/1386-1425/review\\_speed](http://journalinsights.elsevier.com/journals/1386-1425/review_speed)

What happens next?

Here are the steps that you can expect as your manuscript progresses through the editorial process in the Editorial Manager (EM).

Currently your manuscript is with the Managing Editor for technical checks.

1. It will be assigned to an Editor and you will be sent a unique reference number that you can use to track it throughout the process. During this stage, the status in EM will be "With Editor".
  2. If your manuscript matches the scope and satisfies the criteria of Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, the Editor will identify and contact reviewers who are acknowledged experts in the field. Since peer-review is a voluntary service, it can take some time but please be assured that the Editor will regularly remind reviewers if they do not reply in a timely manner. During this stage, the status will appear as "Under Review".
- Once the Editor has received the minimum number of expert reviews, the status will change to "Required Reviews Complete".
3. It is also possible that the Editor may decide that your manuscript does not meet the journal criteria or scope and that it should not be considered further. In this case, the Editor will immediately notify you that the manuscript has been rejected and may recommend a more suitable journal.

For a more detailed description of the editorial process, please see Paper Lifecycle from Submission to Publication: [http://help.elsevier.com/app/answers/detail/a\\_id/160/p/8045/](http://help.elsevier.com/app/answers/detail/a_id/160/p/8045/)

How can I track the progress of my submission?

You can track the status of your submission at any time at <http://ees.elsevier.com/SAA>

Once there, simply:

1. Enter your username: Your username is: andipermana

If you need to retrieve password details, please go to: <https://www.editorialmanager.com/saa/l.asp?i=711227&l=KI4J3UIN>

2. Click on [Author Login]. This will take you to the Author Main Menu
3. Click on [Submissions Being Processed]

Many thanks again for your interest in Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy.

Kind regards,

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

If you require further assistance, you are welcome to contact our Researcher Support team 24/7 by live chat and

email or 24/5 by phone: <http://support.elsevier.com>

#AU\_SAA#

To ensure this email reaches the intended recipient, please do not delete the above code

---

In compliance with data protection regulations, you may request that we remove your personal registration details at any time. (Use the following URL: <https://www.editorialmanager.com/saa/login.asp?a=r>). Please contact the publication office if you have any questions.

**BUKTI  
REVIEW  
DARI  
REVIEWERS**



Andi Dian Permana &lt;andi.dian.permana@farmasi.unhas.ac.id&gt;

---

## Your Submission SAA-D-22-02512

2 messages

---

**Spectrochimica Acta Part A** <em@editorialmanager.com>  
Reply-To: Spectrochimica Acta Part A <support@elsevier.com>  
To: Andi Dian Permana <andi.dian.permana@farmasi.unhas.ac.id>

Sat, Dec 3, 2022 at 2:13 AM

Ms. Ref. No.: SAA-D-22-02512

Title: Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: Application to in-vitro release and ex-vivo dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria.

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

Dear Dr. Andi Dian Permana,

Reviewers have now commented on your paper. You will see that they are advising that you revise your manuscript. If you are prepared to undertake the work required, I would be pleased to reconsider my decision.

For your guidance, reviewers' comments are appended below.

If you decide to revise the work, please submit a list of changes or a rebuttal against each point which is being raised when you submit the revised manuscript.

To submit a revision, please go to <https://www.editorialmanager.com/saa/> and login as an Author.

Your username is: andipermana

If you need to retrieve password details, please go to:

[http://ees.elsevier.com/saa/automail\\_query.asp](http://ees.elsevier.com/saa/automail_query.asp)

NOTE: Upon submitting your revised manuscript, please upload the source files for your article. For additional details regarding acceptable file formats, please refer to the Guide for Authors at: <http://www.elsevier.com/journals/spectrochimica-acta-part-a-molecular-and-biomolecular-spectroscopy/1386-1425/guide-for-authors>

When submitting your revised paper, we ask that you include the following items:

Manuscript and Figure Source Files (mandatory)

We cannot accommodate PDF manuscript files for production purposes. We also ask that when submitting your revision you follow the journal formatting guidelines. Figures and tables may be embedded within the source file for the submission as long as they are of sufficient resolution for Production. For any figure that cannot be embedded within the source file (such as \*.PSD Photoshop files), the original figure needs to be uploaded separately. Refer to the Guide for Authors for additional information.

<http://www.elsevier.com/journals/spectrochimica-acta-part-a-molecular-and-biomolecular-spectroscopy/1386-1425/guide-for-authors>

Highlights (mandatory)

Highlights consist of a short collection of bullet points that convey the core findings of the article and should be submitted in a separate file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point). See the following website for more information

<http://www.elsevier.com/highlights>

Graphical Abstract (mandatory)

Graphical Abstracts should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership online. Refer to the following website for more information: <http://www.elsevier.com/graphicalabstracts>

On your Main Menu page is a folder entitled "Submissions Needing Revision". You will find your submission record there.

Finally, I would appreciate if you could submit your revised paper by Jan 31, 2023.

Note: While submitting the revised manuscript, please double check the author names provided in the submission so that authorship related changes are made in the revision stage. If your manuscript is accepted, any authorship change will involve approval from co-authors and respective editor handling the submission and this may cause a significant delay in publishing your manuscript.

Include interactive data visualizations in your publication and let your readers interact and engage more closely with your research. Follow the instructions here: <https://www.elsevier.com/authors/author-services/data-visualization> to find out about available data visualization options and how to include them with your article.

#### Research Elements (optional)

This journal encourages you to share research objects - including your raw data, methods, protocols, software, hardware and more – which support your original research article in a Research Elements journal. Research Elements are open access, multidisciplinary, peer-reviewed journals which make the objects associated with your research more discoverable, trustworthy and promote replicability and reproducibility. As open access journals, there may be an Article Publishing Charge if your paper is accepted for publication. Find out more about the Research Elements journals at [https://www.elsevier.com/authors/tools-and-resources/research-elements-journals?dgcid=ec\\_em\\_research\\_elements\\_email](https://www.elsevier.com/authors/tools-and-resources/research-elements-journals?dgcid=ec_em_research_elements_email).

Yours sincerely,

Christian Huck  
Editor  
Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

#### Reviewers' comments:

Reviewer #1: This manuscript describes a simple and effective way of quantifying chloramphenicol in biological samples. The information provided in this manuscript is novel and the results are discussed appropriately. I think that the manuscript should be accepted for publication after minor revision.

#### Specific comments:

1. Please change the colors of the UV spectra for "Blank skin tissue" and "CHL in skin tissue" it is difficult to differentiate them.
2. Could you please provide a brief explanation about the release mechanism from PCL particles? I assume that bacteria will accelerate PCL hydrolysis accelerating drug release. Is this correct or it might be related with the pH of the solution. Could you please provide some explanation about this? This could have been described before if so please include the citation to the relevant work.
3. I think that the selected mathematical models are not a good model for the presented release curves. They show a sigmoidal shape and these models are not good for this type of curves. I recommend to remove the mathematical modelling section from this paper as it is not adding any valuable information.
4. Figure 5 is missing the "A" in the legend caption. Please correct this typo

#### Reviewer #3:

Please go through the manuscript and review how the numbers are stated. Commas should be replaced by periods

Abstract - spell out PVA/PSB/TSB/LLOQ

Line 48-49 - check sentence, it seems to be missing a word

Line 66-67 - check sentence for correctness

Line 84 - spell out MP

Line 106-109 - this sentence needs to be clarified

Line 118 - missing "to" - was added to

Line 156 -clarify how it was dried

Line 179-180 - r<sup>2</sup> is the coefficient of determination, not correlation

Line 184 - the term gradient is correct but for consistency with the linearity section, gradient should be replaced by slope.

Sections 2.6.3 and 2.6.4 should be made more similar. This is then same equation, just with a different weight of the standard deviation of the blank. However, the descriptions are very different and thus confusing

Line 201 - "in in" one too many  
Section 2.7.5 title - check spelling of title "preparation"  
Line 247 - check sentence "put into for" is not clear  
Line 279 - add state for Microsoft (WA)  
Line 280-280. Remove period and include line 281 in the previous sentence  
Line 290. Check the sentence as it is missing a verb  
Line 302-304 - check sentences. Capital letter at And, "in in", "while ...)" is not a correct sentence.  
Figure 1. - the green color in the legend does not match the color on all the plots. The blank skin tissue and CHL in skin tissue is not present in plots B and D. Is it normal? The blank skin tissue and CHL in skin tissue have the same color.  
Line 326 - "will be concise" does not make sense here  
Table 4 - concentration difference. Do you mean range?  
Line 338 - r2 is determination, not coefficient  
Line 341 - what is exactly meant by variation coefficient?  
Table 5 to 9 - what does %RE mean?  
Line 427 - spell out LQC, MQC, HQC  
Table 10 - why are recoveries only 90%? When the method is used in later section, recovery of 100% is reached. Why? Discussing is needed here to explain why authors state the method is accurate when 10% active is missing  
Line 445 - after F2, replace comma by period  
Line 462 - does authors mean PCL composition in MPs?  
Line 496 - ". In Figure 4." THIs is not a sentence  
Figure 5 - add (A) at the beginning of the legend  
Line 520 - please confirm "please and dispersal" are correct terms  
Line 531 - what does CAB stand for?  
Line 545 - what does DOX stand for?

For further assistance, please visit our customer support site at <http://help.elsevier.com/app/answers/list/p/7923>. Here you can search for solutions on a range of topics, find answers to frequently asked questions and learn more about EM via interactive tutorials. You will also find our 24/7 support contact details should you need any further assistance from one of our customer support representatives.

#AU\_SAA#

To ensure this email reaches the intended recipient, please do not delete the above code

---

In compliance with data protection regulations, you may request that we remove your personal registration details at any time. (Use the following URL: <https://www.editorialmanager.com/saa/login.asp?a=r>). Please contact the publication office if you have any questions.

---

**Andi Dian Permana** <[andi.dian.permana@farmasi.unhas.ac.id](mailto:andi.dian.permana@farmasi.unhas.ac.id)>  
To: Mukarram Mudjahid <[mukarramfarmasist@gmail.com](mailto:mukarramfarmasist@gmail.com)>

Thu, Dec 15, 2022 at 5:52 PM

[Quoted text hidden]

**BUKTI  
SUBMIT  
HASIL  
REVIEW**



Andi Dian Permana &lt;andi.dian.permana@farmasi.unhas.ac.id&gt;

---

**Submission Confirmation for SAA-D-22-02512R1**

1 message

---

**Spectrochimica Acta Part A** <em@editorialmanager.com>  
Reply-To: Spectrochimica Acta Part A <support@elsevier.com>  
To: Andi Dian Permana <andi.dian.permana@farmasi.unhas.ac.id>

Fri, Dec 16, 2022 at 7:54 AM

Ms. Ref. No.: SAA-D-22-02512R1

Title: Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: application to in vitro release and ex vivo dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria.

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

Dear Dr. Andi Dian Permana,

Your revised manuscript was received for reconsideration for publication in Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy.

You may check the status of your manuscript by logging onto the Editorial Manager as an Author at <https://www.editorialmanager.com/saa/>.

Your username is: andipermana

If you need to retrieve password details, please go to:

[http://ees.elsevier.com/saa/automail\\_query.asp](http://ees.elsevier.com/saa/automail_query.asp)

Kind regards,

Editorial Manager  
Spectrochimica Acta Part A: Molecular and Biomolecular SpectroscopyFor further assistance, please visit our customer support site at <http://help.elsevier.com/app/answers/list/p/7923>. Here you can search for solutions on a range of topics, find answers to frequently asked questions and learn more about EM via interactive tutorials. You will also find our 24/7 support contact details should you need any further assistance from one of our customer support representatives.

#AU\_SAA#

To ensure this email reaches the intended recipient, please do not delete the above code

---

In compliance with data protection regulations, you may request that we remove your personal registration details at any time. (Use the following URL: <https://www.editorialmanager.com/saa/login.asp?a=r>). Please contact the publication office if you have any questions.

# Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

## Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: application to in vitro release and ex vivo dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria.

--Manuscript Draft--

<b>Manuscript Number:</b>	SAA-D-22-02512R1
<b>Article Type:</b>	Full Length Article
<b>Section/Category:</b>	Analytical Spectroscopy and New Methods
<b>Keywords:</b>	Bacterially Sensitive Microparticles; Cellulitis; Chloramphenicol; Microneedle; UV-vis Spectrophotometry; Validation
<b>Corresponding Author:</b>	Andi Dian Permana Universitas Hasanuddin Fakultas Farmasi Makassar, INDONESIA
<b>First Author:</b>	Mukarram Mudjahid
<b>Order of Authors:</b>	Mukarram Mudjahid Sulistiawati Sulistiawati Rangga Meidianto Asri Firzan Nainu Andi Dian Permana
<b>Abstract:</b>	<p>Cellulitis is a common dermis/subcutaneous tissue skin infection and shared global disease burden, with a higher incidence for males and people aged 45–64 years. Application therapy of chloramphenicol (CHL) has been hindered because of its toxicity and limited penetration into the skin. In this research, CHL was developed into a bacterially sensitive microparticles which were further incorporated into a microneedle system to increase penetration. To support this formulation, in this study, UV–visible spectrophotometry method was validated in methanol, polyvinyl alcohol (PVA) 1%, phosphate buffered saline (PBS), tryptic soy broth (TSB) (fluid-mimicking infection), and skin tissue to quantify amount of CHL. The developed analytical method was subsequently validated according to ICH guidelines. The results obtained showed that the correlation coefficients were linear <math>\geq 0.9934</math>. The values of LLOQ inside the methanol, PVA 1%, PBS, TSB, and skin tissue were 7.20 <math>\mu\text{g/mL}</math>, 4.40 <math>\mu\text{g/mL}</math>, 8.18 <math>\mu\text{g/mL}</math>, 387.48 <math>\mu\text{g/mL}</math>, and 7.27 <math>\mu\text{g/mL}</math>, respectively. The accuracy and precision of the developed method were prominent. These methods were successfully applied to quantify the amount of CHL in microparticle and microneedle system in fluid and tissue skin infection. The result showed the high drug release microparticle sensitive bacteria, and high drug retention in ex-vivo dermatokinetic evaluation in rat skin tissue containing bacterial infection. This was due to the presence of Staphylococcus aureus bacteria culture that produced lipase enzymes, playing a role in lysing microparticle matrix to develop selectively delivery antimicrobials. A further analytical method needs to be matured to quantify CHL inside the in vivo studies.</p>



MINISTRY OF EDUCATION CULTURE OF INDONESIA  
HASANUDDIN UNIVERSITY  
FACULTY OF PHARMACY

Alamat Jalan Perintis kemerdekaan Km.10, Makassar 90245  
Telepon (0411) 588556, Faksimili (0411) 590663  
Laman: farmasi@unhas.ac.id

---

The Editor

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy.

15<sup>th</sup> December 2022

Dear Sir/Madam,

I wish you to consider our manuscript entitled: **“Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: application to *in vitro* release and *ex vivo* dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria.”** for publication in journal of Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy.

We have made a few of changes to the manuscript as a result of the reviewers comments. We believe that the manuscript is now substantially improved. Importantly, we have made a great effort to addressed each of the reviewers comments in the response to the reviewer file.

We believe that this work will be of great interest to the readers journal Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy, particularly those working on material and polymeric compounds as medical purposes for selective drug delivery administration. The manuscript has not been previously published in any language anywhere and it is not under simultaneous consideration by another journal. We appreciate your attention. We hope you will now consider publishing our research in journal of Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy and look forward to hearing from you in due course.

Yours sincerely,

Dr. Andi Dian Permana (on behalf of all authors)

Department of Pharmaceutics, Faculty of Pharmacy, Hasanuddin University, Makassar,  
Indonesia

Email: [andi.dian.permana@unhas.ac.id](mailto:andi.dian.permana@unhas.ac.id)

Journal: Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

Manuscript ID: SAA-D-22-02512

Title: "Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: application to *in vitro* release and *ex vivo* dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria."

### **Response to Reviewers**

We are very thankful to the expert reviewers for taking the time to kindly review our manuscript and provide helpful comments for improvement and clarification. We have made some changes to the manuscript as a result of these comments. We believe that the manuscript is now substantially improved. We have addressed each of the reviewers' comments in detail below. Importantly, we have made a great effort to improve the English and the discussion parts of our revised manuscript.

**Reviewer #1:** This manuscript describes a simple and effective way of quantifying chloramphenicol in biological samples. The information provided in this manuscript is novel and the results are discussed appropriately. I think that the manuscript should be accepted for publication after minor revision.

1. Please change the colors of the UV spectra for "Blank skin tissue" and "CHL in skin tissue" it is difficult to differentiate them.

### **Response to Reviewer**

We are very thankful to the Reviewer for taking the time to review this manuscript and for the expert review, providing helpful comments. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 281)

2. Could you please provide a brief explanation about the release mechanism from PCL particles? I assume that bacteria will accelerate PCL hydrolysis accelerating drug release. Is this correct or it might be related with the pH of the solution. Could you please provide some explanation about this? This could have been described before if so please include the citation to the relevant work.

**Response:**

We are very thankful to the Reviewer for taking the time to review this manuscript and for the expert review, providing helpful comments.

PCL has been reported as a good option for a variety of biological and environmental applications. Particularly, PCL's enzymatic biodegradation has been proven for use in controlled drug administration and tissue engineering <sup>1</sup>. It has been also reported, that SA may release a particular enzyme called lipolytic esterase (Lipase), this enzyme can initiate the biocatalytic hydrolysis of PCL <sup>2</sup>.

We have included these in the revised manuscript, as follows:

The results obtained here indicate that CHL encapsulation in PCL MPs may result in an increase in CHL release via two different processes. The PCL outer layer initially aided in the MPs' attachment to the colony infection. Following this, the bacterial strains lipase released into the environment dissolved PCL layers, releasing CHL from MPs. Regarding the impact of pH on kinetical release of CHL in MPs-coated PCL, previous research has demonstrated the profile of nanoparticles (NPs) release utilizing PCL polymers. The release of carvacrol (CAR) in pure form and CAR-PCL NPs was studied at different pH levels in the absence and presence of bacterial lipase enzyme. The result shows that the presence of lipase enzyme has a significant greater impact on the kinetics of polymer breakdown than the different pH levels ( $p < 0,05$ ) <sup>3</sup>. This explains that the effect of pH resulting from bacterial growth factor does not have a significant impact on the kinetics of CHL release from MPs

(The highlight regarding this context can be seen in lines 543-553)

3. I think that the selected mathematical models are not a good model for the presented release curves. They show a sigmoidal shape and these models are not good for this type of curves. I recommend to remove the mathematical modelling section from this paper as it is not adding any valuable information.

#### **Response to Reviewer**

We are very thankful to the Reviewer for taking the time to review this manuscript and for the expert review, providing helpful comments. We have removed this section in the revised manuscript.

4. Figure 5 is missing the "A" in the legend caption. Please correct this typo

#### **Response to Reviewer**

We are very thankful to the Reviewer for taking the time to review this manuscript and for the expert review, providing helpful comments. We have added the "A" in the legend caption in the revised manuscript

(The highlight regarding this context can be seen in lines 491)

**Reviewer #3:** Please go through the manuscript and review how the numbers are stated. Commas should be replaced by periods.

#### **Response to Reviewer**

We are very thankful to the Reviewer for taking the time to review this manuscript and for the expert review, providing helpful comments. We are glad that the Reviewer thinks that our work is clearly been well-planned, carefully executed. We have made a number of key changes to the manuscript as a result of these comments. We believe that the manuscript is now substantially improved. We have addressed each one of the reviewers' comments in detail.

1. Abstract - spell out PVA/PSB/TSB/

**Response:**

We are very thankful to the Reviewer for taking the time to review this manuscript and for the expert review, providing helpful comments. We have spell out these words in the revised manuscript.

(The highlight regarding this context can be seen in lines 21 – 22)

2. Line 48-49 - check sentence, it seems to be missing a word.

**Response:**

We thank the Reviewer for the comment and very useful insight. As a result, we have added several words regarding this content into the revised manuscript, as follows:

These symptoms are also typical in other inflammatory skin conditions, making identification of wound-related cellulitis difficult, while bacterial culture is quickly developing.<sup>4</sup> It is possible to enhance the degree of chronic wounds that were difficult to heal and required treatment management extension.

(The highlight regarding this context can be seen in lines 49 - 52)

3. Line 66-67 - check sentence for correctness.

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript, as follows:

The system can be designed to adapt responsively release the drug to the specific infected area.

(The highlight regarding this context can be seen in lines 66 – 67)

4. Line 84 - spell out MP

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript,

(The highlight regarding this context can be seen in lines 86)

5. Line 106-109 - this sentence needs to be clarified.

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript, as follows:

Finally, the validated approach was used to determine the drug concentration CHL in the MPs system, the drug release profile *in-vitro*, and *ex-vivo* dermatokinetic investigations from DMN loaded MPs sensitive bacteria.

(The highlight regarding this context can be seen in lines 108 - 110)

6. Line 118 - missing "to" - was added to

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript,

(The highlight regarding this context can be seen in lines 119)

7. Line 156 -clarify how it was dried

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript, as follows :

The formed MPs were then collected by centrifugation and washed with distilled water three times, separated and stored in a desiccator until dry.

(The highlight regarding this context can be seen in lines 154-156)

8. Line 179-180 -  $r^2$  is the coefficient of determination, not correlation

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 179 -180)

9. Line 184 - the term gradient is correct but for consistency with the linearity section, gradient should be replaced by slope

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 184)

**10.** Sections 2.6.3 and 2.6.4 should be made more similar. This is then same equation, just with a different weight of the standard deviation of the blank. However, the descriptions are very different and thus confusing

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript, as follows :

LLOQ is determined using the equation. (2) below, that the blank ideal divergence (without specimen) was symbolled by “sy”, and the slope applied through the calibration curve regression equation was symbolled by “b”.<sup>5</sup>.

(The highlight regarding this context can be seen in lines 189-191)

**11.** Line 201 - "in in" one too many

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript

(The highlight regarding this context can be seen in lines 201)

**12.** Section 2.7.5 title - check spelling of title "preparation"

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 229)

**13.** Line 247 - check sentence "put into for" is not clear

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 244)

**14.** Line 279 - add state for Microsoft (WA)

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 256)

**15.** Line 280-281. Remove period and include line 281 in the previous sentence

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 258)

**16.** Line 290. Check the sentence as it is missing a verb

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 267)

**17.** Line 302-304 - check sentences. Capital letter at And, "in in", "while ...)" is not a correct sentence.

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 278-279)

**18.** Figure 1. - the green color in the legend does not match the color on all the plots. The blank skin tissue and CHL in skin tissue is not present in plots B and D. Is it normal? The blank skin tissue and CHL in skin tissue have the same color.

**Response:**

We thank the Reviewer for the comment and very useful insight.

Regarding the blank skin tissue and CHL in skin tissue is not present in plots B and D. This is because on those plots media B (PVA) and D (TSB) are not related to CHL measurements on the biological matrices of rat skin tissue, so that maximum wavelength interference analysis is not carried out on these media.

**19.** Line 326 - "will be concise" does not make sense here

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 301)

**20.** Table 4 - concentration difference. Do you mean range?

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 302)

**21.** Line 338 - r2 is determination, not coefficient

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 313)

**22.** Line 341 - what is exactly meant by variation coefficient?

**Response:**

We thank the Reviewer for the comment and very useful insight.

The coefficient of variation (CV) is the ratio of the standard deviation to the mean. The higher the coefficient of variation will be greater the level of dispersion around the mean.

**23.** Table 5 to 9 - what does %RE mean?

**Response:**

We thank the Reviewer for the comment and very useful insight.

Relative error (RE) when used as a measure of precision is the ratio of the absolute error of a measurement to the measurement being taken. In other words, this type of error is relative to the size of the item being measured. RE is expressed as a percentage and has no units.

**24.** Line 427 - spell out LQC, MQC, HQC

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 401-402)

**25.** Table 10 - why are recoveries only 90%? When the method is used in later section, recovery of 100% is reached. Why? Discussing is needed here to explain why authors state the method is accurate when 10% active is missing

**Response:**

We thank the Reviewer for the comment and very useful insight.

The % recovery measurement in table 10 illustrates how much CHL levels can be extracted from the various extraction procedures employed in biological matrices (rat skin tissue) media, While % Recovery (figure 6) in the following section illustrates how much CHL content can be retrieved in the MPs system that is placed in the DMN system. So it's different meaning and method.

We have included these in the revised manuscript, as follows:

Regarding the %recovery findings in biological matrices obtained average extraction recovery has 90%. The FDA bioanalytical guideline recommendations emphasize the importance of accuracy, consistency, and reproducibility of recovery, acknowledging that the analyte recovery in bioanalytical methods does not need to be 100%. Additionally, the % relative standard deviation (%RSD) of the mean extraction recoveries throughout all QC levels was 2.91% ( $\leq 15\%$ ), indicating the good accuracy, consistency, and reproducibility of the extraction process. As a result, this approach is reliable to measure CHL in rat skin tissue media.

(The highlight regarding this context can be seen in lines 405-410)

26. Line 445 - after F2, replace comma by period

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 424)

27. Line 462 - does authors mean PCL composition in MPs?

**Response:**

We thank the Reviewer for the comment and very useful insight.

Yes, it's mean PCL composition in MPs as seen in table 2. Composition of formulations MPs loaded with CHL.

**Table 2.** Composition of formulations MPs loaded with CHL.

<b>Composition</b>	<b>F1</b>	<b>F2</b>
PCL (mg/7 ml chloroform)	35	70
CHL (mg/3 ml methanol)	35	35
PVA 3% (mL)	25	25

(The highlight regarding this context can be seen in lines 449)

28. Line 496 - ". In Figure 4." This is not a sentence

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 494)

**29.** Figure 5 - add (A) at the beginning of the legend

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

(The highlight regarding this context can be seen in lines 491)

**30.** Line 520 - please confirm "please and dispersal" are correct terms

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

Perhaps we used the incorrect word during the translation process, as the word term explains the drug release from matrices may involve processes of diffusion, erosion, and leaching or dissolution in media. However, we have removed mathematical modelling section from this paper, because suggestions from other reviewers mentioning that it's not adding any valuable information.

**31.** Line 531 - what does CAB stand for?

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

Perhaps we used the incorrect word, as the word term means CHL

(The highlight regarding this context can be seen in lines 515)

32. Line 545 - what does DOX stand for?

**Response:**

We thank the Reviewer for the comment and very useful insight. We have corrected this section in the revised manuscript.

Perhaps we used the incorrect word, as the word term means CHL


(The highlight regarding this context can be seen in lines 531)

**References:**

- (1) Khan, I.; Nagarjuna, R.; Dutta, J. R.; Ganesan, R. Enzyme-Embedded Degradation of Poly( $\epsilon$ -Caprolactone) Using Lipase-Derived from Probiotic *Lactobacillus Plantarum*. *ACS Omega* **2019**, *4* (2), 2844–2852. <https://doi.org/10.1021/acsomega.8b02642>.
- (2) Xiong, M. H.; Bao, Y.; Yang, X. Z.; Wang, Y. C.; Sun, B.; Wang, J. Lipase-Sensitive Polymeric Triple-Layered Nanogel for “on-Demand” Drug Delivery. *J. Am. Chem. Soc.* **2012**, *134* (9), 4355–4362. <https://doi.org/10.1021/ja211279u>.
- (3) Mir, M.; Permana, A. D.; Ahmed, N.; Khan, G. M.; Rehman, A. ur; Donnelly, R. F. Enhancement in Site-Specific Delivery of Carvacrol for Potential Treatment of Infected Wounds Using Infection Responsive Nanoparticles Loaded into Dissolving Microneedles: A Proof of Concept Study. *Eur. J. Pharm. Biopharm.* **2020**, *147* (December 2019), 57–68. <https://doi.org/10.1016/j.ejpb.2019.12.008>.
- (4) Andersen, C. A.; McLeod, K.; Steffan, R. Diagnosis and Treatment of the Invasive Extension of Bacteria (Cellulitis) from Chronic Wounds Utilising Point-of-Care Fluorescence Imaging. *Int. Wound J.* **2021**, No. September, 1–13. <https://doi.org/10.1111/iwj.13696>.
- (5) Silva, J. T. do P.; Silva, A. C. da; Geiss, J. M. T.; de Araújo, P. H. H.; Becker, D.; Bracht, L.; Leimann, F. V.; Bona, E.; Guerra, G. P.; Gonçalves, O. H. Analytical Validation of an

Ultraviolet–Visible Procedure for Determining Lutein Concentration and Application to Lutein-Loaded Nanoparticles. *Food Chem.* **2017**, *230*, 336–342.  
<https://doi.org/10.1016/j.foodchem.2017.03.059>.

# Graphical Abstract



## Highlights

- A spectrophotometric method to quantify chloramphenicol (CHL) was developed
- The analytical method was validated according to ICH guidelines
- Formulation of dissolving microneedle (DMN) loaded microparticles CHL was developed
- Validated method was applied in *in vitro* release study of microparticles CHL
- The validated method was applied in *ex vivo* dermatokinetic study of DMN

[Click here to view linked References](#)

1 **Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat**  
2 **skin tissue mimicking infection environment: application to *in vitro* release and *ex vivo***  
3 **dermatokinetic studies from dissolving microneedle loaded microparticle sensitive**  
4 **bacteria.**

5

6 Mukarram Mudjahid<sup>1</sup>, Sulistiawati<sup>1</sup>, Rangga Meidianto Asri<sup>1</sup>, Firzan Nainu<sup>1</sup>, Andi Dian Per-  
7 mana<sup>1\*</sup>

8 *1. Faculty of Pharmacy, Hasanuddin University, Makassar, 90245, Indonesia*

9

10 **\*Corresponding author:**

11 Andi Dian Permana

12 Faculty of Pharmacy, Hasanuddin University, Indonesia

13 Email: andi.dian.permana@farmasi.unhas.ac.id

14

## Abstract

Cellulitis is a common dermis/subcutaneous tissue skin infection and shared global disease burden, with a higher incidence for males and people aged 45–64 years. Application therapy of chloramphenicol (CHL) has been hindered because of its toxicity and limited penetration into the skin. In this research, CHL was developed into a bacterially sensitive microparticles which were further incorporated into a microneedle system to increase penetration. To support this formulation, in this study, UV–visible spectrophotometry method was validated in methanol, polyvinyl alcohol (PVA) 1%, phosphate buffered saline (PBS), tryptic soy broth (TSB) (fluid-mimicking infection), and skin tissue to quantify amount of CHL. The developed analytical method was subsequently validated according to ICH guidelines. The results obtained showed that the correlation coefficients were linear  $\geq 0.9934$ . The values of LLOQ inside the methanol, PVA 1%, PBS, TSB, and skin tissue were 7.20  $\mu\text{g/mL}$ , 4.40  $\mu\text{g/mL}$ , 8.18  $\mu\text{g/mL}$ , 387.48  $\mu\text{g/mL}$ , and 7.27  $\mu\text{g/mL}$ , respectively. The accuracy and precision of the developed method were prominent. These methods were successfully applied to quantify the amount of CHL in microparticle and microneedle system in fluid and tissue skin infection. The result showed the high drug release microparticle sensitive bacteria, and high drug retention in *ex vivo* dermatokinetic evaluation in rat skin tissue containing bacterial infection. This was due to the presence of *Staphylococcus aureus* bacteria culture that produced lipase enzymes, playing a role in lysing microparticle matrix to develop selectively delivery antimicrobials. A further analytical method needs to be matured to quantify CHL inside the *in vivo* studies.

**Keywords:** Bacterially Sensitive Microparticles; Cellulitis; Chloramphenicol; Microneedle; UV-vis Spectrophotometry; Validation.

## 43 1. Introduction

44 Cellulitis is a type of soft infection in the dermis / subcutaneous tissue characterized by  
45 warmth, erythema, and pain due to the response of cytokines and neutrophils from bacteria that  
46 penetrate the epidermis [1,2]. Cellulitis is a shared global disease burden. Its incidence rate is  
47 24.6/1000 person-years, and it is more common in men and people between the ages of 45 and 64.  
48 [3]. The location of the infection in the cutaneous tissue with no visible wound in the epidermis of  
49 the skin has been often ignored as ordinary inflammation. These symptoms are also typical in other  
50 inflammatory skin conditions, making identification of wound-related cellulitis difficult, while bac-  
51 terial culture is quickly developing. [4]. It is possible to enhance the degree of chronic wounds that  
52 were difficult to heal and required treatment management extension.

53 The microorganisms most commonly associated with cellulitis infection are *Staphylococci*  
54 and -hemolytic *streptococci* bacteria group [5,6]. Multibacterial colonization with different strains  
55 of human skin tissue has been widely reported [7]. This is associated with the interaction of bacteria  
56 with different strains that can synergize the severity of the disease [8]. Accordingly, broad-spectrum  
57 antibiotics are considered in the selection of therapy. Chloramphenicol (CHL) as broad antibiotic is  
58 active against gram- positive, gram-negative bacteria and anaerobic microorganisms [9,10]. Oral  
59 administration of CHL produces systemic effects and has been associated with significant side ef-  
60 fects, hematological disorders (aplastic anemia), spinal cord suppression, and availability of inade-  
61 quate drug concentrations in specific infectious tissues [11]. Therefore, it requires a long time for  
62 treatment. To overcome these shortcomings, CHL could be designed in dermal preparations to re-  
63 duce systemic exposure and increase drug concentrations to adequate target tissues of interest.

64 During the last decade, dermal delivery systems have been developed by designing drugs  
65 into microparticles/nanoparticles [12]. Microparticles are preferred to local infection targets with a  
66 longer drug retention rate in skin tissue than nanoparticles [13,14]. The system can be designed to  
67 adapt responsively release the drug to the specific infected area [15,16]. The tissue's properties re-  
68 sult in lower pH, greater temperature, surface proteins, and increased production of bacterial toxins,  
69 which function as a trigger for infection responsive delivery. [17].

70 Enzymes are considered promising since, under mild conditions, the enzymatic reactions  
71 are more efficient and accurate. Moreover, they are also necessary for the bacterial metabolic and  
72 pathological processes [18]. The chemical synthesis of bacterial lipase-sensitive PCL for specific  
73 delivery of antimicrobials was recently reported [19]. It has been reported, that SA may release a

74 particular enzyme called lipolytic esterase (Lipase), this enzyme can initiate the biocatalytic hydroly-  
75 sis of PCL [19] . Thus, lipase production at infection sites may be utilized to deliver and prove a  
76 practical, safe treatment approach selectively. This principle led to the development of delivery  
77 systems that selectively deliver drugs to specific sites of infection and reduce toxicity which is  
78 minimal when administered dermally.

79 Due to its hydrophobic chemical composition, which prevents effective cutaneous  
80 penetration, CHL is not useful for topical treatment of skin infections. Additionally, because human  
81 skin is thicker, it creates an impenetrable barrier that prevents the transdermal delivery of  
82 hydrophobic drugs. Dissolving microneedles (DMN) have the ability to penetrate biofilms and  
83 necrotic tissue in diseased skin by-passing the primary skin barrier [20] and may dissolve larger  
84 needles that are embedded in the skin. Importantly, the administration of DMN has the benefits of  
85 localized, painless, quick delivery, and patient compliance [21]. Given the potential advantages of  
86 this strategy, adding microparticles (MPs) chloramphenicol to DMNs may enhance the quantity of  
87 CHL that penetrate biofilm and necrotic tissue on diseased skin, which may help the treatment of  
88 burns and chronic wounds.

89 A fundamental part of developing new drug delivery systems in detecting and quantifying  
90 the drug, and therefore, it is crucial to develop an appropriate analytical method. In this research,  
91 the determination of drug content and the amount of drug release concentration from the *in vitro*  
92 method are critical assessments while developing a bacterial-responsive microparticle system. *In*  
93 *vitro* studies are the first phase of evaluating drug release profiles using relevant release media.

94 In measuring CHL in various matrices, numerous systematic methods have been developed  
95 until now. This includes both these prominent research HPLC-UV [22] and HPLC-MS/MS [23].  
96 However, the reported method is unsuitable for a limited budget and time. It also requires sophisti-  
97 cated equipment, making it a challenge to implement this in countries/laboratories with middle to  
98 low-income [24]. In contrast, the promise of UV-visible spectrophotometers utilized in detecting  
99 and quantifying CHL is due to their simplicity and cost-effectiveness. This study developed a CHL  
100 quantification method in artificial infection media and artificial skin fluids. There have been reports  
101 from some previous studies that the ability of UV-visible spectrophotometers to quantify various  
102 types of drugs [23,24]. It was crucial to validate the method used to ensure that the development of  
103 analytical procedures is reliable, comparable, and traceable.

104 This study aimed to develop and validate a method for analyzing CHL in fluid and rat skin  
105 tissue mimicking infection environment where the UV-visible spectrophotometer was utilized. The  
106 next step was to validate the method in accordance with the guidelines of international conference  
107 harmonization (ICH). The evaluated parameters were linearity, accuracy, precision, the limit of de-  
108 tection (LOD), and the limit of quantification (LOQ). Finally, the validated approach was used to  
109 determine the drug concentration CHL in the MPs system, the drug release profile *in vitro*, and *ex*  
110 *vivo* dermatokinetic investigations from DMN loaded MPs sensitive bacteria.

## 111 2. Materials and methods

### 112 2.1. Material

113 Chloramphenicol (CHL) was purchased from Merck (Darmstadt, Germany). Polycaprolactone  
114 (PCL), polyvinyl alcohol (PVA), polyvinyl pyrrolidone (PVP), sodium chloride, tryptic soy broth  
115 (TSB), potassium chloride, disodium phosphate, and potassium dihydrogen phosphate, were  
116 obtained from Sigma-Aldrich (Singapore). All other reagents used were analytical grade.

### 117 2.2. Preparation of CHL stock solution

118 The stock solution of CHL was prepared in 2 concentrations, where 10 mg and 20 mg CHL were  
119 weighed into a 10 mL volumetric flask. After that, to dissolve the CHL, methanol was added to the  
120 solutions with concentrations of 1000 µg/mL and 2000 µg/mL were obtained, respectively.

### 121 2.3. Measurement of the maximum UV absorption wavelength, testing standard, and quality con- 122 trol specimen preparation.

123 UV-visible spectrophotometer (Dynamica, HELLO XB-10) was employed to determine the maxi-  
124 mum wavelength absorbance. The CHL solution was prepared in concentration of 10 µg/mL in  
125 methanol, PVA 1%, PBS media, and 750 µg/mL in TSB media. The solution was scanned at room  
126 temperature at a 200-400 nm wavelength. Subsequently, six concentrations ranging from 2.5 µg/mL  
127 to 40 µg/mL on methanol, PBS, PVA 1%, skin tissue and 125 µg/mL – 1250 µg/mL on TSB media  
128 were prepared as calibration solutions.

129 Moreover, in an attempt to ensure the samples meet the expected criteria, each solvent was used as  
130 quality control (QC) samples at four quality levels, which are lower limit of quantification (LLOQ),  
131 low quality control (LQC), medium quality control (MQC), and high quality control (HQC). All  
132 parameters were measured and prepared in 3 replications.

133 **2.4. Sample preparation and CHL extraction from Skin samples**

134 To prevent any interferences with the measurement, the skin tissue sample was prepared by precip-  
135 itating the proteins and other molecules present in the organ. Methanol and acetonitrile were used  
136 to carry out the CHL extraction procedure. As stated in Table 1, different amounts of acetonitrile  
137 and methanol were employed to extract the drug. In the beginning, 1 g of the CHL spiked with  
138 matrices was combined with the extraction solvent. The mixture was then centrifuged at 14000 rpm  
139 for 15 minutes after being homogenized for 10 minutes using a vortex mixer. The obtained super-  
140 natant was then left at room temperature to allow the organic solvent to evaporate. The dried extract  
141 was then reconstituted with 1 mL of methanol, homogenized, and centrifuged as before. The result-  
142 ing supernatant was then evaluated using UV-visible spectrophotometry.

143 **Table 1.** Quantity of organic solvent required to extract CHL from skin tissue samples  
144

Organic Solvent	Methods	Volume (mL)
Methanol	A	1
	B	3
	C	5
	D	7
Acetonitrile	A	1
	B	3
	C	5
	D	7

145

146 **2.5. Preparation of Chloramphenicol Sensitive Bacterial Microparticles (MPs CHL-PCL) and**  
147 **Microneedle Loaded MPs CHL-PCL**

148 **MPs CHL-PCL** were prepared by a solvent evaporation method using polycaprolactone (PCL) as  
149 polymer and polyvinyl alcohol (PVA) as surfactant. Chloroform and methanol were used as solvents  
150 for the organic phase. Briefly, PCL was dissolved in 7 ml of chloroform, and CHL was dissolved  
151 in 3 ml of methanol, then mixed until homogeneous. The aqueous phase was prepared using PVA  
152 3% solution. The organic phase was added to the aqueous phase and mixed using a mechanical  
153 homogenizer at 500 rpm for 5 minutes. The emulsion system obtained was kept for 5 hours under a  
154 magnetic stirrer to ensure the organic phase evaporated completely. **The formed MPs were then**

155 collected by centrifugation and washed with distilled water three times, separated and stored in a  
156 desiccator until dry.

157 **Table 2.** Composition of formulations MPs loaded with CHL.

Composition	F1	F2
PCL (mg/7 ml chloroform)	35	70
CHL (mg/3 ml methanol)	35	35
PVA 3% (mL)	25	25

158 The dissolving microneedle (DMN) preparations were created. In this investigation, a silicone mold  
159 and the centrifugation process were applied. The mold used possessed several parameters, including  
160 a density of 10x10, pyramidal needles with a height of 700  $\mu\text{m}$ , a base width of 200  $\mu\text{m}$ , and an  
161 interspacing of 200  $\mu\text{m}$ . Each formulation (F1 and F2) contained 1 g of the formula, which contains  
162 MPs CHL-PCL 30%, PVP K-30 25%, PVA 10%, and Aquadest 35%. MPs CHL-PCL was com-  
163 bined with the polymer mixture to create the DMN that hold the active ingredient. They were then  
164 subjected to sonication until a clean, bubble-free dispersion was achieved. The polymer-drug com-  
165 bination was then put on the mold and centrifuged (LC-04S Centrifuge, Zenith Lab (Jiangsu) Co.,  
166 LTD.) for 30 minutes at 3500 rpm. When the centrifugation process was completed, the excess  
167 polymer mixture at the top of the mold was removed and replaced with a fresh polymer mixture.  
168 The MN were then dried for 1 day at room temperature and 1 day at 37°C without being removed  
169 from the mold [27].

## 170 2.6. Validation of analytical methods

### 171 2.6.1 Specificity

172 The comparison of the UV wavelength spectrum of CHL standard solutions in various solvent me-  
173 dia is used as an indicator for the specificity parameter. The assessed parameter was used to identify  
174 the possibility of interference among the analyte and the response of an alternative synthesis at the  
175 appropriate wavelength.

### 176 2.6.2 Linearity calibration curve

177 Parameters of the linear measurement of each medium were assessed with six standard solutions  
178 focusing on the appropriate wavelength spectrum. The results were projected in a calibration curve  
179 consisting of six concentrations with the obtained absorbance results. Parameters of determination  
180 coefficient ( $r^2$ ), slope, and y-intercept were then obtained and analyzed [28].

181 **2.6.3 Limit of detection (LOD)**

182 The parameter represents the minimum analyte concentration that can be measured with a sample  
183 [29]. LOD was determined using the equation. (1) below, that the blank ideal divergence (without  
184 specimen) was symbolled by “sy”, and the slope applied through the calibration curve regression  
185 equation was symbolled by “b”.

186 
$$\text{LOD} = \frac{3.3sy}{b} \quad (1)$$

187 **2.6.4 Lower limit of quantification (LLOQ)**

188 A minimum sample absorption that can be precisely calculated with satisfying precision and accu-  
189 racy is called the parameter of LLOQ [28]. LLOQ is determined using the equation. (2) below, that  
190 the blank ideal divergence (without specimen) was symbolled by “sy”, and the slope applied through  
191 the calibration curve regression equation was symbolled by “b”. [30].

192 
$$\text{LLOQ} = \frac{10sy}{b} \quad (2)$$

193 **2.6.5 Accuracy and precision**

194 The precision and the accuracy of parameters specify the propinquity to the reference value and the  
195 degree of among a sequence of measurements applied from several tests in the scientific method.  
196 These variables were assessed by measuring the inter-day and intra-day QC samples, which are  
197 HQC, QMC, LQC, and LLOQ. Percent relative error (%RE) and relative standard deviation  
198 (%RSD) were determined to express a quality that is accurate along with precise [31].

199 **2.6.6 Extraction recovery**

200 The absolute extraction recovery determination was calculated by comparing the response value of  
201 CHL in quality control samples (LLOQ, LQC, MQC, and HQC) from skin tissue to those obtained  
202 for freshly prepared solutions of the concentrations of the same samples (LLOQ, LQC, MQC, and  
203 HQC) [29].

204 **2.7. Application of analytical methods**

205 **2.7.1. Size Estimation Calculation and Particle Size Distribution**

206 Particle size was calculated using a microscopic method using an optical microscope equipped with  
207 a digital camera for as many as 300 particles. To see the shape and surface of the CHL MPs pro-  
208 duced. Analysis of the particle size distribution by looking at the polydispersity index (PDI) value  
209 using the equation.

210 
$$PDI = \left( \frac{\text{Standard deviation}}{\text{mean particles diameter}} \right)^2 \quad (3)$$

211 **2.7.2. Calculation of % Encapsulation and Drug Loading Efficiency**

212 The MPs suspension was centrifuged, the supernatant was taken, and then analyzed using a UV-Vis  
213 spectrophotometer at a maximum wavelength of 274.8 nm. The calculation of % encapsulation ef-  
214 ficiency (EE) and % drug loading (DL) follows the following equation:

215 
$$\%DL = \frac{\text{Amount of Encapsulated Drug}}{\text{Total Weight}} \times 100 \quad (4)$$

216 
$$\%EE = \frac{\text{Drug Total} - \text{Drug Free}}{\text{Drug Total}} \times 100 \quad (5)$$

217 **2.7.3. Drug content measurement of DMN loaded CHL-PCL**

218 The needle in the **DMN** is taken out by scraping the needle, then the needles is dissolved in water.  
219 After that, the solution was then centrifuged at 5000 rpm for 10 minutes. The precipitated MPs were  
220 then dissolved with an organic solvent (methanol: chloroform) to remove the polymer matrix. The  
221 samples were then analyzed using a UV-Vis spectrophotometer to measure drug content.

222 **2.7.4. In vitro Release Study of Microparticle Sensitive Bacteria in Bacterial Culture**

223 **The In vitro release study of MPs** was conducted to evaluate the sensitivity of bacteria to polymers  
224 by designing MPs on media containing the presence or absence of bacteria. In bacterial culture  
225 media, MPs dispersed in 20 mL of bacterial culture (equivalent to 0.5 Mc Farland) were placed in  
226 an orbital stirrer at 100 rpm, temperature  $37 \pm 1^\circ\text{C}$ . Samples (500  $\mu\text{L}$  volume) were then taken at  
227 predetermined time points (0,5, 2, 4, 6, 8, 12, and 24 hours). The sample was then centrifuged at  
228 10,000 rpm for 15 minutes, and the concentration was calculated with a UV-vis spectrophotometer.

229 **2.7.5. Ex vivo Dermatokinetic studies on infection skin rat model of microneedle **preparation****

230 **2.7.5.1. Preparation of ex vivo model of infection on rat skin**

231 Prior to the experiment, Wistar rat abdominal skins were shaved and adjusted in PBS (pH 7.4). The  
232 rat skins were cleaned in ethanol at a 70 percent concentration for one hour. The ethanol in the skins  
233 was evaporated before use by placing them in a biosafety cabinet for 20 minutes. Briefly, a biopsy  
234 punch was used to make a wound in the skin's surface (Stiefel, Middlesex, UK). Then, with a few  
235 minor adjustments, the earlier published protocols were used to create *ex vivo* models of biofilm on  
236 rat skin [32]. After that, 50  $\mu\text{L}$  of the diluted **SA** bacterial suspensions ( $2 \times 10^5$  CFU/mL) were  
237 dripped to the wound of the skin and distributed evenly after the skins were mounted on TSA plates.

238 The skins were transferred to fresh TSA plates every day for five days while the plates were  
239 incubated at 37 °C to facilitate the growth of the biofilm on the injured skin.

#### 240 **2.7.5.2. *Ex vivo* dermatokinetic studies**

241 The *ex vivo* dermatokinetic studies were also evaluated Using the procedure outlined [20, 32] with  
242 slight modification. *Ex vivo* dermatokinetic experiments of DDMN containing MPs CHL-PCL were  
243 conducted in normal skin and *ex vivo* infection model in rat skins. The skins that had previously  
244 been fastened to the donor compartment of the Franz diffusion cells were manually placed for 30  
245 seconds. The receiver compartment, which contained PBS, was further connected to the donor  
246 compartment (pH 7.4). Afterwards, 5 g of stainless-steel mass was used on top of the DMN to  
247 prevent movement during the trial. The experiment was conducted at 600 rpm and  $37 \pm 1$  °C. The  
248 DMN were separated from the skins in order to assess the concentration of CHL in the skins at  
249 various specified time intervals. The skins were cleaned three times, put into glass vials, and given  
250 2.5 mL of water. The mixtures were centrifuged at 3000 rpm for 15 minutes after being vortexed  
251 for 30 minutes. AAS was used to measure the silver content in the supernatant. PKSolver (China  
252 Pharmaceutical University, Nanjing, China) was used to construct the dermatokinetic profiles [34].

#### 253 **2.8. Statistical Analysis**

254 All results were reported as means  $\pm$  standard deviation (SD). The mean, SD, linear regression  
255 analysis, %RSD and %CV of each sample in the validation method were processed using  
256 Microsoft<sup>®</sup> Excel 2016 (Microsoft Corporation, Redmond, Washington, USA). GraphPad Prism<sup>®</sup>  
257 version 6 (GraphPad Software, San Diego, California, USA) was utilised to statistically analyse the  
258 results where a  $p$ -value  $< 0.05$  denoted a significant difference.

### 259 **3. Results and Discussion**

#### 260 **3.1. Selection of sample preparation method and drug extraction**

261 In this study, methanol and acetonitrile, two organic solvents, were used to extract CHL from skin  
262 tissue. Table 3. displays the outcomes of every extraction technique. The findings demonstrated that  
263 the extraction efficiency might be increased by using more solvent during the extraction procedure.  
264 Methanol offered a greater extraction recovery than acetonitrile, it was also discovered. The find-  
265 ings indicated that techniques C and D had the greatest extraction recovery percentages for extract-  
266 ing CHL from methanol, with respective values of  $91.54 \pm 2,67$  percent and  $94.77 \pm 0,96$  percent.  
267 Furthermore, both of these approaches were not a significantly difference ( $p > 0.05$ ). Because it

268 required the least quantity of solvent while still providing the best extraction recovery, method C (5  
269 mL methanol) was selected as the best extraction technique for this investigation.

270


271 **Table 3.** Mean extraction recovery of CHL of each method with methanol and acetonitrile from skin tissue (n = 3)

Organic Solvent	Methods	Volume (mL)	%Extraction Recovery $\pm$ SD	%RSD
Methanol	A	1	18.52 $\pm$ 1.90	10.26
	B	3	41.78 $\pm$ 1.24	2.97
	C	5	91.54 $\pm$ 2.67	2.91
	D	7	94.77 $\pm$ 0.96	1.01
Acetonitrile	A	1	12.18 $\pm$ 0.82	6.72
	B	3	12.45 $\pm$ 3.11	24.96
	C	5	39.92 $\pm$ 1.90	4.74
	D	7	62.38 $\pm$ 3.27	5.24

272

### 273 3.2. Selectivity of UV-Vis spectrophotometric method

274 In order to prevent interferences between CHL and other chemicals contained in microparticle, mi-  
275 croneedle, and vaginal tissue during the examination utilizing UV-Visible spectrophotometry, a  
276 specificity test was conducted [29, 33]. As was already noted, the content of CHL in the formulation  
277 was determined using the analytical technique in methanol. Additionally, the analytical method's  
278 development in PBS, PVA, and TSB was done in order to calculate the CHL *in vitro*, while skin  
279 tissue sample was used to calculate the CHL in *ex vivo* studies. Figure 1 show representative UV  
280 spectra of CHL and several blank in various media.



281  
 282 **Figure 1.** Representative UV spectra of CHL and several blank in Methanol (A), PVA (B), PBS (C), TSB (D).

283  
 284 Based on the analysis results, in all media, the measurement of the blank spectra of MPs, DMN,  
 285 and skin tissue, there was no peak that appeared at the wavelength of CHL in each medium. These  
 286 findings showed that the inclusion of additional **MPs**, **DMN**, and tissue skin ingredients did not  
 287 cause any interference [35].

288 After measurements were made, there was a shift in the optimum absorption, but it was still in the  
 289 optimum absorption area of CHL in the medium methanol, PBS, and PVA 1%. There was a shift in  
 290 the absorption of a longer wavelength in TSB media. These results are also consistent with those of  
 291 Tseplin et al. who found that the particular impact of the polar solvent, which presents itself here as  
 292 a bathochromic shift of one of the  $\pi-\pi^*$  bands, is generated by the creation of hydrogen bonds

293 between solvent molecules and the molecule as a result, by a decrease in the energy gap between  
294 the corresponding occupied ( $\pi$ ) and empty ( $\pi^*$ ) molecular orbitals [36]. Based on the results ob-  
295 tained, the method developed in this study was specific at the appropriate wavelength.

### 296 3.3. Linearity, LOD, and LLOQ

297 In assessing the linearity parameters and determining the rates of LOD and LLOQ from the scien-  
298 tific form, the measurement graph was produced by calculating the CHL standard solution concen-  
299 tration of each medium (methanol, PBS, PVA 1%, and TSB) utilizing the advanced UV-vis spec-  
300 trophotometry. The spectra of standard solutions across various media are depicted in Figure 2.  
301 Linearity, LOD, and LLOQ settings of the CHL can be seen in Table 4 and Figure 3.


302 **Table 4.** Calibration curve properties for CHL analysis with the quality of LOD and LLOQ.

Media	Concentration Range ( $\mu\text{g/mL}$ )	$r^2$	LOD ( $\mu\text{g/mL}$ )	LLOQ ( $\mu\text{g/mL}$ )
MeOH	2.5 – 40	0.9981	2.37	7.20
PVA 1%	2.5 - 40	0.9993	1.45	4.40
PBS	2.5 - 40	0.9975	2.70	8.18
TSB	125 - 1250	0.9934	127.87	387,48
Skin Tissue	2.5 - 40	0,9981	2,40	7,27

303  
304 Based on the data in Table 6, Each calibration curve comprised six concentration levels: 2.5, 5, 10,  
305 20, 30, 40  $\mu\text{g/mL}$  in MeOH, PVA 1%, PBS, and skin tissue media. While the TSB media obtained  
306 six concentrations of calibration curves, namely, 125, 250, 500, 750, 1000, and 1250. Since there  
307 was no concentration data in the matrix or a similar reference to find out about CHL levels, the  
308 concentration points in the calibration curve were used in the calculation.

309 The slopes and their coefficient of variation as the parameter for analyzing linear equations were  
310 determined because they were the most prominent indicator of the sensitivity of analytical methods  
311 and their quantification capabilities. Linear relationships were obtained between absorbance and  
312 concentration in 2.5 – 40  $\mu\text{g/mL}$  for CHL-MeOH, CHL-PVA 1%, CHL-PBS, CHL-Skin Tissue and  
313 125 – 1250  $\mu\text{g/mL}$  for CHL-TSB. The value of the determination coefficient ( $r^2$ ) of the four CHL  
314 regression equations on Methanol, PVA 1%, PBS, skin tissue, and TSB were 0.9981, 0.9993,

315 0.9975, 0.9981 and 0.9934, respectively. It was shown from the results that the calibration curves  
316 determination coefficients ( $r^2$ ) were more than 0.99, and their variation coefficients were less than  
317 25%. Hence, it can be concluded that we satisfied the acceptance criteria, and a high sensitivity was  
318 achieved in the four-study media. Therefore, a range of 2.5 – 40  $\mu\text{g/mL}$  values for CHL-MeOH,  
319 CHL-PVA 1%, CHL-PBS, and 125 – 1250  $\mu\text{g/mL}$  for CHL-TSB can be used for LOD and LLOQ  
320 calculations. Quality LOD and LLOQ of CHL inside the methanol were 2.37 and 7.20  $\mu\text{g/mL}$ , on  
321 PVA 1% were 1.45 and 4.40  $\mu\text{g/mL}$ , on PBS 2.70 and 8.18, on skin tissue 2.40 and 7.27, and on  
322 TSB medium were 127.87 and 387.48  $\mu\text{g/mL}$ .



323

324 **Figure 2.** Spectrum of CHL standard solutions in of Methanol (A), PVA (B), PBS (C), TSB (D), Skin Tissue (E).

325

326


327

328

329

330

331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359



**Figure 3.** Calibration Curves CHL-Methanol (A), CHL-PVA (B), CHL-PBS (C), CHL-TSB (D), CHL-Skin Tissue (E), (mean  $\pm$  SD, n= 3)

360 **3.4. Accuracy and precision**

361 Determination of accuracy in intra-day and inter-day amounts using the advanced method exhibited  
362 accurate values for PVA 1% (Table 5), methanol (Table 6), PBS (Table 7), TSB (Table 8), skin  
363 tissue (9). The measurement of ratio error values is under 15%, which complied the preconditions  
364 of the ICH guidelines. Intra-day and inter-day precision were also examined to be adequate. Intra-  
365 day and inter-day precision for all solvents determined %RSD values ranging from 0.23% - 13.61%  
366 and 0.3% - 3.38% which obeyed the limits of the ICH guidelines (15%) [37]. Thus, the method  
367 developed using UV-vis spectrophotometry to determine CHL was found to be accurate and  
368 precise.

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

**Table 5.** The Results of evaluation of precision and accuracy of CHL analysis in PVA 1% (mean  $\pm$  SD, n= 3)

<b>Interday Precision and Accuracy</b>				
<b>Replication</b>	<b>Concentration added (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	4.40	4.29 $\pm$ 0.070	1.63	2.53
	7.5	7.36 $\pm$ 0.05	0.72	1.75
	30	30.28 $\pm$ 0.12	0.39	0.96
	15	14.52 $\pm$ 0.069	0.48	3.17
<b>2</b>	4.40	4.16 $\pm$ 0.11	2.66	5.37
	7.5	7.44 $\pm$ 0.07	1.01	0.76
	30	30.09 $\pm$ 0.14	0.48	0.3
	15	14.43 $\pm$ 0.07	0.48	3.78
<b>3</b>	4.40	4.43 $\pm$ 0.09	2.16	-0.59
	7.5	7.67 $\pm$ 0.04	0.59	-2.32
	30	30.77 $\pm$ 0.07	0.23	2.59
	15	14.79 $\pm$ 0.16	1.08	1.34
<b>Intraday Precision and Accuracy</b>				
<b>Day</b>	<b>Concentration added (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	4.40	3.77 $\pm$ 0.046	1.21	14.32
	7.5	6.89 $\pm$ 0.05	0.72	8.11
	30	30.08 $\pm$ 0.09	0.31	0.29
	15	13.49 $\pm$ 0.09	0.67	10
<b>2</b>	4.40	3.85 $\pm$ 0.02	0.67	12.59
	7.5	7.32 $\pm$ 0.09	1.30	2.36
	30	29.92 $\pm$ 0.09	0.30	-0.26
	15	13.43 $\pm$ 0.14	1.06	10.42
<b>3</b>	4.40	3.91 $\pm$ 0.04	1.17	11.21
	7.5	7.36 $\pm$ 0.14	1.90	1.76
	30	30.28 $\pm$ 0.09	0.30	0.96
	15	13.74 $\pm$ 0.07	0.51	8.37

**Table 6.** The Results of evaluation of precision and accuracy of CHL analysis in MeOH (mean  $\pm$  SD, n= 3)

<b>Interday Precision and Accuracy</b>				
<b>Replication</b>	<b>Concentration gained (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	7.2	7.30 $\pm$ 0.11	1.51	-1.48
	10	10.73 $\pm$ 0.08	0.79	-7.36
	30	32.55 $\pm$ 0.11	0.34	8.51
	20	22.18 $\pm$ 0.08	0.39	-10.93
<b>2</b>	7.2	7.35 $\pm$ 0.10	1.43	-2.08
	10	10.86 $\pm$ 0.08	0.78	-8.63
	30	32.76 $\pm$ 0.11	0.34	9.21
	20	22.27 $\pm$ 0.09	0.41	-11.36
<b>3</b>	7.2	7.40 $\pm$ 0.14	2.02	-2.87
	10	10.79 $\pm$ 0.13	1.22	-7.93
	30	32.27 $\pm$ 0.23	0.71	7.57
	20	22.23 $\pm$ 0.12	0.57	-11.15
<b>Intraday Precision and Accuracy</b>				
<b>Day</b>	<b>Concentration gained (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	7.2	7.87 $\pm$ 0.13	1.65	-9.30
	10	10.61 $\pm$ 0.15	1.45	-6.1
	30	31.14 $\pm$ 0.10	0.32	3.82
	20	21.41 $\pm$ 0.14	0.69	-7.06
<b>2</b>	7.2	8.03 $\pm$ 0.12	1.55	-11.62
	10	10.75 $\pm$ 0.10	0.98	-7.5
	30	31.28 $\pm$ 0.30	0.97	4.27
	20	21.66 $\pm$ 0.26	1.22	-8.31
<b>3</b>	7.2	8.16 $\pm$ 0.04	0.55	-13.37
	10	10.86 $\pm$ 0.12	1.15	-8.66
	30	31.2 $\pm$ 0.17	0.54	4
	20	21.48 $\pm$ 0.10	0.49	-7.41

**Table 7.** The Results of evaluation of precision and accuracy of CHL analysis in PBS (mean  $\pm$  SD, n= 3)

<b>Interday Precision and Accuracy</b>				
<b>Replication</b>	<b>Concentration added (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	8.27	8.62 $\pm$ 1.17	13.61	-4.23
	12.5	14.02 $\pm$ 0.06	0.43	-12.21
	30	31.43 $\pm$ 0.13	0.42	4.78
	20	21.96 $\pm$ 0.14	0.66	-9.8
<b>2</b>	8.27	8.05 $\pm$ 0.15	1.91	2.66
	12.5	13.86 $\pm$ 0.36	2.60	-10.88
	30	30.51 $\pm$ 0.74	2.42	1.71
	20	21.65 $\pm$ 0.60	2.77	-8.25
<b>3</b>	8.27	7.98 $\pm$ 0.16	2.09	3.50
	12.5	13.77 $\pm$ 0.45	3.28	-10.18
	30	30.93 $\pm$ 0.73	2.38	3.11
	20	21.66 $\pm$ 0.82	3.81	-8.31
<b>Intraday Precision and Accuracy</b>				
<b>Day</b>	<b>Concentration added (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	8.27	7.82 $\pm$ 0.15	1.92	5.36
	12.5	13.48 $\pm$ 0.34	2.58	-7.84
	30	30.85 $\pm$ 0.67	2.18	2.83
	20	21.38 $\pm$ 0.12	0.58	-6.91
<b>2</b>	8.27	8.19 $\pm$ 0.08	1.08	0.96
	12.5	13.61 $\pm$ 0.11	0.81	-8.93
	30	31.2 $\pm$ 0.12	0.38	4
	20	21.39 $\pm$ 0.06	0.31	-6.98
<b>3</b>	8.27	8.06 $\pm$ 0.17	2.12	2.49
	12.5	13.57 $\pm$ 0.26	1.95	-8.61
	30	31.27 $\pm$ 0.12	0.38	4.23
	20	20.87 $\pm$ 0.40	1.94	-4.38

**Table 8.** The Results of evaluation of precision and accuracy of CHL analysis in TSB (mean  $\pm$  SD, n= 3)

<b>Interday Precision and Accuracy</b>				
<b>Replication</b>	<b>Concentration added (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	388	397.66 $\pm$ 15.06	3.78	-2.49
	450	457.66 $\pm$ 10.10	2.20	-1.70
	938	922.66 $\pm$ 20.81	2.25	-1.63
	600	608.5 $\pm$ 16.39	2.69	-1.41
<b>2</b>	388	366 $\pm$ 13.91	3.80	5.67
	450	416.83 $\pm$ 15.06	3.61	7.37
	938	912.66 $\pm$ 5.77	0.63	-2.70
	600	579.33 $\pm$ 7.21	1.24	3.44
<b>3</b>	388	369.33 $\pm$ 8.77	2.37	4.81
	450	421 $\pm$ 9.01	2.14	6.44
	938	894.33 $\pm$ 19.41	2.17	-4.65
	600	572.66 $\pm$ 10.10	1.76	4.55
<b>Intraday Precision and Accuracy</b>				
<b>Day</b>	<b>Concentration added (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	388	363.08 $\pm$ 10.77	2.96	6.42
	450	426.83 $\pm$ 14.21	3.33	5.14
	938	861.83 $\pm$ 18.76	2.17	-8.12
	600	580.16 $\pm$ 15.27	2.63	3.30
<b>2</b>	388	340.16 $\pm$ 11.54	3.39	12.32
	450	397.66 $\pm$ 11.54	2.90	11.62
	938	841 $\pm$ 17.32	2.05	-10.34
	600	555.16 $\pm$ 14.43	2.59	7.47
<b>3</b>	388	346 $\pm$ 10.89	3.14	10.82
	450	401 $\pm$ 10	2.49	10.88
	938	850.16 $\pm$ 28.75	3.38	-9.36
	600	557.66 $\pm$ 17.01	3.05	7.05

**Table 9.** The Results of evaluation of precision and accuracy of CHL analysis in skin tissue (mean  $\pm$  SD, n= 3)

<b>Interday Precision and Accuracy</b>				
<b>Replication</b>	<b>Concentration added (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	7.28	7.61 $\pm$ 0.24	3.18	4.47
	10	9.93 $\pm$ 0.71	7.11	-0.70
	20	19.42 $\pm$ 0.91	4.66	-2.91
	30	31.21 $\pm$ 0.67	2.14	4.05
<b>2</b>	7.28	7.80 $\pm$ 0.35	4.48	7.08
	10	9.74 $\pm$ 0.17	1.71	-2.55
	20	19.64 $\pm$ 0.48	2.47	-1.81
	30	32.37 $\pm$ 0.46	1.43	7.91
<b>3</b>	7.28	7.84 $\pm$ 0.39	4.98	7.69
	10	10.47 $\pm$ 0.46	4.41	4.69
	20	20.08 $\pm$ 0.51	2.52	0.39
	30	32.08 $\pm$ 0.71	2.23	6.93
<b>Intraday Precision and Accuracy</b>				
<b>Day</b>	<b>Concentration added (<math>\mu\text{g/mL}</math>)</b>	<b>Found concentration (<math>\mu\text{g/mL}</math>) <math>\pm</math> SD</b>	<b>Precision (%RSD)</b>	<b>Accuracy (%RE)</b>
<b>1</b>	7.28	7.43 $\pm$ 0.24	3.20	2.00
	10	9.52 $\pm$ 0.16	1.71	-4.77
	20	19.22 $\pm$ 0.47	2.47	-3.89
	30	31.42 $\pm$ 0.70	2.23	4.73
<b>2</b>	7.28	7.66 $\pm$ 0.38	5.00	5.15
	10	9.18 $\pm$ 0.66	7.19	-8.22
	20	18.66 $\pm$ 0.47	2.53	-6.70
	30	30.57 $\pm$ 0.65	2.14	1.91
<b>3</b>	7.28	7.61 $\pm$ 0.34	4.49	4.56
	10	9.68 $\pm$ 0.43	4.45	-3.19
	20	19.01 $\pm$ 0.89	4.67	-4.97
	30	31.71 $\pm$ 0.45	1.43	5.69

### 400 3.5. Extraction Recovery

401 The quantities of lower limit of quantification (LLOQ), low quality control (LQC), medium quality  
402 control (MQC), and high quality control (HQC) were extracted from skin tissue were compared to  
403 the concentrations obtained by testing samples at the same concentrations to determine the extrac-  
404 tion recovery from the technique utilized in this investigation. Table 10 shows the outcomes of the  
405 typical extraction recovery. The FDA bioanalytical geudeline recommendations emphasize the im-  
406 portance of accuracy, consistency, and reproducibility of recovery, acknowledging that the analyte  
407 recovery in bioanalytical methods does not need to be 100%. Additionally, %RSD of the mean  
408 extraction recoveries throughout all QC levels was  $\leq 15\%$ , indicating the good accuracy, con-  
409 sistency, and reproducibility of the extraction process. As a result, this approach is reliable to meas-  
410 ure CHL in rat skin tissue media.

411 **Table 10.** Average extraction recovery of CHL in skin tissue (n = 3)

Sample	Concentration ( $\mu\text{g/mL}$ )	%Extraction Re- covery $\pm$ SD	%RSD
Skin Tissue	LLOQ (5,13)	91.47 $\pm$ 1.40	1.53
	QC (7,5)	90.93 $\pm$ 1.42	1.56
	MQC (15)	90.84 $\pm$ 1.29	1.42
	HQC (24)	91.72 $\pm$ 1.73	1.88

412

### 413 3.6. Application of the analytical method

#### 414 3.6.1. The Properties of analysis and In- vitro release studies of Microparticle Sensitive Bacteria

415 A UV-visible spectrophotometric method form that has been thoroughly validated was then applied  
416 to confirm the quantity of CHL in the microparticle system. The characteristics of the microparticles  
417 can be seen in Table 11, and the MPs release profile can be seen in Figure 5.


418 **Table 11.** Results of Measurement of Particle Size, PDI, %EE, and %DL MPs CHL (mean  $\pm$  SD, n= 3)

Formulation	Particle Size	PDI	% EE	% DL
F1	0,59 $\pm$ 0,02	0,117 $\pm$ 0,01	40.94 $\pm$ 2.88	20.47 $\pm$ 1.44
F2	1,29 $\pm$ 0,09	0,121 $\pm$ 0,01	39.61 $\pm$ 3.23	13.20 $\pm$ 1.07

419

420 The results showed that an analysis from CHL-carried MPs detected by SEM will be visualized in  
421 Figure 4. As shown in SEM images, all formulations were approximately spherical. Notably, the

422 sizes discovered in this study were closely correlated with the findings from SEM analysis and were  
423 found to be  $0.59 \pm 0.02 \mu\text{m}$  and  $1.29 \pm 0.09 \mu\text{m}$  for F1 and F2, respectively. It was found that the  
424 particle size of F1 were significantly smaller ( $p < 0.05$ ) compared to F2. It has been postulated that  
425 increase in the amount of polymer will increase the particle size due to a rise in the polymer's  
426 viscosity, which causes the creation of bigger emulsion droplets and subsequently larger sizes of  
427 the microparticle [38][39]. For PDI values, it was found that the values were in the range of  $0.117$   
428  $\pm 0.01$  and  $0.121 \pm 0.01$ , demonstrating the generally homogeneous and monodispersed profile of  
429 these MPs [40].



439 **Figure 4.** SEM Visualization Image of F1 (A) and F2 (B) (The black scale bar represents a length of  $10\mu\text{m}$  in  
440 each case).

441 Regarding the EE, it was found that the EE of MPs F1 were not significantly ( $p > 0.05$ ) compared  
442 to MPs F2. These results propose that there can be a relatively high amount of CHL that might be  
443 entrapped in the PCL MPs. The lower EE may be due to the higher hydrophobicity nature of PCL  
444 than CHL. Percentage EE will depend a lot on the polymer type and polymer drug affinity [41].  
445 Therefore, the encapsulation of CHL were low in this polymer. The percentage of encapsulation  
446 efficiency in F1 and F2 were  $40.94 \pm 2.88\%$  and  $39.61 \pm 3.23$ , respectively. In terms of DL, the  
447 values F1 and F2 were  $20.47 \pm 1.44$  and  $13.20 \pm 1.07$ , respectively. Based on the results obtained,  
448 the increase in the number of polymers (2: 1) in formula 2 decreased the percentage of DL values.  
449 This occurred due to the increase of MPs (PCL polymer) composition while retaining the quantity  
450 of CHL in the MPs, resulting in a decreasing DL value.

451 Validated analytical methods also determined the cumulative amount of CHL after *in vitro* release  
452 studies on various media. Figure 5. (A) shows PBS media's *in vitro* release profile. The results  
453 showed that the % CHL measured after 24 hours in F1 and F2 were  $12.75 \pm 1.46$  and  $9.46 \pm 1.13$ ,  
454 respectively. Figure 5. (B) shows the MPs release profile *in-vitro* on TSB media. The results  
455 showed that the % CHL measured after 24 hours in F1 and F2 were  $13.41 \pm 1.55$  and  $10.02 \pm 1.27$ ,  
456 respectively. In both media, it was found that the lower levels of CHL were measured in F1 and F2  
457 when compared to pure CHL due to the absence of a bacterial enzyme stimulus that lyses the PCL  
458 polymer so that the drug remains stable in the MPs polymer matrix. The statistical analysis results  
459 showed that the release profiles of F1, F2, and CHL, compared to both PBS and TSB media, were  
460 not significantly different ( $p > 0.05$ ). This indicated that the media without bacteria did not affect  
461 drug release in the bacterial responsive MPs system. Figure 5. (C) shows *in vitro* release profile of  
462 MPs on TSB media and bacteria. The results showed that the % CHL measured after 24 hours in  
463 F1 and F2 were  $99.29 \pm 11.10$  % and  $79.45 \pm 9.34$  % respectively. In TSB, media containing bacteria  
464 showed high drug release. This was due to the SA bacteria culture that produced lipase enzymes  
465 that played a role in lysing the MPs matrix to measure CHL levels in the media [17,38].

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485


486

487

488

489

490



491

492

**Figure 5.** (A) *in vitro* MPs-CHL release in PBS media, (B) *in vitro* MPs-CHL release in TSB media, (C) *in vitro* MPs-CHL release in TSB + Bacteria media (mean  $\pm$  SD, n= 3)

493

494

495

496

497

498

499

500

501

The statistical analysis results showed significantly different release profile results between bacterial TSB media with PBS and TSB without bacteria. In figure 5, shows the release profile of pure CHL in all test media showed a rapid drug release profile with levels reaching almost 100% at 3 hours, while the microparticle release profile in media containing bacterial stimuli almost reached 100% at 24 hours. Figure 5 shows that the drug release profile of F1 is higher than F2. It is shown that drug release is affected by particle size when drug loading is high (40%). Chen et al also found the same results as we observed regarding the difference in particle size affecting the time and % drug release [43]. The larger surface area of smaller MPs results in a higher concentration of drug molecules at the surface of microspheres, ready for a faster release.

502

503

Thus, in contrast to sterile media, the higher release of CHL from PCL MPs in the bacterial system was possible to be an indicator that the bacteria are releasing the enzymes. Additionally, this result


504 was consistent with that of Wu et al., who observed that the addition of an enzyme multiplies PCL  
505 degradation by a factor of 1000 compared to degradation in an aqueous medium alone. [44]. A stark  
506 contrast between the release of CHL when the bacteria is present or absent demonstrated that the  
507 proposed method has the potential for selective delivery at infection sites. Therefore, it was proven  
508 from these results that loading CHL into a responsive MPs system can avoid release at non-specific  
509 sites.

### 510 ***3.6.2. The Properties of analysis and Ex- vivo dermatokinetic studies of Microneedle Loaded MPs*** 511 ***CHL-PCL***

512 For improved penetration profile, the MPs were incorporated into DMN (Figure 6A and 6B). The  
513 quantity of CHL in the microneedle loaded MPs CHL-PCL was also determined using the approved  
514 spectrophotometry UV-visible technique. Figure 6C shows the results, which revealed that  $97.64 \pm$   
515  $3.21$ ,  $98.09 \pm 2.43$ , and  $97.12 \pm 2.76$  percent of the CHL were recovered from the F1, F2, and Pure  
516 CHL in DMN, respectively. These results demonstrated that the concentration of CHL was unaf-  
517 fected by the incorporation of CHL into DMN preparation. All formulations had acceptable recov-  
518 ery percentages, which fell between 95 and 105 percent in accordance with the ICH standard for  
519 acceptable recovery percentage [37].

520

521



522  
 523 **Figure 6.** The microscope images of DMNs containing F1 (A) and F2 (B). CHL recovery (%) from DMN F1, F2, and  
 524 CHL (C).

525 The release kinetics of CHL from MPs after DMN treatments were then examined in a dermatoki-  
 526 netic investigation. The *ex vivo* infection model in rat skins and uninfected skin were used in this  
 527 investigation. Only choramphenicol released by MPs were measured in order to demonstrate that  
 528 the presence of bacterial infection had no other effect on the release of CHL in the skin. The skin  
 529 samples were vortexed with water at each interval to achieve this, to see how much CHL was re-  
 530 tained in the skin. Our findings showed that this approach would only extract CHL released from  
 531 MPs since no CHL was found in the supernatant of MP dispersion.

532 In this study, we explain to compared the dermatokinetic characteristics of our method to DMNs F1  
 533 and F2 with CHL but no MP formulation (DMN-CHL). Figure 9 shows the concentration of CHL  
 534 released from MPs in the skin vs application time for the kinetic profiles of CHL in normal skin and  
 535 *ex vivo* infection models following application of DMNs. Table 13 has shown lists the dermatoki-  
 536 netic profiles of CHL following application of DMNs, including C<sub>max</sub>, T<sub>max</sub>, T<sub>1/2</sub>, AUC, and  
 537 MRT. As shown, DMNs-MPs released CHL much less ( $p < 0.05$ ) than DMN- CHL in the absence  
 538 of bacterial growth in normal skin. The T<sub>max</sub> values of CHL from MN-MPs F1 and F2 were found  
 539 to be considerably lower ( $p < 0.05$ ) compared to the other formulation due to the explanations pro-  
 540 vided for the C<sub>max</sub> results..This suggested that the inclusion of CHL into MPs would make it

541 possible to avoid the non-specific release of CHL. The release of CHL from DMN-MPs, in contrast  
542 to DMN- CHL, was considerably increased ( $p < 0.05$ ) in *ex vivo* infection models made from SA.  
543 The results obtained here indicate that CHL encapsulation in PCL MPs may result in an increase in  
544 CHL release via two different processes. The PCL outer layer initially aided in the MPs' attachment  
545 to the colony infection. Following this, the bacterial strains lipase released into the environment  
546 dissolved PCL layers, releasing CHL from MPs. Regarding the impact of pH on kinetical release of  
547 CHL in MPs-coated PCL, previous research has demonstrated the profile of nanoparticle release  
548 utilizing PCL polymers. The release of carvacrol (CAR) in pure form and CAR-PCL nanoparticles  
549 (NPs) was studied at different pH levels in the absence and presence of bacterial lipase enzyme.  
550 The result shows that the presence of lipase enzyme has a significant greater impact on the kinetics  
551 of polymer breakdown than the different pH levels ( $p < 0.05$ ) [45]. This explains that the effect of  
552 pH resulting from bacterial growth factor does not have a significant impact on the kinetics of CHL  
553 release from MPs. Specifically, when compared to the DMN-F2 formula, DMN-F1 had a higher  
554 drug release due to the smaller particle size in F1 compared to the F2 formula. In the case of smaller  
555 MPs, the greater surface area produces a higher number of drug molecules at the surface of micro-  
556 spheres, ready for faster release.

557

558


559

560 **Table 13.** Lists the dermatokinetic characteristics of CHL in uninfected rat skin as well as ex infection models created by SA after the administration of DDMN-  
 561 F1, DMN-F2, and DMN-CHL. (means SD, n = 3)

Parameter	Unit	DMN + F1		DMN + F2		DMN + CHL	
		Normal	Infection	Normal	Infection	Normal	Infeccion
t1/2	h	4.35 ± 0.47	34.20 ± 3.76	3.28 ± 0.36	28.50	4.04 ± 0.44	4.14 ± 0.45
Tmax	h	7 ± 0.91	4 ± 0.52	7 ± 0.91	4 ± 0.52	2 ± 0.26	2 ± 0.26
Cmax	µg/cm <sup>3</sup>	15.02 ± 1.86	353.98 ± 43.89	10.87 ± 1.34	298.34 ± 36.99	387.35 ± 48.03	371.23 ± 46.03
AUC 0-t	µg/cm <sup>3</sup> .h	111.65 ± 13.50	3355.96 ± 406.07	79.20 ± 9.53	2444.10 ± 295.73	1583.74 ± 191.63	1587.71 ± 192.11
AUC 0-inf_obs	µg/cm <sup>3</sup> .h	117.12 ± 15.46	8203.39 ± 1082.84	80.52 ± 10.62	5246.17 ± 692.49	1634.65 ± 215.77	1642.79 ± 216.84
MRT 0-inf_obs	h	9.80 ± 1.16	47.67 ± 5.67	8.59 ± 1.02	39.58 ± 4.71	5.36 ± 0.63	5.56 ± 0.66

562

563 Regarding the AUC value, the AUC values of CHL from DMN-MPs especially F1 in *ex vivo* infec-  
 564 tion models was discovered to be considerably higher compared to other formulations, demonstrat-  
 565 ing a good *ex vivo* skin bioavailability of our technique. It was discovered that the DMN containing  
 566 free CHL showed MRT less than 6 h in terms of retention duration in the infected skin. The MRT  
 567 values of CHL from DMN-MPs were found to be considerably higher ( $p < 0.05$ ) than those of  
 568 DMN-CHL, in the case of the retention period in the skin. A lower application time for CHL in the  
 569 treatment of skin infections may be possible because to the increased MRT. As a result, it can lead  
 570 to patients accepting this strategy. The combination of responsive MPs and DMNs might success-  
 571 fully transfer chlramphenicol into an *ex vivo* biofilm model, according to our findings.



592 **Figure 9.** The *ex vivo* concentrations and time profiles of CHL following the application of DMN-F1, DMN-F2, and  
 593 DMN-CHL in non-infected rat skin (a), as well as *ex vivo* infection models formed by SA (b)

594 **4. Conclusion**

595 This research aimed to mature and confirm a CHL analysis procedure that utilizes a UV-visible  
596 spectrophotometric method. This study's proposed method went through a validation process with  
597 these five parameters: accuracy, precision, linearity, LOD, LLOQ, and extraction recovery. The  
598 outcomes indicated which total recognition specifications were right defined and satisfied all of the  
599 qualifications from the ICH instructions. In addition, an accepted scientific method was successfully  
600 used to calculate the characteristics of the entrapment efficiency, drug loading, percentage recovery,  
601 drug release profile of MPs CHL-PCL, and *ex vivo* dermatokinetic of microneedle MPs CHL-PCL  
602 . To conclude, the method that has been validated has numerous applications for CHL studies to  
603 formulate and characterize bacterial responsive microparticle and microneedle systems.

604 **Disclosure of interest**

605 The authors declare no conflicts of interest.

606 **Acknowledgment**

607 Mukarram Mudjahid is thankful for the scholarship provided by German Academic Exchange Ser-  
608 vice (Deutscher Akademischer Austausch Dienst/DAAD). This study was supported by Penelitian  
609 Tesis magister, Ministry of Education, Culture, Research, and Technology of Republic of In-  
610 doonesia (090/E5/PG.02.00/PT/2022).

611 **Author Contribution**

612 **Mukarram Mudjahid:** Conceptualization; Data curation; Investigation; Methodology; Project ad-  
613 ministration; Validation; Roles/Writing - original draft; Writing - review & editing,  
614 **Sulistiawati:** Methodology; Software; Validation; Visualization; Writing - review & editing,  
615 **Rangga Meidianto Asri:** Validation; Writing - review & editing, **Firzan Nainu:** Validation; Su-  
616 pervision, **Andi Dian Permana:** Conceptualization; Funding acquisition; Project administration;  
617 Supervision; Writing - review & editing.

618

619 **References :**

- 620 [1] A. B. Raff and D. Kroshinsky, "Cellulitis A Review," *Jama*, vol. 316, no. 3, pp. 325–337,  
621 2016, doi: 10.1001/jama.2016.8825.
- 622 [2] R. Rrapi, S. Chand, and D. Kroshinsky, "Cellulitis: A Review of Pathogenesis, Diagnosis,

- 623 and Management,” *Med. Clin. North Am.*, vol. 105, no. 4, pp. 723–735, 2021, doi:  
624 10.1016/j.mcna.2021.04.009.
- 625 [3] S. M. Ellis Simonsen *et al.*, “Cellulitis incidence in a defined population,” *Epidemiol.*  
626 *Infect.*, vol. 134, no. 2, pp. 293–299, 2006, doi: 10.1017/S095026880500484X.
- 627 [4] C. A. Andersen, K. McLeod, and R. Steffan, “Diagnosis and treatment of the invasive  
628 extension of bacteria (cellulitis) from chronic wounds utilising point-of-care fluorescence  
629 imaging,” *Int. Wound J.*, no. September, pp. 1–13, 2021, doi: 10.1111/iwj.13696.
- 630 [5] E. L. A. Cross *et al.*, “Route and duration of antibiotic therapy in acute cellulitis: A  
631 systematic review and meta-analysis of the effectiveness and harms of antibiotic  
632 treatment,” *J. Infect.*, vol. 81, no. 4, pp. 521–531, 2020, doi: 10.1016/j.jinf.2020.07.030.
- 633 [6] O. Uddin, J. Hurst, T. Alkayali, and S. A. Schmalzle, “Staphylococcus hominis cellulitis  
634 and bacteremia associated with surgical clips,” *IDCases*, vol. 27, p. e01436, 2022, doi:  
635 10.1016/j.idcr.2022.e01436.
- 636 [7] J. L. P. Gemeinder *et al.*, “Gentamicin encapsulated within a biopolymer for the treatment  
637 of Staphylococcus aureus and Escherichia coli infected skin ulcers,” *J. Biomater. Sci.*  
638 *Polym. Ed.*, vol. 32, no. 1, pp. 93–111, 2021, doi: 10.1080/09205063.2020.1817667.
- 639 [8] S. Sartini *et al.*, “Current state and promising opportunities on pharmaceutical approaches  
640 in the treatment of polymicrobial diseases,” *Pathogens*, vol. 10, no. 2, pp. 1–31, 2021, doi:  
641 10.3390/pathogens10020245.
- 642 [9] S. Kalita *et al.*, “Chloramphenicol encapsulated in poly- $\epsilon$ -caprolactone-pluronic composite:  
643 Nanoparticles for treatment of MRSA-infected burn wounds,” *Int. J. Nanomedicine*, vol.  
644 10, pp. 2971–2984, 2015, doi: 10.2147/IJN.S75023.
- 645 [10] M. Abdollahi and S. Mostafalou, “Chloramphenicol,” *Encycl. Toxicol. Third Ed.*, vol. 1,  
646 pp. 837–840, 2014, doi: 10.1016/B978-0-12-386454-3.00709-0.
- 647 [11] P. Shukla, F. W. Bansode, and R. K. Singh, “Chloramphenicol Toxicity : A Review,” *J.*  
648 *Med. Med. Sci.*, vol. 2, no. 13, pp. 1313–1316, 2011.
- 649 [12] I. S. Ferreira *et al.*, “Activity of daptomycin- and vancomycin-loaded poly-epsilon-

- 650 caprolactone microparticles against mature staphylococcal biofilms,” *Int. J. Nanomedicine*,  
651 vol. 10, pp. 4351–4366, 2015, doi: 10.2147/IJN.S84108.
- 652 [13] T. W. Prow *et al.*, “Nanoparticles and microparticles for skin drug delivery,” *Adv. Drug*  
653 *Deliv. Rev.*, vol. 63, no. 6, pp. 470–491, 2011, doi: 10.1016/j.addr.2011.01.012.
- 654 [14] A. D. Permana *et al.*, “Selective delivery of silver nanoparticles for improved treatment of  
655 biofilm skin infection using bacteria-responsive microparticles loaded into dissolving  
656 microneedles,” *Mater. Sci. Eng. C*, vol. 120, no. November 2020, p. 111786, 2021, doi:  
657 10.1016/j.msec.2020.111786.
- 658 [15] N. Devnarain *et al.*, “Intrinsic stimuli-responsive nanocarriers for smart drug delivery of  
659 antibacterial agents—An in-depth review of the last two decades,” *Wiley Interdiscip. Rev.*  
660 *Nanomedicine Nanobiotechnology*, vol. 13, no. 1, pp. 1–38, 2021, doi: 10.1002/wnan.1664.
- 661 [16] L. Van Gheluwe, I. Chourpa, C. Gaigne, and E. Munnier, “Polymer-based smart drug  
662 delivery systems for skin application and demonstration of stimuli-responsiveness,”  
663 *Polymers (Basel)*, vol. 13, no. 8, 2021, doi: 10.3390/polym13081285.
- 664 [17] H. Chen *et al.*, “Design of smart targeted and responsive drug delivery systems with  
665 enhanced antibacterial properties,” *Nanoscale*, vol. 10, no. 45, pp. 20946–20962, 2018, doi:  
666 10.1039/c8nr07146b.
- 667 [18] Y. Li, G. Liu, X. Wang, J. Hu, and S. Liu, “Enzyme-Responsive Polymeric Vesicles for  
668 Bacterial-Strain-Selective Delivery of Antimicrobial Agents,” *Angew. Chemie - Int. Ed.*,  
669 vol. 55, no. 5, pp. 1760–1764, 2016, doi: 10.1002/anie.201509401.
- 670 [19] M. H. Xiong, Y. Bao, X. Z. Yang, Y. C. Wang, B. Sun, and J. Wang, “Lipase-sensitive  
671 polymeric triple-layered nanogel for ‘on-demand’ drug delivery,” *J. Am. Chem. Soc.*, vol.  
672 134, no. 9, pp. 4355–4362, 2012, doi: 10.1021/ja211279u.
- 673 [20] A. D. Permana, M. T. C. McCrudden, and R. F. Donnelly, “Enhanced intradermal delivery  
674 of nanosuspensions of antifilaria drugs using dissolving microneedles: A proof of  
675 concept study,” *Pharmaceutics*, vol. 11, no. 7, pp. 1–22, 2019, doi:  
676 10.3390/pharmaceutics11070346.
- 677 [21] A. J. Paredes *et al.*, “Microarray Patches: Poking a Hole in the Challenges Faced When

- 678 Delivering Poorly Soluble Drugs,” *Adv. Funct. Mater.*, vol. 31, no. 1, pp. 1–27, 2021, doi:  
679 10.1002/adfm.202005792.
- 680 [22] D. Karunakaran *et al.*, “Design and Testing of a Cabotegravir Implant for HIV Prevention,”  
681 *J. Control. Release*, vol. 330, pp. 658–668, 2021.
- 682 [23] F. P. Pons-Faudoa *et al.*, “2-Hydroxypropyl- $\beta$ -cyclodextrin-enhanced pharmacokinetics of  
683 cabotegravir from a nanofluidic implant for HIV pre-exposure prophylaxis,” *J. Control.*  
684 *Release*, vol. 306, pp. 89–96, 2019.
- 685 [24] Sulistiawati *et al.*, “Validation of spectrophotometric method to quantify cabotegravir in  
686 simulated vaginal fluid and porcine vaginal tissue in ex vivo permeation and retention  
687 studies from thermosensitive and mucoadhesive gels,” *Spectrochim. Acta - Part A Mol.*  
688 *Biomol. Spectrosc.*, vol. 267, p. 120600, 2022, doi: 10.1016/j.saa.2021.120600.
- 689 [25] S. Gorantla, R. N. Saha, and G. Singhvi, “Spectrophotometric method to quantify  
690 tofacitinib in lyotropic liquid crystalline nanoparticles and skin layers: Application in ex  
691 vivo dermal distribution studies,” *Spectrochim. Acta - Part A Mol. Biomol. Spectrosc.*, vol.  
692 255, p. 119719, 2021.
- 693 [26] A. Mahmood *et al.*, “UV spectrophotometric method for simultaneous estimation of  
694 betamethasone valerate and tazarotene with absorption factor method: Application for in-  
695 vitro and ex-vivo characterization of lipidic nanocarriers for topical delivery,” *Spectrochim.*  
696 *Acta - Part A Mol. Biomol. Spectrosc.*, vol. 235, p. 118310, 2020.
- 697 [27] H. E. Putri *et al.*, “Dissolving Microneedle Formulation of Ceftriaxone: Effect of Polymer  
698 Concentrations on Characterisation and Ex Vivo Permeation Study,” *J. Pharm. Innov.*, no.  
699 0123456789, 2021, doi: 10.1007/s12247-021-09593-y.
- 700 [28] D. Ramadan *et al.*, “A sensitive HPLC-UV method for quantifying vancomycin in  
701 biological matrices: Application to pharmacokinetic and biodistribution studies in rat  
702 plasma, skin and lymph nodes,” *J. Pharm. Biomed. Anal.*, vol. 189, p. 113429, 2020.
- 703 [29] A. D. Permana *et al.*, “New and sensitive HPLC-UV method for concomitant quantification  
704 of a combination of antifilaria drugs in rat plasma and organs after simultaneous oral  
705 administration,” *Anal. Methods*, vol. 13, no. 7, pp. 933–945, 2021.

- 706 [30] J. T. do P. Silva *et al.*, “Analytical validation of an ultraviolet–visible procedure for  
707 determining lutein concentration and application to lutein-loaded nanoparticles,” *Food*  
708 *Chem.*, vol. 230, pp. 336–342, 2017, doi: 10.1016/j.foodchem.2017.03.059.
- 709 [31] M. S. Raghu and K. Basavaiah, “Two charge-transfer complexation reactions for  
710 spectrophotometric determination of pheniramine maleate using  $\pi$ -acceptors,” *J. Sci. Ind.*  
711 *Res. (India)*., vol. 70, no. 10, pp. 851–858, 2011.
- 712 [32] N. Alhusein, I. S. Blagbrough, M. L. Beeton, A. Bolhuis, and P. A. De Bank, “Electrospun  
713 Zein/PCL Fibrous Matrices Release Tetracycline in a Controlled Manner, Killing  
714 *Staphylococcus aureus* Both in Biofilms and Ex Vivo on Pig Skin, and are Compatible with  
715 Human Skin Cells,” *Pharm. Res.*, vol. 33, no. 1, pp. 237–246, 2016, doi: 10.1007/s11095-  
716 015-1782-3.
- 717 [33] A. D. Permana *et al.*, “Solid lipid nanoparticle-based dissolving microneedles: A promising  
718 intradermal lymph targeting drug delivery system with potential for enhanced treatment of  
719 lymphatic filariasis,” *J. Control. Release*, vol. 316, no. September, pp. 34–52, 2019, doi:  
720 10.1016/j.jconrel.2019.10.004.
- 721 [34] Y. Zhang, M. Huo, J. Zhou, and S. Xie, “PKSolver: An add-in program for  
722 pharmacokinetic and pharmacodynamic data analysis in Microsoft Excel,” *Comput.*  
723 *Methods Programs Biomed.*, vol. 99, no. 3, pp. 306–314, 2010, doi:  
724 10.1016/j.cmpb.2010.01.007.
- 725 [35] J. T. do P. Silva *et al.*, “Analytical validation of an ultraviolet–visible procedure for  
726 determining lutein concentration and application to lutein-loaded nanoparticles,” *Food*  
727 *Chem.*, vol. 230, pp. 336–342, 2017, doi: 10.1016/j.foodchem.2017.03.059.
- 728 [36] E. E. Tseplin, S. N. Tseplina, and O. G. Khvostenko, “Mechanisms of specific effects of  
729 polar solvent in optical absorption spectra,” *Opt. Spectrosc. (English Transl. Opt. i*  
730 *Spektrosk.*, vol. 110, no. 6, pp. 903–909, 2011, doi: 10.1134/S0030400X11060166.
- 731 [37] ICH, “International Conference on Harmonisation,” in *Validation of Analytical*  
732 *Procedures: Text and Methodology Q2(R1), 1994 (2005)*, .
- 733 [38] M. Kiliçarslan and T. Baykara, “The effect of the drug/polymer ratio on the properties of

- 734 the verapamil HCl loaded microspheres,” *Int. J. Pharm.*, vol. 252, no. 1–2, pp. 99–109,  
735 2003, doi: 10.1016/S0378-5173(02)00630-0.
- 736 [39] N. Sharma, P. Madan, and S. Lin, “Effect of process and formulation variables on the  
737 preparation of parenteral paclitaxel-loaded biodegradable polymeric nanoparticles: A co-  
738 surfactant study,” *Asian J. Pharm. Sci.*, vol. 11, no. 3, pp. 404–416, 2016, doi:  
739 10.1016/j.ajps.2015.09.004.
- 740 [40] C. Folle *et al.*, “Thymol-loaded PLGA nanoparticles: an efficient approach for acne  
741 treatment,” *J. Nanobiotechnology*, vol. 19, no. 1, pp. 1–21, 2021, doi: 10.1186/s12951-021-  
742 01092-z.
- 743 [41] M. M. Badran, A. H. Alomrani, G. I. Harisa, A. E. Ashour, A. Kumar, and A. E. Yassin,  
744 “Novel docetaxel chitosan-coated PLGA/PCL nanoparticles with magnified cytotoxicity  
745 and bioavailability,” *Biomed. Pharmacother.*, vol. 106, no. July, pp. 1461–1468, 2018, doi:  
746 10.1016/j.biopha.2018.07.102.
- 747 [42] M. Mir, N. Ahmed, A. D. Permana, A. M. Rodgers, R. F. Donnelly, and A. U. Rehman,  
748 “Enhancement in site-specific delivery of carvacrol against methicillin resistant  
749 staphylococcus aureus induced skin infections using enzyme responsive nanoparticles: A  
750 proof of concept study,” *Pharmaceutics*, vol. 11, no. 11, 2019, doi:  
751 10.3390/pharmaceutics11110606.
- 752 [43] W. Chen, A. Palazzo, W. E. Hennink, and R. J. Kok, “Effect of particle size on drug  
753 loading and release kinetics of gefitinib-loaded PLGA microspheres,” *Mol. Pharm.*, vol. 14,  
754 no. 2, pp. 459–467, 2017, doi: 10.1021/acs.molpharmaceut.6b00896.
- 755 [44] C. Wu, T. F. Jim, Z. Gan, Y. Zhao, and S. Wang, “A heterogeneous catalytic kinetics for  
756 enzymatic biodegradation of poly (  $\epsilon$ -caprolactone ) nanoparticles in aqueous solution,”  
757 vol. 41, pp. 3593–3597, 2000.
- 758 [45] M. Mir, A. D. Permana, N. Ahmed, G. M. Khan, A. ur Rehman, and R. F. Donnelly,  
759 “Enhancement in site-specific delivery of carvacrol for potential treatment of infected  
760 wounds using infection responsive nanoparticles loaded into dissolving microneedles: A  
761 proof of concept study,” *Eur. J. Pharm. Biopharm.*, vol. 147, no. December 2019, pp. 57–

762 68, 2020, doi: 10.1016/j.ejpb.2019.12.008.

763

### **Credit Author Statement**

**Mukarram Mudjahid:** Conceptualization; Data curation; Investigation; Methodology; Project administration; Validation; Roles/Writing - original draft; Writing - review & editing, **Sulistiawati:** Methodology; Software; Validation; Visualization; Writing - review & editing, **Rangga Meidianto Asri:** Validation; Writing - review & editing, **Firzan Nainu:** Validation; Supervision, **Andi Dian Permana:** Conceptualization; Funding acquisition; Project administration; Supervision; Writing - review & editing.

**Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

**BUKTI**  
**ACCEPTED**



Andi Dian Permana &lt;andi.dian.permana@farmasi.unhas.ac.id&gt;

---

**Your Submission SAA-D-22-02512R1**

2 messages

---

**Spectrochimica Acta Part A** <em@editorialmanager.com>  
Reply-To: Spectrochimica Acta Part A <support@elsevier.com>  
To: Andi Dian Permana <andi.dian.permana@farmasi.unhas.ac.id>

Fri, Jan 13, 2023 at 2:49 AM



Ms. Ref. No.: SAA-D-22-02512R1

Title: Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: application to in vitro release and ex vivo dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria.

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

Dear Dr. Andi Dian Permana,

I am pleased to confirm that your paper "Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: application to in vitro release and ex vivo dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria." has been accepted for publication in Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy.

Comments from the Editor and Reviewers can be found below.

Your accepted manuscript will now be transferred to our production department and work will begin on creation of the proof. If we need any additional information to create the proof, we will let you know. If not, you will be contacted again in the next few days with a request to approve the proof and to complete a number of online forms that are required for publication.

We encourage authors of original research papers to share the research objects – including raw data, methods, protocols, software, hardware and other outputs – associated with their paper. More information on how our open access Research Elements journals can help you do this is available at [https://www.elsevier.com/authors/tools-and-resources/research-elements-journals?dgcid=ec\\_em\\_research\\_elements\\_email](https://www.elsevier.com/authors/tools-and-resources/research-elements-journals?dgcid=ec_em_research_elements_email).

Thank you for submitting your work to this journal.

With kind regards,

Christian Huck  
Editor  
Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

Comments from the Editors and Reviewers:

Reviewer #1: The authors have addressed all the comments. I think that manuscript should be accepted for publication.

Reviewer #3: Across the manuscript. Change r2 to r2

Line 66-67 - "The system can be designed to adapt responsively release the drug to the specific infected area [15,16]." - correct sentence. Maybe: The system can be designed to adapt release of the drug to the specific infected area.

Lines 89-90 - "A fundamental part of developing new drug delivery systems in detecting and quantifying the drug," - please correct sentence. I think in should be replaced by is

Line 95 - "these prominent research" - not sure what this means. Maybe "research articles"

Line96 - "reported method is unsuitable" - plural. "method are"

Line 195 - "degree of among a sequence" - not sure what this means

Line 218 - "The needle in the DMN is taken out by scraping the needle, then the needles is dissolved in water." - please review this sentence. Is only 1 needle removed? Needle/needles is/are

Line 266 - "91.54 ± 2,67 percent and 94.77 ± 0,96 percent" - use % sign and . instead of , for decimals

Line 279 - "show" - replace by shows

Figure 3 legend. It stated mean +- SD n=3 but I only see 1 point and no error bar. Please clarify

Line 402 - "were extracted from skin tissue were compared" - delete the first "were"

Line 405 - "FDA bioanalytical geudeline recommendations" - guideline

Table 11 - replace , by .

Lines 441-442 - "F1 were not significantly ( $p > 0.05$ ) compared to MPs F2." - significantly what? Different?

Line 494 - "In figure 5, shows" - replace by figure 5 shows

Line 501 - ", ready for a faster release" - it is not clear what this means

Line 532 - "In this study, we explain to compared" - maybe replace by we compared

Line 535 - "Table 13 has shown lists" - please clarify

Line 540 - "results..This" - delete 1 .

Line 550 - "has a significant greater" - significantly

Line 556 - ", ready for a faster release" - it is not clear what this means

Line 563 - "especially" - specifically

For further assistance, please visit our customer support site at <http://help.elsevier.com/app/answers/list/p/7923>. Here you can search for solutions on a range of topics, find answers to frequently asked questions and learn more about EM via interactive tutorials. You will also find our 24/7 support contact details should you need any further assistance from one of our customer support representatives.

#AU\_SAA#

To ensure this email reaches the intended recipient, please do not delete the above code



**Spectrochimica Acta Part A** <em@editorialmanager.com>  
Reply-To: Spectrochimica Acta Part A <support@elsevier.com>  
To: Andi Dian Permana <andi.dian.permana@farmasi.unhas.ac.id>

Fri, Jan 13, 2023 at 2:49 AM



Ms. Ref. No.: SAA-D-22-02512R1

Title: Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: application to in vitro release and ex vivo dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria.

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

Dear Dr. Andi Dian Permana,

I am pleased to confirm that your paper "Validation of spectrophotometric method to quantify chloramphenicol in fluid and rat skin tissue mimicking infection environment: application to in vitro release and ex vivo dermatokinetic studies from dissolving microneedle loaded microparticle sensitive bacteria." has been accepted for publication in Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy.

Comments from the Editor and Reviewers can be found below.

Your accepted manuscript will now be transferred to our production department and work will begin on creation of the proof. If we need any additional information to create the proof, we will let you know. If not, you will be contacted again in the next few days with a request to approve the proof and to complete a number of online forms that are required for publication.

We encourage authors of original research papers to share the research objects – including raw data, methods, protocols, software, hardware and other outputs – associated with their paper. More information on how our open access Research Elements journals can help you do this is available at [https://www.elsevier.com/authors/tools-and-resources/research-elements-journals?dgcid=ec\\_em\\_research\\_elements\\_email](https://www.elsevier.com/authors/tools-and-resources/research-elements-journals?dgcid=ec_em_research_elements_email).

Thank you for submitting your work to this journal.

With kind regards,

Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy

Comments from the Editors and Reviewers:

%EDITOR\_DECISION\_PHRASES%

%COMMENTS\_TO\_AUTHOR%

[Quoted text hidden]