

# A.\_Dian\_Permana-1\_Usulan.pdf

*by*

---

**Submission date:** 06-Dec-2022 10:38AM (UTC+0700)

**Submission ID:** 1972790956

**File name:** A.\_Dian\_Permana-1\_Usulan.pdf (5.58M)

**Word count:** 12228

**Character count:** 64608



ELSEVIER

36

Contents lists available at ScienceDirect

Journal of Controlled Release

journal homepage: [www.elsevier.com/locate/jconrel](http://www.elsevier.com/locate/jconrel)

## Hydrogel-forming microarray patches with cyclodextrin drug reservoirs for long-acting delivery of poorly soluble cabotegravir sodium for HIV Pre-Exposure Prophylaxis

Fabiana Volpe-Zanutto<sup>a,b</sup>, Lalit Kumar K. Vora<sup>a</sup>, Ismaiel A. Tekko<sup>a,c</sup>, Peter E. McKenna<sup>a</sup>, Andi Dian Permana<sup>a,d</sup>, Akmal H. Sabri<sup>a</sup>, Qonita K. Anjani<sup>a</sup>, Helen O. McCarthy<sup>a</sup>, Alejandro J. Paredes<sup>a</sup>, Ryan F. Donnelly<sup>a,\*</sup>

<sup>a</sup> School of Pharmacy, Queen's University Belfast, 97 Lisburn Road, Belfast BT9 7BL, UK

<sup>b</sup> Faculty of Pharmaceutical Sciences, R. Cândido Portinari, 200 - Cidade Universitária, Campinas - SP, 13083-871, University of Campinas, Brazil

<sup>c</sup> Faculty of Pharmacy, Aleppo University, Syria

<sup>d</sup> Department of Pharmaceutics, Faculty of Pharmacy, Hasanuddin University, Makassar, Indonesia

### ARTICLE INFO

#### Keywords:

HIV  
Microneedles  
Cabotegravir sodium  
Cyclodextrin  
PrEP  
Pharmacokinetic

### ABSTRACT

Hydrogel-forming microarray patches (HF-MAPs) offer minimally invasive, pain-free and prolonged drug delivery. These devices are designed to be self-administered and self-disabling, avoiding contaminated sharps waste generation. Cabotegravir sodium (CAB-Na) is a poorly soluble anti-human immunodeficiency virus (HIV) drug for the treatment and pre-exposure prophylaxis of HIV infection that lends itself to depot formation following intradermal delivery but presents significant challenges when delivered via HF-MAPs, whose nature is aqueous. Herein, we have investigated, for the first time, the use of hydroxypropyl- $\beta$ -cyclodextrin (HP- $\beta$ -CD) to enhance the solubility of CAB-Na, and its effect on intradermal delivery via HF-MAPs. Accordingly, tablet reservoirs containing CAB-Na and HP- $\beta$ -CD were formulated. These novel reservoirs were combined with two different HF-MAP formulations (MAP1 (Gantrez S97® + poly (ethylene glycol) 10,000 + Na<sub>2</sub>CO<sub>3</sub>) and MAP2 (poly (vinyl pyrrolidone) 58 kDa + poly (vinyl alcohol) 85–120 kDa + citric acid)) to form fully integrated MAP devices which were tested in both *ex vivo* and *in vivo* settings. *Ex vivo* skin deposition results for MAP1 and MAP2 showed that 141 ± 40 µg and 342 ± 34 µg of CAB-Na was deposited into 0.5 cm<sup>2</sup> of excised neonatal porcine skin after 24 h, respectively. Based on these findings, the *in vivo* pharmacokinetics of MAP2 were investigated over 28 days using a Sprague-Dawley rat model. After 24 h patch application, MAP2 demonstrated an extended drug release profile and an observed C<sub>max</sub> of 53.4 ± 10.16 µg/mL, superior to that of an FDA-approved CAB-nanosuspension administered via intramuscular application (C<sub>max</sub> of 43.6 ± 5.3 µg/mL). Consequently, this tablet integrated MAP device is considered to be a viable option for the intradermal delivery of hydrophobic anti-HIV drugs.

### 1. Introduction

The first cases of acquired immunodeficiency syndrome (AIDS), an illness caused by human immunodeficiency virus (HIV), were reported in 1981 [1]. Since then, 59 million has claimed approximately 36 million lives and it is projected that at the end of 2020 that there were 37.7 million people living with HIV, 25.4 million of which are concentrated in Sub-Saharan Africa, yet only 27.5 million people receiving antiretroviral therapy (ART) [2,3]. Currently, there are an estimated 1.5 million new HIV infections each year and, in 2020 alone, approximately

680,000 deaths were attributed to HIV-related illnesses [4,5]. Current ART strategies can be costly, which limits the accessibility of ART for those living in lower- and middle-income areas within the region of Sub-Saharan Africa [6,7]. Furthermore, in this region, the availability of trained healthcare staff for application of parenteral treatment (who are required for ART in its current form) is low, which negatively impacts the rate of successful treatment. Pre-exposure prophylaxis (PrEP) is a highly effective pharmacological tool used for the purpose of HIV prevention among high-risk groups [6]. Effective use of PrEP has previously demonstrated a significant reduction in new HIV infections among high-

19

\* Corresponding author at: School of Pharmacy, Queen's University Belfast, Medical Biology Centre, 97 Lisburn Road, Belfast BT9 7BL, UK.

E-mail address: [r.donnelly@qub.ac.uk](mailto:r.donnelly@qub.ac.uk) (R.F. Donnelly).

46

<https://doi.org/10.1016/j.jconrel.2022.06.028>

Received 16 March 2022; Received in revised form 31 May 2022; Accepted 16 June 2022

Available online 21 June 2022

0168-3659/© 2022 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

risk individuals from 86% to 44% [8,10].

Cabotegravir (CAB) is one of the most promising ART drugs available for PrEP, due to its high effectiveness and reduced occurrence of collateral side effects when compared to other compounds [8–12]. CAB is an analogue of dolutegravir (DOL) and acts as an integrase strand transfer inhibitor [10]. Aiming to develop a parentally administered, long-acting (LA) PrEP for HIV, CAB-Nanosuspension (CAB-LA) was formulated for intramuscular administration [13]. In 2021, the US FDA (Food and Drug Administration) authorised [9] first long-acting treatment of HIV infection in adults containing the combination of CAB-Nanosuspension (CAB LA) and rilpivirine (RPV LA). [13].

CAB LA overcomes the drawbacks related with oral dosing of CAB, extending its half-life by 25-fold (25–52 days in humans). CAB LA can be administered for PrEP in negative-HIV individuals with high risk of infection via intramuscular injection (IM) monthly (400 mg) or bimonthly (800 mg), thus, reducing dosing frequency and improving patient adherence [8,14]. However, parenteral administration requires the use of a hypodermic needle, which many individuals find unacceptable due to needle phobia [15]. Moreover, parenterally-administered products generate clinical sharps waste, a significant issue, especially in the case of HIV treatment. This presents a substantial HIV transmission risk to both patients and healthcare workers, with those in low and middle income regions having higher incidences of unsafe injection practices [16–18]. Other issues that still need to be addressed include the cost of CAB-LA manufacturing, which is both energy and labour-intensive, and the fact that transport in low- and middle-income regions is often challenging and, as a result, many people simply cannot access the treatment that they need. Whilst CAB-LA greatly improves the experience of a patient receiving treatment, it requires administration by trained healthcare staff. Thus, it faces obstacles to widespread use in low-resource countries with insufficient healthcare workers [5,16,19–22]. Therefore, the development of alternative LA formulations that are inexpensive to manufacture, demonstrate reduced side effect incidence, and that can be self-administered is an exciting concept to be explored [18].

A microarray patch (MAPs) is composed of an array of micron-sized needles arranged on a flat baseplate. Upon application, microneedles (MNs) pierce the *stratum corneum*, the outermost layer of the skin, and deliver therapeutics intradermally without eliciting pain or bleeding [23,24]. Significantly, MAPs are designed to be self-administered i.e., trained healthcare staff are not required for their use. Furthermore, certain MAP types, namely hydrogel-forming and dissolving MAPs, are self-disabling in nature, meaning that they are not capable of reinsertion in the skin after use, thus effectively eliminating the risk of infection transmission [24–26]. Hydrogel-forming MAPs (HF-MAPs) are a type of MAP made from a cross-linked polymer matrix that absorbs interstitial fluid from the skin upon application to form a hydrogel *in situ* [25,55]. In contrast to other MAP types, where the drug is loaded into or onto individual MNs, hydrogel-forming MAPs have a separately-formulated drug reservoir that is connected to the back of the MN array. Upon sufficient swelling of the array, this drug reservoir dissolves, releasing drug into the inserted hydrogel, from which it is delivered intradermally in a rate-controlled manner [27]. Crucially, this two-part design facilitates enhanced drug loading and further expands the range of deliverable molecules to include not only potent low-dose molecules, but also those that require high doses. Although intrinsically well-suited for the delivery of hydrophilic drug molecules, hydrophobic compounds may also be successfully delivered using this platform, when combined with a suitable solubility-enhancing technique [28].

In this work, the aqueous solubility of CAB-Na was enhanced by complexing the drug with cyclodextrin (CD) molecules. CDs are cyclic oligosaccharides that possess a truncated cone structure and are classified as  $\alpha$ ,  $\beta$ , or  $\gamma$  types, which contain six, seven, and eight glucose units, respectively. Importantly, they have a hydrophobic inner cavity which is

surrounded by a hydrophilic outer surface [29]. This structure allows them to interact with and house hydrophobic compounds in their inner cavity, whilst simultaneously solubilising themselves and the drug in an aqueous environment [30]. In the present study, CAB-Na was combined with hydroxypropyl- $\beta$ -cyclodextrin to formulate the suitable tablet reservoir that dissolved rapidly and completely when situated atop a swollen HF-MAP. Intradermal delivery of this anti-HIV therapeutic was achieved over 24 h in rats *in vivo* and drug pharmacokinetics were studied over the following 28 days, revealing that the formulated HF-MAP device had a LA delivery profile. The work outlined within highlights, for the first time, the potential of this technology to overcome many issues commonly faced by patients who require HIV PrEP treatment, particularly in low-resource settings.

## 2. Material and methods

### 2.1. Materials

Cabotegravir sodium (CAB-Na) and raltegravir long-acting (CAB LA) vials for intramuscular injection (CAB LA, 400 mg/2 mL) were provided by ViiV Healthcare (Brentford, UK). Polyvinylalcohol (PVA) molecular weight (m/w) of 85–120 kDa, 87%–89% hydrolysed, polyethyleneglycol (PEG) 10,000, Na<sub>2</sub>CO<sub>3</sub> (sodium carbonate), acetonitrile (ACN) (HPLC grade), fluoracetic acid, heparin, isoflurane, formic acid, citric acid, and phosphate buffered saline (PBS, 5.17.4) tablets were purchased from Sigma-Aldrich (Dorset, U.K.). Gantrez® S-97 (copolymer of methylvinylether and maleic acid 1500 kDa (m/w)), polyvinylidene fluoride (PVDF) K29-32 (58 kDa m/w) and 2-hydroxypropyl- $\beta$ -cyclodextrin (HP- $\beta$ -CD) Cavitron™ HP7 Pharma were gifted from Ashland Inc. (Kidderminster, U.K.). All other materials and chemicals were of analytical reagent grade.

### 2.2. Fabrication of “Super-swelling” HF-MAPs

In this work, two HF-MAP formulations were studied. The first to be studied was a “Super swelling” [31] formulation which was composed of copolymer of methylvinylether and maleic acid (20% w/w), PEG 10,000 (7.5% w/w) and Na<sub>2</sub>CO<sub>3</sub> (3% w/w) dispersed in deionized water [31]. Approximately 600 mg of the hydrogel was carefully dispensed into pre-formed silicone MAP moulds (MNs arranged in a 11 × 11 formation, 900  $\mu$ m height, 300  $\mu$ m needle width and 150  $\mu$ m needle interspacing at the base) [32]. Filled moulds were centrifuged at 3500 RPM for 15 min and allowed to dry for 48 h at room temperature. The HF-MAPs were then crosslinked for 24 h at 80 °C to induce ester-bonds between Gantrez® S-97 and PEG [31,57] (Table 1).

### 2.3. Fabrication of PVA/PVP HF-MAPs

The second HF-MAP formulation was composed of PVA (15% w/w), PVP (10% w/w), citric acid (1.5% w/w) dissolved in water [28]. In a similar fashion to the protocol for the fabrication of “Super-swelling” HF-MAPs, analogous MAP moulds were filled with the prepared aqueous blend, centrifuged at 3500 RPM for 15 min, dried at 25 °C for 48 h, and crosslinked according to the conditions outlined in Table 1.

**Table 1**  
Hydrogel formulations and crosslinking conditions.

Formulation	Hydrogel Composition % (w/w)	Crosslinking Temperature °C	Crosslinking Time (h)
MAP1	Gantrez S97® (20%) + sodium carbonate (3%) + PEG 10,000 (7.5%)	80	24
MAP2	PVA (15%) + PVP (10%) + Citric acid (1.5%)	130	1

#### 2.4. Optical microscopy analysis and scanning electron microscopy (SEM)

The structure and morphology of the formulated MAPs was assessed by optical digital light microscopy (Leica EZ4D, Leica, Wetzlar, Germany) and scanning electron microscopy (SEM) (TM3030, Hitachi, Krefeld, Germany). For SEM analysis, MAPs were attached on an aluminum support with copper and carbon conductive tapes. The SEM images were taken by scanning with backscattered electrons at 15 kV [23].

#### 2.5. Swelling studies of crosslinked MAPs

To investigate the swelling profiles of each MAP formulation, individual MAPs (0.5 cm<sup>2</sup>) had their masses recorded in the dry state ( $m_0$ ). They were then placed in an excess volume of water containing 5% w/w HP- $\beta$ -CD and allowed to swell ( $n = 3$ ). At pre-determined time points, the swollen MAPs were removed from the water, gently dabbed with filter paper to remove surface water and then re-weighed, with their masses recorded as  $m_t$  [14]. Percentage swelling of each formulation was calculated using Eq. (1).

$$\%Swelling = \frac{(m_t - m_0)}{m_0} \times 100 \quad (1)$$

#### 2.6. Rationalisation of CAB-Na/HP- $\beta$ -CD

Phase solubility analysis, as proposed by Higuchi and Connors, was employed to investigate the effect of HP- $\beta$ -CD on the aqueous solubility of CAB-Na [33]. To characterise the solubility enhancement inferred by CD complexation and to optimise the formulation of drug/CD inclusion complexes, an excess of CAB-Na was put into aqueous solutions containing increasing concentrations of HP- $\beta$ -CD (0, 5, 10 and 20% w/w). Samples were maintained at room temperature and gently agitated with magnetic stirrers until equilibrium was reached i.e., until the amount of undissolved CAB-Na remained constant. At this point, the saturation solubility of CAB-Na in each of the CD concentrations tested was deduced by filtering samples through a 0.22  $\mu$ m syringe filter to remove undissolved drug and then quantifying dissolved CAB-Na with HPLC-UV validated method (Section 2.17) [32,34].

#### 2.7. Preparation of directly compressed tablets containing CAB-Na/HP- $\beta$ -CD complexes

Solid state CAB-Na/HP- $\beta$ -CD complexes were formed by lyophilisation of the aqueous solution containing the optimum concentration of HP- $\beta$ -CD and CAB-Na [33,35]. Following lyophilisation using the Freeze Drier System (Virtis Advantage, SP Scientific Ltd., Warminster PA, USA), 250 mg of powdered inclusion complex was placed into a tablet die and compressed into tablets under manual hydraulic press (Specac Ltd., Kent, U.K.) with a compression force of 2 t for 20 s. The tablets produced contained 10 mg of CAB-Na, with a diameter and thickness of 13 mm and 1.4 mm, respectively.

#### 2.8. Tablet characterisation

Characterisation of the formed directly compressed tablet reservoirs (i.e., tablet diameter and thickness) was performed visually using a digital microscope ( $n = 6$ ). Tablet dissolution time was measured by placing each tablet in 20 mL of deionized water with continuous stirring at 1200 RPM ( $n = 6$ ).

#### 2.9. Mechanical characterisation of HF-MAPs

Formulated HF-MAPs were characterised in terms of the percentage of height reduction, insertion efficiency, and insertion depth. To

determine the height reduction, individual MNs were measured after compression for 30 s at 32 N onto an aluminum block utilising a TA.XT2 Texture Analyser (Stable Microsystems, Haselmeere, UK) (force = 32 N) [36,37].

An optical coherence tomography (OCT) (EX1301 microscope, Michelson Diagnostics Ltd., Kent, UK) was employed to obtain cross-sectional images of HF-MAPs inserted into excised full-thickness porcine skin and the penetration depth of the MAPs in the skin ( $n = 6$ ). The OCT configuration parameters were pre-established by Donnelly et al. The skin was removed from stillborn piglets, excised, and stored at  $-20^\circ\text{C}$  until use. Prior to use, the skin tissue was de-frosted and equilibrated in PBS (pH 7.4), at which point hair was carefully removed with help of manual disposable razor.

#### 2.10. X-ray powder diffraction (XRD)

The XRD analyses were performed using a MiniFlex II powder X-Ray diffractometer (Rigaku Corporation, Tokyo, Japan), to evaluate the crystallinity of the drug, CD and drug/CD inclusion complex. Samples of CAB-Na, HP- $\beta$ -CD and CAB-Na/HP- $\beta$ -CD inclusion complex were individually placed into the rotating sample support. The analyses were performed in continuous mode within the range of  $3-60^\circ 2\theta$  (scanning rate of  $2^\circ/\text{min}$ ) in triplicate. The analyses were carried using a current of 15 mA and a voltage of 30 kV. Furthermore, the diffractometer was equipped with a Ni filter, and Cu K $\beta$  radiation was used to conduct the experiment [36].

#### 2.11. Fourier transform infrared spectroscopy (FTIR)

The FTIR analyses of CAB-Na, HP- $\beta$ -CD and the CAB-Na/HP- $\beta$ -CD complex were performed using a FTIR spectrometer Cary 630 (Agilent, Santa Clara, California, USA) to identify the interactions that occur during inclusion complex formation. The IR experiment were scanned and recorded in between 4000 and 400  $\text{cm}^{-1}$  at room temperature using attenuated total reflection (ATR) mode. Resolution was maintained at 4.0  $\text{cm}^{-1}$  throughout all analyses and acquired spectra were the result of an average of 64 scans.

#### 2.12. Identification of complex formation by NMR (methods)

NMR analyses of 1D: <sup>1</sup>H and 2D: ROESY were performed using a spectrophotometer with a nuclear magnetic resonance of 11.7 Tesla (Varian® Inova). The samples were solubilised in the deuterated dimethyl sulfoxide (DMSO- $d_6$ ) (Sigma-Aldrich® Co. Ltd. – Gillingham, Dorset, UK). Tetramethylsilane was applied as an internal standard.

#### 2.13. Drug content

The drug content of directly compressed tablets (CAB-Na/HP- $\beta$ -CD complex) was determined by dissolving each tablet in 10 mL of water ( $n = 6$ ). The solution obtained was filtered using a syringe filter (0.22  $\mu$ m), suitably diluted and analysed by validated HPLC-UV system (Section 2.17).

#### 2.14. Skin drug deposition studies (ex vivo)

Ex vivo studies, which investigated the deposition of CAB-Na from the formulated HF-MAPs, were performed using full thickness natal porcine skin (treated and stored as explained in Section 2.9). The skin was affixed to the donor part of the Franz cell apparatus before the HF-MN array was inserted using manual pressure for 30 s. After insertion, 20  $\mu$ L of PBS was aliquoted onto the baseplate of the array to promote reservoir adhesion and the formulated CAB-Na/HP- $\beta$ -CD tablet was placed on top. To guarantee adequate insertion of the HF-MAP device throughout the study, a cylinder-shaped weight made of stainless-steel (12.0 g) was mounted on top of the CAB-Na/HP- $\beta$ -CD tablet and the

Franz-cell apparatus was then sealed using Parafilm® M. To ensure sink conditions, the receiver medium used was composed of HP-β-CD 5% w/w in deionised water (12 mL), degassed and pre-heated at 32 °C ± 1 °C. After 24 h, the apparatus was disassembled, the HF-MAP was removed, and the skin was separated from the donor compartment. Undeposited CAB-Na, which was visible on the side of the skin, was removed. Deposited drugs within the skin tissue were extracted according to the protocol in Section 2.15. Once again, samples (both from the skin and the donor compartment) were analysed with the HPLC-UV method outlined in Section 2.17.

### 2.15. Drug extraction from skin samples

Skin samples were cut from the MAP application site into tiny pieces using scissors and transferred to 2 mL centrifuge tubes having 500 µL of water + 1 mL acetonitrile. These samples were then homogenised for 15 min with stainless steel beads (5 mm diameter) using a TissueLyserLT (QIAGEN, U.K.) to extract the drug. The tissue samples were transferred to a 15 mL falcon tube, where 3.5 mL of acetonitrile: water (v/v) was added and left in ultrasonic bath for 1 h. Finally, samples were homogenised using vortex (1 min) and 100 µL of the supernatant was collected in a 1.5 mL centrifuge tube containing 900 µL of acetonitrile, vortex 1 min, and centrifuged at 14,800 rpm for 10 min. The resulted supernatant was then transferred to 2 mL HPLC vials and the drug was quantified using HPLC-UV system.

### 2.16. Drug recovery from hydrogel-forming MAPs

Following completion of *ex vivo* skin deposition studies, the quantity of drug remaining in the swollen hydrogel MAPs was calculated by immersing the MAP in 5 mL acetonitrile and incubating samples at 37 °C ± 1 °C for 24 h to extract the drug. After incubation, the acetonitrile, into which CAB-Na had been extracted, was removed, and diluted appropriately with the same solvent. Samples were then vortexed for 6 min, centrifuged at 14,800 rpm for 10 min and subsequently analysed by HPLC-UV method.

### 2.17. HPLC-UV validation method (in vitro analysis)

Analysis of CAB-Na was carried out by using the reversed-phase HPLC (Agilent 1200 equipment, Agilent Technologies UK Ltd., UK). The UV detector was set at 257 nm, and the chromatographic separation was carried out using an Agilent ODS-3 C18 column (5 µm particle size and 150 mm length x 4.6 mm internal diameter) (GL Science Inc., Tokyo, Japan) maintained at 40 °C. Isocratic elution was employed with mobile phase which composed of ACN and 0.1% TFA (70:30 v/v). The injection volume of 40 µL and the flow rate was maintained at 0.8 mL/min. The data was obtained using an Agilent Chem-Station® Software (Version: B.02.01) and the analytical method validation was performed according to the corresponding International Council of Harmonisation (ICH) guidelines [32].

### 2.18. In vivo studies

An *in vivo* pharmacokinetic study of MAP2 combined with CAB-Na/HP-β-CD directly compressed tablet reservoirs was conducted (Project Licence no. 2794). This *in vivo* experiment was reviewed and approved by the Queen's University Belfast Biological Services Unit Ethical Committee. Prior to study commencement, female rats (Sprague Dawley, 8–12 weeks of age) (205 ± 8 g) were acclimatised for 7 days and fasted overnight the day before study start. The rats were split into 2 groups (n = 6): Group 1 (control) – each rat was treated with an IM injection of CAB LA suspension 10 mg/rat, and Group 2 – each rat received four MAPs (MAP2 combined with CAB-Na/HP-β-CD) – Drug loading per tablet = 10 mg; total drug “dose” = 40 mg/rat. Approximately 24 h before MAP application, hair was removed from the backs of the rats in

Group 2 using Veet® depilatory cream (RB Healthcare, Ltd., Dansom Lane, Hull, UK). Applied MAPs were immobilised on the backs of each rat using adhesive foam and Tegaderm™ (3 M, Bracknell, UK) and kinesiology tape (KT Health, American Fork, USA). For procedures where anaesthesia was required, 2–4% isoflurane in oxygen was used. On day 1, blood samples were taken via tail vein bleeding at t = 1, 2, 4 and 6 h. Subsequent blood samples were collected at 1, 2, 3, 7, 14, 21, 28 and 35 days after treatment with either IM injection or MAPs.

### 2.19. Drug extraction from plasma

Study samples, which were composed of whole rat blood, were taken in 1.5 mL centrifuge tubes containing of heparin (10 µL) and centrifuged at 1509 g for 15 min (4 °C) to separate the plasma component. To extract CAB-Na from plasma, 100 µL of drug-containing plasma was mixed with 300 µL acetonitrile and vortexed at high speed for 15 min [39]. Then, the precipitated samples were centrifuged at 16,000 g for 15 min (4 °C) and the supernatant collected for analysis using the HPLC-MS method outlined in Section 2.21.

### 2.20. Analysis of pharmacokinetic (PK) parameters

Following quantification of CAB-Na in plasma samples, the delivery profile of CAB-Na was plotted *i.e.*, drug concentration in the plasma versus time. Subsequently, the pharmacokinetic parameters of this profile, namely the maximum CAB-Na concentration (C<sub>max</sub>), the time needed to achieve C<sub>max</sub> (T<sub>max</sub>), half-life elimination time (T<sub>1/2</sub>), mean residence time (MRT) and the area under the curve (AUC) were estimated using a non-compartmental PK model [40]. The relative bioavailability of CAB-Na in plasma after the administration of MAP compared to IM injection was calculated using Eq. (2).

$$FR = \frac{AUC_{MAP} \times Dose_{IM}}{AUC_{IM} \times Dose_{MAP}} \quad (2)$$

where, FR is the relative bioavailability, AUC<sub>MAP</sub> is the area under curve obtained from the integrated MAP2 formulation and AUC<sub>IM</sub> is the area under curve obtained from CAB LA (IM) injection.

### 2.21. HPLC-MS analysis for CAB-Na (in vivo analysis)

The development of a HPLC-MS method with isocratic elution was achieved using a HPLC with diode array detector (DAD), containing a quaternary pump, multi-sampler, multi-column thermostat, attached to a single quadrupole mass spectrometer (API 6400, Agilent Technologies, UK) operated in a positive electrospray ionisation mode was utilised for species detection. CAB and CAB-Na detected at m/z 406 and 428, respectively. The drying temperature was maintained at 300 °C, capillary voltage at 4 kV, nebuliser pressure at 15 psi and drying gas flow at 11 L/min. Chromatographic separation was achieved using a ZORBAX™ Eclipse XDB-C18 column (1.8 µm particle size, 50 mm length and 4.6 mm internal diameter), with column temperature kept at 40 °C. The mobile phase was composed of 70% v/v ACN and 30% v/v ultra-purified water containing 0.1% v/v formic acid. The solvent flow rate was held at 0.3 mL/min for 6 min, and 10 µL of the sample injection volume [34].

### 2.22. Statistical analysis

Where applicable, the data was interpreted using one way analysis of variance (ANOVA) along with the Turkey-Kramer *post hoc* test (GraphPad Prism® software, version 9.1.0, San Diego, California, USA). In all cases, p < 0.05 denoted significance. Linear regression was applied for solubility studies using GraphPad Prism® software where p > 0.0037 denoted significance. Non-compartmental pharmacokinetic profile parameters were evaluated using the PKSolver add-in program in

Microsoft Excel™ (Microsoft Corporation, USA).

80

### 3. Results and discussion

#### 3.1. Microscopic analysis of HF-MAPs

The morphology of formulated HF-MAPs was analysed using both digital light microscopy and SEM (Fig. 1). The chosen MN geometry was an 11 × 11 array (total MNs per array = 121) with MNs that were 900 μm high, composed of a pyramidal tip (600 μm) and cuboidal base (300 μm). Interspacing of MNs was 300 μm and the theoretical needle volume of each array was 5.45 mm<sup>3</sup> [32]. In this work, two HF-MAP types were investigated, namely the “Super-swelling” MAP1 formulation (Fig. 1A) and the PVA/PVP MAP2 formulation (Fig. 1B).

#### 3.2. Swelling studies of crosslinked MAPs

Swelling studies were conducted to evaluate the capacity of swelling for MAP1 (“Super-swelling”) and MAP2 (PVA-PVP) formulations. This work served as an indicator for the capacity of each type of cross-linked hydrogel to absorb interstitial fluid and swell when inserted into the skin [27]. As water containing 5% w/w HP-β-CD was used as the receiver compartment media during *ex vivo* skin deposition studies, it was also used here as the swelling media. It can be seen in Fig. 2 that the “Super swelling” formulation based on Gantrez® S-97 had a swelling capacity that was approximately 9-fold greater than the PVA/PVP hydrogel. Anjani et al. reported similar swelling profiles for both hydrogel formulations when analogous swelling studies were performed in PBS (pH 7.4) [28]. An important observation during this study was that MAPs prepared from the “Super swelling” formulation became structurally [103] after approximately 30 min of swelling, whereas, PVA/PVP MAPs remained intact throughout the duration of the study.

#### 3.3. Mechanical characterisation of HF-MAPs

The ability of HF-MAPs to penetrate and insert into neonatal porcine skin was examined using OCT, as described previously [34]. Analysis of the OCT images obtained (Fig. 3) demonstrated that the penetration depth of MNs from MAP1 and MAP2 were 608.15 ± 43.95 μm for MAP1 and 724.41 ± 32.05 μm, respectively. These insertion depth values were




Fig. 2. Percentage swelling capacity of MAP1 and MAP2 in water containing 5% w/w HP-β-CD (mean ± S.D., n = 3).

higher than values previously reported [27,28]. A possible explanation for this could be related to the geometry of the MNs used in this study *i. e.*, cuboidal bases and pyramidal tips. This MN geometry may enhance skin penetration, compared to those that are conically shaped, due to the sharper edges of the cuboidal base more readily breaching the *stratum corneum* [41]. Additionally, it can be assumed that all MAPs were inserted accounting for approximately 68% and 80% of MAP height for MAP1 and MAP2, respectively.

These results have demonstrated that both MAP1 and MAP2 formulations have the capacity to effectively penetrate the *stratum corneum* (10–20 μm), excluding soles of feet and the palms (400–600 μm) [42]. The viable epidermis in humans, which is 100 μm thick, makes up the next layer of skin [42]. Beyond the epidermis is the dermis, a roughly 1200 μm thick where the majority of systemic drug are absorbed and it occurs due to the high level of vascularization in this layer [42]. The results from the simulated skin [81] sersion studies suggest the ability of MAPs 1 and 2 of penetrating an approximate depth of 600–725 μm, which indicated the capability of the formulated MAPs to easily bypass the *stratum corneum* (~20 μm) as well as epidermis (~100 μm) layers. Provided that the loaded drug can successfully diffuse through the cross-linked hydrogel, it is reasonable to assume that drug deposition in the dermal layer, followed by systemic absorption via the dermal micro-circulation, would occur [31].




Fig. 1. Digital light microscope images of HF-MAPs: (A)1 - MAP1 x8 magnification, (B)1 - MAP2 x8 magnification, (A)2 - MAP1 x35 magnification and (B)2 - x35 magnification. SEM images of HF-MAPs:




Fig. 3. (A) Percentage insertion of MNs on formulated HF-MAPs into full-thickness skin at a force of 32 N/ 30 s/array (means  $\pm$  S.D.,  $n = 5$ ). (B) Reduction in MN height (%) noted following compression onto an aluminum block (32 N/ 30 s/ array) (means  $\pm$  S.D.,  $n = 5$ ). (C) Representative OCT images of HF-MAPs inserted in full-thickness neonatal porcine skin - (C) 1 = MAP1 and (C) 2 = MAP2 (The scale bars represent a length of 500  $\mu$ m).

To ensure penetration of the *stratum corneum* and subsequent drug deposition into lower layers of the skin, MNs should be strong enough to puncture the skin without breaking [40]. The height reduction after compression force application of HF-MNs was evaluated by compressing HF-MAPs using a force of 32 N/ 30 s and calculating the percentage MN height reduction (Fig. 3 (B)). Observed percentage MN height reduction was  $4.57 \pm 1.39\%$  and  $5.26 \pm 1.55\%$  for MAP1 and MAP2, respectively. These values were consistent with those previously reported and confirmed the suitability of both formulations in terms of mechanical strength [28,43].

### 3.4. Rationalisation of CAB-Na/HP- $\beta$ -CD inclusion complex formation and tablet characterisation

The first objective of this work was to formulate a water-soluble drug reservoir using biocompatible material that would enhance the solubility of CAB-Na. Various surfactants and polymers were tested to improve CAB-Na solubility. However, only HP- $\beta$ -CD was able to solubilise CAB-Na successfully (Table S1). To examine the relationship between concentration of HP- $\beta$ -CD and the observed solubility enhancement, an excess of CAB-Na was added to aqueous solutions containing increasing different concentrations of HP- $\beta$ -CD. Fig. 4 (A) illustrates the influence of CD concentration on the aqueous solubility of CAB-Na. A linear relationship existed between the two components *i.e.*, as the concentration of HP- $\beta$ -CD was increased, the solubility of CAB-Na also increased. From this data, it was then possible to estimate the ratio of HP- $\beta$ -CD to CAB-Na required for successful inclusion complex formation using Eqs. (3) and (4), where the moles of CAB-Na and HP- $\beta$ -CD were compared.

$$CAB-Na + HP\beta CD = CAB-Na - HP\beta CD \quad (3)$$

$$K = \frac{[CAB-Na - HP\beta CD]}{[CAB-Na]_x [HP\beta CD]} \quad (4)$$

Based on Eqs. (3) and (4), the stoichiometric molar ratio between CAB-Na: HP- $\beta$ -CD was found to be 1:1 in the inclusion complex (Fig. 4 (B)). This indicated that successful inclusion complex formation occurred between one molecule of CAB-Na and one molecule of HP- $\beta$ -CD. Furthermore, based on the maximum observed solubility of CAB-Na ( $8.6 \pm 0.22$  mg/mL), which occurred in the solution containing HP- $\beta$ -CD 20% w/w, an approximate 19-fold solubility enhancement could be ascribed to the formation of CAB-Na/HP- $\beta$ -CD inclusion complexes. Of those tested, 20% w/w HP- $\beta$ -CD was carefully chosen as the most suitable concentration for the formation of CAB-Na/HP- $\beta$ -CD inclusion complexes, as it demonstrated the greatest improvement in the solubility of the drug. Higher CD concentrations were not investigated, as they would greatly increase the bulk of the final dosage form, which may then have affected the performance of the final integrated MAP device.

To obtain CAB-Na/HP- $\beta$ -CD inclusion complexes in the solid state, an aqueous solution of HP- $\beta$ -CD (20% w/w) was saturated with CAB-Na and the resultant solution was then lyophilised. By removing water in this manner, the remaining powder consisted of CAB-Na complexed with HP- $\beta$ -CD in a mass ratio of 4:96 w/w. To convert this lyophilised powder into a suitable drug reservoir for use with a HF-MN array, it was processed into directly compressed tablets as depicted in Fig. 4 (C) and (D). Each tablet was composed of 250 mg of CAB-Na/HP- $\beta$ -CD inclusion complex powder (drug content =  $10.32 \pm 0.03$  mg or  $4.22 \pm 0.03\%$ ), measured  $13 \pm 0.11$  mm in diameter, and had a thickness of  $1.40 \pm 0.02$  mm. Moreover, the formulated reservoir was distinctly hygroscopic in nature - dissolving in approximately 90 s during dissolution studies.

### 3.5. X-ray powder diffraction (XRD) analyses

The crystalline nature of CAB-Na, HP- $\beta$ -CD, physical mixture (PM) of CAB-Na and HP- $\beta$ -CD PM, and powdered CAB-Na/HP- $\beta$ -CD inclusion complexes were examined using XRD (Fig. 5). The diffractogram obtained for CAB-Na was typical of a crystalline substance, with intense peaks observed at  $7.3, 19.0, 20.6$  and  $24.1^\circ$  theta [44]. When compared




Fig. 4. (A) Observed solubility of CAB-Na when solubilised in aqueous condition containing HP-β-CD at concentrations varying from 5 to 20% w/w (means + S.D.,  $n = 12$ ). (B) Solubilisation of CAB-Na in HP-β-CD solutions. Solubility of CAB-Na is linearly reliant on the concentration of HP-β-CD. The proportion found for the inclusion complex was 1:1 (CAB-Na: HP-β-CD) stoichiometrically. (C) Microscopic image of CAB-Na/HP-β-CD tablet upper surface and (D) Microscopic image of CAB-Na/HP-β-CD tablet lateral surface.

to the diffractogram for HP-β-CD, which had two broad peaks at 11.9 and 19.9° theta and was bereft of any intense crystalline peaks, it was evident that the CD demonstrated extensive amorphous content [45]. The XRD of the CAB-Na and HP-β-CD physical mixture resulted in a diffractogram that overlapped with the patterns of these individual components, verifying that there was no chemical interaction between CD and drug, and that both components had retained their original physical characteristics. Finally, analysis of CAB-Na/HP-β-CD inclusion complexes provided a diffractogram that was similar to that of HP-β-CD alone and revealed a disappearance of the intense crystalline peaks observed during analysis of the pure drug. This finding indicated the conversion of the crystalline structure of CAB-Na to one that was amorphous, most likely due to successful CD/drug inclusion complex formation [45].

### 3.6. NMR and FTIR results

The complexation that took place between HP-β-CD and CAB-Na was confirmed by monitoring the proton chemical shifts using <sup>1</sup>H NMR. <sup>1</sup>H NMR spectra of the host HP-β-CD (host) and CAB-Na (guest) molecules and corresponding complexes were compared as shown in Fig. 6 (A, B and C). Upon complexation, it was clear that some degree of  $\Delta\delta$  (proton chemical shift) had occurred for H-3 and H-5. These are the protons that reside within the cavity of the CD. In the presence of aromatic molecules such as those present on CAB-Na, the effect of resonance would induce an anisotropic effect on the H-3 and H-5 protons, resulting in proton chemical shifts,  $\Delta\delta$  [46]. In addition, by monitoring the degree  $\Delta\delta$  between H-3 and H-5, it can be deduced that there was partial or complete drug inclusion into the CD [47]. Table 3 shows that  $\Delta\delta$  for H-5 > H-3, indicating total drug inclusion into the HP-β-CD cavity. In addition, FTIR

analysis was also conducted to complement the data obtained via <sup>1</sup>H NMR about the chemical interaction between HP-β-CD and CAB-Na. It can be seen from the FTIR analysis in Fig. 6 (D), that a broad peak at 1028 cm<sup>-1</sup> was observed for the CAB-Na/HP-β-CD complex. This was not apparent for free HP-β-CD and CAB-Na + HP-β-CD physical mixture. The broad peak at 1028 cm<sup>-1</sup> was assigned to the C-O stretching vibrations of the cyclodextrin molecule when complexation occurs between the cyclic oligosaccharides and CAB-Na [48]. The interaction between CAB-Na and HP-β-CD to form the complex of inclusion occurred in the H-subject (CAB-Na) to the OH (HP-β-CD), clearly represented at 3363 cm<sup>-1</sup> band and confirmed at the 1371 cm<sup>-1</sup> stretching, suggesting a hydrogen bond interaction between CAB-Na and HP-β-CD occurred [48]. Collectively the <sup>1</sup>H NMR analysis in tandem with FTIR spectra indicate successful CAB-Na complexation with the ring of HP-β-CD.

### 3.7. Ex vivo drug deposition studies

The results shown in Table 4 and Fig. 7 illustrate the amount of CAB-Na delivered into and through neonatal porcine skin (full-thickness, ~ 2 mm) after application of previously formulated CAB-Na/HP-β-CD tablets in combination with MAP1 and MAP2. Additionally, the delivery of CAB-Na from CAB-Na/HP-β-CD tablets without MAPs was investigated i.e., tablets were applied directly to the skin. As detailed in Table 2, the drug content of the formulated CAB-Na/HP-β-CD tablets was approximately 10,325 ± 35.82 μg per tablet.

CAB delivery from the integrated MAP1 containing the CAB-Na/HP-β-CD as a drug reservoir (Fig. 8 (A)) was evaluated employing a Franz-cell set-up with full-thickness neonatal porcine skin. The total drug delivered was 208.9 ± 51.93 μg/0.5 cm<sup>2</sup>/CAB-Na/24 h, corresponding




Fig. 5. XRD diffractograms of CAB-Na, HP- $\beta$ -CD, a physical mixture of CAB-Na and HP- $\beta$ -CD (CAB-Na + HP- $\beta$ -CD PM), and CAB-Na/HP- $\beta$ -CD inclusion complexes.

to  $2.02 \pm 0.52\%$ . The extent of drug deposited in the skin after 24 h was  $141.63 \pm 40.32 \mu\text{g}$  ( $1.41 \pm 0.40\%$ ) of CAB-Na per  $0.5 \text{ cm}^2$  skin, while  $67.27 \pm 11.61 \mu\text{g}$  ( $0.67 \pm 0.11\%$ ) of CAB-Na permeated across the skin into the receptor compartment of the Franz-cells within 24 h.

For MAP1, most of the drug was deposited into the swollen polymeric network, from where  $6650 \pm 395.35 \mu\text{g}$  ( $64.40 \pm 4.99\%$ ) of unreleased CAB-Na was recovered. The recovery from donor compartments and skin surfaces was also studied, with a combined total of  $2.64 \pm 0.28\%$  of CAB-Na found here.

In the same *ex vivo* skin delivery experiment, MAP2 was able to deliver  $456.60 \pm 90.51 \mu\text{g}/0.5 \text{ cm}^2$  of CAB-Na after 24 h. This corresponded to  $4.41 \pm 0.65\%$  of total CAB-Na loaded and was 2.23-fold greater than observed with MAP1. From this total delivery,  $324.00 \pm 38.04 \mu\text{g}/0.5 \text{ cm}^2$  ( $3.13 \pm 0.43\%$ ) was deposited in the skin and  $132.60 \pm 52.47 \mu\text{g}/0.5 \text{ cm}^2$  ( $1.28 \pm 0.22\%$ ) permeated through the skin. Regarding recovery data, a large amount of drug was also found in the swollen polymeric matrix of MAP2, accounting for  $4407 \pm 650.62 \mu\text{g}$  ( $42.68 \pm 8.22\%$ ) of CAB-Na. In addition, a total of  $4542 \pm 1028.71 \mu\text{g}$  ( $43.99 \pm 12.99\%$ ) of CAB-Na was recovered from the donor compartment and skin surface.

The third group tested in this experiment was the CAB-Na/HP- $\beta$ -CD tablet applied to intact skin without MAP application ('No MAP'). Drug deposition in the skin with the No MAP group was greatly reduced, with  $21.13 \pm 2.30 \mu\text{g}$  ( $0.20 \pm 0.02\%$ ) delivered after 24 h. For this group, the drug was not detected in the receiver compartment, highlighting the importance of MAP application to enhance the intradermal and transdermal delivery of CAB-Na. Accordingly, CAB-Na was only recovered from the donor compartment and skin surface after 24 h, where a total of  $6775 \pm 5.68 \mu\text{g}$  ( $65.16 \pm 0.05\%$ ) of CAB-Na was found.

The successful delivery of CAB-Na via MAP2 can be explained by crosslinking difference between the two formulations [28]. As reported by Anjani et al [28], the super-swelling MAP, formulated with Gantrez®

S97, PEG 10,000 and  $\text{Na}_2\text{CO}_3$  has a polymer network formed with bigger pores when compared with the PVA-PVP formulation. In addition, there is the possibility that Gantrez® S97 interacts strongly with the drug and/or the CD as well, explaining a marked difference observed between samples from MAP1 and MAP2 after the skin deposition experiment. Fig. 8 (B) shows the MAP1 (super-swelling) surface, with a large amount of drug deposited into the MAP1 polymeric network ( $\sim 64\%$ ), which resulted in the formation of holes at the tips after the saturation of the hydrogel network with CAB-Na/HP- $\beta$ -CD complex. Fig. 8 (C) shows the image of the skin surface that was in contact with MAP1, after 24 h of experiment. It is possible to see the drug precipitated on the skin surface. This phenomenon occurred due to the super-swelling formulation's high capacity of swelling and getting out of the skin after some period, which explains the amount of drug accumulated on the skin surface. Unlike MAP1, MAP2 remained fully formed, with no apparent precipitation of the drug in the polymeric network - Fig. 8 (D) -, and the skin surface (E) was not covered with the drug.

The CAB-Na/HP- $\beta$ -CD complex tablet was able to release the hydrophobic CAB-Na through the cross-linked HF-MAPs, because these types of hydrogels are inherently aqueous in nature [27,43]. The water-soluble CAB-Na/HP- $\beta$ -CD complex was capable of delivering the hydrophobic CAB-Na through the HF-MAPs due to the drug solubility enhancement achieved by complexation using HP- $\beta$ -CD.

### 3.8. In vivo studies

Pharmacokinetic investigation of the delivery of CAB-Na from the formulated integrated MAP (MAP2 arrays combined with CAB-Na/HP- $\beta$ -CD) tablet reservoirs were carried out on female Sprague-Dawley rats (aged between 8 and 12 weeks). The delivery of this anti-HIV therapeutic using MAP technology was compared to that of an FDA-approved long-acting nanosuspension of CAB (CAB-LA - average particle size =  $200 \text{ nm}$ ) delivered via IM injection. The aim of this study was to deduce if minimally invasive HF MAP-mediated delivery using the formulated MAP device was a suitable alternative to the marketed injectable formulation. After removing MAP2, we observed no inflammation or erythema at the side application as previously reported when applied in the human skin (*in vivo*) [25,49]. Fig. 9 and Table 5 depict the pharmacokinetic profile of the delivery of CAB-LA administered via IM injection and that of CAB-Na delivered via MAPs.

The observed  $C_{\text{max}}$  for the CAB LA (IM) group was  $43.76 \pm 5.36 \mu\text{g}/\text{mL}$  and was achieved after 6 days ( $T_{\text{max}}$ ). The  $C_{\text{max}}$  observed following treatment with the formulated MAP device was  $58.98 \pm 9.95 \mu\text{g}/\text{mL}$ , with a  $T_{\text{max}}$  of 3 days. Exposure of rats to CAB was estimated using AUC as a parameter. The  $\text{AUC}_{0-28}$  values for the IM and MAP groups were  $644.76 \pm 55.43 \mu\text{g}/\text{mL}\cdot\text{day}$  and  $529.58 \pm 86.42 \mu\text{g}/\text{mL}\cdot\text{day}$ , respectively. Interestingly, the AUC values between these groups were not statistically different ( $p > 0.05$ ). The MRT following administration of CAB-Na via MAPs intradermally was approximately 10 days ( $9.71 \pm 0.48$  days), whereas this value was just over 17 days for CAB-LA administered via IM. Additionally, the relative bioavailability of CAB following MAP administration was compared to IM injection, which revealed a bioavailability of  $4.9 \pm 0.8\%$ . Crucially, despite the observed differences in several pharmacokinetic parameters between the two groups, both cohorts demonstrated statistically similar plasma concentrations across the 28-day study ( $p > 0.05$ ). Considering that the therapeutic plasma concentration of CAB ranges from  $0.664 \mu\text{g}/\text{mL}$  ( $4 \times \text{PA-IC}_{90}$ ) to  $0.1664 \mu\text{g}/\text{mL}$  ( $\text{PA-IC}_{90}$ ) [34,48], the present study shows that the formulated MAP device was able to deliver CAB at  $4 \times$  concentrations above the clinically-relevant plasma protein-adjusted concentration. This plasma level inhibits virus replication by  $90\%$  ( $4 \times \text{PA-IC}_{90}$ ), guaranteeing the drug release into tissues at relevant concentrations and consequently maintaining therapeutic levels of CAB *in vivo* for 28 days for HIV prevention [34,48]. Considering the observed plasma concentration at the 28 day time point, which was approximately 22-fold higher than the  $\text{PA-IC}_{90}$ , additional *in vivo* investigation is necessary to




Fig. 6. Proton nuclear magnetic resonance ( $^1\text{H}$  NMR) spectra of (A) HP- $\beta$ -CD, (B) Cabotegravir sodium and (C) CAB-Na/HP- $\beta$ -CD Complex; (D) FTIR spectra of CAB-Na, CAB-Na/HP- $\beta$ -CD Complex, a physical mixture of CAB-Na and HP- $\beta$ -CD (CAB-Na + HP- $\beta$ -CD PM) and HP- $\beta$ -CD.

evaluate <sup>55</sup> formulation over longer time periods. The present data indicates the potential for sustained release of CAB-Na via HF-MAP for long-acting PrEP delivery, deserving further investigation. The initial

burst release from the formulated HF-MAP device may be attributed to the greatly enhanced solubility of CAB-Na brought about by inclusion complex formation. The rapid delivery and, subsequent, absorption of





Fig. 6. (continued).




Fig. 7. (A) Concentration of CAB-Na deposited into full-thickness skin after 24 h (B) Concentration of CAB-Na permeated through the skin after 24 h. (Means  $\pm$  S.D.,  $n = 6$ ) ND\* = drug was not detected.

**Table 2**  
Amount of CAB-Na present in lyophilised CAB-Na/HP- $\beta$ -CD inclusion complex powder (100 mg) and in formulated CAB-Na/HP- $\beta$ -CD tablets (250 mg) (means  $\pm$  S.D.,  $n = 12$ ).

CAB-Na content		
Formulation	mg	% (w/w)
CAB-Na in 100 mg of CAB-Na/HP- $\beta$ -CD	4.13 $\pm$ 0.02	4.13 $\pm$ 0.02
CAB-Na (Tablet 250 mg)	10.32 $\pm$ 0.03	4.22 $\pm$ 0.03

**Table 3**  
 $^1\text{H}$  NMR chemical shift ( $\Delta\delta$ ) data of HP- $\beta$ -CD and the CAB-Na/HP- $\beta$ -CD complexes.  $\Delta\delta = \delta(\text{complex}) - \delta(\text{free})$ .

$^1\text{H}$ -assignment	HP- $\beta$ -CD	CAB-Na/HP- $\beta$ -CD Complexes	$\Delta\delta$ /ppm
H-1	5.890	5.887	-0.003
H-2	3.601	3.601	0.000
H-3	4.832	4.833	0.001
H-4	3.295	3.292	-0.003
H-5	3.750	3.742	-0.008
H-6	3.750	3.742	-0.008
H-8	4.558	4.551	-0.007
H-9	1.018	1.019	-0.001

**Table 4**  
Mass transfer of CAB-Na from formulated tablets using MAP1, MAP2 and No MAP. The percentage values (% w/w) were calculated in relation to 10.3 mg of CAB-Na present in 250 mg CAB-Na/HP- $\beta$ -CD tablet. (Means  $\pm$  S.D.,  $n = 6$ ).

Type of MAP	MAP1 - "Super-Swelling"		MAP2 - PVA-PVP		No MAP	
	$\mu\text{g}$	%	$\mu\text{g}$	%	$\mu\text{g}$	%
Drug deposited skin/24 h	141.63 $\pm$ 40.32	1.37 $\pm$ 0.50	324.00 $\pm$ 38.04	3.13 $\pm$ 0.43	21.13 $\pm$ 2.30	0.20 $\pm$ 0.02
Drug permeated/24 h	67.27 $\pm$ 11.61	0.65 $\pm$ 0.02	132.60 $\pm$ 52.47	1.28 $\pm$ 0.22	ND	ND
Recovery MNs array	6650 $\pm$ 395.35	64.40 $\pm$ 4.99	4407 $\pm$ 650.62	42.68 $\pm$ 8.22	*	*
Recovery weight/donor compartment	273.20 $\pm$ 20.05	2.64 $\pm$ 0.28	4542 $\pm$ 1028.71	43.99 $\pm$ 12.99	6775 $\pm$ 5.68	65.16 $\pm$ 0.05
Recovery total	7132 $\pm$ 442.02	69.07 $\pm$ 5.59	9405 $\pm$ 588.74	91.08 $\pm$ 7.43	6796 $\pm$ 7.22	65.82 $\pm$ 0.06

CAB-Na observed following MAP application provided a faster  $T_{\text{max}}$  and higher  $C_{\text{max}}$  compared to the slow releasing CAB LA IM injection. Encouragingly, the formulated MAP device also demonstrated an


**Table 5**  
CAB pharmacokinetic parameters in animals and their corresponding dose based on the body weight-normalised, following administration of CAB-Na delivered via MAPs (MAP2 formulation) and CAB-LA administered via IM injection (Means  $\pm$  S.D.,  $n = 6$ ).

Pharmacokinetic Parameters	CAB LA (IM)	CAB-Na (MAP2+CAB-Na/HP- $\beta$ -CD tablet)
Applied dose (mg/rat)	2.5	40
Applied dose(mg/kg)	11.8 $\pm$ 0.5	189.4 $\pm$ 9.7
$C_{\text{max}}$ ( $\mu\text{g/mL}$ )	43.76 $\pm$ 5.36	58.98 $\pm$ 9.95
$T_{\text{max}}$ (day)	5.83 $\pm$ 2.05	2.83 $\pm$ 1.17
$C_{28 \text{ days}}$ ( $\mu\text{g/mL}$ )	11.2 $\pm$ 3.3	3.6 $\pm$ 0.8
$AUC_{0-28}$ ( $\mu\text{g/mL}\cdot\text{day}$ )	644.76 $\pm$ 55.43	529.59 $\pm$ 86.42
$AUC_{0\text{inf}}$	816.48 $\pm$ 106.83	559.94 $\pm$ 92.44
MRT (day)	17.27 $\pm$ 4.09	9.71 $\pm$ 0.48
$T_{1/2}$ (day)	10.12 $\pm$ 2.73	5.79 $\pm$ 0.20
FR <sup>2</sup>	1.00	0.049 $\pm$ 0.008

extended delivery profile. This could be attributed to the combination of rate-controlled drug delivery through a cross-linked hydrogel matrix and the formation of drug  $\beta$ -CD inclusion complexes to the dermal layer of the skin, the binding constant between CAB and HP- $\beta$ -CD greatly influences drug release [48]. As a result, it is unlikely that all of the delivered inclusion complexes will remain intact, rather it is probable that at least some dissociate into CAB-Na and HP- $\beta$ -CD [48]. This intradermal dissociation may lead to the formation of micro-depots of the poorly soluble drug CAB-Na in the skin, from which extended drug release occurs over a period of at least one month.

In this study, HF-MAPs have been successfully utilised for the long-acting delivery of the poorly soluble anti-HIV drug CAB. These MAPs, which absorb interstitial fluid from the skin upon application, swell to form an inserted hydrogel matrix *in situ* through which therapeutics can be delivered to the skin. Transcutaneous delivery in this manner is inherently well-suited for the delivery of hydrophilic molecules, due to the aqueous nature of a swollen hydrogel. However, the delivery of hydrophobic drugs using this technique has proven to be problematic [25,28]. Through the complexation of CAB-Na with the solubility enhancing oligosaccharide HP- $\beta$ -CD and subsequent formulation into a soluble tablet reservoir, the previously elusive delivery of a hydrophobic compound via HF-MAPs was achieved. This finding will serve to expand the library of drugs deliverable using HF-MAPs to include those that display poor aqueous solubility.

Regarding the use of MAP technology to deliver anti-HIV therapeutics, only four drugs have been successfully delivered previously, rilpivirine (RPV) [50], etravirine [51], cabotegravir [34], and tenofovir alafenamide [18] using dissolving MAPs or long-acting implantable poly



106

Fig. 8. (A) Schematic representation of HF-MAP mediated delivery of a drug from a tablet reservoir. Digital light microscopy images of (B) MAP1 after skin deposition experiment (x12.5 magnification), (C) skin surface where MAP1 was applied during skin deposition experiment (x12.5 magnification), (D) MAP2 after skin deposition experiment (x8 magnification), and (E) skin surface where MAP2 was applied during skin deposition experiment (x12.5 magnification).

(d,l-lactide-co-glycolide) (PLGA) MAPs [18,58]. In all cases, formulation and successful delivery in a long-acting manner was achieved by forming intradermal micro-depots in the skin using dissolving MAPs. To guarantee the successful delivery of a hydrophobic anti-HIV drug for the first-time using HF-MAPs, it was important to confirm their ability to penetrate the skin and maintain their swollen form after influx of interstitial skin fluid. In this work, a new MAP design which was comprised of individual MNs measuring 900  $\mu\text{m}$  in height, with a cuboidal base (300  $\mu\text{m}$ ) and a pyramidal tip (600  $\mu\text{m}$ ), arranged in an 11  $\times$  11 formation with 300  $\mu\text{m}$  interspacing between MNs, was employed. The formulated MAPs were able to penetrate the skin and swell effectively without exhibiting expulsion from the skin [32]. The formulated MAP2 associated with drug reservoir CAB-Na/HP- $\beta$ -CD had 5% relative bioavailability, corresponding to 500  $\mu\text{g}$  of CAB-Na delivered per 0.5  $\text{cm}^2$  in 24 h from the drug reservoir. Similar amounts of drug were delivered in *ex vivo* experiment using Franz-cell set ups ( $\sim$  456.60  $\mu\text{g}/0.5 \text{ cm}^2/24 \text{ h}$ ) with full-thickness skin, reaffirming that the *in vitro* methodology to evaluate the new formulations can predict the expected

*in vivo* delivery.

Not only is the work reported here the first to deliver a hydrophobic compound via HF-MAPs, but it is also the first example of the delivery of an anti-HIV therapeutic using the same platform. Additionally, we have also demonstrated the successful delivery of a long-acting depot using HF-MAP for the first time. This phenomenon has not been reported previously and may hold great potential for the delivery of a numerous types of molecules for a range of indications. The recently FDA-approved CAB LA and RPV LA injections, which are the first and only marketed formulations that provide long-acting for HIV PrEP in adults [52]. Whilst these products represent a milestone in the evolution of HIV therapy, they require monthly or bimonthly administration by a trained healthcare professional using invasive IM injections. During Phase II and III clinical trials, >95% of individuals receiving CAB-LA via IM injection reported pain upon injection [53,56]. Furthermore, the generation of potentially-dangerous sharps waste is considered a major issue when dealing with a disease such as HIV, particularly in low and middle income settings, where the majority of HIV cases exist [21]. Therefore,




Fig. 9. Pharmacokinetic profiles following the administration of CAB-LA administered via IM injection and CAB-Na delivered via MAP2 + CAB-Na/HP-β-CD (Means ± S.D.,  $n = 6$ ). Statistical analysis using an unpaired *t*-test was used to compare groups at each time point. Accordingly, ns denoted no significant difference ( $p > 0.05$ ), \* denoted  $p \leq 0.05$ ; \*\* denoted  $p \leq 0.01$  and \*\*\* denoted  $p \leq 0.0001$ .

MAPs represent a promising alternative to the currently available oral and injectable treatments due to their ability to deliver HIV therapeutics at clinically relevant levels in a long-acting and minimally-invasive manner, without generating infectious sharps waste. Furthermore, MAPs are designed to be self-administered by the patient [54], thus relieving the burden placed on healthcare professionals in terms of treatment delivery.

#### 4. Conclusion

This work describes, for the first time, the successful delivery of a poorly soluble anti-HIV drug using HF-MAPs. Cyclodextrin complexation has been proven to sufficiently solubilise the hydrophobic drug CAB-Na to facilitate its transdermal delivery across the aqueous channels of a swollen hydrogel MAP. This successful transdermal delivery can be attributed to the enhanced solubility exhibited by CAB-Na/HP-β-CD inclusion complexes which are able to traverse the swollen polymer matrix of an implanted HF-MAP. Following successful intradermal deposition, these complexes are sufficiently diluted in the skin and begin to spontaneously dissociate leading to intradermal drug depot formation and an extended drug release profile. The utilisation of effective HP-β-CD complexation teamed with the rate-controlled delivery of high drug loads provided by the HF-MAP platform facilitated the delivery of clinically relevant doses of CAB-Na in an *in vivo* setting over a 28-day period following a single administration. Further work is now required to optimise the formulated MAP device in terms of delivery efficiency. Nevertheless, the findings of this work demonstrate that the current long-acting delivery profile is similar to that of the FDA-approved CAB-LA (IM injection) marketed for PrEP and HIV treatment. Alternative minimally invasive delivery systems such as HF-MAPs have the potential to unlock benefits for those who are affected by HIV PrEP. Realisation of this potential will surely improve the quality of life of millions across the globe.

#### CRedit authorship contribution statement

**Fabiana Volpe-Zanutto:** Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources, Software, Validation, Writing – original draft, Writing – review & editing. **Lalitkumar K. Vora:** Data curation, Conceptualization, Formal analysis, Investigation, Methodology, Writing – review & editing. **Ismael A. Tekko:** Data curation, Formal analysis, Conceptualization, Investigation. **Peter E. McKenna:** Writing –

original draft, Writing – review & editing. **Andi Dian Permana:** Data curation, Formal analysis, Conceptualization, Writing – original draft. **Akmal H. Sabri:** Data curation, Methodology, Formal analysis, Visualization, Writing – original draft. **Qonita K. Anjani:** Investigation, Methodology, Formal analysis, Writing – original draft. **Helen O. McCarthy:** Investigation, Methodology. **Alejandro J. Paredes:** Writing – review & editing. **Ryan F. Donnelly:** Supervision, Resources, Funding acquisition, Writing – original draft, Writing – review & editing.

#### Acknowledgements

Thanks to ViiV Healthcare for supplying CAB LA injectable formulation, and the micronized cabotegravir sodium. The authors would also like to thank the PATH team members who contributed to the strategic direction of this work and/or provided review and copyediting support, including Courtney Jarrahan, Abra Greene, Jessica Mistilis, Maggie Kilbourne-Brook, Annie Rein-Weston, Jill Sherman-Konkle, and Darin Zehrung. This project was made possible by the generous support of the American people through the United States Agency for International Development (USAID) through the United States President's Emergency Plan for AIDS Relief (PEPFAR), under the terms of Cooperative Agreement #AID-OAA-A-17-00015. The contents are the responsibility of QUB and PATH and do not necessarily reflect the views of USAID, PEPFAR, or the United States government. Also, this work was supported in part by EPSRC grant EP/S028919/1 and Wellcome Trust grant WT094085MA.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jconrel.2022.06.028>.

#### References

- R.K.R. Rajoli, Z.R. Demkovich, C. Flexner, A. Owen, M. Sic, Predicting pharmacokinetics of a tenofovir alafenamide subcutaneous implant using PBPK modelling, *Antimicrob. Agents Chemother.* (2020) 1–19, <https://doi.org/10.1128/AAC.00155-20>.
- UNAIDS, Global HIV & AIDS Statistics vol. 2, 2021. <https://www.unaids.org/en/resources/fact-sheet> (accessed August 11, 2021).
- W. World Health Organization, HIV/AIDS, 30 Novemb. 2021, 2021.
- W. World Health Organization, HIV/AIDS, 30 Novemb. 2021. <https://www.who.int/news-room/fact-sheets/detail/hiv-aids>, 2021 (accessed December 1, 2021).
- A.J. Paredes, I.K. Ramöller, P.E. McKenna, M.T.A. Abbate, F. Volpe-Zanutto, L. K. Vora, M. Kilbourne-Brook, C. Jarrahan, K. Moffatt, C. Zhang, I.A. Tekko, R. F. Donnelly, Microarray patches: breaking down the barriers to contraceptive care and HIV prevention for women across the globe, *Adv. Drug Deliv. Rev.* 173 (2021) 331–348, <https://doi.org/10.1016/j.addr.2021.04.002>.
- J. Ghosh, B. Taiwo, S. Seedat, B. Autran, C. Katlama, Seminar HIV, *Lancet.* 392 (2018), [https://doi.org/10.1016/S0140-6736\(18\)31311-4](https://doi.org/10.1016/S0140-6736(18)31311-4).
- J.M. Molina, I. Charreau, B. Spire, L. Cotte, J. Chas, C. Capitant, C. Tremblay, D. Rojas-Castro, E. Cua, A. Pasquet, C. Bernaud, C. Pintado, C. Delaugerre, L. Sagaon-Teyssier, S. Le Mestre, C. Chidiac, G. Pialoux, D. Ponscarne, J. Fonsart, D. Thompson, M.A. Wainberg, V. Doré, L. Meyer, Efficacy, safety, and effect on sexual behaviour of on-demand pre-exposure prophylaxis for HIV in men who have sex with men: an observational cohort study, *Lancet HIV* 4 (2017) e402–e410, [https://doi.org/10.1016/S2352-3018\(17\)30089-9](https://doi.org/10.1016/S2352-3018(17)30089-9).
- A.N. Nyaku, S.G. Kelly, B.O. Taiwo, Long-acting Antiretrovirals: where are we now? *Curr. HIV/AIDS Rep.* 14 (2017) 63–71, <https://doi.org/10.1007/s11904-017-0353-0>.
- R.K.R. Rajoli, D.J. Back, S. Rannard, C.F. Meyers, C. Flexner, A. Owen, M. Siccardi, In silico dose prediction for long-acting Rilpivirine and Cabotegravir administration to children and adolescents, *Clin. Pharmacokinet.* 57 (2018) 255–266, <https://doi.org/10.1007/s40262-017-0557-x>.
- T. Whitfield, A. Torkington, C. van Halsema, Profile of cabotegravir and its potential in the treatment and prevention of HIV-1 infection: evidence to date, *HIV/AIDS - Res. Palliat. Care.* 8 (2016) 157–164, <https://doi.org/10.2147/HIV.S97920>.
- C. Trezza, S.L. Ford, W. Spreen, R. Pan, S. Piscitelli, Formulation and pharmacology of long-acting cabotegravir, *Curr. Opin. HIV AIDS* 10 (2015) 239–245, <https://doi.org/10.1097/COH.0000000000000168>.
- S.L. Letendre, A. Mills, D. Hagins, S. Swindells, F. Felizarta, J. Devente, C. Bettacchi, Y. Lou, S. Ford, K. Sutton, J.S. Shaik, H. Crauwels, R. D'Amico, P. Patel, Pharmacokinetics and antiviral activity of cabotegravir and rilpivirine in cerebrospinal fluid following long-acting injectable administration in HIV-infected

- adults, *J. Antimicrob. Chemother.* 75 (2020) 648–655, <https://doi.org/10.1093/jac/dkz504>.
- [13] A.J. Paredes, I.K. Ramöller, P.E. McKenna, M.T.A. Abbate, F. Volpe-Zanutto, L. Vora, M. Kilbourne-Brook, C. Jarrhian, K. Moffatt, C. Zhang, I.A. Tekko, R. F. Donnelly, Microarray patches: breaking down the barriers to contraceptive care and HIV prevention for women across the globe, *Adv. Drug Deliv. Rev.* 173 (2021) 331–348, <https://doi.org/10.1016/j.addr.2021.04.002>.
- [14] D. Rial-crestelo, A. Pinto-martínez, F. Pulido, D. Rial-crestelo, A. Pinto-martínez, F. Pulido, Expert Review of Anti-infective Therapy Cabotegravir and rilpivirine for the treatment of HIV Cabotegravir and rilpivirine for the treatment of HIV, *Expert Rev. Anti-Infect. Ther.* 00 (2020) 1–12, <https://doi.org/10.1080/14787210.2020.1736561>.
- [15] Jennifer McLenon, Mary A.M. Rogers, The fear of needles: a systematic review and meta-analysis, *J. Adv. Nurs.* 75 (2019) 30–42, <https://doi.org/10.1111/jan.13818>.
- [16] T. Hayashi, Y.J.F. Hutin, M. Bulterys, A. Altaf, B. Allegranzi, Injection practices in 2011–2015: a review using data from the demographic and health surveys (DHS), *BMC Health Serv. Res.* 19 (2019) 1–10, <https://doi.org/10.1186/s12913-019-4366-9>.
- [17] F. Volpe-Zanutto, B. Fonseca-Santos, P.E. McKenna, A.J. Paredes, J.L. Dávila, M.T.C. McCrudden, M.M.P. Tangerina, M. Ceccheto Figueiredo, W. Vilegas, A. Brisebe, M. Akira D'vila, R.F. Donnelly, M. Chorilli, M.A. Foglio, Novel transdermal bioadhesive surfactant-based system for release and solubility improvement of antimalarial drugs artemether-lumefantrine, *Biomed. Mater.* 16 (2021), <https://doi.org/10.1088/1748-605X/ac2885>.
- [18] A.J. Paredes, F. Volpe-Zanutto, L.K. Vora, I.A. Tekko, A.D. Permana, C.J. Picco, H. O. McCarthy, R.F. Donnelly, Systemic delivery of tenofovir alafenamide using dissolving and implantable microneedle patches, *Mater. Today Bio.* 13 (2022), 100217, <https://doi.org/10.1016/j.mtbio.2022.100217>.
- [19] F.W. Hughes, G.D. Puaone, T.R. Clark, B.L. Wondwossen, T.L. Johnson, Prevalence and Predictors of Traditional Medicine utilization among persons living with AIDS (PLWA) on Antiretroviral (ARV) and Prophylaxis treatment in both rural and urban areas South Africa, *Afr. J. Tradit. Complement. Altern. Med.* 9 (2012) 470–484.
- [20] E. Sicuri, A. Vieta, L. Lindner, C. Sauboin, Economic costs of malaria in children in three Sub-Saharan countries: Ghana, Tanzania and Kenya, *Tropical Med. Int. Health* 16 (2011) 117. <http://ovidsp.ovid.com/ovidweb.cgi?T=JS&PAGE=reference&D=emed13&NEWS=N&AN=70589204>.
- [21] F.V. Zanutto, E. McAlister, M. Marucci, P. Tangerina, B. Fonseca-Santos, H. C. Salles, M.O. Souza, A. Brisebe, W. Vilegas, M. Chorilli, M. Akira D'Avila, R. F. Donnelly, M.A. Foglio, Semisynthetic derivative of *Artemisia annua*-loaded transdermal bioadhesive for the treatment of uncomplicated malaria caused by *Plasmodium falciparum* in children, *J. Pharm. Sci.* 108 (2019) 1177–1188, <https://doi.org/10.1016/j.xphs.2018.10.007>.
- [22] F. Volpe-Zanutto, L. Tiburcio, A. Dian, M. Kirky, A.J. Paredes, L.K. Vora, A. P. Bonfanti, I. Charlie-silva, C. Raposo, M.C. Figueiredo, I.M.O. Sousa, A. Brisebe, R. F. Donnelly, M. Ann, F. Trindade, Artemether and lumefantrine dissolving microneedle patches with improved pharmacokinetic performance and antimalarial efficacy in mice infected with *Plasmodium yoelii*, *J. Control. Release* 333 (2021) 298–315, <https://doi.org/10.1016/j.jconrel.2021.03.036>.
- [23] A.J. Paredes, F. Volpe-Zanutto, A. Dian Permana, A.J. Murphy, C.J. Picco, L. K. Vora, J.A. Coulter, R.F. Donnelly, Novel tip-loaded dissolving and implantable microneedle array patches for sustained release of finasteride, *Int. J. Pharm.* (2021), 120885, <https://doi.org/10.1016/j.ijpharm.2021.120885>.
- [24] A.J. Paredes, P.E. McKenna, I.K. Ramöller, Y.A. Naser, F. Volpe-Zanutto, M. Li, M. T.A. Abbate, L. Zhao, C. Zhang, J.H. Abu-Ershaid, X. Dai, R.F. Donnelly, Microarray patches: poking a hole in the challenges faced when delivering poorly soluble drugs, *Adv. Funct. Mater.* 2005792 (2020) 1–27, <https://doi.org/10.1002/adfm.202005792>.
- [25] R.F. Donnelly, T.R.R. Singh, M.J. Garland, K. Migalska, R. Majithiya, C. M. McCrudden, P.L. Kole, T.M.T. Mahmood, H.O. McCarthy, A.D. Woolfson, Hydrogel-forming microneedle arrays for enhanced transdermal drug delivery, *Adv. Funct. Mater.* 22 (2012) 4879–4890, <https://doi.org/10.1002/adfm.201200864>.
- [26] L.K. Vora, K. Moffatt, I.A. Tekko, A.J. Paredes, F. Volpe-Zanutto, D. Mishra, K. Peng, R. Raj Singh Thakur, R.F. Donnelly, Microneedle array systems for long-acting drug delivery, *Eur. J. Pharm. Biopharm.* 159 (2021) 44–76, <https://doi.org/10.1016/j.ejpb.2020.12.006>.
- [27] D. Ramadan, A.D. Permana, A.J. Courtenay, M.T.C. McCrudden, I.A. Tekko, E. McAlister, Q.K. Anjani, E. Utomo, H.O. McCarthy, R.F. Donnelly, Development, evaluation, and pharmacokinetic assessment of polymeric microarray patches for transdermal delivery of vancomycin hydrochloride, *Mol. Pharm.* 17 (2020) 3353–3368, <https://doi.org/10.1021/acs.molpharmaceut.0c00431>.
- [28] Q.K. Anjani, A.D. Permana, A. Cárcamo-Martínez, J. Domínguez-Robles, I. A. Tekko, E. Larrañeta, L.K. Vora, D. Ramadan, R.F. Donnelly, Versatility of hydrogel-forming microneedles in vitro transdermal delivery of tuberculosis drugs, *Eur. J. Pharm. Biopharm.* 158 (2021) 294–312, <https://doi.org/10.1016/j.ejpb.2020.12.003>.
- [29] A. Ryzhakov, T. Do Thi, J. Stappaerts, L. Bertolotti, K. Kimpe, A.R. Sá Couto, P. Saokham, G. Van den Mooter, P. Augustijns, G.W. Somsen, S. Kurkov, S. Inghelbrecht, A. Arien, M.I. Jimidar, K. Schrijnemakers, T. Loftsson, Self-assembly of Cyclodextrins and their complexes in aqueous solutions, *J. Pharm. Sci.* 105 (2016) 2556–2569, <https://doi.org/10.1016/j.xphs.2016.01.019>.
- [30] C.W. Yong, C. Washington, W. Smith, Structural behaviour of 2-hydroxypropyl-β-cyclodextrin in water: molecular dynamics simulation studies, *Pharm. Res.* 25 (2008) 1092–1099, <https://doi.org/10.1007/s11095-007-9506-y>.
- [31] R.F. Donnelly, M.T.C. McCrudden, A.Z. Alkilani, E. Larrañeta, E. McAlister, A. J. Courtenay, M.C. Kearney, T.R. Raj Singh, H.O. McCarthy, V.L. Kett, E. Caffarel-Salvador, S. Al-Zahrani, A.D. Woolfson, Hydrogel-forming microneedles prepared from “super swelling” polymers combined with lyophilised wafers for transdermal drug delivery, *PLoS One* 9 (2014), <https://doi.org/10.1371/journal.pone.0111547>.
- [32] A.S. Cordeiro, I.A. Tekko, M.H. Jomaa, L. Vora, E. McAlister, F. Volpe-Zanutto, M. Nethery, P.T. Baine, N. Mitchell, D.W. McNeill, R.F. Donnelly, Two-photon polymerisation 3D printing of microneedle Array templates with versatile designs: application in the development of polymeric drug delivery systems, *Pharm. Res.* 37 (2020) 174, <https://doi.org/10.1007/s11095-020-02887-9>.
- [33] K.A. Higuchi, T. Connors, Phase-solubility techniques, in: C.N. Reilly (Ed.), *Adv. Anal. Chem. Instrum.*, Wiley-Interscience, New York, NY, 1965, pp. 117–122.
- [34] I.A. Tekko, L.K. Vora, F. Volpe-Zanutto, K. Moffatt, C. Jarrhian, H.O. McCarthy, R. F. Donnelly, Novel bilayer microarray patch-assisted long-acting Micro-depot Cabotegravir intradermal delivery for HIV pre-exposure prophylaxis, *Adv. Funct. Mater.* (2021) 2106999, <https://doi.org/10.1002/adfm.202106999>.
- [35] J. Siepmann, N.A. Peppas, Higuchi equation: derivation, applications, use and misuse, *Int. J. Pharm.* 418 (2011) 6–12, <https://doi.org/10.1016/j.ijpharm.2011.03.051>.
- [36] A.D. Permana, A.J. Paredes, F. Volpe-Zanutto, Q.K. Anjani, E. Utomo, R. F. Donnelly, Dissolving microneedle-mediated dermal delivery of itraconazole nanocrystals for improved treatment of cutaneous candidiasis, *Eur. J. Pharm. Biopharm.* 154 (2020) 50–61, <https://doi.org/10.1016/j.ejpb.2020.06.025>.
- [37] A.J. Paredes, F. Volpe-Zanutto, A.D. Permana, A.J. Murphy, C.J. Picco, L.K. Vora, J. A. Coulter, R.F. Donnelly, Novel tip-loaded dissolving and implantable microneedle array patches for sustained release of finasteride, *Int. J. Pharm.* 606 (2021), 120885, <https://doi.org/10.1016/j.ijpharm.2021.120885>.
- [38] R.F. Donnelly, M.J. Garland, D.I.J. Morrow, K. Migalska, T. Raghu, R. Singh, R. Majithiya, A.D. Woolfson, Optical coherence tomography is a valuable tool in the effects of microneedle geometry on skin penetration characteristics and in-skin dissolution, *J. Control. Release* 147 (2010) 333–341.
- [39] I.K. Ramöller, M.T.A. Abbate, L.K. Vora, A.R.J. Hutton, K. Peng, F. Volpe-Zanutto, I.A. Tekko, K. Moffatt, A.J. Paredes, H.O. McCarthy, R.F. Donnelly, HPLC-MS method for simultaneous quantification of the antiretroviral agents rilpivirine and cabotegravir in rat plasma and tissues, *J. Pharm. Biomed. Anal.* 213 (2022), 114698, <https://doi.org/10.1016/j.jpba.2022.114698>.
- [40] A.D. Permana, A.J. Paredes, F.V. Zanutto, M.N. Amir, I. Ismail, M.A. Bahar, S. D. Sumarheni, R.F. Donnelly Palma, Albendazole nanocrystal-based dissolving microneedles with improved pharmacokinetic performance for enhanced treatment of cystic echinococcosis, *ACS Appl. Mater. Interfaces* 13 (2021) 38745–38760, <https://doi.org/10.1021/acsami.1c11179>.
- [41] E.Z. Loizidou, N.T. Inoue, J. Ashton-Barnett, D.A. Barrow, C.J. Allender, Evaluation of geometrical effects of microneedles on skin penetration by CT scan and finite element analysis, *Eur. J. Pharm. Biopharm.* 107 (2016) 1–6, <https://doi.org/10.1016/j.ejpb.2016.06.023>.
- [42] L.G.C. Burkhart, D. Morrell, *Dermatological Pharmacology*, in: B.C.K.L.L. Brunton, A.B. Chabner (Eds.), *Goodman Gilman's Pharmacol. Basis Ther.*, 12<sup>th</sup> ed, McGraw-Hill Medical, New York, NY, 2012.
- [43] E.M. Migdadi, A.J. Courtenay, I.A. Tekko, M.T.C. McCrudden, M. Kearney, E. McAlister, H.O. McCarthy, R.F. Donnelly, Hydrogel-forming microneedles enhance transdermal delivery of metformin hydrochloride, *J. Control. Release* 285 (2018) 142–151, <https://doi.org/10.1016/j.jconrel.2018.07.009>.
- [44] A. Adamer, Verena; Thaler, *Crystalline Forms of Cabotegravir Sodium*, WO 2018149608 (A1), 2018.
- [45] D. Han, Z. Han, L. Liu, Y. Wang, S. Xin, H. Zhang, Z. Yu, Solubility enhancement of myricetin by inclusion complexation with heptakis-(2-hydroxypropyl)-β-cyclodextrin: a joint experimental and theoretical study, *Int. J. Mol. Sci.* 21 (2020) 1–11, <https://doi.org/10.3390/ijms21030766>.
- [46] H.-J. Schneider, F. Hacket, V. Rudiger, H. Ikeda, NMR studies of Cyclodextrins and Cyclodextrin complexes, *Chem. Rev.* 2665 (1998), <https://doi.org/10.1021/cr970019t>.
- [47] D. Greatbanks, R. Pickford, Cyclodextrins as chiral complexing agents in water, and their application to optical purity measurements, *Magn. Reson. Chem.* 25 (1987) 208–215, <https://doi.org/10.1002/mrc.1260250306>.
- [48] F.P. Pons-Faudoa, A. Sizovs, N. Di Trani, J. Paez-Mayorga, G. Bruno, J. Rhudy, M. Manohar, K. Gwenden, C. Martini, C.Y.X. Chua, G. Varchi, M.A. Marzinke, A. Grattoni, 2-Hydroxypropyl-β-cyclodextrin-enhanced pharmacokinetics of cabotegravir from a nanofluidic implant for HIV pre-exposure prophylaxis, *J. Control. Release* 306 (2019) 89–96, <https://doi.org/10.1016/j.jconrel.2019.05.037>.
- [49] R. Al-Kasasbeh, A.J. Brady, A.J. Courtenay, E. Larrañeta, M.T.C. McCrudden, D. O'Kane, S. Liggett, R.F. Donnelly, Evaluation of the clinical impact of repeat application of hydrogel-forming microneedle array patches, *Drug Deliv. Transl. Res.* 10 (2020) 690–705, <https://doi.org/10.1007/s13346-020-00727-2>.
- [50] M.T.C. McCrudden, E. Larrañeta, A. Clark, C. Jarrhian, A. Rein-Weston, S. Lachau-Durand, N. Niemeijer, P. Williams, C. Haec, H.O. McCarthy, D. Zehring, R.F. Donnelly, Design, formulation and evaluation of novel dissolving microarray patches containing a long-acting rilpivirine nanosuspension, *J. Control. Release* 292 (2018) 119–129, <https://doi.org/10.1016/j.jconrel.2018.11.002>.
- [51] S. Rojekar, L.K. Vora, I.A. Tekko, F. Volpe-Zanutto, H.O. McCarthy, P.R. Vavia, R. F. Donnelly, Etravirine-loaded dissolving microneedle arrays for long-acting delivery, *Eur. J. Pharm. Biopharm.* (2021), <https://doi.org/10.1016/j.ejpb.2021.04.024>.
- [52] M.L. Murray, M. Markowitz, I. Frank, R.M. Grant, K.H. Mayer, K.J. Hudson, B. S. Stancil, S.L. Ford, P. Patel, A.R. Rinehart, W.R. Spreen, D.A. Margolis, Satisfaction and acceptability of cabotegravir long-acting injectable suspension for

- prevention of HIV: patient perspectives from the ECLAIR trial, *HIV Clin. Trials*. 19 (2018) 129–138, <https://doi.org/10.1080/15284336.2018.1511346>.
- [53] C. Orkin, K. Arasteh, M. Górgolas Hernández-Mora, V. Pokrovsky, E.T. Overton, P.-M. Girard, S. Oka, S. Walmsley, C. Bettaachi, C. Brinson, P. Philibert, J. Lombaard, M. St. H. Clair, S.L. Crauwels, P. Ford, V. Patel, R. Chounta, S. D'Amico, D. Vanveggel, A. Dorey, S. Cutrell, D.A. Griffith, P.E. Margolis, W. Williams, K. Y. Parys, W.R. Spreen Smith, Long-acting Cabotegravir and Rilpivirine after Oral induction for HIV-1 infection, *N. Engl. J. Med.* 382 (2020) 1124–1135, <https://doi.org/10.1056/NEJMoa1909512>.
- [54] J.J. Norman, J.M. Arya, M.A. McClain, P.M. Frew, M.I. Meltzer, M.R. Prausnitz, Microneedle patches: usability and acceptability for self-vaccination against influenza, *Vaccine*. 32 (2014) 1856–1862, <https://doi.org/10.1016/j.vaccine.2014.01.076>.
- [55] P. Ranjan Yadav, M. Iqbal Nasiri, L.K. Vora, E. Larrañeta, R.F. Donnelly, S. K. Pattanayek, D. Bhusan Das, Super-swelling hydrogel-forming microneedle based transdermal drug delivery: Mathematical modelling, simulation and experimental validation, *Int. J. Pharmaceutics* 622 (2022) 121835, <https://doi.org/10.1016/j.ijpharm.2022.121835>.
- [56] A. Rein-Weston, I. Tekko, L. Vora, C. Jarrahian, B. Spreen, T. Scott, R. Donnelly, D. Zehrung, LBS. Microarray Patch Delivery of Long-Acting HIV PrEP and Contraception, *Open Forum Infectious Diseases* 6 (2019) S996, <https://doi.org/10.1093/ofid/ofz415.2491>.
- [57] K. Peng, L.K. Vora, J. Domínguez-Robles, Y.A. Naser, M. Li, E. Larrañeta, R. F. Donnelly, Hydrogel-forming microneedles for rapid and efficient skin deposition of controlled release tip-implants, *Materials Sci. Eng.: C* 127 (2021) 112226, <https://doi.org/10.1016/j.msec.2021.112226>.
- [58] L.K. Vora, K. Moffatt, R.F. Donnelly, 9 - Long-lasting drug delivery systems based on microneedles. *Long-Acting Drug Delivery Systems*, Woodhead Publishing, 2022, pp. 249–287.

ORIGINALITY REPORT

---

19%  
SIMILARITY INDEX

%  
INTERNET SOURCES

19%  
PUBLICATIONS

%  
STUDENT PAPERS

---

PRIMARY SOURCES

---

- 1** Donnelly, R.F.. "Bioadhesive patch-based delivery of 5-aminolevulinic acid to the nail for photodynamic therapy of onychomycosis", *Journal of Controlled Release*, 20050321  
Publication 1%
- 2** Maelíosa T. C. Mc Crudden, Eneko Larrañeta, Annie Clark, Courtney Jarrahian et al. "Design, Formulation, and Evaluation of Novel Dissolving Microarray Patches Containing Rilpivirine for Intravaginal Delivery", *Advanced Healthcare Materials*, 2019  
Publication 1%
- 3** Satish Rojekar, Lalitkumar K. Vora, Ismaiel A. Tekko, Fabiana Volpe-Zanutto et al. "Etravirine-loaded dissolving microneedle arrays for long-acting delivery", *European Journal of Pharmaceutics and Biopharmaceutics*, 2021  
Publication 1%
- 4** Neng Qiu, Beilisha Shen, Xuebing Li, Xiaotong Zhang, Zitai Sang, Tao Yang, Lianying An, <1%

Junda Liu, Lijuan Chen, Lan Wang. "Inclusion complex of magnolol with hydroxypropyl- $\beta$ -cyclodextrin: characterization, solubility, stability and cell viability", Journal of Inclusion Phenomena and Macrocyclic Chemistry, 2016

Publication

---

5

Delly Ramadon, Andi Dian Permana, Aaron J. Courtenay, Maelíosa T. C. McCrudden et al. "Development, Evaluation, and Pharmacokinetic Assessment of Polymeric Microarray Patches for Transdermal Delivery of Vancomycin Hydrochloride", Molecular Pharmaceutics, 2020

Publication

---

<1 %

6

Ismaiel A. Tekko, Andi Dian Permana, Lalitkumar Vora, Taher Hatahet, Helen O. McCarthy, Ryan F. Donnelly. "Localised and sustained intradermal delivery of methotrexate using nanocrystal-loaded microneedle arrays: Potential for enhanced treatment of psoriasis", European Journal of Pharmaceutical Sciences, 2020

Publication

---

<1 %

7

Alejandro J. Paredes, Fabiana Volpe-Zanutto, Andi Dian Permana, Aidan J. Murphy et al. "Novel tip-loaded dissolving and implantable microneedle array patches for sustained

<1 %

release of finasteride", International Journal of  
Pharmaceutics, 2021

Publication

---

8

Ke Peng, Lalitkumar K. Vora, Ismaiel A. Tekko,  
Andi Dian Permana et al. "Dissolving  
microneedle patches loaded with  
amphotericin B microparticles for localised  
and sustained intradermal delivery: Potential  
for enhanced treatment of cutaneous fungal  
infections", Journal of Controlled Release,  
2021

Publication

---

<1 %

9

Li Wang, Shen Li, Hongda Cai, Xiaohu Liu,  
Tongtong Feng, Xiuhua Zhao. "Preparation  
And Characterisation Of  
Ecdysterone/Hydroxypropyl-B-Cyclodextrin  
Inclusion Complex With Enhanced Oral And  
Transdermal Bioavailability", Journal of  
Microencapsulation, 2022

Publication

---

<1 %

10

Qonita Kurnia Anjani, Andi Dian Permana,  
Álvaro Cárcamo-Martínez, Juan Domínguez-  
Robles et al. "Versatility of hydrogel-forming  
microneedles in in vitro transdermal delivery  
of tuberculosis drugs", European Journal of  
Pharmaceutics and Biopharmaceutics, 2021

Publication

---

<1 %

11

Muhammad Iqbal Nasiri, Lalitkumar K. Vora, Juhaina Abu Ershaid, Ke Peng, Ismaiel A. Tekko, Ryan F. Donnelly. "Nanoemulsion-based dissolving microneedle arrays for enhanced intradermal and transdermal delivery", Drug Delivery and Translational Research, 2021

Publication

&lt;1 %

12

Jef Stappaerts, Thao Do Thi, Elena Dominguez-Vega, Govert W. Somsen, Guy Van den Mooter, Patrick Augustijns. "The impact of guest compounds on cyclodextrin aggregation behavior: A series of structurally related parabens", International Journal of Pharmaceutics, 2017

Publication

&lt;1 %

13

Ebru Altuntaş, Ismaiel A. Tekko, Lalitkumar K. Vora, Narender Kumar et al. "Nestorone nanosuspension-loaded dissolving microneedles array patch: A promising novel approach for "on-demand" hormonal female-controlled peritcoital contraception", International Journal of Pharmaceutics, 2021

Publication

&lt;1 %

14

Andi Dian Permana, Ismaiel A. Tekko, Maelíosa T.C. McCrudden, Qonita Kurnia Anjani et al. "Solid lipid nanoparticle-based dissolving microneedles: A promising

&lt;1 %

intradermal lymph targeting drug delivery system with potential for enhanced treatment of lymphatic filariasis", Journal of Controlled Release, 2019

Publication

---

15

Ismaiel A. Tekko, Gaoyun Chen, Juan Domínguez-Robles, Raghu Raj Singh Thakur et al. "Development and characterisation of novel poly (vinyl alcohol)/poly (vinyl pyrrolidone)-based hydrogel-forming microneedle arrays for enhanced and sustained transdermal delivery of methotrexate", International Journal of Pharmaceutics, 2020

Publication

---

<1 %

16

Lalit K. Vora, Kurtis Moffatt, Ismaiel A. Tekko, Alejandro J. Paredes et al. "Microneedle array systems for long-acting drug delivery", European Journal of Pharmaceutics and Biopharmaceutics, 2021

Publication

---

<1 %

17

Ana Sara Cordeiro, Ismaiel A. Tekko, Mohamed H. Jomaa, Lalitkumar Vora et al. "Two-Photon Polymerisation 3D Printing of Microneedle Array Templates with Versatile Designs: Application in the Development of Polymeric Drug Delivery Systems", Pharmaceutical Research, 2020

Publication

---

<1 %

18

Eneko Larrañeta, Sarah Stewart, Steven J. Fallows, Lena L. Birkhäuser et al. "A facile system to evaluate in vitro drug release from dissolving microneedle arrays", *International Journal of Pharmaceutics*, 2016

Publication

&lt;1 %

19

Qonita Kurnia Anjani, Akmal Hidayat Bin Sabri, Natalia Moreno-Castellanos, Emilia Utomo et al. "Soluplus®-based dissolving microarray patches loaded with colchicine: towards a minimally invasive treatment and management of gout", *Biomaterials Science*, 2022

Publication

&lt;1 %

20

Heba Abd-El-Azim, Ismaiel A. Tekko, Ahlam Ali, Alyaa Ramadan et al. "Hollow microneedle assisted intradermal delivery of hypericin lipid nanocapsules with light enabled photodynamic therapy against skin cancer", *Journal of Controlled Release*, 2022

Publication

&lt;1 %

21

Zhaojun Xu, Junxia Liu. "Research on Temperature Control of Liposome High Pressure Homogenizer Based on Genetic Algorithm Optimization PID", *Procedia Computer Science*, 2022

Publication

&lt;1 %

22

Cindy Kristina Enggi, Hansel Tridatmojo Isa, Sulistiawati Sulistiawati, Komang Agus Rai Ardika et al. "Development of thermosensitive and mucoadhesive gels of cabotegravir for enhanced permeation and retention profiles in vaginal tissue: A proof of concept study", *International Journal of Pharmaceutics*, 2021

Publication

&lt;1 %

23

Emma McAlister, Bridie Dutton, Lalitkumar K. Vora, Li Zhao et al. "Directly Compressed Tablets: A Novel Drug - Containing Reservoir Combined with Hydrogel - Forming Microneedle Arrays for Transdermal Drug Delivery", *Advanced Healthcare Materials*, 2020

Publication

&lt;1 %

24

Irina V. Terekhova, Tatyana V. Volkova, German L. Perlovich. "Experimental Analysis of Complex Formation of Niflumic Acid with  $\beta$ -Cyclodextrins", *Journal of Inclusion Phenomena and Macrocyclic Chemistry*, 2006

Publication

&lt;1 %

25

Kellici, Tahsin F, Dimitrios Ntountaniotis, Georgios Leonis, Maria Chatziathanasiadou, Alexandra V. Chatzikonstantinou, Johanna Becker-Baldus, Clemens Glaubitz, Andreas G. Tzacos, Kyriakos Viras, Petros Chatzigeorgiou, Stavros Tzimas, Evangelia Kefala, Georgia

&lt;1 %

Valsami, Helen Archontaki, Manthos G. Papadopoulos, and Thomas Mavromoustakos. "Investigation of the interactions of silibinin with 2-hydroxypropyl- $\beta$ -cyclodextrin through biophysical techniques and computational methods", *Molecular Pharmaceutics*, 2015.

Publication

---

26

Andi Dian Permana, Alejandro J. Paredes, Fabiana Volpe-Zanutto, Qonita Kurnia Anjani, Emilia Utomo, Ryan F. Donnelly. "Dissolving microneedle-mediated dermal delivery of itraconazole nanocrystals for improved treatment of cutaneous candidiasis", *European Journal of Pharmaceutics and Biopharmaceutics*, 2020

Publication

---

27

Andrew W. Brooks, Kevin D. Kohl, Robert M. Brucker, Edward J. van Opstal, Seth R. Bordenstein. "Phylosymbiosis: Relationships and Functional Effects of Microbial Communities across Host Evolutionary History", *PLOS Biology*, 2016

Publication

---

28

Madan, Jitender, Bharat Baruah, Mulpuri Nagaraju, Mohamed O. Abdalla, Clayton Yates, Timothy Turner, Vijay Rangari, Donald Hamelberg, and Ritu Aneja. "Molecular

<1 %

<1 %

<1 %

Cycloencapsulation Augments Solubility and Improves Therapeutic Index of Brominated Noscipine in Prostate Cancer Cells", Molecular Pharmaceutics, 2012.

Publication

---

29

Beltran, A.. "Synthesis and application of a carbamazepine-imprinted polymer for solid-phase extraction from urine and wastewater", Analytica Chimica Acta, 20070730

Publication

---

30

Prajapati Jagruti B., Sawant Krutika K., Bhramanand Dubey. "Oral Bioavailability Enhancement of Amisulpride: Complexation and its Pharmacokinetics and Pharmacodynamics Evaluations", Drug Metabolism Letters, 2020

Publication

---

31

María Beatrice Bianchi, Chunyang Zhang, Elise Catlin, Giuseppina Sandri et al. "Bioadhesive eutectogels supporting drug nanocrystals for long-acting delivery to mucosal tissues", Materials Today Bio, 2022

Publication

---

32

Yapeng Lu, Siyuan Liu, Yu Zhao, Li Zhu, Shuqin Yu. "Complexation of Z-ligustilide with hydroxypropyl- $\beta$ -cyclodextrin to improve stability and oral bioavailability", Acta Pharmaceutica, 2014

Publication

<1 %

<1 %

<1 %

<1 %

---

33 Andi Dian Permana, Alejandro J. Paredes, Fabiana Volpe Zanutto, Muh. Nur Amir et al. "Albendazole Nanocrystal-Based Dissolving Microneedles with Improved Pharmacokinetic Performance for Enhanced Treatment of Cystic Echinococcosis", ACS Applied Materials & Interfaces, 2021

<1 %

Publication

---

34 Meina Liu, Wen Cao, Yinghua Sun, Zhonggui He. "Preparation, characterization and in vivo evaluation of formulation of repaglinide with hydroxypropyl- $\beta$ -cyclodextrin", International Journal of Pharmaceutics, 2014

<1 %

Publication

---

35 Proceedings of the Ninth International Symposium on Cyclodextrins, 1999.

<1 %

Publication

---

36 "Corrigendum to "Books received" [J. Control. Release. Volume 126 (2008) 281-282]", Journal of Controlled Release, 20090319

<1 %

Publication

---

37 Ruiping Kong, Xingyi Zhu, Elizaveta S. Meteleva, Alexander V. Dushkin, Weike Su. "Physicochemical characteristics of the complexes of simvastatin and atorvastatin calcium with hydroxypropyl-  $\beta$  -cyclodextrin produced by mechanochemical activation", Journal of Drug Delivery Science and

<1 %

38

Nathir A.F. Al-Rawashdeh, Khaled S. Al-Sadeh, Mohammad-Bassam Al-Bitar.

"Physicochemical study on microencapsulation of hydroxypropyl- $\beta$ -cyclodextrin in dermal preparations", Drug Development and Industrial Pharmacy, 2010

Publication

---

<1 %

39

Nutsarun Wangsawangrung, Chasuda Choipang, Sonthaya Chairwut, Pongpol Ekabutr et al. "Quercetin/Hydroxypropyl- $\beta$ -Cyclodextrin Inclusion Complex-Loaded Hydrogels for Accelerated Wound Healing", Gels, 2022

Publication

---

<1 %

40

Shengfeng Wang, Yanfei Ding, Yao Yao. "Inclusion complexes of fluorofenidone with  $\beta$ -cyclodextrin and hydroxypropyl- $\beta$ -cyclodextrin", Drug Development and Industrial Pharmacy, 2009

Publication

---

<1 %

41

Shoulian Wei, Jianwen Li, Yong Liu. "Colourimetric assay for  $\beta$ -estradiol based on the peroxidase-like activity of Fe O @mSiO @HP- $\beta$ -CD nanoparticles ", RSC Advances, 2015

Publication

---

<1 %

42

Álvaro Cárcamo-Martínez, Brónach Mallon, Qonita Kurnia Anjani, Juan Domínguez-Robles et al. "Enhancing intradermal delivery of tofacitinib citrate: Comparison between powder-loaded hollow microneedle arrays and dissolving microneedle arrays", International Journal of Pharmaceutics, 2020

Publication

<1 %

43

Akmal Hidayat Bin Sabri, Qonita Kurnia Anjani, Ryan F. Donnelly. "Synthesis and characterization of sorbitol laced hydrogel-forming microneedles for therapeutic drug monitoring", International Journal of Pharmaceutics, 2021

Publication

<1 %

44

Alejandro J. Paredes, Peter E. McKenna, Inken K. Ramöller, Yara A. Naser et al. "Microarray Patches: Poking a Hole in the Challenges Faced When Delivering Poorly Soluble Drugs", Advanced Functional Materials, 2020

Publication

<1 %

45

Emma McAlister, Mary-Carmel Kearney, E. Linzi Martin, Ryan F. Donnelly. "From the laboratory to the end-user: a primary packaging study for microneedle patches containing amoxicillin sodium", Drug Delivery and Translational Research, 2021

Publication

<1 %

46

Milena Giulia Gonçalves. "História natural da infecção por variantes genéticas de HPV-16 no canal anal de homens participantes do estudo internacional multicêntrico HIM (HPV Infection in Men)", Universidade de Sao Paulo, Agencia USP de Gestao da Informacao Academica (AGUIA), 2022

Publication

&lt;1 %

47

Mingshan Li, Lalitkumar K. Vora, Ke Peng, Ryan F. Donnelly. "Trilayer microneedle array assisted transdermal and intradermal delivery of dexamethasone", International Journal of Pharmaceutics, 2021

Publication

&lt;1 %

48

Patricia González-Vázquez, Eneko Larrañeta, Maelíosa T.C. McCrudden, Courtney Jarrahan et al. "Transdermal delivery of gentamicin using dissolving microneedle arrays for potential treatment of neonatal sepsis", Journal of Controlled Release, 2017

Publication

&lt;1 %

49

V. Concha - Herrera, J. R. Torres - Lapasió, G. Vivó - Truyols, M. C. García - Álvarez - Coque. "Separation of Proteic Primary Amino Acids under Several Reversed - Phase Liquid Chromatographic Conditions", Journal of Liquid Chromatography & Related Technologies, 2011

&lt;1 %

50

Victor David, Andrei Medvedovici, Florin Albu. "Retention Behavior of Metformin and Related Impurities in Ion - Pairing Liquid Chromatography", Journal of Liquid Chromatography & Related Technologies, 2007

Publication

---

<1 %

51

Zhiqiang Yan, Wen Xu, Jin Sun, Xiaohong Liu, Yanping Zhao, Yinghua Sun, Tianhong Zhang, Zhonggui He. "Characterization and In Vivo Evaluation of an Inclusion Complex of Oridonin and 2-hydroxypropyl- $\beta$ -cyclodextrin", Drug Development and Industrial Pharmacy, 2008

Publication

---

<1 %

52

Chirag Desai, Bala Prabhakar. "Nano-amorphous composites of cilostazol-HP- $\beta$ -CD inclusion complexes: physicochemical characterization, structure elucidation, thermodynamic studies and in vitro evaluation", Journal of Inclusion Phenomena and Macrocyclic Chemistry, 2014

Publication

---

<1 %

53

Delly Ramadan, Aaron J. Courtenay, Andi Dian Permana, Ismaiel A. Tekko et al. "A sensitive HPLC-UV method for quantifying vancomycin in biological matrices: Application to

<1 %

pharmacokinetic and biodistribution studies in rat plasma, skin and lymph nodes", Journal of Pharmaceutical and Biomedical Analysis, 2020

Publication

---

54

Donnelly, Ryan F., Desmond I. J. Morrow, Maelíosa T. C. McCrudden, Ahlam Zaid Alkilani, Eva M. Vicente-Pérez, Conor O'Mahony, Patricia González-Vázquez, Paul A. McCarron, and A. David Woolfson. "Hydrogel-Forming and Dissolving Microneedles for Enhanced Delivery of Photosensitizers and Precursors", Photochemistry and Photobiology, 2013.

Publication

---

<1 %

55

Fernanda P. Pons-Faudoa, Antons Sizovs, Nicola Di Trani, Jesus Paez-Mayorga et al. "2-Hydroxypropyl- $\beta$ -cyclodextrin-enhanced pharmacokinetics of cabotegravir from a nanofluidic implant for HIV pre-exposure prophylaxis", Journal of Controlled Release, 2019

Publication

---

<1 %

56

Xianghe Meng, Qiuyue Pan, Yun Liu. "Preparation and properties of phytosterols with hydroxypropyl  $\beta$ -cyclodextrin inclusion complexes", European Food Research and Technology, 2012

Publication

---

<1 %

57

Dongxu Han, Zhongbao Han, Liyan Liu, Ying Wang, Shigang Xin, Hongbo Zhang, Zhan Yu. "Solubility Enhancement of Myricetin by Inclusion Complexation with Heptakis-O-(2-Hydroxypropyl)- $\beta$ -Cyclodextrin: A Joint Experimental and Theoretical Study", International Journal of Molecular Sciences, 2020

Publication

---

<1 %

58

Jef Stappaerts, Patrick Augustijns. "Displacement of itraconazole from cyclodextrin complexes in biorelevant media: In vitro evaluation of supersaturation and precipitation behavior", International Journal of Pharmaceutics, 2016

Publication

---

<1 %

59

Omar Sued, Tomás M. Grosso. "Pathophysiology of HIV and strategies to eliminate AIDS as a public health threat", Elsevier BV, 2023

Publication

---

<1 %

60

Yu Wu, Lalitkumar K. Vora, Deepakkumar Mishra, Muhammad Faris Adrianto et al. "Nanosuspension-loaded dissolving bilayer microneedles for hydrophobic drug delivery to the posterior segment of the eye", Biomaterials Advances, 2022

Publication

---

<1 %

61

"Supramolecules in Drug Discovery and Drug Delivery", Springer Science and Business Media LLC, 2021

Publication

---

<1 %

62

Alyaa A. Albadr, Ismaiel A. Tekko, Lalitkumar K. Vora, Ahlam A. Ali, Garry Laverty, Ryan F. Donnelly, Raghu Raj Thakur. "Rapidly Dissolving Microneedle Patch of Amphotericin-B for Intracorneal Fungal Infections.", Research Square Platform LLC, 2021

Publication

---

<1 %

63

Benjámín Gyarmati, Gergő Dargó, Barnabás Áron Szilágyi, Anna Vincze et al. "Synthesis, complex formation and corneal permeation of cyclodextrin-modified, thiolated poly(aspartic acid) as self-gelling formulation of dexamethasone", European Journal of Pharmaceutics and Biopharmaceutics, 2022

Publication

---

<1 %

64

Jin-Ki Kim. "Thermo-reversible flurbiprofen liquid suppository with HP- $\beta$ -CD as a solubility enhancer: improvement of rectal bioavailability", Journal of Inclusion Phenomena and Macrocyclic Chemistry, 03/13/2009

Publication

---

<1 %

65

Martin Holt, Toby Lea, Limin Mao, Johann Kolstee et al. "Community-level changes in condom use and uptake of HIV pre-exposure prophylaxis by gay and bisexual men in Melbourne and Sydney, Australia: results of repeated behavioural surveillance in 2013–17", *The Lancet HIV*, 2018

Publication

&lt;1 %

66

Nathan Smith, Aditya N. Bade, Dhruvkumar Soni, Nagsen Gautam et al. "A long acting nanoformulated lamivudine ProTide", *Biomaterials*, 2019

Publication

&lt;1 %

67

Da-Wei Wang, Can-Bin Ouyang, Qi Liu, Hao-Liang Yuan, Xiao-Hui Liu. "Inclusion of quineestrol and 2,6-di-O-methyl- $\beta$ -cyclodextrin: Preparation, characterization, and inclusion mode", *Carbohydrate Polymers*, 2013

Publication

&lt;1 %

68

Eneko Larrañeta, Laura Barturen, Michael Ervine, Ryan F. Donnelly. "Hydrogels based on poly(methyl vinyl ether-co-maleic acid) and Tween 85 for sustained delivery of hydrophobic drugs", *International Journal of Pharmaceutics*, 2018

Publication

&lt;1 %

69

Juma Adinan, Bridgit Adamou, Caroline Amour, Aisa Shayo, Paulo Lino Kidayi, Levina

&lt;1 %

Msuya. "Feasibility of home-based HIV counselling and testing and linking to HIV services among women delivering at home in Geita, Tanzania: A household longitudinal survey", Research Square Platform LLC, 2019

Publication

---

70

Karla Helena-Bueno, Chinenye L. Ekemezie, Charlotte R. Brown, Arnaud Baslé, James N. Blaza, Chris H. Hill, Sergey V. Melnikov.

"Structure of dormant bacterial ribosomes reveals an elusive translation factor with evolutionary connections to eukaryotic protein synthesis factors", Cold Spring Harbor Laboratory, 2022

Publication

---

<1 %

71

L. Harivardhan Reddy, K. Vivek, Nishant Bakshi, R.S.R. Murthy. "Tamoxifen Citrate Loaded Solid Lipid Nanoparticles (SLN™): Preparation, Characterization, In Vitro Drug Release, and Pharmacokinetic Evaluation", Pharmaceutical Development and Technology, 2008

Publication

---

<1 %

72

Maelíosa T.C. Mc Crudden, Eneko Larrañeta, Annie Clark, Courtney Jarrahan et al. "Design, formulation and evaluation of novel dissolving microarray patches containing a long-acting

<1 %

rilpivirine nanosuspension", Journal of  
Controlled Release, 2018

Publication

---

73

Mark A Marzinke, Beatriz Grinsztejn, Jessica M Fogel, Estelle Piwowar-Manning et al.

"Characterization of Human Immunodeficiency Virus (HIV) Infection in Cisgender Men and Transgender Women Who Have Sex With Men Receiving Injectable Cabotegravir for HIV Prevention: HPTN 083",  
The Journal of Infectious Diseases, 2021

Publication

---

74

Siepmann, J., and F. Siepmann. "Mathematical modeling of drug dissolution", International Journal of Pharmaceutics, 2013.

Publication

---

75

Aaron J. Courtenay, Maelíosa T.C. McCrudden, Kathryn J. McAvoy, Helen O. McCarthy, Ryan F. Donnelly. "Microneedle Mediated Transdermal Delivery of Bevacizumab",  
Molecular Pharmaceutics, 2018

Publication

---

76

Akmal Hidayat Bin Sabri, Qonita Kurnia Anjani, Emilia Utomo, Anastasia Ripolin, Ryan F. Donnelly. "Development and characterization of a dry reservoir-hydrogel-forming microneedles composite for minimally

<1 %

<1 %

<1 %

<1 %

invasive delivery of cefazolin", International Journal of Pharmaceutics, 2022

Publication

---

77

Andi Dian Permana, Maria Mir, Emilia Utomo, Ryan F. Donnelly. "Bacterially sensitive nanoparticle-based dissolving microneedles of doxycycline for enhanced treatment of bacterial biofilm skin infection: A proof of concept study", International Journal of Pharmaceutics: X, 2020

Publication

---

<1 %

78

B. Malaekheh-Nikouei. "Enhancement of cyclosporine aqueous solubility using  $\alpha$ - and hydroxypropyl  $\beta$ -cyclodextrin mixtures", Journal of Inclusion Phenomena and Macrocyclic Chemistry, 11/07/2007

Publication

---

<1 %

79

Benoît S. Assogba, Pascal Milesi, Luc S. Djogbénou, Arnaud Berthomieu, Patrick Makoundou, Lamine S. Baba-Moussa, Anna-Sophie Fiston-Lavier, Khalid Belkhir, Pierrick Labbé, Mylène Weill. "The ace-1 Locus Is Amplified in All Resistant Anopheles gambiae Mosquitoes: Fitness Consequences of Homogeneous and Heterogeneous Duplications", PLOS Biology, 2016

Publication

---

<1 %

80

Boyd, Peter, Susan M. Fetherston, Clare F. McCoy, Ian Major, Diarmaid J. Murphy, Sandeep Kumar, Jonathon Holt, Andrew Brimer, Wendy Blanda, Brid Devlin, and R. Karl Malcolm. "Matrix and reservoir-type multipurpose vaginal rings for controlled release of dapivirine and levonorgestrel", *International Journal of Pharmaceutics*, 2016.

Publication

&lt;1 %

81

Essyrose Mathew, Giulia Pitzanti, Ana L. Gomes dos Santos, Dimitrios A. Lamprou. "Optimization of Printing Parameters for Digital Light Processing 3D Printing of Hollow Microneedle Arrays", *Pharmaceutics*, 2021

Publication

&lt;1 %

82

Ester Caffarel-Salvador, Aaron J. Brady, Eyman Eltayib, Teng Meng, Ana Alonso-Vicente, Patricia Gonzalez-Vazquez, Barbara M. Torrisi, Eva Maria Vicente-Perez, Karen Mooney, David S. Jones, Steven E. J. Bell, Colin P. McCoy, Helen O. McCarthy, James C. McElnay, Ryan F. Donnelly. "Hydrogel-Forming Microneedle Arrays Allow Detection of Drugs and Glucose In Vivo: Potential for Use in Diagnosis and Therapeutic Drug Monitoring", *PLOS ONE*, 2015

Publication

&lt;1 %

83

Laura Benítez-Gutiérrez, Vicente Soriano, Silvia Requena, Ana Arias, Pablo Barreiro, Carmen de Mendoza. "Treatment and prevention of HIV infection with long-acting antiretrovirals", Expert Review of Clinical Pharmacology, 2018

Publication

&lt;1 %

84

Maali Saad Mokhtar, FakhrEldin O. Suliman, Abdalla A. Elbashir. "Experimental and molecular modeling investigations of inclusion complexes of imazapyr with 2-hydroxypropyl( $\beta/\gamma$ ) cyclodextrin", Journal of Molecular Liquids, 2018

Publication

&lt;1 %

85

Mary-Carmel Kearney, Ester Caffarel-Salvador, Steven J. Fallows, Helen O. McCarthy, Ryan F. Donnelly. "Microneedle-mediated delivery of donepezil: Potential for improved treatment options in Alzheimer's disease", European Journal of Pharmaceutics and Biopharmaceutics, 2016

Publication

&lt;1 %

86

Qonita Kurnia Anjani, Akmal Hidayat Bin Sabri, Juan Domínguez-Robles, Natalia Moreno-Castellanos et al. "Metronidazole nanosuspension loaded dissolving microarray patches: An engineered composite pharmaceutical system for the treatment of

&lt;1 %

skin and soft tissue infection", Biomaterials  
Advances, 2022

Publication

---

87

Raghu Raj Singh Thakur, Ismaiel A. Tekko, Farhan Al-Shammari, Ahlam A. Ali, Helen McCarthy, Ryan F. Donnelly. "Rapidly dissolving polymeric microneedles for minimally invasive intraocular drug delivery", Drug Delivery and Translational Research, 2016

Publication

---

<1 %

88

Rui Tian, Hui Wang, Yi Xiao, Pengyue Hu, Ruofei Du, Xiufeng Shi, Zhengtao Wang, Yan Xie. "Fabrication of Nanosuspensions to Improve the Oral Bioavailability of Total Flavones from Hippophae rhamnoides L. and their Comparison with an Inclusion Complex", AAPS PharmSciTech, 2020

Publication

---

<1 %

89

Usanee Detamornrat, Emma McAlister, Aaron R. J. Hutton, Eneko Larrañeta, Ryan F. Donnelly. "The Role of 3D Printing Technology in Microengineering of Microneedles", Small, 2022

Publication

---

<1 %

90

Anne Melhuish, Penny Lewthwaite. "Natural history of HIV and AIDS", Medicine, 2022

Publication

---

<1 %

91 Burcin Yavuz, Jessica L. Morgan, Laura Showalter, Katti R. Horng et al. "Pharmaceutical Approaches to HIV Treatment and Prevention", *Advanced Therapeutics*, 2018  
Publication

---

92 Craig R MacKenzie, J.Paul Fawcett, David W Boulton, Ian G Tucker. "Formulation and evaluation of a propanidid hydroxypropyl- $\beta$ -cyclodextrin solution for intravenous anaesthesia", *International Journal of Pharmaceutics*, 1997  
Publication

---

93 Gilbert Lazarus, Vincent Kharisma Wangsaputra, Christianto, Melva Louisa, Vivian Soetikno, Raph L. Hamers. "Safety and Pharmacokinetic Profiles of Long-Acting Injectable Antiretroviral Drugs for HIV-1 Pre-Exposure Prophylaxis: A Systematic Review and Meta-analysis of Randomized Trials", *Frontiers in Pharmacology*, 2021  
Publication

---

94 Jean-Michel Molina, Isabelle Charreau, Bruno Spire, Laurent Cotte et al. "Efficacy, safety, and effect on sexual behaviour of on-demand pre-exposure prophylaxis for HIV in men who have sex with men: an observational cohort study", *The Lancet HIV*, 2017

95

Jian Li, Yifei Huang, Lei Huang, Lei Ye, Zhou Zhou, Guangya Xiang, Li Xu. "Determination of imatinib mesylate and related compounds by field amplified sample stacking with large volume sample injection capillary electrophoresis", Journal of Pharmaceutical and Biomedical Analysis, 2012

Publication

---

<1 %

96

Jos I Pérez-Martínez, Juan M Gínez, Esmeralda Morillo, María L González-Rodríguez, José R Moyano Méndez. "Improvement of the desorption of the pesticide 2,4-D via complexation with HP- $\beta$ -cyclodextrin", Pest Management Science, 2000

Publication

---

<1 %

97

Junior, Osvaldo Valarini, João Henrique Dantas, Carlos Eduardo Barão, Everton Fernando Zanoelo, Lúcio Cardozo-Filho, and Flávio Faria de Moraes. "Formation of inclusion compounds of (+)catechin with  $\beta$ -cyclodextrin in different complexation media: Spectral, thermal and antioxidant properties", The Journal of Supercritical Fluids, 2017.

Publication

---

<1 %

98

Kloprogge, J.T.. "Spectroscopic studies of synthetic and natural beidellites: A review",

<1 %

99

Koji Yamaguchi, Tetsuya Mitsui, Yoshinori Aso, Kenji Sugibayashi. "Analysis of in vitro skin permeation of 22-oxacalcitriol from ointments based on a two- or three-layer diffusion model considering diffusivity in a vehicle", International Journal of Pharmaceutics, 2007

Publication

---

<1 %

100

Kurtis Moffatt, Ismaiel A. Tekko, Lalitkumar Vora, Fabiana Volpe-Zanutto et al. "Development and Evaluation of Dissolving Microarray Patches for Co-administered and Repeated Intradermal Delivery of Long-acting Rilpivirine and Cabotegravir Nanosuspensions for Paediatric HIV Antiretroviral Therapy", Pharmaceutical Research, 2022

Publication

---

<1 %

101

Laura Ferreira, Joana Campos, Francisco Veiga, Catarina Cardoso, Ana Cláudia Paiva-Santos. "Cyclodextrin-based delivery systems in parenteral formulations: A critical update review", European Journal of Pharmaceutics and Biopharmaceutics, 2022

Publication

---

<1 %

102

Lidija Kovač, Zdenko Časar. "A literature review of the patent application publications on cabotegravir – an HIV integrase strand

<1 %

transfer inhibitor", Expert Opinion on  
Therapeutic Patents, 2020

Publication

---

- 103 Linying Li, Gregory J. Gatto, Rhonda M. Brand, Sai Archana Krovi et al. "Long-acting biodegradable implant for sustained delivery of antiretroviral (ARV) and hormones", Journal of Controlled Release, 2021 <1 %
- Publication
- 

- 104 Mahmoud M. Al Omari, Mohammad B. Zughul, J. Eric D. Davies, Adnan A. Badwan. "Effect of Buffer Species on the Complexation of Basic Drug Terfenadine with  $\beta$ -Cyclodextrin", Journal of Inclusion Phenomena and Macrocyclic Chemistry, 2006 <1 %
- Publication
- 

- 105 Makoto Anraku, Ryo Tabuchi, Miwa Goto, Daisuke Iohara et al. "Design and Evaluation of An Extended-Release Olmesartan Tablet Using Chitosan/Cyclodextrin Composites", Pharmaceutics, 2019 <1 %
- Publication
- 

- 106 Maurizio Ventre, Valerio Coppola, Maria Iannone, Paolo A. Netti et al. "Nanotechnologies for tissue engineering and regeneration", Elsevier BV, 2018 <1 %
- Publication
-

107 Mayuri Dalvi, Pratik Kharat, Pradip Thakor, Valamla Bhavana, Shashi Bala Singh, Neelesh Kumar Mehra. "Panorama of dissolving microneedles for transdermal drug delivery", Life Sciences, 2021

Publication

<1 %

---

108 Păduraru, Oana Maria, Andreea Bosînceanu, Gladiola Țântaru, and Cornelia Vasile. "Effect of Hydroxypropyl- $\beta$ -Cyclodextrin on the Solubility of an Antiarrhythmic Agent", Industrial & Engineering Chemistry Research, 2013.

Publication

<1 %

---

109 Qonita Kurnia Anjani, Akmal Hidayat Bin Sabri, Emilia Utomo, Juan Domínguez-Robles, Ryan F. Donnelly. "Elucidating the Impact of Surfactants on the Performance of Dissolving Microneedle Array Patches", Molecular Pharmaceutics, 2022

Publication

<1 %

---

110 Subhraseema Das, Usharani Subuddhi. "Controlled Delivery of Ibuprofen from Poly(vinyl alcohol)-Poly(ethylene glycol) Interpenetrating Polymeric Network Hydrogels", Journal of Pharmaceutical Analysis, 2018

Publication

<1 %

---

111 Teng Meng Sheng, Palanirajan Vijayaraj Kumar. "A New Approach for  $\beta$ -cyclodextrin Conjugated Drug Delivery System in Cancer Therapy", Current Drug Delivery, 2022

Publication

<1 %

---

112 Trapani, G., A. Cutrignelli, A. Latrofa, M. Franco, M. Serra, M. G. Pisu, G. Biggio, and G. Liso. "Valproic Acid - Hydrophilic Cyclodextrin Complexes and Valproic Acid - Solid Dispersions: Evaluation of Their Potential Pharmaceutical Use", Drug Development and Industrial Pharmacy, 2004.

Publication

<1 %

---

113 Vuyolwethu Khwaza, Buhle Buyana, Xhamla Nqoro, Sijongesonke Peter et al. "Strategies for delivery of antiviral agents", Elsevier BV, 2023

Publication

<1 %

---

114 Wang, Fen, Bo Yang, Yulin Zhao, Xiali Liao, Chuanzhu Gao, Ruijian Jiang, Bin Han, Jian Yang, Man Liu, and Rongguang Zhou. "Host-guest inclusion system of scutellarein with 2-hydroxypropyl-beta-cyclodextrin: preparation, characterization, and anticancer activity", Journal of Biomaterials Science Polymer Edition, 2014.

Publication

<1 %

---

115 Yali Huang, Honghong Yu, Shiling Lu, Liangjun Zou, Zonggui Tang, Tao Zeng, Juan Tang. "Effect and mechanism of ferulic acid inclusion complexes on tyramine production by *Enterobacter hormaechei* MW386398 in smoked horsemeat sausages", *Food Bioscience*, 2021  
Publication

---

116 Bibby, D.C.. "Mechanisms by which cyclodextrins modify drug release from polymeric drug delivery systems", *International Journal of Pharmaceutics*, 20000320  
Publication

---

117 Donnelly, R.F.. "Influence of solution viscosity and injection protocol on distribution patterns of jet injectors: Application to photodynamic tumour targeting", *Journal of Photochemistry & Photobiology, B: Biology*, 20071214  
Publication

---

118 Jianping Deng. "Photo-induced polymerization of methyl methacrylate/cyclodextrin complex in aqueous solution", *Polymers for Advanced Technologies*, 2008  
Publication

---

119 Ta Chen Chen, Song-Cu Yu, Chin-Mu Hsu, Fuu-Jen Tsai, Yuhsin Tsai. "Minoxidil-2-hydroxypropyl- $\beta$ -cyclodextrin inclusion

complexes: characterization and in vivo evaluation of an aqueous solution for hair growth in rats", Journal of Inclusion Phenomena and Macrocyclic Chemistry, 2017

Publication

---

120

Ebru BAŞARAN, A. Alper ÖZTÜRK, Behiye ŞENEL, Müzeyyen DEMİREL, Şenay SARICA. "Quercetin, Rutin And Quercetin-Rutin Incorporated Hydroxypropyl  $\beta$ -Cyclodextrin Inclusion Complexes", European Journal of Pharmaceutical Sciences, 2022

Publication

---

<1 %

121

Sharif Abdelghany, Ismaiel A. Tekko, Lalitkumar Vora, Eneko Larrañeta, Andi Dian Permana, Ryan F. Donnelly. "Nanosuspension-Based Dissolving Microneedle Arrays for Intradermal Delivery of Curcumin", Pharmaceutics, 2019

Publication

---

<1 %

Exclude quotes  On

Exclude matches  < 4 words

Exclude bibliography  On