

PAPER • OPEN ACCESS

The digestive and physiological visceral organs of male Bali cattle were fed with cocoa bean shell

To cite this article: H Hikmah *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **492** 012063

View the [article online](#) for updates and enhancements.

You may also like

- [A unified perspective of complex band structure: interpretations, formulations, and applications](#)
Matthew G Reuter
- [Fundamentals of the gut for capsule engineers](#)
Lavinia Barducci, Joseph C Norton, Sunandita Sarker et al.
- [Accurate potential energy curve and spectroscopic properties of \$S_2\(b^1_g^+\)\$ via extrapolation to the complete basis set limit](#)
Lu-Lu Zhang, Jing Zhang, Qing-Tian Meng et al.



The Electrochemical Society
Advancing solid state & electrochemical science & technology

243rd ECS Meeting with SOFC-XVIII

Boston, MA • May 28 – June 2, 2023

**Abstract Submission Extended
Deadline: December 16**

[Learn more and submit!](#)

The digestive and physiological visceral organs of male Bali cattle were fed with cocoa bean shell

H Hikmah¹, G Alam², J A Syamsu¹, S Salengke³ and N Nahariah¹

¹Faculty of Animal Husbandry, Universitas Hasanuddin, 90245, Indonesia

²Faculty of Pharmacy, Universitas Hasanuddin, 90245, Indonesia

³Faculty of Agriculture, Universitas Hasanuddin, 90245, Indonesia

E-mail: hikmah2unhas@gmail.com, hikmah@unhas.ac.id

Abstract. Cocoa bean shell (CBS) is a waste from cocoa bean processing, and recently used as supplements in ruminants feed. However, a CBS contains theobromines that well known for its anti-nutrient effect in the animal digestive system. This study examined the effect of CBS supplementation on some digestive organs in Bali cattle bulls. The total of 12 Bali bulls (average initial body weight, 159.25 ± 15.68 kg) aged 16 months age were fed with four levels of CBS concentrate in the diet for 16 weeks (0, 3, 6, and 9%, respectively). At the end of the experiment, digestive tract and some visceral organs weight (e.g., rumen, digest tracts, lymph, liver, lungs, heart, and kidney) were measured and calculated as the percentage of body weight (BW). The results showed that the percentage of spleen, liver and kidney weights were increased ($p < 0.05$) according to the level of CBS supplementation. The effect of CBS feeding on the percentage of spleen weight was occurred at level 9%, and significantly higher ($p < 0.05$) compared to the level of 3%, 6% and without cocoa shell. The liver and kidney percentages were higher ($p < 0.05$) at levels 6% and 9% compared to 0% and 3%. There was no effect of supplementation on the digestive tract, lungs and heart. The study concluded that CBS supplementation at 6% and 9% levels had an effect on physiological visceral organs of Bali cattle bulls.

1. Introduction

Increasing meat production can be done through an increasing population and production efficiency of livestock. Using feed from agricultural and industrial waste is an alternative to increase environmentally friendly livestock production. Cocoa bean shell (CBS) is one of the agricultural industries wastes were potentially to use as feed supplements for livestock. The potential of CBS in Indonesia is very large. BPS data for 2017 shows that cocoa production in 2017 reached 658 tons [1], those an estimated production of CBS of 112 tons per year. CBS contains 68.8% dry matter, 6.8% fat, 25.1% fiber and 16.7% crude protein. It is also has a high palatability for cattle with total digestible nutrient (TDN) is equivalent to grass, 52.3% [2], CBS also contains other beneficial polyphenol compounds as in cocoa, especially in the form of catechins, flavonol glycosides, anthocyanins and procianidin [3], theobromine is the main responsible factor for cocoa's effects on body weight gain as well as on lipid and glucose metabolism [4], event they also had potency to be an alternative antibiotic for animal farm [5].

However, there are weaknesses in the content of anti-nutrition substances such as tannins and several types of alkaloids including theobromin. In our previous study [6], we found that high concentration of theobromine effected on nitrogen urea in blood plasma, in other hand in limited use, the presence of anti-nutrients can help in body metabolism as an antioxidant. Theobromine and some phlavanoids work



in energy metabolism, including the process of forming, storing and utilizing muscle fat and glycogen. However, the study on CBS as feed supplementation, and its depth effects on digestive and metabolism are needed. This study aimed to measure the effect of providing various levels of CBS (as supplement) on the development of some visceral organs of Bali cattle after fattening for 4 months. The research is expected to find the right formulation of CBS as supplementation for Bali cattle in forages basal feed.

2. Research method

The study was conducted in cattle fattening pen, CV Akbar Jaya, Makassar. The study used 12 1.5-year-old bulls with an average body weight of 159.25 ± 15.68 kg, body condition score was 2 on a scale of 5. Slaughtering of cows was done after 12 hours fasting, carried out at Tamangapa Animal Slaughterhouse, Antang-Makassar.

The study was arranged according to a completely randomized design with 4 treatments of CBS supplemented concentrate, A = 0%; B = 3%; C = 6%; and D = 9%. Each treatment was repeated three times. Further tests for multiple comparisons use the Tukey-HSD test. The administration of worming was carried out during the feed adaptation period in the initial 2 weeks of the study. Weighing was done every 14 days, for 3 months. Measurement of the final weight was done prior the cattle were slaughtered; the end of fasting for 12 hours. Measurement of the weight of digestive organs and visceral organs was carried out after slaughtering cattle at the end of the study. The parameters measured include body weight, rumen, intestine, liver, heart, lungs, kidneys, and spleen. The percentage of organ weight was obtained from the distribution of organ weight to cow's body weight before slaughter.

3. Result and discussion

The analysis showed an increase ($p < 0.05$) in the percentage of spleen, liver, and kidney weights, while the digestive tract, rumen, lungs and heart showed no significant differences. The complete results of the study are presented in Table 1.

Table 1. Weight percentage of visceral organs of Bali bulls after 4 months fattening with cocoa beans shell supplemented concentrate feed

Visceral organs	Cocoa beans shell levels				SE	P
	0	3	6	9		
Rumen	3.06	2.98	2.97	2.95	0.057	0.086
Digestive tract	1.71	1.69	1.69	1.66	0.023	0.109
Spleen	0.53 ^a	0.55 ^a	0.58 ^{ab}	0.68 ^b	0.055	0.037
Liver	2.71 ^a	2.86 ^{ab}	2.89 ^b	3.02 ^b	0.136	0.018
Lungs	0.86	0.90	0.86	0.90	0.037	0.462
Heart	0.79	0.82	0.82	0.86	0.040	0.126
Kidney	0.35 ^a	0.38 ^{ab}	0.43 ^{bc}	0.47 ^c	0.056	0.011

Least squares means within a row with a common superscript letter differ ($P < 0.05$)

Analysis of variance showed that there was no difference in rumen percentage due to treatment. This indicated that a CBS increase of up to 9% in feed has no direct effect on rumen structure or on the rumen epithelial layer. However, there was a decrease in rumen weight percentage in line with the CBS level increase. Likewise, in the digestive organs, lungs, and heart, which did not show significant differences. The CBS treatment has a significant effect on the Spleen, liver, and kidney. The percentage of these organs increases with CBS level.

The relationship of metabolic rate to body size has become an interesting topic for doctors, especially pediatricians. It has been learned that many quantitative functions vary during growth in relation to metabolic rate, not body size. Example from this was cardiac output, glomerular filtration rate, oxygen

consumption and drug dosage. This phenomenon can reflect direct or possible cause and effect relationships coincidence parallels between a relatively slow increase in metabolic rate compared to body size and function in question. Metabolism decreases and many other measures of physiological function in relation to the unit of body size are widely observed. This phenomenon can be demonstrated by comparing between species, as well as within a species during growth or among members of mature species of varying sizes [7]. Especially in bovine, cow have a basal metabolic rate per kg (BMR/kg) about $69.8 W^{0.75}$; Cattle, beef heifer $75.4 W^{0.75}$ and $117.0 W^{0.75}$ in dairy cattle [8]. In the case of humans during growth, babies have BMR / kg more than double that of normal adults. Normal adults may have a BMR / kg one and a half times that of obese adults. Holliday [7] suggested that high BMR is caused, in part, by a relatively high proportion of body weight taken by all internal organs, including the brain - not because they are large, but their supporting structure.

Spleen, Liver and Kidney were appearing their weight increase depend on the CBS supplementation levels. Spleen is known has important role for antibacterial and antifungal immune reactivity [9]. In the case of responses on anti-nutrient or anti-oxidant affect in metabolic system, various insignificant negative effects on some species were reported widely. Kuku [10] reported that laboratory animals histopathological analysis indicated that the spleen and small intestines were adversely affected, but could have deleterious effects in animals if ingested without adequate processing. Spleen abnormalities, such as large depositions of hemosiderin and melanin pigments and proliferation of melano-macrophage centers, lymphocytic depletion of the white pulp areas (hypocellularity), and presence of vacuoles and necrotic in observed areas were similar to the effects of vitamin E and/or vitamin C deficiencies observed in fish [11]. Some anti-nutritional factors have been observed to inhibit absorption of nutrients and their subsequent utilization and assimilation by animals [12]. Besides, they cause some level of damages to some organs such as liver, kidney and spleen [13].

The condition of hepatotoxicity is a concern in the treatment of CBS supplementation, the high content of active compounds contained therein needs to be a particular concern, because it can affect the performance of organ metabolism nutrition, as theobromine has responsibility for some effect in metabolism [14]. The severity of liver abscesses was evaluated by [15] in individually feeder cows, the report show that concentrate level in the diet of 77.7% affects the severity score of liver abscess, final live weight ($p < .10$), hot carcass weight ($p < .0001$), intake of dry matter ($p < .10$), increase in daily body weight. In this study, no significant differences between all CBS treatments but significantly higher compared to the control, and the means that the range of 3% of CBS supplementation did not have significant effects on heart size development in bulls. However, [16] workload determines organ size, but dietary factors influencing workload clearly vary for each organ.

The kidneys play to filter the blood, remove wastes, control the body's fluid balance, and keep the right levels of electrolytes. In cattle, which have a higher metabolic rate per unit of metabolic BW than do most other domestic ruminant, but has poorly developed mechanism in the kidney and gut for water retention [17]. In basal metabolism rate (BMR) the proportion of kidney organ was 608 ± 25 g, for cattle with $277,000 \pm 20,800$ BW [18] or about 0.20 – 0.21% BW. The recent study found that CBS on control bulls had no significant differences than those supplemented by 3% of CBS (0.35 vs 0.38), and significantly increase for 0.43 and 0.47% at 6 and 9% of CBS, a greater value than the reference indicates the differences of the cattle breed. Greater value in Bali Bulls may due to their body condition which about two or three times smaller than European breed bulls. Depends on cauture et al. [19], larger mammals have lower tissue metabolic rates (TMR; $\text{mmol O}_2 \cdot \text{g wet wt}^{-1} \cdot \text{min}^{-1}$) at 37 degrees C, yielding the equations $\text{TMR} = 3.6 M - 0.21$ for liver slices and $\text{TMR} = 3.2 M - 0.11$ for kidney cortex slices. Increased the kidney mass can also be influenced by protein intake, Hammod et al [20] concluded that liver, kidney and stomach mass increased with protein intake rate, while digestive tract and other vital organ masses increased only in response to increased energy intake rate. There has not been found any influence or relationship between theobromine or other methylxanthine derivatives directly on kidney performance, but the indirect effect of changes in metabolic activity is very likely to occur.

4. Conclusion

In conclusion, the supplementation of CBS concentrate up to 9% level in the rations did not affect the rumen, gastrointestinal, lung, and cardiac aspiration; but has an effect on the increasing of spleen, liver and, kidneys percentage. Although there was no significant effect on the percentage of the rumen and the digestive tract, there was a tendency in decreasing the percentage of the such two organs weight; indicated the occurrence of degradation conditions for the long time during the research fattening period.

References

- [1] Kementerian Petanian 2018 Statistik Pertanian Nasional Tahun 2018
- [2] Ozung P O, Oko O O K and Agiang E A 2016 Chemical composition of differently treated forms of cocoa POD husk meal (CPHM) *Asian J. Agric. Sci.* **8** 5–9
- [3] Utami R R, Armunanto R, Rahardjo S and Supriyanto 2016 Effects of cocoa bean (*Theobroma cacao* L.) fermentation on phenolic content, antioxidant activity and functional group of cocoa bean shell *Pakistan J. Nutr.* **15** 948–53
- [4] Camps-Bossacoma M, Garcia-Aloy M, Saldana-Ruiz S, Cambras T, González-Domínguez R, Franch À, Pérez-Cano F J, Andres-Lacueva C and Castell M 2019 Role of theobromine in cocoa's metabolic properties in healthy rats *J. Agric. Food Chem.* **67** 3605–14
- [5] Huang Q, Liu X, Zhao G, Hu T and Wang Y 2018 Potential and challenges of tannins as an alternative to in-feed antibiotics for farm animal production *Anim. Nutr.* **4** 137–50
- [6] Ali H M, Alam G, Syamsu J A, Salengke S and Asja M A 2014 Average daily gain, AST and blood nitrogen urea (BUN) responses of bali beef on cocoa waste extract supplement *J. Adv. Agric. Technol.* **1** 5–9
- [7] Holliday M A, Potter D, Jarrah A and Bearg S 1967 The relation of metabolic rate to body weight and organ size: a review *Pediat. Res.* **1** 185–95
- [8] Tombropoulos E G and Kleiber M 1961 The metabolism of specifically labelled glucose in bovine ketosis *Biochem. J.* **80** 414–20
- [9] Mebius R E and Kraal G 2005 Structure and function of the spleen *Nat. Rev. Immunol.* **5** 606–16
- [10] Kuku A, Etti U J and Ibrionke I S 2014 Processing of fluted pumpkin seeds, *Telfairia occidentalis* (Hook F) as it affects growth performance and nutrient metabolism in rats *African J. Food, Agric. Nutr. Dev.* **14** 1992–2014
- [11] Martinez-Garcia C G, Rayas-Amor A A, Anaya-Ortega J P, Martinez-Castaneda F E, Espinoza-Ortega A, Prospero-Bernal F and Arriaga-Jordan C M 2015 Performance of small-scale dairy farms in the highlands of central Mexico during the dry season under traditional feeding strategies. *Trop. Anim. Health Prod.* **47** 331–7
- [12] Tope A K 2014 Effect of fermentation on nutrient composition and anti-nutrient contents of ground Lima bean seeds fermented with *Aspergillus fumigatus*, *Rhizopus stolonifer* and *Saccharomyces cerevisiae* *Int. J. Adv. Res.* **2** 1208–15
- [13] Emiola I A, Ologhobo A D and Gous R M 2007 Performance and histological responses of internal organs of broiler chickens fed raw, dehulled, and aqueous and dry-heated kidney bean meals *Poult. Sci.* **86** 1234–40
- [14] Zbaida S, Kariv R, Fischer P and Gilhar D 1987 Reactions of theophylline, theobromine and caffeine with Fenton's reagent-simulation of hepatic metabolism *Xenobiotica* **17** 617–21
- [15] Brink D R, Lowry S R, Stock R A and Parrott J C 1990 Severity of liver abscesses and efficiency of feed utilization of feedlot cattle. *J. Anim. Sci.* **68** 1201
- [16] Sainz R D and Bentley B E 1997 Visceral organ mass and cellularity in growth-restricted and refed beef steers *J. Anim. Sci.* **75** 1229–36
- [17] Maltz E and Silanikove N 1996 Kidney function and nitrogen balance of high yielding dairy cows at the onset of lactation *J. Dairy Sci.* **79** 1621–6
- [18] Couture P and Hulbert A J 1995 Membrane fatty acid composition of tissues is related to body mass of mammals. *J. Membr. Biol.* **148** 27–39
- [19] Couture P and Hulbert A J 1995 Relationship between body mass, tissue metabolic rate, and

sodium pump activity in mammalian liver and kidney *Am. J. Physiol. - Regul. Integr. Comp. Physiol.* **268**

- [20] Hammond K A and Janes D N 1998 The effects of increased protein intake on kidney size and function. *J. Exp. Biol.* **201** 2081 LP – 2090