

PAPER • OPEN ACCESS

The effect of temperature levels on antioxidant activity in chicken eggs

To cite this article: N Nahariah and H Hikmah 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **788** 012099

View the [article online](#) for updates and enhancements.

You may also like

- [Antioxidant Activity and Stability of Radish Bulbs \(*Raphanus sativus* L.\) Crude Extract](#)
Eveline and R L Pasau
- [Effect of combination of poultry eggs and acids on the color profile and consumers' acceptance on mayonnaise](#)
I Amalia, N Nahariah and H Hikmah
- [An Evaluation of a Systematic Series of Cobalt-Free Ni-Rich Core-Shell Materials as Positive Electrode Materials for Li-Ion Batteries](#)
Yulong Liu, Dongxu Ouyang, Divya Rathore et al.



The Electrochemical Society
Advancing solid state & electrochemical science & technology

243rd ECS Meeting with SOFC-XVIII

Boston, MA • May 28 – June 2, 2023

**Abstract Submission Extended
Deadline: December 16**

[Learn more and submit!](#)

The effect of temperature levels on antioxidant activity in chicken eggs

N Nahariah and H Hikmah

Faculty of Animal Science, Hasanuddin University, Jl. Perintis Kemerdekaan Km 10, Makassar, South Sulawesi, Indonesia

E-mail: nahariah11@gmail.com

Abstract. Generally, egg heating aims to inhibit microbial contamination, especially on the surface of eggshells and liquid egg products. The application of heating temperature to eggs was carried out at temperatures below 100°C. However, the heating temperature can have an impact on antioxidant activity and the characteristics of egg protein consumption. The treatment uses 3 temperature levels i.e., 45°C, 55°C, and 65°C, respectively. Each treatment was repeated 5 times, a total of 15 units. Each sample was heated for 3 minutes. Parameters were antioxidant activity, total protein, and dissolved protein. The results showed that the different heating temperatures had a significant effect ($P < 0.01$) on antioxidant activity, total protein, and dissolved protein. Value of antioxidant activity and total protein decreased with the increasing of heating temperature. However, increasing the heating temperature can increase the dissolved protein value of chicken eggs. The heating temperature of 45°C on chicken eggs can optimize the antioxidant activity, total protein, and dissolved protein value.

1. Introduction

Eggs are highly nutritious animal food. Besides, eggs contain bioactive compounds that are beneficial to health called functional foods [1,2]. Functional food is food that provides health effects to its users. The potential of eggs as a functional food is quite high [3], due to the nature of poultry as a bio-convert from feed nutrition to egg nutrition [4,5]. Also, the functional properties of eggs can be obtained through processing mechanisms [6], either by heating or pasteurization [7].

Egg pasteurization is done by heating eggs at a temperature below 100°C [8]. Pasteurization or heating of eggs aims to inhibit microbial contamination, especially on the surface of eggshells and liquid egg products. The pasteurization temperature used depends on the types of raw material. According to the USDA (1980) that heating the eggs aims for pasteurization [8]. The heating can be done at 60°C for 3.5 minutes [9]. Meanwhile, according to Cotterill [10], the heating process is 56.7°C for 3.5 minutes or 55.6°C for 6.2 minutes. The maximum pasteurization temperature for eggs is 65°C. The high temperature was reported as a cause of a decrease in egg characteristics [6,11].

Research by Ubba *et al.*, (2018) and Nahariah *et al.*, (2014) were revealed that the heating process increased the antioxidant activity of eggs [12,13]. The temperature and heating time in eggs changed the protein structure of the eggs. Changes in egg protein structure from complex to simple. The change in protein structure resulted in the formation of peptide bonds such as dipeptides or oligopeptides. The ability of oligopeptides to inhibit oxidative reactions is indicated as antioxidants. Research by Nahariah *et al.*, (2014), explained that egg whites that were oven-dried at 45°C for 39 hours showed antioxidant



activity of 26.85% [13]. The use of an incubation temperature of 38°C in infertile eggs from the hatchery industry can provide an inhibitory action against oxidation of 77.99% [14].

The heating process in eggs has an impact on changes in their antioxidant activity [7,15]. Likewise, heating of eggs can also result in changes in total protein and dissolved protein. However, the studies of this examined are limited. Therefore, the objectives of this study were to examine the effect of optimal heating temperature on consumption eggs on antioxidant activity, total protein, and dissolved protein.

2. Materials and Methods

The materials were one day old chicken eggs, aquades, aquabidistila, alcohol, methanol, phenolic acid, H₂SO₄, ice water, HCl, Folin Ciocalteu, 10% Na₂CO₃, K₂SO₄, HgO, NaOH-thiosulfate, Lowrey reagent, Folin reagent, Calcium Permanganate (PK), 2,2-diphenyl-1-picrylhydrazyl (DPPH), Tricarboxylic Acid (TCA), and Bovine Serum Albumine (BSA). The equipment was beaker glass, waterbaths, test tubes, *Graduated Cylinder*, Kjeldahl flasks, stopwatches, ovens, UV-VIS spectrophotometers, analytical scales, 1000µm micropipettes, vortex, mixer, erlenmeyer flask, distillators and incubators, hot plate, spoits, and centrifuges.

This study used a completely randomized design (CRD) with 3 treatments and 5 replications. The total treatment was 15 units. The treatment of the research was the heating temperature of the eggs i.e., 45°C, 55°C, and 65°C, respectively. Furthermore, all the treatments were heating for 3 minutes. The parameters measured were antioxidant activity, total protein, and dissolved protein. Measurement of the antioxidant activity by DPPH method [16], while measurement of total protein by AOAC (1984) and dissolved protein by Wikandari *et al.*, (2011), respectively [17,18].

3. Results and Discussions

The results showed that the antioxidant activity and characteristics of egg protein changed due to the differences in heating temperature. The results of the study are presented in Table 1.

Table 1. Antioxidant activity and characteristics of egg protein with different heating temperatures.

Description (%)	Temperature (°C)			Mean±SD or SE
	45	55	65	
Antioxidant activity	92.69±1.43 ^a	86.89±1.04 ^b	87.47±1.80 ^c	89.02±3.04
Total protein	12.42±0.13 ^a	11.53±0.09 ^b	11.46±0.54 ^b	11.80±0.54
Dissolved protein	0.24±0.01 ^a	0.24±0.02 ^b	0.31±0.15 ^c	0.27±0.03

Different superscripts on the same line show significant differences (P <0.01). Data±SD or SE?

3.1. Antioxidant activity in eggs by different heating temperature treatment

The results showed (Table 1), that different heating temperatures had a significant effect (P <0.01) on the antioxidant activity of the eggs.

The LSD test showed that the antioxidant activity was significantly different with the increasing heating temperature. This is probably caused by changes in the protein structure of the eggs due to high temperature. By Miryanti *et al.*, (2011), using high temperatures in food can have an impact on the damage to its antioxidant compounds [19]. Heating of foodstuffs can cause an increase in the chain of initiation and propagation of oxidation reactions, also decreased the antioxidant activity in ingredients [20,21]. High heating temperatures can cause oxidative stress on foodstuffs. The application of high temperatures in the heating process of eggs can cause oxidative stress. Oxidative stress occurs in ingredients or eggs that requires antioxidants to inhibit or reduces the oxidation of ingredients or eggs [7].

Table 1 showed the antioxidant activity of eggs at a heating temperature of 45°C for 3 minutes at 92.69% and decreased with the increase heating temperature. The value of egg antioxidant activity obtained in this study was higher than the results of previous studies. Research by Nirmalaratne and Wu (2015) found that antioxidants for raw whole eggs were 86.95% [3]. However, high temperatures can also damage antioxidants, so it was necessary to control the temperature to maintain the antioxidant activity of eggs [7,19]. The application of proper heating temperature to eggs has a positive effect on changes in egg protein structure. The proper heating mechanism can inhibit oxidative reactions. However, it is expected to optimize the antioxidant content [1].

3.2. Total protein in eggs by different heating temperature treatment

The results (Table 1) showed that heating temperature had a significant effect ($P < 0.01$) on the total eggs protein.

Heating eggs at a temperature of 45°C was significantly different ($P < 0.01$) from reducing the protein of the total egg due to the increased heating temperature used in this study. However, total egg protein did not significant difference at heating at 55°C nor 65°C. This is probably due to the utilization of temperatures above 45°C in the egg heating process which has resulted in changes in protein structure. Heating at high temperatures causes protein to decrease [7,22]. Materials that experience high heating result in changes to the secondary, tertiary, and quaternary structures, respectively [23]. These changes result in the denaturation of the protein material [23]. Heating causes the breaking of the hydrogen bonds that support the secondary and tertiary structures of a protein [7,23,24]. This change causes the hydrophobic part of the polypeptide side group to open and release water [1,23].

Relevant research showed that heating eggs using a water bath at 63°C for 3 minutes and heating by oven at 70°C for 60 minutes makes egg protein and eggshell membrane clump [25]. Likewise, the research of Novia *et al.*, (2011) showed that the protein content of salted eggs at an oven temperature of 70°C was 37.54% and decreased to 28.40% at an oven temperature of 80°C [26].

3.3. Value of dissolved protein in eggs by different heating temperature treatment

The results presented in Table 1 indicated that different heating temperatures have a significant effect ($P < 0.01$) on egg dissolved protein.

The LSD test showed that the soluble egg protein significantly increased with the increase in heating temperature. This is probably due to the changes in protein structure because the heating breaks down into simpler structures. Eggs are hydrated and their protein structure breaks down and denatures [6,27]. This condition causes the protein to dissolve in solvents such as water [27]. [27] explains that heat-denatured protein material can decrease its water binding ability. Heat energy causes the interruption of non-covalent interactions that exist in the natural structure of proteins. However, the low energy does not break the covalent bonds, namely the peptide bonds [6].

The high heating temperature can be increasing the kinetic energy and it is causing the building blocks of protein to move very fast. This can be resulting in broken protein molecular bonds [28]. Protein components are broken down by heat into simpler compounds and dissolve in water [13]. This is relevant to Tyl *et al.*, (2020), which stated that changes in protein structure can result a decrease in soluble protein [24].

4. Conclusion

There were changes in antioxidant activity and protein characteristics in eggs due to different heating temperatures. Heating eggs at 45°C for 3 minutes increased the antioxidant activities and protein total. Applying high temperatures to egg heating increase dissolved protein. However, high temperatures decreased the antioxidant activities and protein total.

References

- [1] Nahariah, N Hikmah H and Yuliati F N 2020 The evaluation of changes in organoleptic flavor of fermented egg whites at different levels and types of fruit. In *IOP Conf. Ser. Earth and*

- Environ. Sci.* **492** 012041
- [2] Godbert S R, Guyet N and Nys Y 2019 The Golden Egg Nutritional, value, bioactivities and emerging benefits for human healthy *Nutrients* **11** 648
- [3] Nimalaratne C and Wu J 2015 Hen egg as an antioxidant food commodity Review *Nutrients* **7** 8274–93
- [4] Nirmalaratne C, Lutz D L, Schieber A and Wu J 2011 Free aromatic amino acids in egg yolk show antioxidant properties *Food Chem.* **129** 155–61
- [5] Kartina, Nahariah N, Fatma and Hikmah H 2020 Egg chip quality with different types dan levels of fillers *Hasanuddin J. Anim. Sci.* **2** 9–16
- [6] Nahariah N, Legowo A M, Abustam E, Hintono A 2015 Angiotensin I-Converting enzyme inhibitor activity on egg albumen fermentation *Asian Australas J. Anim. Sci.* **28** 855–61
- [7] Yuliati, F N and Ali H M 2019 Effects honey on different levels of antioxidant activity and chemical of pasteurized eggs *IOP Conf. Ser. Earth. Environ. Sci.* **247** 012067
- [8] USDA 1980 *Regulation governing inspection of egg products* (Albany: USDA)
- [9] USDA 1969 *Egg Pasteurization Manual* (Albany: USDA)
- [10] Cotterill O J 1968 Equivalent pasteurization temperatures toll kill Salmonella in liquid egg white at various pH levels *Poult. Sci.* **47** 352–65
- [11] Irianto K 2010 *Mikrobiologi Menguak Dunia Mikroorganisme* (Bandung: Yrama Widya)
- [12] Ubba E T, Abustam E dan Nahariah N 2018 Antioxidant activity of infertile egg residue from the hatching industry by giving chitosan during different storage periods *Advan. Environ. Biol.* **12**(10) 1–4
- [13] Nahariah, Legowo A M, Abustam E, Hintono A, Bintoro V P and Pramono Y B 2014 Evaluasi potensi aktivitas ACE-Inhibitor endogenous pada putih telur dari jenis unggas yang berbeda. *Prosiding Seminar Nasional Optimasi Sumber Daya Lokal pada Peternakan Rakyat Berbasis Teknologi* 9-10 Oktober 2014 (Makassar: Faculty of Animal Science, Universitas Hasanuddin) pp 207-213.
- [14] Ubba E 2014 *Potensi Antioksidan pada Telur Infertil Hasil Seleksi Berdasarkan Waktu Pengeraman yang Berbeda* Skripsi (Makassar: Fakultas Peternakan Universitas Hasanuddin Makassar)
- [15] Nimalaratne C, Schieber A, Wu J 2016 Effects of storage and cooling on the antioxidant capacity of laying hen eggs *Food Chem.* **1** 111–16
- [16] Gasic U, Keckes S, Dabic D, Trifkovic J, Oспенica D M, Natie M and Tesic Z 2014 Phenolic profile and antioxidant activity of Serbian polyforal honeys *Food Chem.* **145** 599–07
- [17] AOAC 1984 *Official Methods of Analysis* 12th Ed. (Washington: DC Association of Official Analysis Chemist)
- [18] Wikandari, P R, Suparmo, Y Marsono and E S Rahayu 2011 Potensi bekasem sebagai Sumber Angiotensin I-converting Enzyme Inhibitory *Biota* **16** 145–52
- [19] Miryanti Y I P, Sapei L, Budiono K and Indra S 2011 *Ekstraksi Antioksidan dari Kulit Buah Manggis (Garcinia mangostana L.)* Laporan Penelitian (Bandung: Lembaga Penelitian dan Pengabdian Kepada Masyarakat Universitas Katolik Parahyangan)
- [20] Pokorny J, Yanishlieva N and Gordon M 2001 *Antioxidant in Food* (New York: CRC Press)
- [21] Raharjo S 2006 *Kerusakan Oksidatif pada Makanan* (Yogyakarta: Gadjah Mada University Press)
- [22] Sundari 2015 Pengaruh proses pemasakan terhadap komposisi zat gizi bahan pangan sumber protein *Media Penelitian dan Pengembangan Kesehatan* **25** 235–42.
- [23] Zulfikar 2008 *Kimia Kesehatan Jilid 3* (Jakarta: Departemen Pendidikan Nasional)
- [24] Tyl C, Marti A and Ismail B P 2020 Changes in protein structural characteristic upon processing of gluten free millet pasta *Food Chemistry* **327** 1–7
- [25] Rizal B, Hintono A and Nirwantoro 2012 Microbial growth in post-heating eggs *Animal Agriculture Journal* **1** 208–18
- [26] Novia D, Melia S and Ayuza N Z 2011 Study of oven temperature on protein content and

- organoleptic value of salted eggs *Jurnal Peternakan* **8** 70–6
- [27] Ophart C E 2003 *Virtual Chembook* (Illinois: Elmhurst College Press)
- [28] Winarno F G and Koswara S 2002 *Telur, Penanganan dan Pengolahannya* (Bogor: M- BRIO Press)