

ENDOSYMBIONT BACTERIA  
OF PHERETIMA SP.  
EARTHWORMS (ANNELIDA:  
OLIGOCHAETA) POSSESSES  
ANTIBACTERIAL ACTIVITY  
*by*

---

FILE	AJ-17.PDF (140.85K)	WORD COUNT	3851
TIME SUBMITTED	07-MAY-2019 08:37AM (UTC+0700)	CHARACTER COUNT	21185
SUBMISSION ID	1126122625		

## ENDOSYMBIONT BACTERIA OF *PHERETIMA* SP. EARTHWORMS (ANNELIDA: OLIGOCHAETA) POSSESSES ANTIBACTERIAL ACTIVITY

DIRAYAH RAUF HUSAIN, FITRIANI, ZOHRA HASYIM, ASADI ABDULLAH AND SULFAHRI\*

Departement of Biology, Faculty of Mathematics and Natural Sciences,  
Hasanuddin University, Makassar, Indonesia

(Received 28 April, 2018; accepted 25 June, 2018)

**Key words :** *Pheretima* sp. earthworm, Endosymbiont bacteria, Times of incubation, *Salmonella typhi*,  
*Staphylococcus aureus*.

**Abstract**– Earthworms are used as drugs, not only the compound earthworms but also endosymbiont microorganisms in earthworms suspected of having antibiotic activities. This research is aimed to identify the ability of several endosymbiont bacteria in *Pheretima* sp. earthworms to inhibit the growth of the bacteria *Salmonella typhi* and *Staphylococcus aureus*. Isolation of bacteria from *Pheretima* sp. earthworms was carried out on NA (nutrient agar medium). The observed characteristics of bacterial isolates from *Pheretima* sp. earthworms consisted of colony morphology, gram staining, and the presence of endospores. This research demonstrates the variety of colors, shapes, edges, and elevations of the isolates. The result showed that there are 2 isolate that was isolation form *Pheretima* sp. namely *Bacillus choshinensis* and *Bacillus brevis*. Based on the test results of isolation and inhibition of *Pheretima* sp. earthworms, all isolate effectively inhibited the growth of the bacteria *S. typhi* and *S. aureus*, and they have the potential to act as antibiotics. Isolates *Bacillus choshinensis* had the largest inhibitory zone diameter of 21.32 mm against *S. typhi* after 15 days of incubation. Inhibitory properties of all isolates that were isolated from *Pheretima* sp. indicate that bacteria could be described as bactericidal.

### INTRODUCTION

Earthworms (Phylum: Annelida, Order: Oligochaeta, Class: Clitellata) are known to be the most important soil fauna biomass in humid soils of temperate and tropical regions (Leveque *et al.*, 2013; Singh *et al.*, 2016). These animals are most often found in moist soil and contain numerous beneficial organic compounds and minerals from natural sources or waste disposal bins, their natural habitats (Chachina *et al.*, 2016). Earthworm species consist of three major groups, namely Polychaeta, Oligochaeta, and Harudinea, as well as two small groups, namely Aeolosamata and Branchiobdella (Klarica *et al.*, 2012; Carnovale *et al.*, 2015).

Polychaeta is the largest group of invertebrates, comprising approximately 8000 species; the largest group is found in the sea (Fiege *et al.*, 2010; Bohlen, 2001; Metcalfe and Glasby, 2008). The typical form of Polychaeta is characterized by a jointed body shape, and on every segment, there is a pair of parapodia

(Bohlen, 2001; Omena *et al.*, 2012; Choi *et al.*, 2011). The potential of polychaeta earthworms as medicinal raw materials and ingredients for healing various diseases has been expressed by many researchers (Li *et al.*, 2011; Cho *et al.*, 2014).

Polychaeta earthworms are used as drugs, not only the compound polychaeta earthworms but also endosymbiont microorganisms in polychaeta earthworms suspected of having antibiotic activities (Fiolka *et al.*, 2010). As reported in earlier studies, earthworms are used as drugs, not only the compound earthworms but also endosymbiont microorganisms in earthworms suspected of having antibiotic activities. Therefore, further screening of the potential bacterial endosymbionts contained in the worm *Pheretima* sp. needs to be done. This study is aimed to investigate antibiotic compounds from endosymbionts of *Pheretima* sp. earthworms (polychaetes) as inhibitors of the growth of *Salmonella typhi* and *Staphylococcus aureus*.

---

\*Corresponding author's email: sulfahri@unhas.ac.id

## MATERIAL AND METHODS

### Isolation of Bacteria

Samples of *Pheretima* sp. earthworms were obtained from garbage and washed with sterile distilled water until the soil and dirt attached to the skin surface of the earthworms was washed away. The earthworms were then slowly killed by washing with 70% alcohol. Earthworms were measured, crushed using a mortar, and serially diluted by adding 0.1 mL to 0.9 mL of diluent to a maximum dilution of 10<sup>-6</sup>. A total of 1 mL of each dilution was then inoculated on NA (nutrient agar) medium by using the casting method, then incubated at 37°C for 48 h. Each colony that grew was selected for the purification step.

### Purification of Bacterial Culture

This stage of purification began with selecting individual colonies. Round, sterile needles were touched to the surface of bacterial colonies and then inoculated on the surface of NA medium by the scratch method to obtain isolated colonies. It took several attempts to obtain pure colonies, which were then incubated at 37°C for 48 h. The purification step can be performed 2–3 times to ensure the complete purity of the selected colonies.

### Bacterial Identification

#### Observation of Morphology

In the morphological observation, each colony formed after purification was observed. Observations made include colony form, color, edge, center, diameter, and surface morphology, as well as cell morphology and gram-positive or negative status of bacteria and their spores to classify isolates.

#### Gram Staining

Observations of cell morphology were made using Gram staining techniques. First, bacteria were mounted and fixed on glass slides. A total of 2–3 drops of stain A (crystal violet) were dropped onto bacterial colonies and incubated for 60 s. Preparations were then washed under running water and dried. A total of 2–3 drops of stain B (lugol) were dripped over the preparation and left for 60 s. The mixture was washed under running water and dried. Preparations were then etched with 2–3 drops of stain C (alcohol-acetone), incubated for approximately 30 s, then washed again and dried. Further, 2–3 drops of spilled stain D

(safranin) were added and allowed to stand for 60 s, then washed and dried. Observations were then made under light microscopy.

#### Endospore Staining

This test is used to determine the presence of bacterial spores in soil worms. A smear of bacteria from pure cultures was fixed on a sterile glass microscope slide, flooded with 5% malachite green, and placed on a hot plate that had been preheated to 200°C for 10 min. After 10 min, the slide was cooled and then cleaned under running water. The smear of bacteria was flooded with safranin for 1 min and washed under running water. The slide was drained of excess water on absorbent paper and then observed under a microscope. Spores appeared green in color or refractive.

#### Biochemical Identification

The isolated bacteria strains from *Pheretima* sp. gut were identified using biochemical tests API 20 E (Bio-Merieux).

#### Bacterial Isolates of Preculture Stage Symbionts

Isolates obtained were subsequently used to inoculate a preculture/starter culture in NB (nutrient broth) medium and incubated with shaking for 24 h at 37°C. After it grew, the preculture was inoculated into NB (nutrient broth) medium at a 1:10 ratio.

#### Testing Antibiotics

Antibiotic activity was tested using MHA medium (Muller Hinton order) by the agar diffusion method using a paper disk. Each paper disk was immersed in the supernatant of bacterial symbionts for 15 min and a positive control solution of chloramphenicol. A 1 mL suspension of test bacteria was added to a sterile petri dish and MHA medium was added at 45°C and allowed to solidify. Paper disks that had been soaked in MHA media solidified with a distance of 20 mm between each paper disk and then incubated at 37°C. Observations were made after 24 h, then the diameter of the inhibition zone formed was measured, and incubation was resumed for 48 h to determine the nature of the antibiotic compounds contained in the worm.

## RESULTS

### Microscopic Observation and Identification of Bacterial Isolates from *Pheretima* sp.

Based on the results of microscopic observation and

biochemical identification, there are 2 isolates: *Bacillus choshinensis* and *Bacillus brevis*. Based on the results of Gram staining, both of bacteria obtained were in the shape of gram-negative bacilli (rods). Endospore tests of the stained slides revealed that both isolates capable of forming endospores. Isolate *Bacillus choshinensis* in the form of single bacilli, while isolate *Bacillus brevis* in the form of bacilli with two chain (Table 1).

#### Inhibitory Power of Earthworm Test Against Bacterial Pathogens *Staphylococcus aureus* (gram-positive) and *Salmonella typhi* (gram-negative)

Earthworm play a critical role in the decomposition of organic matter. In organic materials, there are many types of bacterial, including pathogenic bacteria. This type of bacteria can affect the composition of the endosymbiont bacterial on earthworm. Inhibition test of bacteria is a method for determining the ability of bacteria to inhibit other bacteria. Table 2 shows the effect of incubation time on the inhibition zone on both test bacteria, *S. typhi* and *S. aureus*. After an incubation time of 15

days, 15.27 mm and 21.32 mm diameter zones of inhibition occurred around *Bacillus choshinensis* in the test bacteria *S. typhi* and *S. aureus*, respectively. The appearance of an inhibition zone around *Bacillus choshinensis* greater than 14 mm in diameter indicated effective inhibition of bacterial growth, in accordance with Cappuccino & Sherman [16]. Zones of inhibition with diameters of 10–11 mm and 9 mm are considered less effective and ineffective, respectively (Cappuccino and Sherman, 2011).

#### DISCUSSION

The inhibition zones around bacteria isolated from *Pheretima* sp. worms were greater than those around the commercial antibiotic Amoxicillin (Table 2). This suggests that the bacteria isolated from *Pheretima* sp. worms have potential as an antibiotic. The same phenomenon was reported by Fiolka *et al.* (2010), Brito-Vega and Espinosa-Victoria (2009), and Byzov *et al.* (2007) that some worm species have bacterial endosymbionts that can inhibit the growth of some types of pathogenic bacteria.

**Table 1.** Characterization of bacterial colony morphology earthworms *Pheretima* sp. and Gram's staining and endospore staining

Isolates	Forms colonies				Gram Staining		Endospore
	Color	Form	Edge	Elevation	Form	Gram	
<i>Bacillus choshinensis</i>	White	Circular	Entire	Flat	Single basil	Positive	exist
<i>Bacillus brevis</i>	White	Circular	Entire	Convex	Basil two chain	Positive	exist

**Table 2.** Inhibitory Zone of isolate bacteria that was isolated from *Pheretima* sp.

No.	Isolate	Incubation Time	Inhibitory Zone			
			<i>Salmonella typhi</i>		<i>Staphylococcus aureus</i>	
			24 hours	48 hours	24 hours	48 hours
1.	<i>B. choshinensis</i>	5 days	13,09 mm	13,60 mm	12,20 mm	12,42 mm
		10 days	14,24 mm	14,63 mm	13,20 mm	13,57 mm
		15 days	20,05 mm	21,32 mm	15,56 mm	16,88 mm
		20 days	13,40 mm	13,55 mm	13,01 mm	13,15 mm
2.	<i>B. brevis</i>	5 days	11,90 mm	12,60 mm	9,90 mm	10,10 mm
		10 days	9,97 mm	10,40 mm	10,20 mm	10,31 mm
		15 days	15,13 mm	15,25 mm	12,20 mm	12,40 mm
		20 days	10,70 mm	10,79 mm	10,20 mm	10,35 mm
3.	Amoxicillin	5 days	8,40 mm	9,05 mm	13,80 mm	13,98 mm
		10 days	8,40 mm	9,05 mm	13,90 mm	13,98 mm
		15 days	8,40 mm	9,05 mm	13,80 mm	13,98 mm
		20 days	8,40 mm	9,05 mm	13,80 mm	13,98 mm
4.	Chloramphenicol	5 days	25,00 mm	25,10 mm	13,09 mm	13,10 mm
		10 days	25,00 mm	25,10 mm	14,11 mm	14,50 mm
		15 days	25,20 mm	25,45 mm	14,15 mm	14,50 mm
		20 days	25,20 mm	25,45 mm	14,15 mm	14,50 mm

The results showed that the inhibitory zone around chloramphenicol was larger than those around bacteria isolated from *Pheretima* sp. worms (Table 2). This can occur because chloramphenicol is an antibiotic that is commonly used in the treatment of typhoid and contains active compounds that have been tested. However, the use of chloramphenicol is limited because it can damage the ribosomes of mitochondria in mammalian cells [19]. As with several compounds utilized in medicine that have a nitro group component, chloramphenicol may be toxic to animals and plants (Madigan *et al.*, 2012).

Based on observations of inhibitory activity, the test showed that the bacterial inhibition isolated from *Pheretima* sp. earthworms (*Bacillus choshinensis* and *Bacillus brevis*) was capable of inhibiting bacterial growth after a 24-h incubation. After incubation for 48 h, the diameter of the inhibitory

zone had increased from the from the first 24 h. According to Madigan *et al.*, (2012), when the inhibition zone formed after an incubation period of 48 h remains the same or expansion of the inhibition zone diameter occurs, then the bacteria could be described as bactericidal. This is the case with bacteria isolated from *Pheretima* sp.

This result shows that incubation for 15 days effectively assessed inhibition of the growth of pathogenic bacteria, marked by the largest inhibition zone. These results agree with Madigan *et al.*, (2012), that secondary metabolites (antimicrobial) are produced by microorganisms in the late stationary phase of growth. This is because secondary metabolites are usually synthesized by the end of the growth cycle of the cell, namely the stationary phase, as the population remains stable because the number of cells that grow equal the

**Table 3.** Comparison of Isolated bacteria that was Isolated form Various Earthworm

Earthworm	Location	Identification Method	Isolated Bacteria	References
<i>Dendrobaena veneta</i> (Annelida)	Poland	Biochemical identification (API 20 E Bio-Merieux)	<i>Raoultella ornithinolytica</i>	Fiolka <i>et al.</i> (2010) [15]
<i>Libyodrilus violaceus</i> (Annelida)	Nigeria	Biochemical identification (Manual)	<i>Alcaligans faecalis</i> , <i>Bacillus brevis</i> , <i>Bacillus ceveus</i> , <i>Bacillus lalerosporus</i> , <i>Bacillus licheniformis</i> , <i>Bacillus maceraus</i> , <i>Bacillus cereus</i>	Owa <i>et al.</i> (2013)[24]
<i>Lampito mauritii</i> (Annelida)	India	16S rRNA sequencing		Biswas <i>et al.</i> (2014)[25]
<i>Lumbricus terrestris</i> (Annelida)	Ireland	Automated ribosomal intergenic spacer analysis of 16S and 23S genes	Proteobacteria, Firmicutes and an actinobacterium	Thakuria <i>et al.</i> (2010)[26]
<i>Perionyx excavatus</i> (Annelida)	India	Biochemical identification (Manual)	<i>Bacillus</i> , <i>Staphylococcus</i> , <i>Enterococci</i> , <i>Micrococcus</i> , <i>Enterobacter</i> , and <i>Citrobacter</i>	Samanta & Das (2016)[27]
<i>Eisenia fetida</i> (Annelida)	China	16S rRNA sequencing	<i>Bacillus</i> , <i>Balneimonas</i> , <i>Cupriavidus</i> , <i>Hylemonella</i> , <i>Kaistobacter</i> , <i>Lysobacter</i> , <i>Thermomonas</i> , <i>Flaviumibacter</i> , <i>Flavisolibacter</i> , <i>Pseudomonas</i> , <i>Stenotrophomonas</i> , and <i>Achromobacter</i>	Ma <i>et al.</i> (2017)[28]
<i>Aporrectodea caliginosa</i> (Annelida)	Russia	16S rRNA sequencing	<i>Aeromonas encheleia</i> , <i>Bacillus thuringiensis</i> , <i>Acinetobacter</i> sp., <i>Aeromonas</i> sp. and <i>Streptomyces</i> sp.	Tikhonov <i>et al.</i> (2016)[29]

number of cells that die. The synthesis of secondary metabolites is initiated when some of the nutrients in the microbial growth medium have been exhausted. Limitation of these nutrients causes the accumulation of secondary metabolites, enzyme inducers.

The observation that 15 days is the optimum incubation time for bacteria in this research to inhibit pathogenic bacteria is related to secondary metabolites. One of the factors that affects the production of secondary metabolites is the duration of growth of the bacteria. An indicator of the optimum time of production of anti-microbials is the time at which anti-microbial compounds produced are maximally characterized by the formation of the largest inhibition zone at a certain incubation time on the growth of gram-negative bacteria *S. thypi* and gram-positive bacteria *S. aureus*.

Based on the bacteria that have been isolated from *Pheretima* sp. earthworms, there are antibiotics that can inhibit the growth of pathogenic bacteria because they produce a large inhibition zone. This is in accordance with the study of Fiolka *et al.* (2010), who showed that a bacterium isolated from the midgut of earthworms has antimycobacterial activity and has shown activity against four strains of fast-growing mycobacteria. According to Li *et al.* (2011) and Fiolka *et al.* (2010), microbial soil can routinely produce antibiotics through a chemical process called secondary metabolism. It is believed that the production of antibiotics by microorganisms in the soil increases when they are likely to compete with other microbes in soil. This occurs in the form of competition for space or food (Omena *et al.*, 2012).

In this study, two bacterial isolates from *Pheretima* sp. worms namely *Bacillus choshinensis* and *Bacillus brevis*. Table 1 shows various types of bacteria that was isolated by several studies from various worms. Table 1 shows the different types of worms that have different endosymbiont bacteria. Endosymbiont bacteria that can be isolated by several studies dominated by the genus *Bacillus* (Owa *et al.*, 2013; Biswas *et al.*, 2014; Samanta and Das, 2016; Ma *et al.*, 2017; Tikhonov *et al.*, 2016). These results are in accordance with a recent study succeeded in isolating two species of bacteria of the genus *Bacillus*. This suggests that genus *Bacillus* is a genus generally be endosymbiont bacteria in the earthworm. According to Rahman *et al.* (2014), the genus *Bacillus* is commonly found in soil bacteria that is often found as endosymbiont on earthworm.

Two bacterial isolates from *Pheretima* sp. worms

namely *Bacillus choshinensis* and *Bacillus brevis* were shown to inhibit the growth of the bacteria *S. thypi* and *S. aureus*. Therefore, detailed studies of the role of bacterial *Bacillus choshinensis* and *Bacillus brevis* on the lives of worms *Pheretima* sp. worms should be further investigated to uncover benefits and reasons why it is endosymbiont with the worm *Pheretima* sp.

## CONCLUSION

Based on the test results of isolation and inhibition of *Pheretima* sp. earthworms, it was concluded that *Bacillus choshinensis* and *Bacillus brevis* effectively inhibited the growth of the bacteria *S. thypi* and *S. aureus*, and they have the potential to act as antibiotics. Isolates *Bacillus choshinensis* had the largest inhibitory zone diameter of 21.32 mm against *S. thypi* after 15 days of incubation. Inhibitory properties of all isolates that were isolated from *Pheretima* sp. indicate that bacteria could be described as bactericidal.

## Significance Statements

As reported in earlier studies, earthworms are used as drugs, not only the compound earthworms but also endosymbiont microorganisms in earthworms suspected of having antibiotic activities. Therefore, further screening of the potential bacterial endosymbionts contained in the worm *Pheretima* sp. Our findings revealed two bacterial isolates from *Pheretima* sp. worms namely *Bacillus choshinensis* and *Bacillus brevis*. It showed significant inhibitory activities as inhibitors of the growth of *Salmonella thypi* and *Staphylococcus aureus*. We postulate that *Bacillus choshinensis* and *Bacillus brevis* that was isolated from *Pheretima* sp. earthworm have great potentials to become a source of therapeutic agent for pathogenic bacterial infection.

## ACKNOWLEDGEMENT

The authors gratefully acknowledge to Department of Biology, Faculty of Mathematics and Natural Sciences, Hasanuddin University.

## Conflict of interest

No potential conflict of interest was reported by the authors.

## REFERENCES

- Ali, Rahman, M.M., Rahman, A., Basaglia, M., Rahman, M.M., Sultana, T. and Casella, S. 2014. Isolation of

- Bacillus* spp. from Soil and an Evaluation of Their Sensitivity towards Different Extracts and Essential Oils of Cumin (*Cuminum cyminum* L.). *J. Agr. Sci. Tech.* 16: 623-633.
- Biswas, S., Lahiri, P. and Das, S. 2014. Isolation of predominant bacterium from gut of earthworm *Lampito mauritii* for effective use in soil fertility. *Current Science.* 107(1) : 105-109.
- Bohlen. 2001. Earthworm. *Encyclopedia of Soil Science.* USA: Archboid Biological Station.
- Brito-Vega, H. and Espinosa-Victoria, D., 2009. Bacterial diversity in the digestive tract of earthworms (Oligochaeta). *Journal of Biological Sciences.* 9 : 192-199.
- Byzov, B.A., Khomyakov, N.V., Kharin, S.A., Kurakov, A.V., 2007. Fate of soil bacteria and fungi in the gut of earthworms. *European Journal of Soil Biology.* 43 : 149-156.
- Cappuccino J. G., dan Sherman N. 2011. Microbiology a Laboratory Manual. Ed. 9. San Francisco. Benjamin Cummings.
- Carnovale, D., Baker, G., Bissett, A. and Thrall, P. 2015. Earthworm composition, diversity and biomass under three land use systems in south-eastern Australia. *Applied Soil Ecology.* 88 : 32-40.
- Chachina S.B., Voronkova N.A., Baklanova O.N. 2016. Biological remediation of the petroleum and diesel contaminated soil with earthworms *Eisenia fetida*. *Procedia Engineering.* 152 : 122-133.
- Cho, J.H., Park, C.B., Yoon, Y.G., dan Kim S.C. 2014. Lumbricin I, a novel praline-rich antimicrobial peptide from the Earthworm :purification,cDNA cloning and molecular characterization. *Biochim Biophys Acta.* 1408 : 67-76.
- Choi, J., Oh, H., Koo, B.J., Ryu, J. and Lee, S. 2011. Spatial polychaeta habitat potential mapping using probabilistic models. *Estuarine, Coastal and Shelf Science.* 93 : 98-105.
- Fiege, D., Patricia, A., Ramey, Ebbe, B. 2010. Diversity and distributional patterns of Polychaeta in the deep South Atlantic. *Deep-Sea Research I.* 57 : 1329-1344.
- Fiolka, M.J., Zagaja, M.P., Piersiak, T.D., Wróbel, M. and Pawelec, J. 2010. Gut bacterium of *Dendrobaena veneta* (Annelida: Oligochaeta) possesses antimycobacterial activity. *Journal of Invertebrate Pathology.* 105 : 63-73.
- Hada, N., Shida, Y., Shimamura, H., Sonoda, Y., Kasahara, T., Sugita, M. and Takeda, T. 2008. Synthetic studies on glycosphingolipids from Protostomia phyla: syntheses and biological activities of amphoteric glycolipids containing a phosphocholine residue from the earthworm *Pheretima hilgendorfi*. *Carbohydrate Research.* 343 : 2221-2228.
- Hanekamp, J.C. and Bast, A. 2015. Antibiotics exposure and health risks: Chloramphenicol. *Environmental Toxicology and Pharmacology.* 39 : 213-220.
- Klarica, J., Kloss-Brandstätter, A., Traugott, M. and Juen, A. 2012. Comparing four mitochondrial genes in earthworms : Implications for identification, phylogenetics, and discovery of cryptic species. *Soil Biology & Biochemistry.* 45 : 23-30.
- Leveque, T., Capowiez, Y., Schreck, E., Mazzia, C., Auffan, M., Foucault, Y., Austruy, A. and Dumat, C. 2013. Assessing ecotoxicity and uptake of metals and metalloids in relation to two different earthworm species (*Eisenia hortensis* and *Lumbricus terrestris*). *Environmental Pollution.* 179 : 232-241.
- Li, W., Li, S., Zhong, J., Zhu, Z., Liu, J. and Wang, W. 2011. A novel antimicrobial peptide from skin secretions of the earthworm, *Pheretima guillelmi* (Michaelson). *Peptides.* 32 : 1146-1150.
- Ma, L., Xie, Y., Han, Z., Giesy, J.P. and Zhang, X. 2017. Responses of earthworms and microbial communities in their guts to Triclosan. *Chemosphere.* 168 : 1194-1202.
- Madigan, M.T., Martinko, J.M., Sthal, D.A. and Clark, D.P. 2012. Brock, Biology of Microorganisms, Thirteenth Edition. Pearson Education. San Francisco.
- Metcalfe, K.N. and Glasby, C.J. 2008. Diversity of Polychaeta (Annelida) and other worm taxa in mangrove habitats of Darwin Harbour, northern Australia. *Journal of Sea Research.* 59 : 70-82.
- Monari, M., Foschi, J., Cortesi, P., Rosmini, R., Cattani, O. Serrazanetti, G.P. 2008. Chloramphenicol influence on antioxidant enzymes with preliminary approach on microsomal CYP1A immunopositive-protein in *Chamelea gallina*. *Chemosphere.* 73 : 272-280.
- Omena, E.P., Lavrado, H.P., Paranhos, R. and Silva, T.A. 2012. Spatial distribution of intertidal sandy beach polychaeta along an estuarine and morphodynamic gradient in an eutrophic tropical bay. *Marine Pollution Bulletin.* 64 : 1861-1873.
- Owa, S. O., Olowoparija, S.B., Aladesida, A. and Dedeke, G.A. 2013. Enteric bacteria and fungi of the Eudrilid earthworm *Libyodrilus violaceus*. *African Journal of Agricultural Research.* 8(17): 1760-1766.
- Pollack, R.A, Findlay, L., Mondschein, W. and Modesso, R.R. 2009. Laboratory Exercises in Microbiology Third edition. United States of America
- Samanta, T.T. and Das, A. 2016. Isolation, identification, and characterization of gut microflora of *Perionyx excavatus* collected from Midnapore, West Bengal. *Journal of Basic Microbiology.* 56 : 286-293.
- Singh, S., Singh, J. and Vig, A.P. 2016. Effect of abiotic factors on the distribution of earthworms in different land use patterns. *The Journal of Basic & Applied Zoology.* 74 : 41-50.
- Tikhonov, V., Zavgorodnyaya, J., Demin, V. and Byzov, B. 2016. Transformation of soil humic acids by *Aporrectodea caliginosa* earthworm: Effect of gut fluid and gut associated bacteria. *European Journal of Soil Biology.* 75 : 47-53.
- Verma, M.K., Xavier, F., Verma, Y.K. and Sobha, K. 2013. Evaluation of cytotoxic and anti-tumor activity of partially purified serine protease isolate from the Indian earthworm *Pheretima posthuma*. *Asian Pacific Journal of Tropical Biomedicine.* 3 : 896-901.

# ENDOSYMBIONT BACTERIA OF PHERETIMA SP. EARTHWORMS (ANNELIDA: OLIGOCHAETA) POSSESSES ANTIBACTERIAL ACTIVITY

---

## ORIGINALITY REPORT

---

% **6**

SIMILARITY INDEX

% **1**

INTERNET SOURCES

% **5**

PUBLICATIONS

% **3**

STUDENT PAPERS

---

## PRIMARY SOURCES

---

**1**

Sharanpreet Singh, Jaswinder Singh, Adarsh Pal Vig. "Effect of abiotic factors on the distribution of earthworms in different land use patterns", *The Journal of Basic & Applied Zoology*, 2016

Publication

% **1**

**2**

M. Monari, J. Foschi, P. Cortesi, R. Rosmini, O. Cattani, G.P. Serrazanetti. "Chloramphenicol influence on antioxidant enzymes with preliminary approach on microsomal CYP1A immunopositive-protein in *Chamelea gallina*", *Chemosphere*, 2008

Publication

% **1**

**3**

[centaur.vri.cz](http://centaur.vri.cz)

Internet Source

<% **1**

**4**

[www.jove.com](http://www.jove.com)

Internet Source

<% **1**

**5**

M.A. Haxhiu, A.D. Loewy. "Central connections

of the motor and sensory vagal systems innervating the trachea", Journal of the Autonomic Nervous System, 1996

Publication

<% 1

6

Submitted to University of Warwick

Student Paper

<% 1

7

www.omicsonline.org

Internet Source

<% 1

8

apjtb.com

Internet Source

<% 1

9

Vladimir Tikhonov, Julia Zavgorodnyaya, Vladimir Demin, Boris Byzov. "Transformation of soil humic acids by Aporetodea caliginosa earthworm: Effect of gut fluid and gut associated bacteria", European Journal of Soil Biology, 2016

Publication

<% 1

10

Chenglong Ma, Dongdong Gu, Donghua Dai, Wenhua Chen, Fei Chang, Pengpeng Yuan, Yifu Shen. "Aluminum-based nanocomposites with hybrid reinforcements prepared by mechanical alloying and selective laser melting consolidation", Journal of Materials Research, 2015

Publication

<% 1

11

Submitted to Hoggard High School

Student Paper

<% 1

12

Submitted to Canakkale Onsekiz Mart  
University

Student Paper

<% 1

13

Submitted to Chulalongkorn University

Student Paper

<% 1

14

A. Prysor Williams, Paula Roberts, Lisa M.  
Avery, Ken Killham, David L. Jones.

"Earthworms as vectors of Escherichia coli  
O157:H7 in soil and vermicomposts", FEMS  
Microbiology Ecology, 2006

Publication

<% 1

15

Submitted to National University of Singapore

Student Paper

<% 1

16

Submitted to CSU, San Marcos

Student Paper

<% 1

17

Anurag Gupta, Pushpa V. Hazarey, Om P.  
Kharbanda, Virender S. Kohli, Amit Gunjal.

"Stress distribution in the temporomandibular  
joint after mandibular protraction: A 3-  
dimensional finite element study. Part 2",  
American Journal of Orthodontics and  
Dentofacial Orthopedics, 2009

Publication

<% 1

18

Armin Tschermak. "Über Veränderung der  
Form, Farbe und Zeichnung von Kanarieneiern  
durch Bastardierung", Pflügers Archiv -

<% 1

# European Journal of Physiology, 10/1912

Publication

---

---

EXCLUDE QUOTES ON

EXCLUDE  
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5  
WORDS