

Antiinflammation effect of
garciana mangostana pericarp
extract in albino mice skin
induced by 12-0-
tetradecanoylphorbol-13-acetate
by Khairuddin Djawad

FILE	JURNAL_KIRANA_PRASANTY_MOKOAGOW_2020.PDF (502.54K)		
TIME SUBMITTED	19-DEC-2020 12:14PM (UTC+0700)	WORD COUNT	4446
SUBMISSION ID	1479144127	CHARACTER COUNT	24940



Antiinflammation effect of *garciana mangostana* pericarp extract in albino mice skin induced by 12-O-tetradecanoylphorbol-13-acetate

Kirana Prasanty Mokoagow^{1*}, Khairuddin Djawad¹, Siswanto Wahab¹,
Burhanuddin Bahar², Upik Andriani Miskad³, Nurelly Waspodo¹

¹ Department of Dermatology and Venereology, Faculty of Medicine, Hasanuddin University, INDONESIA

² Department of Public Health, Faculty of Public Health, Hasanuddin University, INDONESIA

³ Department of Pathology and Anatomical, Faculty of Medicine, Hasanuddin University, INDONESIA

*Corresponding author: kiranakiranaw@gmail.com

Abstract

Background: Mangosteen (*Garcinia mangostana* Linn.) is a tropical fruit that is most commonly found in Southeast Asia. Mangosteen skin has been reported to contain various beneficial bioactive components such as anti-inflammatory effects. The skin not only serves as a physical and chemical barrier but also as a competent immune system that elicits both innate and adaptive immune responses to pathogens or other inflammatory damages. 12-O-tetradecanoylphorbol-13-acetate (TPA) is a protein kinase activator that induces inflammatory skin reactions. In this study, we investigated the effect of mangosteen pericarp extract as an inhibitory agent for acute inflammatory responses on TPA-induced albino mice skin inflammation.

Materials and Methods: Hematoxylin & Eosin staining histopathology was used to analyze neutrophil infiltration, epidermal thickness, edema, and vascular permeability.

Results: *Garcinia mangostana* pericarp extract cream showed reduced TPA-induced neutrophil infiltration and epidermal thickness. The best concentration is both at 20% and 10%, respectively.

Conclusion: Our study suggests that the naturally derived *Garcinia mangostana* pericarp extract cream provides an acute anti-inflammatory effect on the skin.

Keywords: *garciana mangostana*, histopathology, TPA-induced acute inflammation

Mokoagow KP, Djawad K, Wahab S, Bahar B, Miskad UA, Waspodo N (2020) Antiinflammation effect of *garciana mangostana* pericarp extract in albino mice skin induced by 12-O-tetradecanoylphorbol-13-acetate. Eurasia J Biosci 14: 5815-5821.

© 2020 Mokoagow et al.

This is an open-access article distributed under the terms of the Creative Commons Attribution License.

INTRODUCTION

Natural products are traditionally utilized as a therapeutic agent against many conditions. *Garcinia Mangostana* (GM) is one of those naturally derived medications that have been used to treat gastrointestinal infection, urinary tract infection, anti-scorbutic, laxative, and as an antipyretic. In recent times, mangosteen is believed to be able to treat an assortment of infectious diseases, such as inflammation, diarrhea, abdominal pain, fever, acne, food allergy, and arthritis. (Ovalle-Magallanes et al., 2017, Maione et al., 2016) Secondary metabolites from mangosteen, especially on the skin, have been reported to possess extensive pharmacological properties, such as anti-inflammatory, antioxidant, anti-proliferative, anti-microbial, anti-cancer, hepatoprotective, cardioprotective, and neuroprotective on Alzheimer's disease. (Pedraza-Chaverri et al., 2008, Chen et al., 2018) In dermatological use, a slew of inflammatory conditions, both acute and chronic, requires medication associated with several side effects.

Mangosteen, also known as *Garcinia mangostana* Linn. is a tropical plant belonging to the *Clusiaceae* family that has been widely cultivated as food in Southeast Asia (Indonesia, Malaysia, Thailand, the Philippines, and India). This fruit, known as the "Queen of fruits", consists of white-colored fruit divided into septa and covered by dark purple skin. The mangosteen tree grows very slowly and is in the shape of a pyramidal crown. The tree can reach a height of between 20 and 82 feet (6-25 m), dark brown or almost black peeling bark with the inner bark containing a lot of rubbery, yellow, and bitter sap. The leaves are green with short stalks. Mangosteen skin is reported to contain various bioactive components that have potential as therapeutic agents such as xanthenes, tannins, flavonoids, saponins, quinines, and other bioactive components. (Anwar et al., 2016, Suttirak et al., 2014) *Xanthone*, as a

Received: January 2020

Accepted: April 2020

Printed: November 2020

Significant component is a hydrophobic compound composed of a tricyclic aromatic ring system, which is a structure of isoprenyl, hydroxyl, and methoxy substitution. α -mangostin and γ -mangostin are the most common xanthone found and has the most prominent therapeutic role. Other xanthones include β -mangostin, gartanin, 8-deoxygartanin, garcinones A, B, C, D and E, mangostinone, 9-hydroxycalabaxanthone, and isomangostin consistently show that xanthones have antioxidant, anti-proliferative, proapoptotic, antimicrobial, anti-cancer, and anti-inflammatory activities. (Guzmán-Orozco et al., 2013)

Acute inflammation is the initial immune response to maintain tissue homeostasis against trauma or infection by eliminating potential pathogens. (Loynes et al., 2018) The inflammatory process in the skin occurs due to a tissue reaction that elicits cell defense mediators and proteins to the site of infection and damaged tissue. (Abbas et al., 2015) Inflammatory mediators such as cyclooxygenase-2 (COX-2), interleukin-1 beta (IL-1 β), interleukin-6 (IL-6), and tumor necrosis factor-alpha (TNF- α) play a role in inflammatory diseases. (Lee et al., 2012). The release of inflammatory mediators such as nitric oxide and prostaglandins also increases the permeability of blood vessels, allowing the migration of leukocytes, especially neutrophils, to inflamed tissues. Cyclooxygenase 2 (COX-2) is an enzyme that regulates the prostaglandin/eicosanoid pathway induced by proinflammatory cytokines in an NF- κ B-dependent manner. (Wei et al., 2011)

Phorbol ester, 12-O-tetradecanoylphorbol-13-acetate (TPA) also known as tetradecanoyl phorbol acetate, or phorbol 12-myristate 13-acetate (PMA) is a protein kinase-C (PKC) activator. TPA is a potent phorbol diester and tumor promoter often used in biomedical research. TPA is also used to induce other

biological effects in low concentrations such as skin irritation (Madsen et al., 2016)

We conducted a study assessing the anti-inflammatory effect of *Garcinia mangostana* Linn skin extract on TPA-induced mice skin. The histopathological analysis will be conducted based on neutrophil infiltration, epidermal thickness, edema, and increased tissue permeability.

MATERIAL AND METHODS

GM skin extract was made in the Phytochemical Pharmacognosy Laboratory of the Hasanuddin University Faculty of Pharmacy. The method used is the maceration method with 96% ethanol as a solvent, a ratio of 1:2, and evaporation time of 20 hours. GM skin extract cream was formulated and made at the Laboratory of the Faculty of Pharmacy, University of Surabaya, Surabaya, East Java. The TPA substance used is made by Sigma-Aldrich, USA. The empirical formula for TPA is C₃₆H₅₆O₈, with a molecular weight of

Table 1. Anti-inflammatory effect of GM skin extract

Component	Concentration	Mean (SD)	
		Neutrophil Infiltration	Epidermal Thickness
Control	-	1,40 (0,548)	56,571 (23,766)
TPA	-	3,40 (0,548)	107,907 (48,734)
TPA + Base Cream	-	3,80 (0,447)	155,807 (31,573)
TPA + GM	2,5%	2,80 (0,447)	64,830 (25,985)
TPA + GM	5%	2,40 (0,548)	67,994 (11,059)
TPA + GM	10%	1,40 (0,548)	62,994 (18,701)
TPA + GM	20%	1,20 (0,447)	67,591 (11,019)

616.83 g mol⁻¹. Each glass bottle packaging contains 5 mg of TPA, 5 mg of TPA is dissolved in 50 ml of acetone as a solvent to become homogeneous. The dose of TPA required in this study is 2 μ g dissolved in 20 μ l acetone.

Animal

The Samples used were female albino mice, aged 6-9 weeks, weighing 20-30 grams, healthy and derived from one brood. Animals are housed in a veterinary care facility in the veterinary laboratory of Hasanuddin University Faculty of Medicine with a light-dark cycle of 12/12 hours (dark from 8 PM to 8 AM and light from 8 AM to 8 PM). Animals were given free access to their food and water and were left in this state for 1 week prior to the study. Mice were divided into 7 groups. Mice ears were smeared with 2 μ g TPA dissolved in 20 μ l acetone, then 12 hours and 24 hours later after TPA application, the mice ears were applied topical cream base, GM skin extract in four different concentrations, namely 2.5%, 5%, 10% and 20%.

Group A: The control group was 5 mice without treatment.

Group B: 5 mice with TPA applied on the outer ear.

Group C: 5 mice that were applied with base cream after inflammation induced by TPA.

Group D: 5 mice were applied with a 2.5% GM extract cream after inflammation induced by TPA.

Group E: 5 mice were applied with 5% GM extract cream after inflammation induced by TPA.

Group F: 5 mice were applied with 10% GM extract cream after inflammation induced by TPA.

Group G: 5 mice were applied with 20% GM extract cream after inflammation induced by TPA.

The mice were then biopsied 6 hours after the application of the second GM skin extract cream. Indicators of inflammation include neutrophil infiltration in the epidermal dermis, epidermal thickness, edema, and vascular permeability through a microscope (Olympus, Japan). Epidermal thickness was measured using ImageJ software.

Histology

Tissue samples were fixed in a 10% formalin buffer. All tissues were placed in paraffin blocks and cut to a

Table 2. Effect on edema and increased vascular permeability by *Garciana Mangostana* in TPA-induced mice

Group	N	Edema ^a		Asymptotic Significance ^b	Vascular Permeability ^a		Asymptotic Significance ^b
		Yes	No		Yes	No	
Control	5	0	5	0.000	0	5	0.000
TPA	5	5	0		5	0	
TPA+Base Cream	5	5	0		5	0	
TPA+GM 2.5%	5	5	0		5	0	
TPA+GM 5%	5	5	0		5	0	
TPA+GM 10%	5	5	0		5	0	
TPA+GM 20%	5	5	0		5	0	

^aEdema and increased vascular permeability were examined histologically and categorized into scores, namely: loose stroma and erythrocyte extravasation (present), dense stroma and no erythrocyte extravasation (absent)

^bStatistical test was performed with Crosstab. The reading is based on the Chi-Square Tests, where 14 cells (100%) are estimated to have a count less than 5. The minimum expected count is 0.71.

thickness of 4 - 5 μ m. Each part that was cut was deparaffinated with *xylene* and divided to scale with serial alcohol to water then stained with *hematoxylin-eosin* (HE) for standard evaluation using an Olympus CX21 microscope.

Histopathologi⁴¹ preparations were obtained from excisional biopsy, the ear tissue of mice. Each specimen was fixed with a buffer of formalin, placed on a flat plane. The slide was taken⁴⁰ from the middle area then cut perpendicularly with a thickness of 4 μ m then stained with HE.


Statistical Analysis

The data is collected from all the results of the research, then edited, tabulated, and entered into a computer program, then a descriptive and analytic analysis is carried out. The results of the analysis will be presented in tables or graphs accompanied by an explanation using SPSS version 25. A normality test is conducted from the seven treatment groups. Due to the small sample size (5 mice) per group, One way ANOVA and Crosstab tests were used as statistical tests. Kruskal-Wallis test is performed if the data distribution is not normal. The test result is significant if the p-value <0.05.

RESULT

Anti-inflammatory effect of GM extract cream against TPA-induced neutrophil infiltration

Acute inflammatory processes are measured through a set of parameters, such as Increased neutrophil infiltration, epidermal thickness, vascular permeability, and edema. Neutrophil infiltration is calculated using a microscope, with neutrophils measured as scores, such as single neutrophil infiltration/sparse 1-1 cells (score 1), focal neutrophil cell infiltration (score 2), group of neutrophil cell infiltration (score 3), and diffuse neutrophil cell infiltration (score 4). Measurements were carried out 6 hours after interventions had been performed, including the second GM extract application. TPA exposure was found to increase neutrophil infiltration at 12 o'clock position. More neutrophil infiltration was seen after the base cream application. However, GM extract cream application reduces the neutrophil infiltration, and the

**Fig. 1.** Neutrophil infiltration on all groups

greater the concentration, the lower the neutrophil infiltration.

The ability of GM extract cream to reduce neutrophil infiltration is evident in this study, with the best result obtained with a concentration of 20%.

Anti-inflammatory effect of GM extract on epidermal thickness induced by TPA

Epidermal¹⁰ thickness is the first sign of local inflammation. The thickness of the epidermis was measured from the basal cell layer up to the stratum corneum using ImageJ software. The data is then reported in micrometers. Based on the measurement results, the thickness of the epidermis was found to be the thickest in group C with a TPA-induced inflammation and given base cream afterward. Meanwhile, the application of GM skin extract cream is decreased in the GM group with a concentration of 5%, then 20%, 2.5%; the best result was in the group with a concentration of 10%. A concentration of 10% GM skin extract cream had the best anti-inflammatory effect compared to other formulations.

Anti-inflammatory effect of GM extract cream against edema and TPA-induced vascular permeability improvement

Histopathological examination did not show GM extract cream to have any effect on edema and vascular permeability. This can be seen in the Kruskal-Wallis test, where there is no significant difference. Furthermore, Post Hoc test with Mann-Whitney, showed that there




Fig. 2. Epidermal thickness on all groups




Fig. 3. Histopathology of the skin of albino mice with 100x magnification. Slides of Hematoxylin and Eosin staining on normal mouse skin (A), TPA-induced inflammation (B), Base cream applied to the skin of albino mice after being induced to TPA 24 hours prior (C), GM skin extract cream with 2.5% concentration Applied to the skin of albino mice after being induced to TPA 24 hours prior (D), 5% concentration of GM skin extract cream that was applied to the skin of albino mice after being induced to TPA 24 hours prior (E), 10% concentration GM skin extract cream which Applied to the skin of albino mice after being induced to TPA 24 hours prior (F), 20% concentration of GM skin extract cream was applied to the skin of albino mice after being induced to TPA 24 hours prior (G). The picture shows a representative of 7 mice from each group

was no edema and increased vascular permeability in the untreated group. Whereas in the TPA-induced or on groups treated with GM extract cream, edema, and increased vascular permeability were found.

DISCUSSION

The skin is the largest organ of the human body; it forms a protective barrier, which is essential in protecting individuals from the external environment. The skin not only serves as a physical and chemical barrier but also as a competent immune system that elicits both innate and adaptive immune responses to various pathogens and inflammatory damages. Due to the presence of exogenous factors such as foreign pathogens, ultraviolet radiation, and chemical irritants, the innate immune cells (granulocytes, mononuclear

phagocytes, natural killer cells, keratinocytes) elicit various types of immune responses such as (1) release of anti-microbial agents; (2) induction of inflammatory mediators such as cytokines, chemokines, neuropeptides, and eicosanoids; and (3) initiation and modulation of adaptive immune responses. (Xie et al., 2015, Modlin et al., 2012)

Acute inflammation is a rapid response to infection or tissue damage that lasts only for a short time, ranging from a few minutes to several days. Excessive inflammation or inadequate immune response in the skin may cause various skin diseases such as allergic reactions, psoriasis, and skin cancer. (Wei et al., 2011) TPA-induced inflammatory responses in mouse models have been widely used in several studies to test the effectiveness of various drugs or natural substances as anti-inflammatory agents. TPA application results in the recruitment of inflammatory cells and the massive production of proinflammatory cytokines. (Khan et al., 2013) Multiple studies have used TPA to assess the effects of inflammation in mouse models. (Lee et al., 2012, Park et al., 2016, Choi et al., 2009, Wei et al., 2011, Huang et al., 2006, Yasukawa et al., 2009) Epidermis exposure to TPA causes keratinocytes and Langerhans cells to be stimulated to activate protein kinase C (PKC), resulting in the activation of P13K/Akt, Erk and NF- κ B signaling transduction pathways. Akt and NF- κ B play an essential role in activating key inflammatory enzymes such as inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2). One study showed that COX-2 and iNOS expression was detected in the epidermis after TPA stimulation, indicating that COX-2 and iNOS are released from epidermal cells (keratinocytes). (Wei et al., 2011)

Along with the development of science and technology, naturally derived products are continuously being developed. The anti-inflammatory effect of *Garciana mangostana* is one of them. Our study shows that the anti-inflammatory capabilities of *Garciana mangostana* can suppress the TPA-induced inflammation in the skin of a mouse model. *Garciana mangostana*'s anti-inflammatory mechanism of action is by inhibiting the cyclooxygenase (COX) enzyme in the arachidonic acid pathway, which results in decreased inflammation. Xanthone also inhibits inflammation through the NF- κ B signaling cascade. This inhibition is due to the inactivation of cyclic adenosine monophosphate (cAMP) and cyclic guanosine monophosphate (cGMP) cyclase in vascular endothelial cells due to the absence of prostanoid production, thus inhibiting inflammatory reactions. Other studies have found that mangosteen contains several identical components as nonsteroidal anti-inflammatory drugs, an anti-inflammatory agent that targets the COX enzyme. (Marzaimi and Aizat, 2019)

It is evident in *in vivo* studies that α - and γ -mangostin inhibit histamine release and PGE2 induction. Previous

in vitro studies [13](#) showed that *Garciana mangostana* L. could inhibit COX-1 and COX-2. (Nakatani et al., [22](#)) Furthermore, α -mangostin shows effectiveness as an antimetastatic agent against the expression of TPA-induced matrix metalloproteinase-2 (MMP-2) and MMP-9 on [18](#) adenocarcinoma cells A549. Data indicate that α -mangostin can inhibit the activation of α v β 3 integrin, focal adhesion kinase (FAK), extracellular signal-regulated kinase $\frac{1}{2}$ (ERK $\frac{1}{2}$); all of these are involved in the downregulating activity of enzymes and NF κ B. (Shih et al., 2010) In vivo effect of isogarcinol (mangosteen) using xylene on acute inflammation in the animal model [31](#) in the reduction of edema in the ear of the mouse. Oral administration of 100 mg kg⁻¹ of isogarcinol has an aspirin-like effect on xylene-induced ear edema. Meanwhile, in vitro studies have shown that isogarcinol inhibits COX-2 mRNA and iNOS expression; Isogarcinol also suppresses NO production in supernatant RAW 264.7 [24](#) cells. The investigators concluded that isogarcinol significantly reduced serum levels of IL-1 β , TNF- α , IL-6, and IL-17. (Fu et al., 2014) α -mangostin exhibit [7](#) a significant effect in inhibiting lipopolysaccharide production and cytotoxicity in mice stimulated by monocyte macrophage cell line (RAW 264.7 cells). At 3 to 25 μ M of α -mangostin, the amount of NO production was measured [1](#) continuously and the IC50 value was found to be 12.4. Production of lipopolysaccharide-activated PGE2 in RAW 264.7 cells was also significantly reduced by α -mangostin, with [47](#) IC50 value of 11.08 μ M. The concentration of α -mangostin was found to reduce the induction of iNOS. 1 μ g mL⁻¹ lipopolysaccharide was used to activate RAW 264.7 cells for 12 hours, iNO [1](#) activity in activated macrophages was inhibited after treatment with 5 μ g mL⁻¹ α -mangostin in 24 hours. Carrageenan-induced edema in mice was used to evaluate the anti-inflammatory effect of α -mangostin, which showed potent inhibition 3 hours after treatment. (Ibrahim et al., 2016)

Histopathological examination showed that topical administration of TPA induces inflammation, as evidenced by the increase of epidermal thickness, leukocyte infiltration, vascular permeability, and edema. The release of mediators such as nitric oxide and prostaglandins raises vascular permeability, resulting in fluid accumulation [2](#) the vicinity of inflammation (edema) and allows the migration of leukocytes, especially neutrophils, in the inflammatory tissue. Epidermal hyperplasia is the result of increased inflammatory mediators. (Wei et al., 2011) Based on our preliminary

study, the administration of 2 μ g topical TPA has shown increased signs of inflammation at 12 hours after TPA induction as evidenced by increased neutrophil infiltration, vascular permeability, and edema. Inflammatory markers are significantly increased in a linear correlation with more prolonged TPA exposure. Histopathological examination was [10](#) performed using an Olympus microscope, where the thickness of the epidermis was measured vertically from the basal cell layer to the corneum layer using ImageJ software. Neutrophil infiltration was assessed using histopathological scores.

In this study, the group induced by inflammation and given base cream showed higher inflammation than the group that was only exposed to TPA. This may be [21](#) used by the base cream components such as alcohol, propylene glycol, sorbitan oleate esters, cetylalcohol, and isopropyl myristate, all of which have the potential to cause allergic contact dermatitis. (Goossens and Clinics, 2014) Our study showed that GM skin extract cream could inhibit neutrophil infiltration and hamper epidermal thickness in all groups. Meanwhile, based on edema measurements and increased vascular permeability, it appears that GM cream extract did not show any anti-inflammatory effect on both categories.

CONCLUSION

Topical application of GM skin extract cream with concentrations ranging from 2.5%, 5%, 10%, and 20% can inhibit TPA-induced inflammatory reactions in the skin of mice. The topical anti-inflammatory effect of GM skin extract cream is evident by the decrease in neutrophil infiltration and epidermal thickness. In comparison, edema and increased vascular permeability were unchanged. The group treated with a topical administration of 20% GM skin extract cream exhibit the most significant effect on neutrophil infiltration. Meanwhile, a concentration of 10% provides an anti-inflammatory effect, as shown by the lowest [37](#) epidermal thickness compared to other groups. The results of this study indicate that GM skin extract cream can be developed as an alternative topical treatment for inflammatory skin diseases.

ACKNOWLEDGEMENTS

We are grateful to all parties for their assistance in the literature search.

REFERENCES

- Abbas, A. K., Lichtman, A. H. & Pillai, S. 2015. Cellular and Molecular Immunology, Philadelphia, ELSEVIER.
- Anwar, A. I., Djawad, K. & Fitri, E. W. J. P. J. O. N. 2016. Effectiveness of Topical Mangosteen Pericarp Extract Against Angiogenesis in Mice Exposed to Ultra Violet B. 15, 745-751.
- Chen, G., Li, Y., Wang, W. & Deng, L. J. E. O. T. P. 2018. Bioactivity and pharmacological properties of α -mangostin from the mangosteen fruit: a review. 28, 415-427.

- Choi, G., Yoon, T., Cheon, M. S., Choo, B. K. & Kim, H. K. 2009. Anti-inflammatory activity of *Chrysanthemum indicum* extract in acute and chronic cutaneous inflammation. *Journal of Ethnopharmacology*, 123, 149-154.
- Fu, Y., Zhou, H., Wang, M., Cen, J., Wei, Q. J. J. O. A. & Chemistry, F. 2014. Immune regulation and anti-inflammatory effects of isogarcinol extracted from *Garcinia mangostana* L. against collagen-induced arthritis. 62, 4127-4134.
- Goossens, R. A. J. I. & Clinics, A. 2014. Allergic contact dermatitis from the vehicle components of topical pharmaceutical products. 34, 663-670.
- Gutierrez-Orozco, F., Chitchumroonchokchai, C., Lesinski, G. B., Suksamrarn, S., Failla, M. L. J. J. O. A. & Chemistry, F. 2013. α -Mangostin: anti-inflammatory activity and metabolism by human cells. 61, 3891-3900.
- Huang, M. T., Liu, Y., Ramji, D., Lo, C. Y., Ghai, G., Dushenkov, S., Ho, C. T. J. M. N. & Research, F. 2006. Inhibitory effects of black tea theaflavin derivatives on 12-O-tetradecanoylphorbol-13-acetate-induced inflammation and arachidonic acid metabolism in mouse ears. 50, 115-122.
- Ibrahim, M. Y., Hashim, N. M., Mariod, A. A., Mohan, S., Abdulla, M. A., Abdelwahab, S. I. & Arbab, I. A. J. A. J. O. C. 2016. α -Mangostin from *Garcinia mangostana* Linn: an updated review of its pharmacological properties. 9, 317-329.
- Khan, A. Q., Khan, R., Qamar, W., Lateef, A., Rehman, M. U., Tahir, M., Ali, F., Hamiza, O. O., Hasan, S. K., Sultana, S. J. E. & Pathology, M. 2013. Geraniol attenuates 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced oxidative stress and inflammation in mouse skin: possible role of p38 MAP Kinase and NF- κ B. 94, 419-429.
- Lee, S. H., Kim, D. W., Eom, S. A., Jun, S.-Y., Park, M., Kim, D.-S., Kwon, H. J., Kwon, H. Y., Han, K. H. & Park, J. J. B. R. 2012. Suppression of 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced skin inflammation in mice by transduced Tat-Annexin protein. 45, 354-359.
- Loynes, C. A., Lee, J. A., Robertson, A. L., Steel, M. J., Ellett, F., Feng, Y., Levy, B. D., Whyte, M. K. & Renshaw, S. A. J. S. A. 2018. PGE2 production at sites of tissue injury promotes an anti-inflammatory neutrophil phenotype and determines the outcome of inflammation resolution in vivo. 4, eaar8320.
- Madsen, M., Hansen, P. R., Nielsen, L. B., Hartvigsen, K., Pedersen, A. E., Christensen, J. P., Aarup, A. & Pedersen, T. X. J. B. D. 2016. Effect of 12-O-tetradecanoylphorbol-13-acetate-induced psoriasis-like skin lesions on systemic inflammation and atherosclerosis in hypercholesterolaemic apolipoprotein E deficient mice. 16, 1-10.
- Maione, F., Russo, R., Khan, H. & Mascolo, N. J. N. P. R. 2016. Medicinal plants with anti-inflammatory activities. 30, 1343-1352.
- Marzaimi, I. N. & Aizat, W. M. 2019. Current review on mangosteen usages in antiinflammation and other related disorders. *Bioactive Food as Dietary Interventions for Arthritis and Related Inflammatory Diseases*. Elsevier.
- Modlin, Miller & Bangert 2012. Innate and Adaptive Immunity in the Skin. In: Goldsmith, Katz, Gilchrist & Leffell (eds.) *Fitzpatrick's Dermatology In General Medicine*. 8 ed.: Mc Graw Hill.
- Nakatani, K., Atsumi, M., Arakawa, T., Oosawa, K., Shimura, S., Nakahata, N. & Ohizumi, Y. 2002. Inhibitions of histamine release and prostaglandin E2 synthesis by mangosteen, a Thai medicinal plant. *Biological and Pharmaceutical Bulletin*, 25, 1137-1141.
- Ovalle-Magallanes, B., Eugenio-Perez, D., Pedraza-Chaverri, J. J. F. & Toxicology, C. 2017. Medicinal properties of mangosteen (*Garcinia mangostana* L.): a comprehensive update. 109, 102-122.
- Park, S.-H., Seo, W., Eun, H. S., Kim, S. Y., Jo, E., Kim, M.-H., Choi, W.-M., Lee, J.-H., Shim, Y.-R., Cui, C.-H. J. B. & Communications, B. R. 2016. Protective effects of ginsenoside F2 on 12-O-tetradecanoylphorbol-13-acetate-induced skin inflammation in mice. 478, 1713-1719.
- Pedraza-Chaverri, J., Cerdenas-Rodriguez, N., Orozco-Ibarra, M., Perez-Rojas, J. M. J. F. & Toxicology, C. 2008. Medicinal properties of mangosteen (*Garcinia mangostana*). 46, 3227-3239.
- Shih, Y.-W., Chien, S.-T., Chen, P.-S., Lee, J.-H., Wu, S.-H., Yin, L.-T. J. C. B. & Biophysics 2010. α -Mangostin suppresses phorbol 12-myristate 13-acetate-induced MMP-2/MMP-9 expressions via α v β 3 integrin/FAK/ERK and NF- κ B signaling pathway in human lung adenocarcinoma A549 cells. 58, 31-44.
- Suttirak, W., Manurakchinakorn, S. J. J. O. F. S. & Technology 2014. In vitro antioxidant properties of mangosteen peel extract. 51, 3546-3558.
- Wei, W.-C., Lin, S.-Y., Chen, Y.-J., Wen, C.-C., Huang, C.-Y., Palanisamy, A., Yang, N.-S. & Sheu, J.-H. J. J. O. B. S. 2011. Topical application of marine briarane-type diterpenes effectively inhibits 12-O-tetradecanoylphorbol-13-acetate-induced inflammation and dermatitis in murine skin. 18, 94.

Xie, Z., Sintara, M., Chang, T., Ou, B. J. F. S. & Nutrition 2015. Daily consumption of a mangosteen-based drink improves in vivo antioxidant and anti-inflammatory biomarkers in healthy adults: a randomized, double-blind, placebo-controlled clinical trial. 3, 342-348.

Yasukawa, K., Iida, T., Fujimoto, Y. J. J. O. P. & Pharmacology 2009. Relative inhibitory activity of bile acids against 12-O-tetradecanoylphorbol-13-acetate-induced inflammation, and chenodeoxycholic acid inhibition of tumour promotion in mouse skin two-stage carcinogenesis. 61, 1051-1056.

www.ejobios.org

Antiinflammation effect of garciana mangostana pericarp extract in albino mice skin induced by 12-0- tetradecanoylphorbol-13-acetate

ORIGINALITY REPORT

%24 SIMILARITY INDEX	%16 INTERNET SOURCES	%21 PUBLICATIONS	%7 STUDENT PAPERS
--------------------------------	--------------------------------	----------------------------	-----------------------------

PRIMARY SOURCES

- 1** Mohamed Yousif Ibrahim, Najihah Mohd Hashim, Abdalbasit Adam Mariod, Syam Mohan et al. " α -Mangostin from Garcinia mangostana Linn: An updated review of its pharmacological properties", Arabian Journal of Chemistry, 2016
Publication **%2**
- 2** www.pubmedcentral.nih.gov
Internet Source **%2**
- 3** www.ncbi.nlm.nih.gov
Internet Source **%1**
- 4** plasticsurgerykey.com
Internet Source **%1**
- 5** preview-jbiomedsci.biomedcentral.com
Internet Source **%1**
- 6** www.mdpi.com
Internet Source **%1**
- 7** Submitted to King Saud University

% 1

8

Ili Natasya Marzaimi, Wan Mohd Aizat. "Current Review on Mangosteen Usages in Antiinflammation and Other Related Disorders", Elsevier BV, 2019

Publication

% 1

9

"Plant-derived Bioactives", Springer Science and Business Media LLC, 2020

Publication

% 1

10

www.frontiersin.org

Internet Source

% 1

11

Submitted to Udayana University

Student Paper

% 1

12

fjfsdata01prod.blob.core.windows.net

Internet Source

% 1

13

repository.unhas.ac.id

Internet Source

% 1

14

Panarat Arunrattiyakorn, Mayuso Kuno, Thammarat Aree, Surat Laphookhieo et al. " Biotransformation of β -Mangostin by an Endophytic Fungus of to Furnish Xanthenes with an Unprecedented Heterocyclic Skeleton ", Journal of Natural Products, 2018

Publication

% 1

15

peerj.com

Internet Source

<% 1

16

www.deepdyve.com

Internet Source

<% 1

17

Manoranjan Sharma, Jogeswar Mohapatra, Anil Argade, Shrikalp S. Deshpande, Gaurang B. Shah, Abhijit Chatterjee, Mukul R. Jain.

"Chemopreventive effect of a novel, selective TACE inhibitor on DMBA- and TPA-induced skin carcinogenesis", Immunopharmacology and Immunotoxicology, 2014

Publication

<% 1

18

Chiung-Man Tsai, Gow-Chin Yen, Fang-Ming Sun, Shun-Fa Yang, Chia-Jui Weng.

"Assessment of the Anti-invasion Potential and Mechanism of Select Cinnamic Acid Derivatives on Human Lung Adenocarcinoma Cells", Molecular Pharmaceutics, 2013

Publication

<% 1

19

Al-Sha'er, Mahmoud A., Iman Mansi, Inas Almazari, and Nancy Hakooz. "Evaluation of novel Akt1 inhibitors as anticancer agents using virtual co-crystallized pharmacophore generation", Journal of Molecular Graphics and Modelling, 2015.

Publication

<% 1

- | | | |
|----|--|------|
| 20 | Advances in Experimental Medicine and Biology, 2014.
Publication | <% 1 |
| 21 | "Contact Dermatitis", Springer Science and Business Media LLC, 2021
Publication | <% 1 |
| 22 | Submitted to Padjadjaran University
Student Paper | <% 1 |
| 23 | Rohman Abdul, Arifah Fitriana Hayyu, Irnawati Irnawati, Alam Gemini, Muchtaridi Muchtaridi, Rafi Muhammad. "A review on phytochemical constituents, role on metabolic diseases, and toxicological assessments of underutilized part of Garcinia mangostana L. fruit", Journal of Applied Pharmaceutical Science, 2020
Publication | <% 1 |
| 24 | insights.ovid.com
Internet Source | <% 1 |
| 25 | www.hospitalpharmacyeurope.com
Internet Source | <% 1 |
| 26 | Khan, Abdul Quaiyoom, Rehan Khan, Wajhul Qamar, Abdul Lateef, Muneeb U Rehman, Mir Tahir, Farrah Ali, Oday O. Hamiza, Syed Kazim Hasan, and Sarwat Sultana. "Geraniol attenuates 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced oxidative stress and | <% 1 |

inflammation in mouse skin: Possible role of p38 MAP Kinase and NF-κB", Experimental and Molecular Pathology, 2013.

Publication

27

bmcdermatol.biomedcentral.com

Internet Source

<% 1

28

www.researchsquare.com

Internet Source

<% 1

29

Submitted to Surabaya University

Student Paper

<% 1

30

eprints.utar.edu.my

Internet Source

<% 1

31

Fu, Yanxia, Hailing Zhou, Mengqi Wang, Juren Cen, and Qun Wei. "Immune Regulation and Anti-inflammatory Effects of Isogarcinol Extracted from *Garcinia mangostana* L. against Collagen-Induced Arthritis", *Journal of Agricultural and Food Chemistry*

Publication

<% 1

32

Nelson, Randy J.. "An Introduction to Behavioral Endocrinology", Oxford University Press

Publication

<% 1

33

Edward B. Walker. "HPLC analysis of selected xanthenes in mangosteen fruit", *Journal of Separation Science*, 06/2007

Publication

<% 1

34

thieme-connect.de

Internet Source

<% 1

35

www.scribd.com

Internet Source

<% 1

36

stormpdf.com

Internet Source

<% 1

37

Kamarza Mulia, Fitria Hasanah, Elsa A Krisanti. "The effect of heating temperature on cytotoxicity and α -mangostin yield: Mangosteen pericarp juice and mangosteen extract", IOP Conference Series: Materials Science and Engineering, 2018

Publication

<% 1

38

Yan Weng, Natsumi Mizuno, Jiangxu Dong, Ryosuke Segawa et al. "Induction of thymic stromal lymphopoietin by a steroid alkaloid derivative in mouse keratinocytes", International Immunopharmacology, 2018

Publication

<% 1

39

worldwidescience.org

Internet Source

<% 1

40

In Jun Yang, Dong Ung Lee, Heung Mook Shin. "Anti-inflammatory and antioxidant effects of coumarins isolated from in lipopolysaccharide-stimulated macrophages and 12- - tetradecanoylphorbol-13-acetate-stimulated

<% 1

mice ", Immunopharmacology and
Immunotoxicology, 2015

Publication

41

Sun Phil Choi, Sung Phil Kim, Mi Young Kang,
Seok Hyun Nam, Mendel Friedman. "Protective
Effects of Black Rice Bran against Chemically-
Induced Inflammation of Mouse Skin", Journal of
Agricultural and Food Chemistry, 2010

Publication

<% 1

42

res.mdpi.com

Internet Source

<% 1

43

www.science.gov

Internet Source

<% 1

44

www.stuartxchange.org

Internet Source

<% 1

45

Jing J. Wang, Qing H. Shi, Wei Zhang, Barbara
J.S. Sanderson. "Anti-skin cancer properties of
phenolic-rich extract from the pericarp of
mangosteen (*Garcinia mangostana* Linn.)",
Food and Chemical Toxicology, 2012

Publication

<% 1

46

Yan-Yun Ding, Jia-Jie Luan, Yan Fan, Opeyemi
Joshua Olatunji, Jing Song, Jian Zuo. " α -
Mangostin reduced the viability of A594 cells in
vitro by provoking ROS production through
downregulation of NAMPT/NAD", Cell Stress

<% 1

47

"Bioactive Molecules in Food", Springer Science and Business Media LLC, 2019

Publication

<% 1

48

Borsato, Débora M., Arthur S. Prudente, Patrícia M. Döll-Boscardin, Aurélio V. Borsato, Cynthia F. P. Luz, Beatriz H. L. N. S. Maia, Daniela A. Cabrini, Michel F. Otuki, Marilis D. Miguel, Paulo V. Farago, and Obdulio G. Miguel. "Topical Anti-Inflammatory Activity of a Monofloral Honey of *Mimosa scabrella* Provided by *Melipona marginata* During Winter in Southern Brazil", *Journal of Medicinal Food*, 2014.

Publication

<% 1

49

Gutierrez-Orozco, Fabiola, and Mark Failla. "Biological Activities and Bioavailability of Mangosteen Xanthones: A Critical Review of the Current Evidence", *Nutrients*, 2013.

Publication

<% 1

50

Masa-Aki Shibata, Mariko Harada-Shiba, Eiko Shibata, Hideki Tosa et al. "Crude α -Mangostin Suppresses the Development of Atherosclerotic Lesions in Apoe-Deficient Mice by a Possible M2 Macrophage-Mediated Mechanism", *International Journal of Molecular Sciences*, 2019

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE ON
BIBLIOGRAPHY

EXCLUDE MATCHES < 5
WORDS