

1-

PB_Dewi_Rahmawati,_Apik_Ind
arty_Moedjiono,_Sukri_Palutturi.
pdf

by

FILE	1- PB_DEWI_RAHMAWATI,_APIK_INDARTY_MOEDJIONO,_SUKRI_PALUTTU RI.PDF (165.18K)		
TIME SUBMITTED	04-JUL-2020 09:47AM (UTC+0700)	WORD COUNT	4492
SUBMISSION ID	1353246110	CHARACTER COUNT	24049

DOI: <http://dx.doi.org/10.33846/hn30706>
<http://heanoti.com/index.php/hn>



RESEARCH ARTICLE

URL of this article: <http://heanoti.com/index.php/hn/article/view/hn30706>

The Effect of Health Center Service and Role of Husband Towards Contraceptive Use in Makassar City, Indonesia

Dewi Rahmawati^{1(CA)}, Api⁸ Indarty Moedjiono², Sukri Palutturi³

¹Postgraduate Program of Reproductive Health, Faculty of Public Health, Universitas Hasanuddin, Indonesia; dewrahma¹⁵95@gmail.com (Corresponding Author)

²Department of Biostatistic, Faculty of Public Health, Universitas Hasanuddin, Indonesia

³Departemen of Administration and Health Policy, Faculty of Public Health, Universitas Hasanuddin, Indonesia

ABSTRACT

Contraceptive use is one way to reduce population growth rates. This research is aim to measure the effect of health center¹⁹ service and husband role on contraception use in makassar city. This research is done at the Tamalanrea Health Center and Kassi-Kassi Health Center, Makassar Regency. The research design use¹⁴ as a cross sectional study. Research the sample was 178 people who were selected using the stratified random sampling technique. Data analyzed using the chi-square test and logistic regression. The result shows that public health center services that are related to¹ use of contraception in makassar city are the choice of contraception (p-value = 0.001), information provided (p-value = 0,000), officer technical ability (p-value = 0,000) and husband-communication wife p-value = 0.007). Information provided is the most influential variable on contraceptive use in the city of makassar with a wald value (42,958)

Keywords: contraception; family planning; health center services; husband's role

INTRODUCTION

Background

Family planning is an attempt to outline or determine the amount and distance of assessment by using contraception, to manifest a small, happy family and prosperous⁽¹⁾ so that²⁶ family planning programs it is possible for everyone to decide freely the distance²⁵ between children, the number of children desired and the time of birth through the use of contraception⁽²⁾. The use of contraception can reduce the risk of maternal death, newborns, the number of unwanted births and¹² abortions and can reduce the poverty chain⁽³⁾. in 2002 up to 2013 the birth¹² in Indonesia was 2.6 children per 1000 women of childbearing age so the target of the birth rate was 2.11 per 1000 women of childbearing age in 2015⁽⁴⁾.

Globally, modern contraceptive use increased, from 54% in 1990 to 57.4% in 2015 for Asia, up from 60.9% to 61.8%⁽⁵⁾. In 2017 the number of PUS (couples of childbearing age) in Indonesia continues to increase every year and there is a decline in long-term contraceptive use and an increase in injectable contraceptive users from 2015 to 2017⁽⁶⁾. The province of South Sulawesi is one of the provinces that has continued to decline in the use of MKJP from 2016-2017. The number of MKJP users includes 4.51% IUDs and 13.66% implants and there has been a decrease in 2017 covering 3.28% IUD and mplan 9.10%⁽⁶⁾. Although continued use of MKJP in Makassar City, the increase has not yet reached the MKJP prevalence target set by the 2015-2019 Strategic Plan, wh⁷ is 60.9% in 2017⁽⁷⁾.

One of the factors²⁴ that determine the use of family planning is the place of contraception service, waiting time, contraception that you want to use and information^(8,9). Husband is someone that participated in the use of the couple contraception through the giving of communications and communications with the partners^(10,11).

Based on early survey results in Kassi-Kassi Health Center and Tamalanrea Health Center is a national accredited in Makassar Ci¹³ but the coverage of contraception use has not reached target resntras that has been set. Based on the profile of the Kassi-Kassi Health Center and the Tamalanrea Health Center, the number of

long-term contraceptive users in the Kassi-Kassi Health Center is 69 acceptors and there has been an increase in long-term contraceptive users from 2017 to 2018 and the Tamalanrea Health Center has decreased long-term contraceptive users by 20 acceptors in 2018⁽¹²⁾.

Service quality and husband's role are the main factors in contraceptive use. Health workers have the duty to improve the quality of care for women in the use of family planning services while the key factor that contributes to contraceptive use is the role of the husband^(13,14).

Purpose

This study aims to look at the influence of health center services and the role of husbands on contraceptive use in the City of Makassar, Indonesia.

METHODS

Study Design and setting

The observational study with Cross-Sectional design was carried out in the work area of the Cassium Health Center and Tamalanrea Health Center, Makassar City, South Sulawesi in April - May 2019.

Population and sample

The population in this study were women of childbearing age (WUS) who used contraceptive services at the Kassi-kassi Health Center and Tamalanrea Health Center. Sample size determination is based on. The study sample consisted of 178 contraceptive acceptors selected using the stratified random sampling technique, so that the number of samples obtained in each public health center was 89 acceptors.

Data collection and variables

Data was collected using a structured questionnaire. The enumerator was first given training related to the procedures to be carried out in this study. Questionnaires were given to the respondent's homes which were selected based on data obtained from the contraception user registration book. Contraceptive use is divided into 2 receptors that use injection contraception and acceptors who use long-term contraception. Health Center services are the role of health workers in terms of contraceptive choices, information provided, interpersonal relationships, technical officers' abilities and follow-up (based on Jain theory) while the related roles consist of those who provide assistance to those who receive assistance and communication with their husbands.

Data analysis

Data was taken for suitability and completeness and code and included in Microsoft Excel and SPSS Version 20. Analysis was performed to see the frequency distribution of respondents' characteristics. Bivariate and multivariate analysis was performed to see the relationship of each independent variable to the dependent variable. To measure the relationship between independent and dependent variables with a 95% confidence interval. Only socio-demographic variables that have a value of $p < 0.05$ are included in the multivariate regression logistic model. The final model obtained retaliates with the Hosmer and Lemeshow Test.

RESULTS

Characteristics of Respondents

Table 1. shows a total of 178 respondents consisting of 111 acceptors who use contraceptive services at the Kassi-Kassi Health Center and Tamalanrea Health Center. Based on age, the majority of respondents were 31-35 years old, 27 (15.2%) respondents. Based on the latest education, the highest number of respondents was in the last level of education D1-D3 of 51 (28.7%) respondents while the least was in the junior secondary education level of 33 (18.5%) respondents. Based on the age of married respondents, the age of marriage with the oldest age is at the age of 31-35 years as many as 10 (5.6%) respondents and the highest age of marriage at the age of 21-25 years as many as 83 (46.6%) respondents. Based on the reasons for contraceptive use, the most reason for respondents to maintain a pregnancy distance of 99 (55.6%) respondents while respondents with a reason to postpone pregnancy was 79 (44.4%) respondents. Based on the history of previous contraceptive use,

respondents with the most history of contraceptive use were not replacing contraceptives used as many as 141 respondents (79.2%) and the history of not using contraception was 2 respondents (1.1%). Based on the number of children desired, as many as 43 (24.2%) respondents had children who were not ideal and as many as 135 (75.8%) had children with ideal numbers.

Table 1. Description of the respondents' socio-demographic characteristics

Variable	Respondents (n = 178)	%
Age		
• 15-20	4	2,2
• 21-25	27	15,2
• 26-30	44	24,7
• 31-35	35	19,07
• 36-40	39	21,9
• 41-45	24	13,5
• 46-50	5	2,8
Educational level		
• SLTP	33	18,5
• SMA	49	27,5
• D1 – D3	51	28,7
• S1-S2	37	20,8
Married age		
• 15-20	43	24,2
• 21-25	83	46,6
• 26-30	42	23,6
• 31-35	10	5,6
Reasons for using KB		
• Delaying pregnancy	79	21,9
• Maintain pregnancy distance	99	22,5
KB usage history		
• Pill	20	11,2
• Injection	9	5,1
• MKJP	6	3,4
• Do not use contraception	2	1,1
• Haven't changed the method	141	79,2
Number of children desired		
• Not Ideal	43	24,2
• Ideal	135	75,8
Total	178	100

Effect Of Health Center Service And Husband's Role On Contraceptive Use

Table 2 shows the service variables and the role of the husband include contraceptive choices that injection contraceptive users as many as 38 (42.7%) respondents stated that contraceptive choices were not available and 51 (57.3%) respondents stated that contraceptive choices were available, whereas for contraceptive users Long term (MKJP) of 17 (19.1%) respondents stated that contraceptive choices that respondents wanted to use were not available and as many as 72 (80.9%) respondents stated that the choice of contraception respondents wanted to use was always available. In the information provided, injecting contraceptive users as many as 71 (79.8%) respondents stated that the information provided was lacking and 18 (20.2%) respondents stated that the information provided was sufficient, whereas for long-term contraceptive users (MKJP) there were 15 (16.9%) respondents stated that the information provided was less and as many as 74 (83.1%) respondents stated that the information provided was sufficient. In interpersonal relationships, respondents stated that interpersonal relationships were built less and 47 (52.8%) respondents stated that interpersonal relations were built sufficiently, while in long-term contraceptive users (MKJP) 45 (50.6%) respondents stated that interpersonal relations which was built less and as many as 44 (49.4%) respondents stated that the interpersonal relations provided were sufficient. In the technical abilities of the officers,

respondents stated that the technical abilities of officers were bad and 38 (42.7%) respondents stated that the technical abilities of officers were good, whereas in long-term contraceptive users (MKJP) 10 (11.2%) respondents stated that the technical capabilities of officers bad and as many as 79 (88.8%) respondents stated the technical capabilities of the officers were good. In the follow-up of injecting contraceptive users as many as 53 (59.6%) respondents stated that follow-up or advanced service mechanisms were poor technical officers and 36 (40.4%) respondents stated that the technical abilities of officers were good, while those with term contraception length (MKJP) of 44 (49.4%) respondents stated that the technical abilities of officers were poor and as many as 45 (50.6%) respondents stated that the technical abilities of officers were good.

Table 2. Effect of health center services and the role of husbands on contraceptive use

Variable	Contraceptive				p-value
	Injection (89)		MKJP (89)		
	n	%	n	%	
Health center services					
Choice of contraception					
• Available	38	42.7	17	19.1	0.001
• Not available	51	57.3	72	80.9	
Information provided					
• Less	71	79.8	15	16.9	0.000
• Enough	18	20.2	74	83.1	
Interpersonal relations					
• Less	42	47.2	45	50.6	0.764
• Well	47	52.8	44	49.4	
Officer technical capability					
• Bad	51	57.3	10	11.2	0.000
• Well	38	42.7	79	88.8	
<i>Follow-up</i>					
• Bad	53	59.6	44	49.4	0.229
• Well	36	40.4	45	50.6	
Husband's role					
Husband's Agreement					
• Less	31	34.8	32	50.8	1.000
• Full	58	65.2	57	64.0	
Husband and wife communication					
• Bad	49	55.1	30	33.7	0.007
• Well	40	44.9	59	66.3	

Table 2. In the husband's role variables, namely husband's consent, 32 (34.8%) respondents stated that injection of husband to wife was less and 58 (65.2%) respondents stated that the husband's consent was given in full, while for long-term contraceptive users (MKJP) 32 (36.0%) respondents stated that husband's approval was lacking and as many as 57 (64.0%) respondents stated that the husband's consent was given in full. and husband and wife communication variables indicate that, injecting contraceptive users with poor communication as many as 49 (55.1%) respondents and 40 (44.9%) good respondents while 30 MKJP users (33.7%).) respondents showed poor husband and wife communication and as many as 59 (66.3%) respondents showed good husband and wife communication.

Table 3. Multiple logistic regression analysis

Variable	B	p-value	OR	Wald
Information provided	3.194	0.000	0.041	42.958

The results of bivariate statistical analysis based on the significance of $p < 0.05$ indicate the variables that influence contraceptive use, namely contraceptive choice (p -value = 0.001), information provided (p -value = 0.000), technical ability of the officer (p -value = 0.000), conjugal communication (p -value = 0.007) while interpersonal relations (p -value = 0.764), follow-up (p -value = 0.229) and husband's consent (p -value = 1.000) do not affect contraceptive use (p -value > 0.05) (Table 2). Multiple logistic regression analysis was used to see the variables that most influenced contraceptive use. The results of logistic regression analysis indicate that the most influencing variable is the information provided with a Wald value of 42.958 (Table 3).

DISCUSSION

The concept of a special family planning service standard for infrastructure facilities in the provision of services, the contribution of tools and drugs is very important because there is no contribution of contraceptive tools and drugs so that the implementation of family planning services will not succeed so that the program should offer a method that is sufficient to serve the whole group of people to nest or limit the number of children (15, 16) so the government issued a policy on the availability of various types of contraception but in reality these contraceptives were often limited (17).

Based on the results of the analysis of choice of contraceptive use in the City of Makassar, there is an influence between contraceptive choices on contraceptive use and the value of $p (0.000) < \alpha (0.005)$. The results of this study are consistent with the statement that the availability of contraception to be used will increase the coverage of users of contraception (18). In this thing client satisfaction affects the contraception provided, if the family planning service is able to provide contraceptives properly, the acceptor will experience satisfaction with the quality of the service provided and can encourage other acceptors to take advantage of family planning services at that place (19).

A survey with the theme "Contraception: Getting the Right Facts" involving 9 countries, namely China, South Korea, Thailand, Singapore, Indonesia, India, Pakistan, Taiwan and Malaysia with a total of 1,800 male and female respondents aged 20-35 years. the results show that around 30 percent of correspondents in Asia get false and inaccurate information about contraception (20).

The results of this study indicate p -value (0.000) < α (0.05) so that there is an influence of information provided on contraceptive use in Makassar City. information provided relating to contraceptive use in Ethiopia information about side effects related to information provided by family service providers needed can help individuals and couples to meet their health needs safely and effectively (21, 22).

Interpersonal relationships refer to the level of empathy, trust, guarantee of confidentiality and the sensitivity of provide to meet the needs and expectations of clients which has an impact on the creation of positive relationships including the way they are treated by service providers (23, 24).

In this study shows that there is no influence between interpersonal relationships on contraceptive use in the city of Makassar with a value of $p (0.764) > \alpha (0.05)$. Good interpersonal relationships make it possible to maintain contraceptive use (25) so that interpersonal relations is one of the keys to using contraception (26).

The technical capacity of family planning service providers is a basic procedure that must be obtained by family planning acceptors in using family planning services. The results of this study indicate the value of p -value (0.000) < α (0.05) so that there is an influence between the technical ability of officers to use contraception. In this study, the ability of officers in two health centers to carry out a variety of standard services but some officers have not carried out basic procedures for other acceptors in both long-term contraceptive users and short-term users of contraception.

The quality of services covering (availability of contraception, information provided, ease of service, interpersonal / counseling relationships, mechanism of follow-up and technical abilities of officers) is a determinant of contraceptive use. however, even though the officers have carried out appropriate procedures, they do not affect women of childbearing age in contraceptive use (27, 28).

A person's behavior in complying with every recommendation from a health worker is influenced by several factors such as the level of knowledge, level of education, socio-economic, and employment. In addition, health care facilities, attitudes and behavior of health workers also support and strengthen the formation of a person's behavior (29). Follow-up visits (controls) are considered important but not felt as a routine need to be done, control is carried out if there are serious complaints that arise when using contraception. If

there are no complaints, they do not follow the schedule. Thus, the mechanism of follow-up recommended by officers is considered important or not depending on the type of contraception used⁽³⁰⁾.

Husband's support is verbal or non-verbal information, advice, real assistance or behavior given by an official life partner of a woman (wife)⁽³¹⁾. The results of this study indicate the value of p -value $(1,000) > \alpha$ $(0,05)$ so that there is no influence between husband's agreement on contraceptive use in the city of Makassar, H_1 is rejected and H_0 is accepted.

Husbands who do not participate in contraceptive use for partners cause them to be responsible for their use of contraception⁽³²⁾. Emotional support for wives in family planning can be realized through good communication between husband and wife in reproductive health and participation in using contraception⁽³³⁾.

The results in this study indicate that the p -value $(0,007) < \alpha$ $(0,005)$ indicates that there is an influence between conjugal communication on contraceptive use in Makassar City in 2019.

Increasing knowledge about family planning can be done by counseling or active activities using the game method so that it can be used in contraception⁽³⁴⁾. The relationship between a woman and her partner can also be a factor in determining certain contraceptive choices. In many communities, couples do not communicate with each other about family planning, but women often have to get and use contraception if they want to control their fertility⁽³⁵⁾.

CONCLUSION

Based on the results of the research and discussion with reference to the formulation of the problem and the research hypothesis, it can be concluded that health centers and husbands are a determining factor for someone to use contraception. So that health workers are advised to further enhance their role in terms of service delivery so that acceptors can use contraceptives that are suitable for reproductive purposes to avoid failure rates and prevent maternal deaths from unwanted pregnancies or pregnancies in old age and husbands as the first people to make decisions, contribute well to the choice of contraception that will be used by the wife.

REFERENCES

1. Munandar B. The Role of Information on Family Planning about Perception in Family Planning Practices. *Jurnal swarnabhumi: Jurnal Geografi dan Pembelajaran Geografi*. 2017;2(1):59-50
2. NCANDS. [National Child Abuse And Neglect Data System (NCANDS)]. Childre's Bureau. 2017[2019 Jan 02]. Available from: https://www.acf.hhs.gov/sites/default/files/cb/ncands_glossary.pdf.
3. USAID. [Fast For Family Planning 2012]. JHU Center for Communication Programs.2012 [2019 Jan 02]. Available from: https://www.fphandbook.org/sites/default/files/familyplanning_web.pdf.
4. Ministry Of Health Of The Republic Of Indonesia. Situation Of Family Planning In Indonesia (Situasi Keluarga Berencana Di Indonesia). Jakarta: Ministry Of Health Of The Republic Of Indonesia;2015.
5. WHO. Family Planning/Contraception 2018. [Cited 2019 10 Januari]. Available from:<https://www.who.int/news-room/fact-sheets/detail/family-planning-contraception>.
6. Ministry Of Health Of The Republic Of Indonesia. Indonesian Health Profile (Profil Kesehatan Indonesia). Jakarta : Ministry Of Health Of The Republic Of Indonesia;2017.
7. Makassar City Health Office. Coverage of Contraceptive Users Based on Public Health Centers in Makassar City (Cakupan Pengguna Kontrasepsi Berdasarkan Puskesmas di Kota Makassar). Makassar : Makassar City Health Office 2018.
8. Tessema GA, Gomersall JS, Mahmood MA, Laurence CO. Factors Determining Quality Of Care In Family Planning Services In Africa: A Systematic Review Of Mixed Evidence. *JBIDatabase of Systematic Reviews and Implementation Reports*. 2016;11(11):116-103.
9. Endriyas M, Belachew T, Megerssa B. Missed Opportunities In Family Planning: Process Evaluation Of Family Planning Program In Omo Nada District, Oromia Region, Ethiopia. *Contraception and reproductive medicine*. 2018;3(14):9-1.
10. Nanda G, Schuler SR, Lenzi R. The Influence Of Gender Attitudes On Contraceptive Use In Tanzania: New Evidence Using Husbands'and Wives'survey Data. *Journal of biosocial science*. 2013;45(3):44-331.
11. White K, Ocampo M, Scarinci IC. A Socio-ecological Approach For Examining Factors Related To Contraceptive Use Among Recent Latina Immigrants In An Emerging Latino State. *Journal Women & Health*. 2017;57(7):872-89.
12. Health Center Tamalanrea and Health Center Kassi-Kassi. Daftar Pengguna Kontrasepsi. 2018.

13. Srivastava A, Singh D, Montagu D, Bhattacharyya S. Putting Women At The Center: A Review Of Indian Policy To Address Person-Centered Care In Maternal and Newborn Health, Family Planning and Abortion. *BMC public health*. 2018;18(1):10-1.
14. Moedjiono AI, Kuntoro, K., Notobroto, H B. Indicators Of Husband role in pregnancy and maternity care. *International Journal of Public Health Science (IJPHS)*.2017;6(2):192-6.
15. Imron R, Kes M, Novita Rudiyanti S. Determinants For Using Iud Contraception Tools In Family Planning Acceptors are Worked in The Sidomulyo Health center, Sidomulyo Sub District. *Journal Of Health*. 2011;2(2):312-318.
16. Kahraman K, Göç G, Taşkın S, Haznedar P, Karagözlü S, Kale B, et al. Factors influencing the contraceptive method choice: a university hospital experience. *Journal of the Turkish German Gynecological Association*. 2012;13(2):102.
17. Spagnoletti BRM, Bennett LR, Kermod M, Wilopo SA. 'I Wanted to Enjoy our Marriage First But I got Pregnant Right Away': A Qualitative Study Of Family Planning Understandings and Decisions Of Women In Urban Yogyakarta, Indonesia. *BMC pregnancy and childbirth*. 2018;18(1):353.
18. Charles MM, Abel NM, Simon IK, Henry MS, Baptiste KSZJ, Françoise MK. Availability and Quality of Family Planning Services in the City of Lubumbashi, Democratic Republic of Congo. *Open Access Library Journal*. 2019;6:11-1.
19. Slater AM, Estrada F, Suarez-Lopez L, de la Vara-Salazar E, Campero L. Overall User Satisfaction With Family Planning Services and Associated Quality Care Factors: A Cross-Sectional Analysis. *Reproductive Health Journal*. 2018;15(1):172.
20. McCauley HL, Silverman JG, Decker MR, Agénor M, Borrero S, Tancredi DJ, et al. Sexual and reproductive Health Indicators and Intimate Partner Violence Victimization Among Female Family Planning Clinic Patients Who Have Sex With Women and Men. *Journal of Women's Health*. 2015;24(8):621-8.
21. Jain AK. Information about methods received by contraceptive users in india. *Journal of biosocial science*. 2017;49(6):798-810.
22. Tafese F, Woldie M, Megeressa B. Quality Of Family Planning Services In Primary Health Centers Of Jimma Zone, Southwest Ethiopia. *Ethiopian Journal Of Health Sciences*. 2013;23(3):245-54.
23. Galle A, Vermandere H, Griffin S, de Melo M, Machaieie L, Van Braeckel D, et al. Quality Of Care In Family Planning Services In Rural Mozambique With A Focus On Long Acting Reversible Contraceptives: A Cross-Sectional Survey. *BMC women's health Journal*. 2018;18(1):201.
24. Chavane L, Dgedge M, Bailey P, Loquiha O, Aerts M, Temmerman M. Assessing Women's Satisfaction With Family Planning Services In Mozambique. *J Fam Plann Reprod Health Care*. 2017;43(3):222-8.
25. Dehlendorf C, Henderson JT, Vittinghoff E, Grumbach K, Levy K, Schmittiel J, et al. Association Of The Quality Of Interpersonal Care During Family Planning Counseling With Contraceptive Use. *American Journal Of Obstetrics And Gynecology*. 2016;215(1):78. e1-. e9.
26. Sans-Corrales M, Pujol-Ribera E, Gene-Badia J, Pasarin-Rua MI, Iglesias-Perez B, Casajuana-Brunet J. Family Medicine Attributes Related To Satisfaction, Health And Costs. *Family practice*. 2006;23(3):308-16.
27. Najib N. Knowledge and Quality of Service for Clients as Basics for Choosing Hormone Contraception. *Kesmas: National Public Health Journal*. 2011;6(3):111-6.
28. Setiasih s. Analysis of Factors Affecting the Selection of Long-Term Contraceptive Methods (MKIP) in Fertile Age Couples in the District of Kendal in 2013. *Indonesian Health Promotion Journal*. 2016; 11(2):46-32
29. Andriani D, Lestari T. Analysis Of Factors Related To Risk Sexual Behavior In Adolescents / High School In The District Of Sungai Manau Year 2014. *Jurnal Penelitian Universitas Jambi: Seri Sains*. 2015;17(1):66-55.
30. Budiman B, Kasto K. Quality of Family Planning Services in Unaaha District, Konawe Regency. *Journal Of Gadjah Mada University*.2016(2):154-75.
31. Agustini NNM, Suryani N, Murdani P. Relationship Between Knowledge Level of Mother and Family Support with Antenatal Service Coverage in the Work Area of Buleleng Health Center. *Journal of the Master of Family Medicine*. 2013;1(1):67-79.
32. Mandayani S. Knowledge And Level Of Husband Education Relationship With The Support Of Use Of Contraception Instruments In Iud (Iud) In Lambaro Village, Aceh District. *ETD Unsyiah*. 2014.
33. Muniroh ID, Luthviatin N, Istiaji E. Husband's Social Support for Wives for Using Women's Operation Medical Contraception (MOW) (Qualitative Study on Fertile Age Unmet Need in Puger District, Jember Regency. *e-Jurnal Pustaka Kesehatan*. 2014;2(1):66-71.

34. Moedjiono AI, Awaluddin, Hidayanty H, Abdullah T. Prevention of Delay of Decision Making as Efforts to Improve The Mother Health in Polewali Mandar, West Sulawesi Indonesia. *Indian Journal of Public Health Research & Development*. 2019;10(1):1230-6.
35. Supiani S. Relationship Of Husband's Support To The Use Of Contraceptives In The Womb In The Work Area Of Gamping II Sleman Yogyakarta Health Center: ; 2015.

1-

PB_Dewi_Rahmawati,_Apik_Indarty_Moedjiono,_Sukri_Palutturi..

ORIGINALITY REPORT

% **12**
SIMILARITY INDEX

% **6**
INTERNET SOURCES

% **7**
PUBLICATIONS

% **6**
STUDENT PAPERS

PRIMARY SOURCES

1 www.ijsk.org Internet Source % **1**

2 ovihealth.com Internet Source % **1**

3 journals.plos.org Internet Source % **1**

4 Submitted to Universitas Hasanuddin Student Paper % **1**

5 Submitted to Moulton College Student Paper % **1**

6 Muh. Kahfi, Hasanuddin Ishak, Ridwan M. Thaha. "Analysis of the Environmental Factors and the Efforts of the Community to Avoid the Mosquito Bites Against the Endemic Dengue in Rappocini Sub-District, Makassar City", Proceedings of the International Conference on Healthcare Service Management 2018 - ICHSM '18, 2018
Publication % **1**

-
- 7 worldwidescience.org % 1
Internet Source
-
- 8 Healthy Hidayanty, Devintha Virani, Sabaria Manti, Rahayu Indriasari, Dian Sidik Arsyad. "Inadequate nutrients intake and wasting status among adolescent students in Small Island of Indonesia", *Enfermería Clínica*, 2020 % 1
Publication
-
- 9 Agus Heru Darjono, Ujang Sumarwan, Lilik Noor Yuliati, Hari Wijayanto. "The Influence of Family, Community, and Information Technology Supports on Empowerment of Diabetes Mellitus Patients Using SEM (*Structural Equation Modelling*)", *Asian Journal of Applied Sciences*, 2018 <% 1
Publication
-
- 10 www.researchsquare.com <% 1
Internet Source
-
- 11 Submitted to School of Business and Management ITB <% 1
Student Paper
-
- 12 Prabir C. Bhattacharya. "Determinants of Fertility in Three North Indian States: Caste, Female Autonomy and Economic Development", *Canadian Journal of Development Studies/Revue canadienne* <% 1

13 Citrakesumasari ., Vonny kalsum, Chaidir Masyhuri Majiding, Tenri Andi Ayu Rahman, Yessy Kurniati. "Mineral Concentrations in Breast Milk across Infant Birth Weight", Pakistan Journal of Nutrition, 2019

Publication

14 repositori.usu.ac.id

Internet Source

15 fkm.unhas.ac.id

Internet Source

16 Winda Liftiana Sari, Fuadah Fakhruddiana. "Internal locus of control, social support and academic procrastination among students in completing the thesis", International Journal of Evaluation and Research in Education (IJERE), 2019

Publication

17 Submitted to University of Southampton

Student Paper

18 journals.scholarpublishing.org

Internet Source

19 Muhammad Yunus, Saenab, Andi Ahmad Yani, Sangkala et al. "Health service innovation in City of Makassar", Enfermería Clínica, 2020

20 www.ijsspr.com <% 1
Internet Source

21 Submitted to Stourbridge College <% 1
Student Paper

22 Submitted to University of Nottingham <% 1
Student Paper

23 www.nivel.nl <% 1
Internet Source

24 Elsa Palinggi, Andi Zulkifli, Alimin Maidin, Andi Ulfiana Fitri. "Effect of logotherapy to control depression of drug abuser in Kassi-Kassi and Jumandang Baru health-center", Enfermería Clínica, 2020 <% 1
Publication

25 Submitted to London School of Economics and Political Science <% 1
Student Paper

26 Submitted to Institute of Graduate Studies, UiTM <% 1
Student Paper

27 Anna Galle, Heleen Vermandere, Sally Griffin, Málica de Melo, Lino Machaieie, Dirk Van Braeckel, Olivier Degomme. "Quality of care in family planning services in rural Mozambique with a focus on long acting reversible <% 1

contraceptives: a cross-sectional survey", BMC Women's Health, 2018

Publication

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5

WORDS