

Selection of Dominant and Co-dominant Markers for Red Wood (*Pterocarpus indicus* Willd) Polymorphism from Five Provenances in East Nusa Tenggara

by Siti Halimah Larekeng

Submission date: 18-Mar-2020 01:14PM (UTC+0100)

Submission ID: 1277522027

File name: Ild_Polymorphism_from_Five_Provenances_in_East_Nusa_Tenggara.pdf (696.82K)

Word count: 4622

Character count: 22520

Selection of Dominant and Co-dominant Markers for Red Wood (*Pterocarpus indicus* Willd) Polymorphism from Five Provenances in East Nusa Tenggara

3

Siti Halimah Larekeng

Biotechnology and Tree Breeding Laboratory, Department of Forestry, Faculty of Forestry, Universitas Hasanuddin, Indonesia. 90245

6

{siti5h.82@gmail.com}

Abstract. Information about the genetic diversity of Red wood is needed to improve the quality and quantity of the plant. This information can be provided by analyzing plant DNA using molecular markers which are polymorphic and generate clear and thick bands. The objective of the study was to obtain primers that could generate clear and polymorphically bands from Red wood DNA. The evaluated Red wood was from Alor, Sika, Lembata, Rote, and Timur Tengah Selatan (TTS) provenances. As many as 54 primers screened to polymorphism assay. Those primers consisted of RAPD, SSR, and chloroplast. The SSR primers were developed from *Pterocarpus indicus* (six primers) and *Prosopis alba* (12 primers). Chloroplast primers were divided into two types, four Rubisco Large Sub-unit genes (*rbcL*) and 12 MaturaseK genes (*matK*). SSR primers of *P. alba* were not able to amplify Red wood DNA, whereas those of *P. indicus* could amplify the DNA. Three of each chloroplast primers successfully amplified the DNA, yet, each primer could only amplify one to four of 12 evaluated samples. *rbcL* 2 generated the highest number of sample, and samples from Lembata were amplified the most. Moreover, 16 RAPD primers were able to generate DNA.

Keywords: Genetic marker, *matK*, *RAPD*, *rbcL*, *SSR*, Red wood.

1 Introduction

A genetic marker is a tool to analyze the genetic diversity of an individual including trees. A molecular marker is one of the genetic markers that can distinguish individuals by carrying out an analysis at the level of Deoxyribose Nucleic Acid (DNA). This marker is more accurate than other genetic markers as it is not influenced by the environment.

Molecular markers can be developed from nucleus DNA or cytoplasmic DNA, such as chloroplasts DNA. The developed markers are Random Amplified Polymorphic DNA (RAPD), Simple Sequence Repeat (SSR), Amplified Fragment Length Polymorphic (AFLP), Single Nucleotide Polymorphism (SNP), and Inter-Simple Sequence Repeat (ISSR). Meanwhile, molecular markers developed from chloroplast DNA are *matK*, *rbcL*, *SSRcp* and others. These markers are primers used for DNA amplification in genetic analysis.

10 Primer is a nucleic acid chain that functions as a starting point for synthesizing DNA, which is needed for DNA replication because the enzymes that catalyze this process are DNA polymerase. Primer screening aims to obtain a high polymorphism level, and thus the used primers were selected from several plant species sharing the same family with the studied

species. This selection is done to find random primers that produce bands, both clearness of the band and number of the locus [1].

RAPD marker is dominant, and consequently, the produced DNA fragments cannot distinguish between homozygous (AA) and heterozygotes (Aa) individuals, whereas those without bands are recessive (aa) genotypes [3]. The utilization of co-dominant marker, for instance SSR, is needed in analyzing the genetic diversity because it is able to distinguish homozygous and heterozygous individuals. SSR has a simple sequence consisting one to six bases that are repeated and often found in plant genomes [4]. The variation in the number of repetitions of microsatellite sequences causes it has high polymorphism level and suitable for utilizing in genetic diversity of a population and parental analysis studies [5]. Molecular analysis can also be performed using the chloroplast genome. This analysis is able to identify genetic diversity and phylogenetic between species. *rbcL* and *matK* chloroplast genome markers are recommended primers for analysis in plants [6].

The distribution of Red wood in Indonesia is quite extensive, especially in eastern Indonesia; Sulawesi, Maluku, West Nusa Tenggara, East Nusa Tenggara, and Papua. Even though having widespread distribution, it has experienced a decline in population and genetic diversity hence it has been categorized as Vulnerable species in the International Union for Conservation of Nature (IUCN) Red List of Threatened Species due to high exploitation [7]. Efforts to maintain the existence of Red wood need to be done for the preservation of this species in the future in order to improve the plant characters and plant breeding.

The previous studies on primer screening have been done on *Vitex coffassus* [1], *Elmerillia tsiampacca* [8], *Swietenia mahogany* [9], *Swietenia macrophylla* [10]. Information about the suitable primers that can amplify the Red wood DNA is still limited. Therefore, study on primer screening with several primer types needs to be done. The objective of the study was to obtain primers that could amplify clear and polymorphic bands from Red wood DNA.

2 Material and Methods

This study was conducted in September 2018 up to March 2019 at Biotechnology and Tree Breeding Laboratory, Faculty of Forestry, Hasanuddin University, Makassar, Indonesia. Leaves of Red wood were collected from genetic resources area of Watershed Management Center of East Nusa Tenggara (BPDAS Nusa Tenggara Timur) located in Silu village, Kupang, East Nusa Tenggara, Indonesia. The seedling collected at the study site were from Alor, Rote, Sika, Lembata, dan Timur Tengah Selatan (TTS) provenances. Twelve DNA samples of Red wood were selected for primer screening; three samples from Rote and Lembata, and two samples from Alor, Sika, and TTS, respectively. Leaves were weighed as much as 100 mg and then grounded for the DNA extraction process. The DNA extraction procedure was done using DNA Geneaid extraction kit protocol. The extracted DNA was mixed with RNase and stored in -20°C freezer for 24 hours. After 24 hours, the DNA working solution was prepared by diluting 1 µL DNA master with 9 µL ddH₂O.

Primer screening used RAPD, SSR, and Chloroplast primers. Nine RAPD primers, six SSR primers from *Pterocarpus indicus* [6], four *rbcL* primers, and twelve *matK* primers were screened in this study (Table 1).

DNAs were amplified using PCR Sensoquest Thermal Cycler with one PCR reaction consisted of 2µL of DNA template; 0,625µL of Hotstar PCR mix; and 3µL of ddH₂O. One PCR reaction for RAPD marker contained 1,25 µL of RAPD primer, while that of SSR and chloroplast markers contained 1,25 µL primers (0.625µL of forward and 0.625µL of reverse). The amplification was performed using the following steps: one cycle of preheat at 95°C for 5 minutes, 35 cycles of denaturation at 94°C for 60 seconds, primer annealing at annealing

temperature for each primer for one minute, elongation at 72°C for 60 seconds, and final elongation at 72°C at 10 minutes. The precise annealing temperature for each primer was determined by doing a temperature gradient at $\pm 5^\circ\text{C}$ of each primer temperature.

SSR amplification products were separated on 3% of SFR agarose with 1x TAE buffer at 100 V for 90 minutes. Meanwhile, that of RAPD, RbcL, and MatK were separated on 2% of agarose with 1x TAE buffer at 100 V for 60 minutes. The products were then visualized over UV light and documented using a Gel Doc system.

Table 1. Primer name, sequence, and primer melting temperature

No	Locus Name	Primer Sequence	Melting Temperature (°C)
SSR primers of <i>P. indicus</i>			
1	Pin2_36/GU9 03082	F: TCA AAA TTA GAA TGA AGG TGA G R: GCAATAAGAGAGAATGAAGAAA	51,2
2	Pin2_29/GU9 03081	F: TCGTTTGGAAAGTGTTTCCATT R: TACCACCTTGCCCGAGTTTATG	54,1
3	Pin2_04/GU9 03076	F: ATTGTCAGTTGATGTATCTTGC R: TTTAATCAGCTTGTTTCCTATG	49,9
4	Pin1_23/GU9 03073	F: ATTCCTGGATAGGCTGACAAAT R: CTGAAAGCACGAGAGCATTITTT	56,1
5	Pin1_04/GU9 03071	F: CTC GTA AGC CCA AGT CAA CAT A R: AAT CAG AAA CCT AAA ACG GAA C	52,9
6	Pin1_01/GU9 03070	F: CTAACCTGAATACCAAGAAAACCTCC R: TCTCAAGAATAAAAAGATGTGC	48,4
7	Pin2_39/GU9 03083	F: GCAACACCTCCAACACCAAAAAA R: CTCTAATCCCTCCCTCGCCAAA	57,6
SSR Primers of <i>Prosopis alba</i>			
1	GL3	F: CAC CGA TCTCACAAAGCTGC R: AATGGATCTGGTGTGTCGC	56
2	JX136853	F: CTGGTTGCTGTGATTGGAGG R: CTCCAGGGATCACAAAGACAAAC	55,8
3	GL6 JX136854	F: CAGGTGGGCATGAAGTTTCC R: CCAAGAACAACCTGCCGAAG	56,3
4	GL8	F: ACTCTGCGGGTTAGGTAAGC R: ACCTGGAGCTGACATGGATC	56,7
5	JX136855	F: GAGTGAAGGTCGGGAAGAGG R: CCATTGGACCAAGGCAGAAC	56,7
6	GL9 JX136856	F: GTGTTATGGTCCCAACAGCC R: TGAAGAGGGAGGAATCGCAG	56,4
7	GL12	F: GTTGGATTTACGGAAGGGC R: TCAGCTAAGTGGCCATACGG	56,8
8	JX136857	F: GAGAATCTGGAGCAGCAACG R: AAGGTAGCGTCCAGGTATG	56,2

9	GL15 JX136858	F: ATCTCCGTCACAACCTTGCAC R: ACCCTCACTCCCGAATGATG	56,1
10	GL16 JX136859	F: GTCTTCTCTCCCGTGGATCC R: TGAGGCAAAGGAAGAGCAAC	56,4
11	GL18 JX136860	F: CCTTAATCTCCCTCTCGGCC R: AACCAGGCTCTGCAGAAATG	56,3
12	GL21 JX136861	F: CGAATGTGGATCTTCTGCGG R: TTAAGCGGCCCAAGTTTCTC	55,8
	GL23 JX136862		
	GL24 JX136863		
	GL26 JX136864		
RAPD Primers			
1	OPQ-07	CCCCGATGGT	38,5
2	OPAC-12	GGCGAGTGTG	38,1
3	OPAE-11	AAGACCGGGA	35,5
4	OPAA-20	TTG CCT TCG G	35,6
5	OPA 15	TTCCGAACCC	34,2
6	OPZ 05	TCCCATGCTG	4,3
7	OPG-19	GTCAGGGCAA	34,7
8	OPC-11	AAAGCTGCGG	36,9
9	OPA-09	GGGTAACGCC	37,4
10	OPP-08	ACATCGCCCA	37,6
11	OPD-03	GTCGCCGTCA	40,8
12	OPA-02	TGCCGAGCTG	40,7
13	OPD-20	ACCCGGTCAC	39,1
14	OPY-09	AGCAGCGCAC	42,5
15	OPA 05	AGGGGTCTTG	32,6
16	OPA 11	CAATCGCCGT	32,0
17	OPG 09	CTGACGTCAC	32,0
18	PLC 14	TGCCGTGCTTG	32,0
19	PLR 13	GGACGACAAG	32,0
20	PLW 04	CAGAAGCGGA	32,0
Chloroplast Primers			
1	rbcL 1	F: ATG TCA CCA CAA ACA GAA AC R: TCG CAT GTA CCT GCA GTA GC	54,2
2	rbcL 2	F: TAT GCG TTG GAG AGA CCG TTT C	57,2

3	rbcL 3	R: TCG CAT GTA CCT GCA GTA GC F: ATG TCA CCA CAA ACA GAA AC	54,05
4	rbcL 4	R: TCC TTT TAG TAA AAG ATT GGG CCG AG F: TAT GCG TTG GAG AGA CCG TTT C	54,05
5	matK 1	R: TCC TTT TAG TAA AAG ATT GGG CCG AG F: CAT GCA TTA TGT TAG GTA TCA AGG	50,05
6	matK 2	R: CGA AGT ATA TAY TTT ATT CGA TAC A F: CCA TAG TTC CAA TTA TTC CTC TG	49,65
7	matK 3	R: CGA AGT ATA TAY TTT ATT CGA TAC A F: CCC CCA TAT ATT TGA TAC CTT CTC	51,1
8	matK 4	R: CGA AGT ATA TAY TTY ATT CGA TAC A F: CAT GCA TTA TGT TAG GTA TCA AGG	54,1
9	matK 5	R: GCT TGC ATT TTT CAT TGC ACA CG F: CCA TAG TTC CAA TTA TTC CTC TG	53,7
10	matK 6	R: GCT TGC ATT TTT CAT TGC ACA CG F: CCC CCA YAT ATT TGA TAC CTT CTC	55,1
11	matK 7	R: GCT TGC ATT TTT CAT TGC ACA CG F: CAT GCA TTA TGT TAG GTA TCA AGG	50,65
12	matK 8	R: ATT CGA GTA ATT AAA CGT TTT ACA A F: CCA TAG TTC CAA TTA TTC CTC TG	50,25
13	matK 9	R: ATT CGA GTA ATT AAA CGT TTT ACA A F: CCC CCA YAT ATT TGA TAC CTT CTC	51,7
14	matK 10	R: ATT CGA GTA ATT AAA CGT TTT ACA A F: CAT GCA TTA TGT TAG GTA TCA AGG	52,1
15	matK 11	R: CCA GAA ATT GAC AAG GTA ATA TTT CC	51,7
16	matK 12	R: CCA GAA ATT GAC AAG GTA ATA TTT CC F: CCC CCA TAT ATT TGA TAC CTT CTC R: CCA GAA ATT GAC AAG GTA ATA TTT CC	53,15

3 Results and Discussion

Three primers of the evaluated primers, SSR (developed from *P. indicus*), RAPD and chloroplast primers, could amplify DNA samples. Those of SSR primers were pin2_39, pin2_36, pin2_29, pin2_04, pin1_04, and pin1_01 (Table 2). Each primer amplified two until seven DNA samples from Rote, Lembata, Sika, and Alor. Figure 1 and 2 present the electropherogram of PCR products using six SSR primers. Figure 1 depicts only two of 12 DNA samples could be generated using Pin2-29, sample #1 and 5, which was from Alor.

Table 2. PCR amplification product information using SSR primers

No	Primer name	Number of amplified sample	Annealing temperature	Provenance	Band
1	Pin2_39	6	62,6	Rote,Lembata,Sika,Alor	Thin and clear

2	Pin2_36	4	56,2	Sika,Alor	Thick and clear
3	Pin2_29	2	52,4	Rote,Lembata	Thin and clear
4	Pin2_04	7	51,7	Rote,Lembata,Sika,Alor	Thick and clear
5	Pin1_04	5	58	Rote,Lembata,Sika,Alor	Thick and clear
6	Pin1_01	4	53,5	Rote,Lembata,Sika,Alor	Thin and clear
7	Pin1_23	0	-	-	Unamplified




Fig 1. Electropherogram of DNA amplification products of Red wood using SSR primers of *P. indicus*, Pin2_39, Pin2_36, and Pin2_29. Notes: M = 100 bp marker, 1-12 = DNA amplification products using Pin2-39 Pin2-29, and Pin2-36




Fig 2. Electropherogram of DNA amplification products of Red wood using SSR primers of *P. indicus*, Pin2_04, Pin1_01, and Pin1_04. Notes: M = 100 bp marker, 1-12 = DNA amplification products using Pin2_04, Pin1_01, and Pin1_04.

SSR primers of *P. indicus* that amplified DNA samples were Pin2_39, Pin2_36, Pin2_29, Pin2_04, Pin1_04, and Pin1_01 (Table 2). Each primer was only able to generate one from 12 evaluated samples. As the primers could only amplify one sample, they are eliminated for further analysis on genetic diversity. Genetic diversity analysis requires primers that produce polymorphic bands (many alleles) in order to distinguish the individuals and calculate the parameters of genetic diversity analysis. [11] employed polymorphic primers for measuring parameters of genetic diversity analysis.

Genetic diversity using SSR primer has been widely applied on tree species as it is categorized as a co-dominant primer that able to distinguish between homozygote and heterozygote individuals. This primer is used to analyze the genetic diversity of *Anthocephalus macrophyllus* [12].

P. alba is in the same family with Red wood, Fabaceae. Thus its primers can be used to amplify Red woods DNA. Meanwhile, SSR primers belonged to *P. indicus* were also utilized because of sharing the same Latin name, *Pterocarpus indicus*, with Red wood. SSR primers of *P. indicus* generated more DNA than *P. alba*' due to taxonomic closeness to Red wood. [11] stated that SSR primer screening originated from a species can be employed to its close relatives as they have similar base sequences. However, in this study, none of those evaluated primers could generate DNAs of Red wood (Figure 3).

rbcL 2 generated four DNA samples and had the highest number of successfully amplified samples. Those successfully amplified samples were from Lembata (two samples), Alor and Sika (one sample, respectively). rbcL 1 and matK 1 amplified two samples from Lembata, and rbcL 4, matK 2, and matK 4 amplified one sample from Lembara, respectively (Table 3). Figure 2 depicts the electropherogram of amplified DNA products using rbcL 4.




Fig 3. Electropherogram of DNA amplification products of Red wood using SSR primers of *P. alba*, GL3, GL6, and GL21. Notes: M = 100 bp marker, 1-12 = DNA amplification products

Table 3. PCR amplification product information using chloroplast primer

No	Primer name	Number of amplified sample	Annealing temperature	Note
1	rbcL 1	2	52,4	Thick and clear
2	rbcL 2	4	55,5	Thick and clear
3	rbcL 3	0	-	Unamplified/No product
4	rbcL 4	1	52,4	Thick and clear
5	matK 1	2	48,4	Thin and clear
6	matK 2	1	48,0	Thick and clear
7	matK 3	0	-	Unamplified/No product
8	matK 4	1	52,4	Thick and clear
9	matK 5	0	-	Unamplified/No product
10	matK 6	0	-	Unamplified/No product
11	matK 7	0	-	Unamplified/No product
12	matK 8	0	-	Unamplified/No product
13	matK 9	0	-	Unamplified/No product
14	matK 10	0	-	Unamplified/No product
15	matK 11	0	-	Unamplified/No product
16	matK 12	0	-	Unamplified/No product




Fig 3. Electropherogram of DNA amplification products of Red wood using chloroplast primer, rbcL4. Notes : M = 100 bp marker, 1-12 = DNA amplification products

In general, rbcL (3 of 4 primers) primers amplified more samples than matK (3 of 12 primers). This result was assumed as rbcL primers have more general sequences, yet, lower accuracy in distinguishing individuals between species than matK [13] and [14].

Screening primer using 20 RAPD primers showed 16 primers could amplify the DNA samples. Allele number of amplified samples was ranged one to nine alleles. That of the highest was using OPA 02 (nine alleles), OPD 03 and OPAC 12 were five alleles, and OPA 15, OPA 09, OPY 09, and OPG 09 were three alleles, respectively. Moreover, OPAE 11, OPC 11, OPA 11, PLR 13, and PLW 04 were two alleles, OPQ 07 and PLC 14 were 1 allele (the lowest), and OPAA 20, OPA 05, OPG 19, and OPD 20 were none, respectively. A total number of the band was 49 polymorphic bands, and the highest was obtained by OPA 02 (nine alleles). The monomorphic bands were observed using OPQ 07 and PLC 14. Polymorphism was also detected on the 16 evaluated primers. Polymorphism is the existence of locus or amplicon (present band) with different sizes on one primer. The presence of the band can be polymorphic and monomorphic primer (the existence of locus with the same size on the analyzed samples) [15].

As many as 20 RAPD primers screened by [16] using Mahogany DNA and [17] using DNA *Gigantochloa atter* to obtain polymorphic primers. Those studies only received four polymorphic primers that could be used in genetic diversity analysis. Criteria of proper primers for genetic diversity analysis are good reproducibility, producing polymorphic, bright, clear, stable as well as easy to score bands [18].

This study is in similar to the previous study by [16] who found two RAPD primers which polymorphically amplified DNA *S. mahogany*, OPQ 07 and OPAE 11, but contrast in OPG 19 and OPA 05 primers. In this study, none of both primers could amplify the DNA. Meanwhile, [17] selected OPA 15, OPZ 05, OPA 05, and OPK 20 to generate polymorphically *G. atter* DNA.

Primer selection on RAPD primers influences polymorphism of the produced bands as each primer has specific annealing site. As consequent of the specific annealing site, every primer generates different polymorphic bands both size and number of DNA bands. Polymorphism refers to the visualization of amplification based on different observed DNA fragments and score in order to determine whether the variation is presented by observing the sequence differences.

Table 4. PCR amplification product information using RAPD primer

No	Locus Name	Number of amplified sample	Annealing Temperature (°C)	Provenance	Note	Number of allele
1	OPQ 07	2	40,2	Rote	Clear	1
2	OPA 15	4	37,1	Sika,Lembata,Rote	Clear	3
3	OPZ 05	4	37,1	Lembata,Rote,Sika	Clear	4
4	OPD 03	3	42,5	Lembata,Rote	Clear	5
5	OPA 02	4	46,9	Sika,Lembata,Rote	Thin and clear	9
6	OPP 08	3	33,8	Lembata,Rote	Thin and clear	4
7	OPA 09	4	34,6	Sika,Timur Tengah Selatan,Rote	Clear	3
8	OPAE 11	2	42,1	Rote	Thin and clear	2
9	OPAA 20	0	-	-	Unamplified/No product	0
10	OPA 05	0	-	-	Unamplified/No product	0
11	OPG 19	0	-	-	Unamplified/No product	0
12	OPC 11	2	39,7	Rote	Thin and clear	2
13	OPAC 12	5	33,6	Lembata,Rote	Thin and clear	5
14	OPD 20	0	-	-	Unamplified/No product	0
15	OPY 09	3	43,6	Lembata,Rote	Clear	3
16	OPA 11	3	34,8	Sika,Lembata,Rote	Unclear/Smear	2
17	OPG 09	2	29,2	Lembata,Rote	Unclear/Smear	3
18	PLC 14	1	27,0	Rote	Thin and clear	1
19	PLR 13	4	32,0	Sika,Lembata,Rote	Clear	2
20	PLW 04	2	33,7	Rote	Thin and clear	2



Fig 4. Electropherogram of DNA amplification products of Red wood using RAPD primers, OPD-03. Notes : M = 100 bp marker, 1-12 = DNA amplification products

RAPD is still used to analyze genetic diversity because of its ability to amplify various species, like *Pinus merkusii* [19] in Educational Forest Hasanuddin University, onion (*Allium cepa* L) [20], *Dioscorea* sp [21], and Indonesian Edible Canna (*Canna indica* L.) [22]. The comparison co-dominant and dominant primers have been performed in molecular research, in genetic diversity using RAPD and SSR of *Ustilago hordei* [23], and *Zingiber officinale* [24]. SSR and SNAP marker to calculate the xenia effect in Kopyor coconut [25], and to analysis mating system in Ebony, Jabon Merah and Kopyor coconut [26,27,28].

4 Conclusion

The results concluded that there were sixteen out of twenty RAPD primers showed good amplification products and polymorphisms. Four of 6 SSR primers developed from *P. Indicus* generated DNA samples, while that of originated from *P. alba* could not amplify any DNA. *rbcl* and *matK* successfully amplified the evaluated DNA for Red Wood DNA amplification along with specific PCR temperature in each primer.

Acknowledgment

The author thanks The 2nd Seed/Seedling Forest Office (Sulawesi BPTH) for providing plant materials, Andi Fahriadi Amir, Gunawan, Yuni Fitri Cahyaningsih dan Mirza Arsiaty Arsyad for participating in this research activities.

References

- [1] Pongtuluran, I., Restu M., and Gusmiaty: Primary Selection for the Genetic Diversity of Bitti (*Vitex coffasus*). Jurnal Perennial. Vol. 8 (1), pp. 25-29 (2012)
- [2] Weeden, N. F. G. M., M. Timmerman, B. E. Hemmat, M. A. Kneen, and Lodhi: Inheritance and Reliability of RAPD Markers. In Applications of RAPD Technology to

- Plant Breeding. Symposium Proceedings. Crop Science Society of America, Madison. Pp. 12-17 (1992)
- [3] Cheng, K. T., H. C. Chang, C. H. Su, and F. L. Hsu: Identification of dried rhizomes of *Coptis* species using random amplified polymorphic DNA. *Botanical Bulletin of Academia Sinica*. Vol. 38, pp. 241-244 (1997)
- [4] Tani, N., Y. Tsumura, T. Kado, Y. Taguchi, S. L. Lee, N. Muhammad, S. Numata, S. Nishimura, A. Konuma, and T. Okuda: Paternity analysis based inference of pollen dispersal patterns, male fecundity variation, and influence of flowering tree density and general flowering magnitude in two dipterocarp species. *Annals of Botany*, Vol. 104, pp. 1421-1434 (2009)
- [5] Karakousis, A., A. R. Barr, K. J. Chalmers, G. A. Ablett, T. A. Holton, R. J. Henry, P. Lim, and P. Langridge: Potential of SSR markers for plant breeding and variety identification in Australian barley germplasm. *Australian Journal of Agricultural Research* Vol. 54 (12), pp. 1197-1210 (2003)
- [6] Li, X., Y. Yang, R. J. Henry, M. Rossetto, Y. Wang, and S. Chen: Plant DNA barcoding: from gene to genome. *Biol. Rev.* Doi: 10.1111/brv.12104 (2014)
- [7] Sulistyawati, P., dan Widyatmoko, A. Y. P. B. C.: Keragaman Genetik Populasi Kayu Merah (*Pterocarpus indicus* Willd.) Menggunakan Penanda Random Amplified Polymorphism DNA. Yogyakarta (2017)
- [8] Abubakar, M. H.: Seleksi Primer Untuk Analisis Keragaman Genetik Uru (*Elmerillia tsiampacca*). Skripsi Sarjana Fakultas Kehutanan Universitas Hasanuddin (2012)
- [9] Mangiwa, S. T.: Seleksi Primer untuk Analisis Keragaman Genetik Tanaman Mahoni (*Swietenia mahagoni*). Skripsi Sarjana Fakultas Kehutanan Universitas Hasanuddin (2012)
- [10] Nurutami, K.: Seleksi Primer Mikrosatelit Mahoni (*Swietenia macrophylla*). Skripsi Sarjana Fakultas Kehutanan Universitas Hasanuddin (2018)
- [11] Nurtjahjansih, I. L. G., A. Y. P. B. C. Widyatmoko, P. Sulistyawati dan Rimbawanto, A.: Karakteristik dan Aplikasi Penanda Mikrosatelit pada Beberapa Spesies *Eucalyptus*. Balai Besar Penelitian Bioteknologi dan Pemuliaan Tanaman Hutan. Yogyakarta. Vol. 7, pp. 107-108 (2013)
- [12] Larekeng SH., Restu M., Gusmiaty, Syamsuddin M., Budirman B.: Moderate Level of Genetic diversity in *Anthocephalus macrophyllus* Roxb, an endemic tree of Sulawesi and Its Implication in conservation. *Int. J. Agr. Syst.* Vol (6), pp. 74-81 (2018)
- [13] Watoo, J. I., Saleem, M. Z., Shahzad, M. S., Arif, A., Hameed, A., and Saleem, M. A.: DNA Barcoding: Amplification and sequence analysis of *rbcl* and *matK* genome regions in three divergent plant species. [Short Communication], *Advancements in Life Sciences*. Vol 4(1), pp. 1-5 (2016)
- [14] Gonzalez, M. A., Baraloto, C., Engel, J., Scott A. M., Pascal P., Bernard R., Aure' lien R., Christophe T., Je'ro'me C.: Identification of Amazonian Trees with DNA Barcodes. Vol. 4 (10), pp. 1-7 (2009)
- [15] Berdakci F.: Random amplified polymorphic DNA (RAPD) markers [review]. *Turk J Biol.* Vol. 25, pp. 185-196 (2001)
- [16] Iswanto.: Analisis Keragaman Genetik Jenis Mahoni (*Swietenia mahagoni* (L) Jacq) pada Berbagai Sumber Benih di Sulawesi Selatan. Skripsi Sarjana Fakultas Kehutanan Universitas Hasanuddin (2016)
- [17] Larekeng, S.H., Restu M., Mis'al, Oktavina J, Cahyaningsih YF.b.: Penggunaan Penanda RAPD Untuk Mengevaluasi Keragaman Genetik Bambu Parring (*Gigantochloa*

- atter) di Kabupaten Maros Sulawesi Selatan. Prosiding Seminar Nasional Silvikultur V Lambung Mangkurat University Press. pp. 404-413 (2018)
- [18] Cannon, J., Robert A.S., Lynn C., Wendy H.Y., Joanne C.S.: Optimization of PCR primers to detect phylogenetically diverse *nrfA* genes associated with nitrite ammonification. *Journal of Microbiological Methods* Vol. 160. Pp. 49-59 (2019).
- [19] Gusmiaty, Restu M, Asriyanni, Siti H.L.: Polimorfisme Penanda RAPD untuk Analisis Keragaman Genetik Pinus merkusii di Hutan Pendidikan Unhas. *Jurnal Natur Indonesia* Vol. 12, pp. 47-53 (2016)
- [20] Sudha, G.S., P. Ramesh, A.Chandra S., T.Sail K., Bramhachari, K.Riazunnisa.: Genetic diversity analysis of selected Onion (*Allium cepa* L.) germplasm using specific RAPD and ISSR polymorphism markers. *Biocatalysis and Agricultural Biotechnology*. Vol. 17, pp. 110-118 (2019)
- [21] Purnomo, Daryono BS, Rugayah, Sumardi I, Shiwachi H.: Genetic variability and classification of Indonesian Yams (*Dioscorea* spp.) based on RAPD analysis. *SABRAO J. Breed. Genet.* Vol. 48, pp. 377-390 (2016)
- [22] Sari N, Suryadiantina, B.S. Daryono and Purnomo.: Variability And Intraspecific Classification Of Indonesian Edible Canna (*Canna indica* L.) Based On RAPD Marker Analysis. *SABRAO Journal of Breeding and Genetics*. Vol. 50, pp. 156-167 (2018)
- [23] Zhou,Y., Gui-Mie C., Jia-Jia L., Ming-qi Z., Yang W., Bai-li F.: Genetic diversity of *Ustilago hordei* in Tibetan areas as revealed by RAPD and SSR. *Journal of Integrative Agriculture*. Vol.15 (10), pp. 2299-2308 (2016)
- [24] Baruah, J., Sudin K.P., Twahira B, Neelav S, Manabi P, Mohan L.: Molecular diversity assessed amongst high dry rhizome recovery Ginger germplasm (*Zingiber officinale* Roscoe) from NE-India using RAPD and ISSR markers. *Industrial Crops and Products*. Vol. 129, pp. 463-471 (2019)
- [25] Maskromo I, Siti H.L., Hengky N and Sudarsono S.: Xenia negatively affecting kopyor nut yield in Kalianda Tall kopyor and Pati Dwarf kopyor coconuts. *Emir. J. Food Agric* Vol. 28, pp. 644-52 (2016)
- [26] Restu, M., Gusmiaty, dan Siti H.L.: High Outcrossing Rate And Pollen Dispersal Distance of *Diospyros celebica* Bakh. (Ebenaceae), an Endemic Tree Species In Sulawesi Island, Indonesia. *Biotropia*, Vol. 24, pp. 173 – 181 (2017)
- [27] Larekeng SH, M Restu, A Arif, Y F Cahyaningsih, and J Mukti.: A genetic approach to study mating system on Jabon Merah (*Anthocephalus macrophyllus* Roxb.) from three different provenances in South Sulawesi. *IOP Conf. Ser.: Earth Environ. Sci.*pp. 1-9. (2019)
- [28] Larekeng SH, Purwito A, Mattjik NA and Sudarsono: Microsatellite and SNAP markers used for evaluating pollen dispersal on Pati tall coconuts and Xenia effect on the production of 'Kopyor' fruits *IOP Conf.Ser.: Earth Environ.Sci.* Vol. 157. Pp. 012-042 (2018).

Selection of Dominant and Co-dominant Markers for Red Wood (*Pterocarpus indicus* Willd) Polymorphism from Five Provenances in East Nusa Tenggara

ORIGINALITY REPORT

12%

SIMILARITY INDEX

4%

INTERNET SOURCES

10%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Duangjai, S.. "A multi-locus plastid phylogenetic analysis of the pantropical genus *Diospyros* (Ebenaceae), with an emphasis on the radiation and biogeographic origins of the New Caledonian endemic species", *Molecular Phylogenetics and Evolution*, 200909
Publication 5%
- 2 medcraveonline.com
Internet Source 1%
- 3 journal.biotrop.org
Internet Source 1%
- 4 onlinelibrary.wiley.com
Internet Source <1%
- 5 worldwidescience.org
Internet Source <1%
- 6 S H Larekeng, M Restu, M A Arsyad, Mutia. "
Observation of morphological and physiological <1%

characteristics on Abangares Mahogany (.) In South Sulawesi ", IOP Conference Series: Earth and Environmental Science, 2019

Publication

7	Submitted to University of Zakho Student Paper	<1%
8	szie.hu Internet Source	<1%
9	Submitted to University of Birmingham Student Paper	<1%
10	Submitted to Republic Polytechnic Student Paper	<1%
11	pasca.unhas.ac.id Internet Source	<1%
12	equityhealthj.biomedcentral.com Internet Source	<1%
13	Rosabelle Samuel, Hashendra Kathriarachchi, Petra Hoffmann, Michael H. J. Barfuss et al. "Molecular phylogenetics of Phyllanthaceae: evidence from plastid MATK and nuclear PHYC sequences", American Journal of Botany, 2005 Publication	<1%
14	Submitted to University of Nottingham Student Paper	<1%
15	www.alice.cnptia.embrapa.br	

Internet Source

<1%

16

pdfs.semanticscholar.org

Internet Source

<1%

17

Joyashree Baruah, Sudin Kumar Pandey, Twahira Begum, Neelav Sarma, Manabi Paw, Mohan Lal. "Molecular diversity assessed amongst high dry rhizome recovery Ginger germplasm (*Zingiber officinale* Roscoe) from NE-India using RAPD and ISSR markers", *Industrial Crops and Products*, 2019

Publication

<1%

18

Submitted to University of Edinburgh

Student Paper

<1%

19

www.tandfonline.com

Internet Source

<1%

20

journals.plos.org

Internet Source

<1%

21

Prem Lal Kashyap, Sudheer Kumar, Ravi Shekhar Kumar, Rahul Tripathi et al. "Identification of Novel Microsatellite Markers to Assess the Population Structure and Genetic Differentiation of *Ustilago hordei* Causing Covered Smut of Barley", *Frontiers in Microbiology*, 2020

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Selection of Dominant and Co-dominant Markers for Red Wood (*Pterocarpus indicus* Willd) Polymorphism from Five Provenances in East Nusa Tenggara

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13
