

Transgenic Phalaenopsis plants with resistance to Erwinia carotovora produced by introducing wasabi defensin gene using Agrobacterium method

by Rinaldi Sjahril

Submission date: 17-Aug-2021 09:27PM (UTC+0700)

Submission ID: 1632476631

File name: Transgenic_Phalaenopsis_Plants.pdf (545.19K)

Word count: 2962

Character count: 16732

Short Communication

Transgenic *Phalaenopsis* plants with resistance to *Erwinia carotovora* produced by introducing wasabi defensin gene using *Agrobacterium* method

Rinaldi Sjahril^{1,2}, Dong Poh Chin¹, Raham Sher Khan¹, Saburo Yamamura³, Ikuo Nakamura¹, Yoshimiki Amemiya¹, Masahiro Mii^{1*}

¹ Faculty of Horticulture, Chiba University, 648 Matsudo, Matsudo City, Chiba 271-8510, Japan; ² Faculty of Agriculture and Forestry, Hasanuddin University, Jl. Perintis Kemerdekaan Km 10, Makassar 90245, South Sulawesi, Indonesia; ³ Iwate Biotechnology Research Center, 22-174-4 Narita, Kitakami City, Iwate 024-0003, Japan

*E-mail: miim@faculty.chiba-u.jp Tel & Fax: +81-47-308-8852

Received November 14, 2005; accepted December 22, 2005 (Edited by M. Iwano)

Abstract Transgenic plants over-expressing wasabi defensin gene were successfully produced in *Phalaenopsis* orchid by *Agrobacterium*-mediated transformation method. Embryogenic cell suspension culture of *Phalaenopsis* Wataboushi ‘#6.13’ was infected with *A. tumefaciens* strain EHA101 carrying a plasmid containing wasabi defensin gene and selectable marker *nptII*, *hpt* genes. Plantlets were regenerated through somatic embryogenesis from the calli selected on hygromycin-containing medium. Transformation of plantlets with wasabi defensin gene was confirmed by PCR analysis. Southern blot analysis confirmed successful integration of 1–4 copies of the gene. Production of the 5 kDa wasabi defensin protein with varying levels was confirmed in the leaf extracts of different transgenic clones using Western blot analysis. Most of the transgenic plants showed strong resistance to *Erwinia carotovora*, which causes soft rot disease in the control plant. These results suggest the usefulness of this gene for conferring the resistance to various diseases of *Phalaenopsis* and possibly other orchids.

Key words: *Agrobacterium tumefaciens*, *Erwinia carotovora*, *Phalaenopsis*, wasabi defensin.

Genetic improvement of *Phalaenopsis* for conferring disease resistance by conventional plant breeding (sexual hybridization) is restricted by the lack of available useful genes in the species of *Phalaenopsis* and related genera. Consequently, genetic transformation method is expected to apply for transferring disease resistance genes as an alternative procedure. So far, transformation with direct delivery of marker genes such as GUS and GFP genes into plant cells by particle bombardment and indirectly through the use of *Agrobacterium*-mediated procedure has been reported on orchids (Anzai et al. 1996; Belarmino and Mii 2000; Chia et al. 1994). However, there have been no reports on the transformation of useful genes including plant disease resistance genes into *Phalaenopsis*.

Plants react against pathogen attacks in a variety of defensive way such as hypersensitive cell death, production of phytoalexins and expression of pathogenesis-related proteins including a number of antimicrobial peptides (Kiba et al. 2003). So far, many

antimicrobial proteins such as chitin-binding proteins, thionins, lipid-transfer proteins and defensins have been identified and their activity has been tested against fungi and bacteria (Florack and Stiekema 1994; Broekaert et al. 1995; Terras et al. 1995).

Plant defensin protein families have been detected in different tissues from wide range of plants such as wheat, barley, pepper, mustard and radish. They are cysteine-rich polypeptides of 5 kDa in size and are considered to have defensive role against plant pathogenic bacteria (Bohlmann 1994) and fungi (Thevissen et al. 1996). Integration and over-expression studies of defensin gene has been reported in many transgenic dicotyledonous plants such as *Arabidopsis*, tobacco, potato, mung bean and canola (Eppel et al. 1997; Gao et al. 2000; Saitoh et al. 2001; Kiba et al. 2003) and only few studies were reported on monocotyledonous plants such as rice, wheat (Kanzaki et al. 2002; Liang et al. 1998) and banana (Chakrabarti et al. 2003).

The Japanese horseradish, wasabi (*Wasabia japonica*),

Abbreviations: GFP, green fluorescence protein; GUS, β -glucuronidase; NDM, New Dogashima medium; NDM-10M, New Dogashima medium with 1% maltose; NDM-20S, New Dogashima medium with 2% sucrose; PGR, plant growth regulator; PLB, protocorm-like body; PSA, potato sucrose agar; WjAMP-1, *Wasabia japonica* antimicrobial protein.

This article can be found at <http://www.jspcmb.jp/>

which has a protective role against microorganisms and has been used as a food ingredient, are considered to have some antimicrobial substances. Secondary metabolites of wasabi, such as wasalexin and 6-methylallylhexyl isothiocyanate, have been reported to have antifungal and antibacterial activity, respectively (Pedras et al. 1999; Ono et al. 1998). Recently, antimicrobial protein-encoding gene (WjAMP-1) was isolated from wasabi (Saitoh et al. 2001) and named as wasabi defensin gene by Kanzaki et al. (2002). Antimicrobial protein, coded by WjAMP-1 gene showed a strong expression and inhibitory effect on fungal and bacterial growth in transgenic *Nicotiana benthamiana* (Saitoh et al. 2001). Furthermore, Kanzaki et al. (2002) have reported that growth of rice blast fungus was inhibited in transgenic rice over-expressing the wasabi defensin gene.

In this study, we report an attempt to transfer wasabi defensin gene derived from a dicotyledonous plant, *Wasabia japonica* into monocotyledonous plant, *Phalaenopsis* orchid, and evaluate their resistance to bacterial pathogens.

Fine embryonic cell suspension culture of *Phalaenopsis* Wataboushi '#6.13', which was induced and maintained from meristem culture of axillary bud as described previously (Tokuhara and Mii 1993) was used for the transformation study. Five day-old suspension cells after each subculture were used for the transformation experiments.

The *A. tumefaciens* strain EHA101 (pEKH-WT) used harbored the wasabi defensin gene isolated from *Wasabia japonica* (Figure 1). Inoculation of the suspension cells with the *Agrobacterium* was performed following the method of Belarmino and Mii (2000) with some modifications (Sjahril and Mii in press), by using 2 h of inoculation instead of 10 h recommended. Also, instead of using 500 mg l⁻¹ cefotaxime for bacteria elimination, cells were washed with 10 mg l⁻¹ meropenem for 30 min and then transferred onto new sterilized filter paper placed on 40 ml selective medium, which is 2.5 g l⁻¹ Gelrite-solidified NDM-20S supplemented with 25 mg l⁻¹ hygromycin and 5 mg l⁻¹ meropenem. Meropenem was omitted after the fifth or sixth subculture at 2 week-intervals by visually confirming the absence of *Agrobacterium*.

Most of the cells turned brown and died after 4 to 6 weeks on selection medium. After another month, small creamy white cell colonies with hygromycin-resistance initiated to grow among browned or dead cell population and formed small greenish yellow calli. Approximately 19 hygromycin-resistant callus clones were selected per 1 g of cells co-cultivated with *Agrobacterium* in three separate transformation experiments. The yield of hygromycin-resistant calli was much higher than that obtained in the previous report (Belarmino and Mii




Figure 1. T-DNA region of the binary vector pEKH-WT used for *Phalaenopsis* genetic transformation and molecular analyses of the transgenic plants. (A) T-DNA regions of the binary vector pEKH-WT. The genes for wasabi defensin and hygromycin phosphotransferase (*hpt*) are driven by 35S promoter and the gene for neomycin phosphotransferase (*npII*) is driven by nopaline synthase promoter (*nos-pro*). *RB* and *LB* Right and left border sequences of the T-DNA region, respectively. Bold arrow, PCR-amplification regions for confirming *hpt* gene (1.2 kb) and wasabi defensin gene (0.7 kb). Dotted lined box Regions used as the probe for wasabi defensin gene. 35S-P Cauliflower mosaic virus 35S promoter. Wasabi defensin Wasabi defensin cDNA, *Nos-T* Terminator of the nopaline synthase gene. *HindIII*, *XbaI* Restriction enzyme recognition sites. (B) Detection of *hpt* (1.2 kb) and wasabi defensin (0.7 kb) genes in the putative transgenic *Phalaenopsis* plantlets by PCR. Lane *M* molecular marker (ϕ X174/*HaeIII*), lane *P* plasmid pEKH-WT as positive control. Lane *C* non-transformed control, lanes 1–11 different transgenic *Phalaenopsis* plantlets #1 to #11. (C) and (D) Southern blot analysis of genomic DNA of transgenic plantlets, digested with *HindIII* (C) and *XbaI* (D), respectively. Lane *C* non-transformed control plant, lanes 1–11 transgenic plantlets. Lane *M* digoxigenin (DIG)-labeled molecular marker, lane *N* negative control.

2000), in which 1 g cell clumps or clusters only yielded about five hygromycin-resistant clones after selection with 50 mg l⁻¹ hygromycin. The use of finer cell aggregates and lower concentration of hygromycin in the present study might have resulted in the

higher yield probably because of more chance to make *Agrobacterium* cells contact directly with the *Phalaenopsis* cells. The fine cell aggregates after infection might also be exposed directly to the selection antibiotics, resulting in less escaped calli.

Putative transgenic callus selected were transferred onto 2.5 g l^{-1} Gelrite-solidified NDM-20S containing 25 mg l^{-1} hygromycin for further proliferation. One month after the transfer, the calli were sub-cultured to PGR free Gelrite-solidified NDM containing 10 g l^{-1} maltose (NDM-10M) and the same hygromycin concentration for 1 month for callus greening. Green calli produced were subcultured into NDM-10S containing 10 mg l^{-1} sucrose with 30 mg l^{-1} hygromycin for regeneration of PLBs and plantlets for 3–4 months with monthly subcultures.

Approximately 78% of the hygromycin-resistant callus clones, regenerated into hygromycin-resistant PLBs on PGR-free NDM-10S containing 30 mg l^{-1} hygromycin and about 30 transgenic plantlets were obtained from 1 g fresh weight of embryonic suspension calli, after 7–8 months of culture after infection.

The presence and integration of wasabi defensin gene in the plant genome was indicated by PCR of the extracted genomic DNA from putative transgenic plantlets. Total genomic DNAs were isolated from young leaves of putative transgenic plants and control plant following the CTAB method (Murray and Thompson, 1980). PCR reaction was performed using genomic DNA as a target and oligonucleotide primers (Bex Co. Ltd., Tokyo, Japan) for the wasabi defensin and *hpt* genes. All eleven hygromycin-resistant plantlets showed positive specific amplification of approximately 0.7 kb fragment for wasabi defensin gene (Figure 1B) and approximately 1.2 kb for *hpt* gene, respectively (Figure 1B), suggesting the transgenic nature of the *Phalaenopsis* plantlets analysed.

Integration and the number of insertion sites of the transgenes in the *Phalaenopsis* genome were examined by Southern blot analysis. Genomic DNA samples ($10 \mu\text{g}$) from hygromycin-resistant transgenic plantlets and control plantlet were digested with *Hind*III and *Xba*I and electrophoresed on a 0.8% (w/v) agarose gel, blotted to nylon membrane (Immobilon-NY⁺, Millipore, Bedford, MA, USA) and hybridised with digoxigenin-labeled probe of the wasabi defensin gene following the manufacturer's instruction (Roche Diagnostics GmbH, Mannheim, Germany). The insertion of the wasabi defensin gene into the genome was confirmed in both the digest with *Hind*III (Figure 1C) and *Xba*I (Figure 1D), respectively, whereas it was not detected in non-transformed plant. Digestion with *Xba*I also revealed that the number of inserted T-DNA copies varied from 1 to 4 in the independent 11 transgenic clones (Figure 1D).

For Western blot analysis, total proteins were extracted




Figure 2. Defensin production and resistance of transgenic *Phalaenopsis* plant against *Erwinia carotovora*. (A) Western blot analysis of the expression of wasabi defensin protein in leaves of *Phalaenopsis* plantlets. Arrow indicates the 5 kDa band of the wasabi defensin protein. Note the varying level of expression among different clones. C non-transformed control plantlet, lanes 1, 2, 4, 5, 6, 7, 8, 10 correspond to the different transgenic clone numbers shown in Figure 1B to 1D. (B) Evaluation of transgenic clone #2 of *Phalaenopsis* Wataboushi '6.13' for resistance to bacterial soft rot, one week after the inoculation. Black arrows represent the points of inoculation with the bacteria. Note the soft rot symptom in the control plant (left), whereas no sign of symptom in the transformant (right).

from 50 mg of leaf tissues from non-transgenic and transgenic plantlets, fractionated by 15% SDS-PAGE and electro blotted onto PVDF membrane (Bio-Rad, BioActive Co. Hercules, CA, USA). Detection of the wasabi defensin protein was performed using ECL Advance Western Blotting Detection Kit (Amersham Bioscience, Buckinghamshire, UK). Antibodies used were polyclonal antisera raised in rabbit against the defensin protein (primary antibody at 1 : 500 v/v) and goat-anti-rabbit IgG conjugated to horseradish peroxidase (HRP) as secondary antibody at 1 : 10,000 v/v.

As a result, the 5 kDa peptide which corresponds to the translated form of the wasabi defensin (Kanzaki et al. 2002), was detected as a single band in the leaves of transgenic clones but not in the non-transgenic control plant. The level of wasabi defensin expression varied among the clones (Figure 2A) and clones #2 and #8 showed relatively stronger expression. Similar difference in the expression levels of the same protein has also been reported in other species (Fagard and Vaucheret 2000; Kanzaki et al. 2002). It is interesting to note that the control plant showed no expression of the defensin protein at all, suggesting that *Phalaenopsis* might not have this specific anti-fungal protein at all or maybe very little.

In order to evaluate disease resistance of the

transgenic plant, bacterial strain of *Erwinia carotovora* was isolated from a *Phalaenopsis* plant severely showing symptom of soft rot disease in a greenhouse and grown on potato sucrose agar (PSA; 500 ml of potato broth made from 200 g potato, 20 g sucrose and 15 g agar in 1000 ml Milli-Q water, pH 5.6). For testing resistance of transformant against *E. carotovora*, non-transformed and transgenic plantlets grown on Gelrite-solidified NDM-10S were infected with the bacteria by inoculating at 4 points of one leaf for each plant using sterile needles. One week after the infection treatment, all non-transformant plantlets tested were susceptible to *E. carotovora* and died with soft rot symptom, whereas transformed plantlets showed strong resistance against inoculation of the bacteria (Figure 2B). Out of 15 transgenic clones tested, almost all showed no symptom 1 week after inoculation except for 1 clone (6.7%), which showed some susceptibility and died 4 weeks after the treatment.

We have succeeded in efficiently producing transgenic plants with wasabi defensin gene of *Phalaenopsis* orchid by using the transformation protocol used in the present study. Since almost all the transgenic plants successfully showed strong resistance to *E. carotovora*, they are expected to be resistant to some other important diseases of this orchid.

Acknowledgement

We thank Dr. Ken Tokuhara, Orchid Sanctuary Dogashima, for providing embryonic callus culture used in the study.

References

- Anzai H, Ishii Y, Shichinohe M, Katsumata K, Nojiri C, Morikawa H, Tanaka M (1996) Transformation of *Phalaenopsis* by particle bombardment. *Plant Tissue Culture Lett* 13: 265–272
- Belarmino MM, Mii M (2000) *Agrobacterium*-mediated genetic transformation of a *Phalaenopsis* orchid. *Plant Cell Rep* 19: 435–442
- Bohlmann H (1994) The role of thionins in plant protection. *Crit Rev Plant Sci* 13: 1–16
- Broekaert WF, Terras FRG, Cammue BPA, Osborn RW (1995) Plant defensins: novel antimicrobial peptides as components of the host defense system. *Plant Physiol* 108: 1353–1358
- Chakrabarti A, Ganapathi TR, Mukherjee PK, Bapat VA (2003) MSI-99, a magainin analogue, imparts enhanced disease resistance in transgenic tobacco and banana. *Planta* 216: 587–596
- Chia TF, Chan YS, Chua NH (1994) The firefly luciferase gene as a non-invasive reporter for *Dendrobium* transformation. *Plant J* 6: 441–446
- Epple P, Apel K, Bohlmann H (1997) Overexpression of an endogenous thionin enhances resistance of *Arabidopsis* against *Fusarium oxysporum*. *Plant Cell* 9: 509–520
- Fagard M, Vaucheret H (2000) Systemic silencing signal(s). *Plant Mol Biol* 43: 285–293
- Florack DEA, Stiekema WJ (1994) Thionins: properties, possible biological roles, and mechanisms of action. *Plant Mol Biol* 26: 25–37
- Gao AG, Hakimi SM, Mittanck CA, Wu Y, Woerner BM, Stark DM, Shah DM, Liang J, Rommens CMT (2000) Fungal pathogen protection in potato by expression of a plant defensin peptide. *Nat Biotechnol* 18: 1307–1310
- Kanzaki H, Nirasawa S, Saitoh H, Ito M, Nishihara M, Terauchi R, Nakamura I (2002) Overexpression of the wasabi defensin gene confers enhanced resistance to blast fungus (*Magnaporthe grisea*) in transgenic rice. *Theor Appl Genet* 105: 809–814
- Kiba A, Saitoh H, Nishihara M, Omiya K, Yamamura S (2003) C-terminal domain of a hevein-like protein from *Wasabia japonica* has potent antimicrobial activity. *Plant Cell Physiol* 44: 296–303
- Liang J, Wu Y, Rosenberger C, Hakimi SM, Castro S, Berg J (1998) AFP genes confer disease resistance to transgenic potato and wheat plants. In: Abstract no. L-49. *5th International Workshop on Pathogenesis-related Proteins in Plants: Signalling Pathways and Biological Activities 1998*. Aussois, France
- Murray MG, Thompson WF (1980) Rapid isolation of high molecular weight plant DNA. *Nucl Acids Res* 8: 4321–4325
- Ono H, Tesaki S, Tanabe S, Watanabe M (1998) 6-methylsulfinylhexyl isothiocyanate and its homologues as food-originated compounds with antibacterial activity against *Escherichia coli* and *Staphylococcus aureus*. *Biosci Biotechnol Biochem* 62: 363–365
- Pedras MSC, Sorensen JL, Okanga FI, Zaharia I (1999) Wasalexins A and B, new phytoalexins from wasabi: isolations and synthesis and antifungal activity. *Bioorgan Medic Chem Lett* 9: 3015–3020
- Saitoh H, Kiba A, Nishihara M, Yamamura S, Suzuki K, Terauchi R (2001) Production of antimicrobial defensin in *Nicotiana benthamiana* with a potato virus X vector. *Mol Plant Microbe Interact* 14: 111–115
- Sjahril R, Mii M (in press) High-efficiency *Agrobacterium*-mediated transformation of *Phalaenopsis* using meropenem, a novel antibiotic to eliminate *Agrobacterium*. *J Hort Sci Biotech*
- Terras FRG, Eggermont K, Kovaleva V, Raikhel NV, Osborn RW, Kester A, Rees SB, Torrekens S, Van Leuven F, Vanderleyden J, Cammue BPA, Broekaert WF (1995) Small cysteine-rich antifungal proteins from radish: their role in host defense. *Plant Cell* 7: 573–588
- Thevissen K, Ghazi A, De Samblanx GW, Brownlee C, Osborn RW, Broekaert WF (1996) Fungal membrane responses induced by plant defensins and thionins. *J Biol Chem* 271: 15018–15025
- Tokuhara K, Mii M (1993) Micropropagation of *Phalaenopsis* and *Doratenopsis* by culturing shoot tips of flower stalk buds. *Plant Cell Rep* 13: 7–11

Transgenic Phalaenopsis plants with resistance to Erwinia carotovora produced by introducing wasabi defensin gene using Agrobacterium method

ORIGINALITY REPORT

17%

SIMILARITY INDEX

7%

INTERNET SOURCES

15%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Kushikawa, S.. "Agrobacterium-mediated transformation of Saintpaulia ionantha Wendl.", Plant Science, 2001 10 2%
Publication

- 2** B. López-García, B. San Segundo, M. Coca. "Antimicrobial Peptides as a Promising Alternative for Plant Disease Protection", American Chemical Society (ACS), 2012 1%
Publication

- 3** R. S. Khan. "Production and selection of marker-free transgenic plants of Petunia hybrida using site-specific recombination", Biologia Plantarum, 06/2010 1%
Publication

- 4** M. Ogaki. "Importance of co-cultivation medium pH for successful Agrobacterium-mediated transformation of Lilium × formolongi", Plant Cell Reports, 04/2008 1%
Publication

5

Midori Sugimoto. "Genetic diversity of the entomopathogen *Verticillium lecanii* on the basis of vegetative compatibility", *Phytoparasitica*, 10/2003

Publication

1 %

6

V. Colova-Tsolova. "Progress in Genetic Engineering of Grapevine for Disease & Stress Tolerance", *Grapevine Molecular Physiology & Biotechnology*, 2009

Publication

1 %

7

Submitted to Gyeongsang National University

Student Paper

1 %

8

C.-H. Liau, S.-J. You, V. Prasad, H.-H. Hsiao, J.-C. Lu, N.-S. Yang, M.-T. Chan. "Agrobacterium tumefaciens-mediated transformation of an *Oncidium* orchid", *Plant Cell Reports*, 2003

Publication

1 %

9

Nobuhiko Muramoto, Tomoko Tanaka, Takashi Shimamura, Norihiro Mitsukawa et al. "Transgenic sweet potato expressing thionin from barley gives resistance to black rot disease caused by *Ceratocystis fimbriata* in leaves and storage roots", *Plant Cell Reports*, 2012

Publication

1 %

10

www.tbiomed.com

Internet Source

1 %

11

biozoojournals.ro

Internet Source

<1 %

12

Zhong Hu. "Factors affecting *Agrobacterium tumefaciens*-mediated genetic transformation of *Lycium barbarum* L.", *In Vitro Cellular & Developmental Biology - Plant*, 09/2006

Publication

<1 %

13

www.plantphysiol.org

Internet Source

<1 %

14

Fraiture, Marie-Alice, Nancy H.C. Roosens, Isabel Taverniers, Marc De Loose, Dieter Deforce, and Philippe Herman. "Biotech rice: Current developments and future detection challenges in food and feed chain", *Trends in Food Science & Technology*, 2016.

Publication

<1 %

15

Masahiro Nishihara, Takashi Nakatsuka, Saburo Yamamura. "Flavonoid components and flower color change in transgenic tobacco plants by suppression of chalcone isomerase gene", *FEBS Letters*, 2005

Publication

<1 %

16

Nailish Samanani. "In vitro regeneration and genetic transformation of the berberine-producing plant, *Thalictrum flavum* ssp. *glaucum*", *Physiologia Plantarum*, 9/2002

Publication

<1 %

17	moam.info Internet Source	<1 %
18	www.frontiersin.org Internet Source	<1 %
19	Advances in Photosynthesis and Respiration, 2010. Publication	<1 %
20	K.-C. Cho. "Resistance to Rhizoctonia solani AG-2-2 (IIIB) in creeping bentgrass plants transformed with pepper esterase gene PepEST : Brown patch disease-resistant creeping bentgrass", Plant Pathology, 08/2011 Publication	<1 %
21	bmcplantbiol.biomedcentral.com Internet Source	<1 %
22	ir-library.ku.ac.ke Internet Source	<1 %
23	Abdul Aziz M.A. Mohamed, David B. Hogg, Yong Chul Park, Walter G. Goodman. " An Immunoassay Method for Detecting (Hymenoptera: Braconidae: Euphorinae) Parasitism in the Alfalfa Weevil, (Coleoptera: Curculionidae) ", Biocontrol Science and Technology, 2003 Publication	<1 %

24

Juan Yan, Su-su Yuan, Luan-luan Jiang, Xiu-juan Ye, Tzi Bun Ng, Zu-jian Wu. "Plant antifungal proteins and their applications in agriculture", Applied Microbiology and Biotechnology, 2015

Publication

<1 %

25

K. Hosokawa, R. Matsuki, Y. Oikawa, S. Yamamura. "Production of transgenic gentian plants by particle bombardment of suspension-culture cells", Plant Cell Reports, 2000

Publication

<1 %

26

Kanzaki Hiroyuki. "Regulation of expression of rice thaumatin-like protein: inducibility by elicitor requires promoter W-box elements", Plant Cell Reports, 09/2008

Publication

<1 %

27

Mei-Kuen Yip, Hsiang-En Huang, Mang-Jye Ger, Shih-Hua Chiu, Yuh-Chih Tsai, Chin-I Lin, Teng-Yung Feng. "Production of soft rot resistant calla lily by expressing a ferredoxin-like protein gene (pflp) in transgenic plants", Plant Cell Reports, 2006

Publication

<1 %

28

RICARDO B. FERREIRA. "The role of plant defence proteins in fungal pathogenesis", Molecular Plant Pathology, 9/2007

Publication

<1 %

29

Su-Juan You, Chia-Hui Liao, Hsiang-En Huang, Teng-Yung Feng, Venkatesh Prasad, Hsin-hao Hsiao, Jian-Cheng Lu, Ming-Tsair Chan. "Sweet pepper ferredoxin-like protein (pflp) gene as a novel selection marker for orchid transformation", *Planta*, 2003

Publication

<1 %

30

fjfsdata01prod.blob.core.windows.net

Internet Source

<1 %

31

M. Akutsu, T. Ishizaki, H. Sato. "Transformation of the monocotyledonous *Alstroemeria* by *Agrobacterium tumefaciens*", *Plant Cell Reports*, 2004

Publication

<1 %

32

S. Suzuki, M. Nakano. "Agrobacterium-mediated production of transgenic plants of *Muscari armeniacum* Leichtl. ex Bak.", *Plant Cell Reports*, 2002

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On