

Dr.Dwi_Kesuma_Adachi_2008.p df

by Dwi Kesuma Sari

FILE	DR.DWI_KESUMA_ADACHI_2008.PDF (174.2K)		
TIME SUBMITTED	03-MAR-2020 09:46AM (UTC+0700)	WORD COUNT	4979
SUBMISSION ID	1268120665	CHARACTER COUNT	27006

Development of neutralization antibodies against highly pathogenic H5N1 avian influenza virus using ostrich (*Struthio camelus*) yolk

KAZUHIDE ADACHI¹, EKOWATI HANDHARYANI³, DWI KESUMA SARI³, KENTARO TAKAMA¹, KEIKO FUKUDA¹, ISAKO ENDO¹, RYOHEI YAMAMOTO², MASAKI SAWA², MASARU TANAKA², ITSURO KONISHI² and YASUHIRO TSUKAMOTO¹

¹Laboratory of Veterinary Anatomy, Graduate School of Biology and Environmental Sciences, Osaka Prefecture University, Gakuencho 1-1, Sakai, Osaka 599-8531; ²Ostrich Breeding Farm 'Ostrich-Kobe', Kande-Housei, Nishiku, Kobe-City, Hyogo 651-2321, Japan; ³Departments of Veterinary Pathology and Veterinary Anatomy, Faculty of Veterinary Medicine, Bogor Agricultural University, Jl. Agatis Kampus IPB, Darmaga, Bogor 16680, Indonesia

Received December 3, 2007; Accepted January 4, 2008

Abstract. The rapid outbreak of the highly pathogenic H5N1 avian influenza virus and its transmission to humans have induced world-wide fears of a new influenza pandemic. The most effective method for the reduction of the impact of such a pandemic would be prophylaxis with a safe and effective vaccine, as well as anti-viral materials. In this study, we generated the specific antibodies (immunoglobulin yolk (IgY) from ostrich eggs immunized with a full-length glycosylated recombinant H5 protein of the strain H5N1/Vietnam/1203/2004. Using this simple method, abundant specific antibody (about 200 g) against H5 was successfully produced by one female ostrich in a year. The IgY from the immunized ostrich eggs had strong reactivity to the H5N1 virus as well as to H5 proteins. Furthermore, the antibodies strongly inhibited cytopathic effects in MDCK cells and prevented the death of an embryonated chick after a viral inoculation, indicating strong neutralization activity against H5N1 infections. These findings suggest that the neutralization antibody produced by the H5-immunized ostrich is suitable for industrial purposes, such as the development of antibody-binding filters, which can be applied to a mask or to air-conditioners to prevent the influenza pandemic through antigen-antibody reactions. Of note, the mortality rate of chicks inoculated with the H5N1 virus was dramatically decreased with antibody injection. This indicates that ostrich IgY is a potentially effective therapeutic modality for H5N1 infection.

Correspondence to: Dr Yasuhiro Tsukamoto, Laboratory of Veterinary Anatomy, Graduate School of Biology and Environmental Sciences, Osaka Prefecture University, Gakuencho 1-1, Sakai, Osaka 599-8531, Japan

E-mail: ytsuka@vet.osakafu-u.ac.jp

Key words: antibody, influenza virus, H5N1, pandemic, ostrich, IgY

Introduction

Influenza is recognized as a zoonotic disease, with the most commonly affected animals being humans, pigs, horses and species of aquatic birds (1). Influenza viruses belong to the family *Orthomyxoviridae* and are divided into three types, A, B and C. A type viruses are responsible for major disease problems in humans as well as in avian species (2-4) and are further classified into subtypes on the basis of their antigenic properties, including hemagglutinin (HA) and neuraminidase on the viral particle. There is world-wide concern over the possibility of a new influenza pandemic, especially since the appearance of the highly pathogenic avian influenza viruses H5N1, which have the capacity for lethal infection in man. Since 1997, over 200 people have been infected with H5N1, resulting in a mortality rate of over 50% (5). In addition, the H5N1 virus has been able to spread by means of domestic and wild birds from Hong Kong, where it was first detected, across Asia, Africa, and into northern Europe. Should the H5N1 virus develop the ability to spread efficiently from man to man, there would be a high risk of worldwide pandemic causing considerable mortality and economic disruption (6). It has been discussed that the most effective method to reduce the impact of a pandemic would be prophylaxis with a safe and effective vaccine (6,7), as well as anti-viral materials including masks or facial tissues (8).

Antibodies are increasingly used for research, diagnosis and therapeutic purposes. However, antibodies from experimental mammals, including mice and rabbits, are not adapted for industrial use because of their high production cost. Of late, the avian egg as an antibody source has proven to be attractive for the non-invasive production of antibodies with applications in research, diagnosis and immunotherapy (9-11). In addition, the production of avian antibodies offers many advantages over mammalian antibodies in terms of their specificity for antigens, production cost and uses (9).

The predominant class of immunoglobulin in birds is immunoglobulin yolk (IgY), which is transferred from the serum to the yolk to confer passive immunity to the embryo (12). Although the IgY and IgG of mammals are functionally

equivalent, they differ with respect to certain physicochemical properties (13).

Recently, we developed a convenient method for the mass-production of antibodies using the female ostrich (*Struthio camelus*). The ostrich grows to up to 250 cm in height and 160 kg in weight, and has a life span of about 60 years. Its eggs weigh about 1.5 kg and are 30-fold bigger than chicken eggs. They can lay one hundred eggs every year. We can purify about 2 g of IgY per ostrich egg. Accordingly, about 200 g of IgY is obtained from only one ostrich in the course of a year. Recent developments in breeding have permitted low-cost egg production from the ostrich, because they can be fed the factory waste from bean sprout production. Accordingly, we strongly believe that the ostrich egg may provide an excellent source of antibody for industrial purposes (14).

In the present study, we tried to produce antibodies against H5N1 from ostrich eggs. The HA proteins of the influenza viruses were considered to be suitable antigens for the immunization of the ostrich for the following reasons: i) influenza virus entry is mediated by the receptor binding domain of HA, and adaptation of avian viruses to humans is associated with HA specificity, and ii) the blockage of HA antigens leads to inactivation of the viral infection to host cells. We herein show that a high amount of neutralizing antibody against H5N1 was generated by the ostrich using a simple and economical method, which may lead to the development of anti-viral materials against an influenza pandemic.

Materials and methods

Generation of antibodies against influenza virus HA antigens. A mixture of HA antigens from vaccine strains of the human influenza virus (H1N1/NewCaledonia/20/99, H3N2/Hiroshima/52/2005 and B/Malaysia) (The Kitasato Institute Research Center for Biologicals, Japan) were used as antigens for the immunization of the ostrich, as well as H5 recombinant protein (rH5) purified from insect cells transfected with baculovirus vector inserting H5 gene of H5N1/Vietman/1203/2004 (Protein Sciences, USA).

The laying ostriches were immunized intra-muscularly in the lumbar region at multiple sites with the mixture of HA (30 µg/bird) or rH5 (50 µg/bird) emulsified in Freund's complete adjuvant (FCA). The boosters were administered every other week with both the antigens in Freund's incomplete adjuvant (FIA). Eggs were collected daily from the initial immune day. The yolk was separated from the albumin using egg yolk separator, the volume of the yolk was recorded and it was stored at -20°C. Serum samples were also collected every week and stored at -20°C.

The yolk was separated from the albumin of the eggs and diluted to 5-fold with TBS-buffer [0.02 M Tris/HCl (pH 7.5), 0.15 M NaCl], increased 1- to 10-fold with 30% dextran sulfate in TBS and 2- to 3-fold with 2.5 M CaCl₂ in TBS, then stored at 4°C for at least 4 h. The supernatant containing IgY was collected by centrifugation (10000 x g at 4°C for 15 min) and precipitated with 45% saturated ammonium sulfate. The solution was centrifuged again at 10000 x g at 4°C for 15 min. The precipitate was then redissolved in TBS and dialyzed against PBS. Finally, the purified antibody solutions were verified by 10% SDS-PAGE under non-reducing or

reducing conditions and stained with Coomassie Brilliant Blue (CBB).

Generation of rabbit polyclonal antibodies against ostrich IgY. An experimental rabbit (Japanese-White) was immunized subcutaneously in the lumbar region at multiple sites with purified ostrich IgY (50 µg/body) emulsified in FCA. The boosters were given every other week with the same antigen in FIA. At the hyper-immunized stage, confirmed by ELISA, the serum was collected and the IgG purified with protein G as a routine procedure. The purified antibody fractions were verified by 10% SDS-PAGE under non-reducing conditions and stained with CBB. Finally, the purified IgG was conjugated with horseradish peroxidase (HRP) or FITC using the appropriate labeling kits (Dojin, Japan), then used as a secondary antibody (14).

Western blot analysis. rH5 proteins were separated with 10% SDS-PAGE in reducing conditions and electroblotted onto a nitrocellulose membrane (Bio-Rad). The membrane was air dried for 5-10 min, then blocked with blocking reagent (2% skim-milk in PBS) for 1 h. After washing with a washing buffer (0.01 M PBS, pH 7.4 with 0.05% Tween-20), the membrane was incubated with the ostrich IgY against rH5 (1:4000) in the same blocking buffer at room temperature for 1 h on a horizontal shaker. Next, the membrane was washed and incubated with 1:5000 dilutions in blocking buffer of HRP-conjugated rabbit IgG against ostrich IgY at room temperature for 1 h. After thorough washing, the membrane was placed in peroxidase chromogenic substrate solution (0.01 M PBS, pH 7.4 with 0.05% diaminobenzidine and 0.03% H₂O₂) for 5 min, then immersed in excess distilled water to stop the color development. Finally, the Western blot membrane was digitally recorded using a gel documentation system (Fujifilm, Japan).

Enzyme linked immunosorbent assay (ELISA) using ostrich IgY. Each well of polystyrene ELISA plates (Sumitomo Bakelite, Japan) was coated with 0.2 µg HA antigens or rH5 protein in phosphate-buffered saline (PBS) and the plate was incubated overnight at 4°C. Each of the following incubation steps were preceded by washing the wells twice with PBS containing 0.05% Tween-20. The wells were blocked for non-specific binding by the addition of a commercial blocking buffer (DS Pharma Biomedical, Japan) and incubated at 37°C for 2 h. The serial dilutions of purified IgY were added vertically to the wells and kept for incubation at 37°C for 1 h. HRP-conjugated rabbit IgG diluted (1:5000) in PBS was dispensed into each well. The plate was incubated for 1 h at 37°C. Later, a substrate buffer containing TMB (Sumitomo Bakelite) was added to each well and kept for incubation at 37°C for 15 min. The reaction was terminated by the addition of a stopping reagent (1.25 M sulfuric acid). The absorbance was recorded at 450 nm using the ELISA plate reader (DS Pharma Biomedical).

Viral infectivity (TCID₅₀ and EID₅₀) assays. The H5N1/Indonesian strain cloned from affected chickens was used throughout this study. The viral solutions were titered using a cell culture system (MDCK cells) onto 96-well microtiter

2

13 MOLECULAR MEDICINE REPORTS 1: 203-209, 2008

205

plates or embryonated chick eggs by serial 10-fold dilutions of the samples as a routine procedure.

Immunocytochemistry. MDCK cells seeded at a density of 5×10^4 on the well of the microtiter plate were infected with H5N1 virus (100TCID_{50}) for 2-3 days at 35°C . The cultures showing cytopathic effect (CPE) were fixed with 10% buffered formalin for immunocytochemistry. After washing in PBS, the cells were incubated with the ostrich IgY (1:4000) against HA or rH5 for 1 h at 37°C , and then incubated with FITC-conjugated rabbit IgG (1:4000) against ostrich IgY following a sufficient number of washes in PBS. Finally, the specific signal was observed under fluorescence microscopy.

Neutralization assays for H5N1 infection. For the standard neutralization assay on cell cultures, serial dilutions in 2-fold step ostrich IgY were mixed at a ratio of 1:1 with H5N1 virus (100TCID_{50}), incubated for 1 h at 37°C , and then transferred to a microtiter plate with a MDCK monolayer. After 2-3 days incubation at 35°C , the cultures were inspected for CPE. The neutralizing titer, expressed as the reciprocal of the IgY dilution at which virus growth is 50% inhibited, was calculated by the number of virus negative wells and the IgY dilution according to the report by Reed *et al* (15).

For the assay on the embryonated eggs, serial dilutions in a 4-fold step of ostrich IgY were mixed at a ratio of 1:1 with H5N1 virus (100EID_{50}), incubated for 1 h at 37°C , and injected into the allantoic cavity of the embryonic chicken eggs (10 days). After 2-3 days incubation at 35°C , the embryos were inspected for their death. The neutralizing titer, expressed as the reciprocal of the IgY dilution at which virus growth is 50% inhibited, was calculated by the number of survivors and the IgY dilution.

Antibody injection into H5N1-infected chicks. At 10 days of age, the chicks were intra-nasally inoculated with H5N1 virus at a dose of $10^{7.8}\text{TCID}_{50}$. They were injected intra-muscularly with PBS or preimmune IgY ($10 \mu\text{g}/\text{bird}$) or IgY against rH5 ($10 \mu\text{g}/\text{bird}$) at 1 h post-inoculation with H5N1. Then all chicks were boarded in individual cages in a BSL3 laboratory in Indonesia (Vaksindo Satwa Nusantara). At 5-days post-viral-challenge, the number of dead chicks was counted and the results were expressed as the mean of dead birds in each experimental group (over 4 individuals in each group).

Results

Ostrich antibodies against HA antigen of influenza viruses. The IgY from the immunized ostrich with influenza virus HA antigens was purified from the yolk and visualized by SDS-PAGE. The molecular weight of ostrich IgY is around 200 kDa (heavy chain, 64 kDa; light chain, 28 kDa) and is heavier than that of rabbit IgG (Fig. 1). In a Western blot analysis, using a lysate of H5N1, the purified IgY from the rH5-immunized ostrich recognized the doublet bands of 45 and 25 kDa, corresponding to HA₁ and HA₂ of the H5 antigens, respectively (Fig. 2). This indicated that the antibody against the H5 antigen of H5N1 was generated in the ostrich eggs.




Figure 1. SDS-PAGE analysis of rabbit IgG and ostrich IgY. (A) Molecular weight of Rabbit IgG and ostrich IgY. Purified rabbit IgG and ostrich IgY were loaded under non-reducing conditions. The intact ostrich IgY is around 200 kDa - heavier than rabbit IgG, which is about 150 kDa. A faint band under 82 kDa in the ostrich IgY sample seems to be a heavy chain of IgY. (B) The heavy and light chains of ostrich IgY. The samples were heated at 95°C for 2 min under reducing conditions. The doublet bands of 64 and 28 kDa correspond to a heavy and light chain of IgY, respectively. M, molecular weight marker.




Figure 2. Western blot analysis of HA antigen of the H5N1 virus. The reaction of ostrich antibodies to baculovirus-derived H5 recombinant protein (rH5) was examined by Western blotting. The HA (45 kDa) and HA₂ (25 kDa) proteins were strongly recognized by the IgY from rH5 immunized ostrich and slightly by HA (H1, H2, B) immunized ostrich. In contrast, no reaction was found by preimmune IgY.

The IgY against each HA antigen was titered by ELISA in time course from the primary immunization day. The titers were increased dramatically in both serum and yolk from 2 weeks after the primary immunization time and reached the maximal peak at 8 weeks (data not shown). At the hyper-immune stage from 8 weeks on, about 2 g of IgY could be purified from one egg, indicating that it is possible to collect 200 g of IgY against the influenza virus from one female ostrich in one year. The ELISA titers of IgY against HA antigens are shown in Table I. The IgY from eggs immunized with HA-mixtures of antigens from the vaccine strains showed high reactivity to H1N1, H3N2 and B strains, but reacted

Table I. ELISA titers of ostrich IgY against influenza viruses HA antigens.

Antibodies against indicated HA antigens	Concentration (mg/ml)	Reciprocal IgY titer against indicated influenza virus HA antigens			
		H1N3	H3N2	B	H5N1
H1, H3, B	2	102400	204800	102400	3200
rH5	2	800	1600	800	51200

The IgY activities to each HA antigens of influenza viruses were measured by ELISA. The antibody titer was defined as the reciprocal of the highest IgY dilutions that produced an ELISA signal twice as intense as the signal from equivalently diluted preimmune IgY.




Figure 3. Immunocytochemistry of H5N1-infected MDCK cells. MDCK cells were reacted with ostrich IgY, and visualized by FITC-conjugated secondary antibody. Normal MDCK cells have no immunoreactivity to all antibodies. The H5N1-infected cells are not stained with preimmune IgY. In contrast, the cytoplasm of infected cells are stained strongly with the IgY from H5 immunized ostrich, but only slight with IgY from mixed HA (H1, H3, B) immunized bird. Bar, 50 μ m.

more weakly to the H5N1 strain. In contrast, IgY from rH5-immunized eggs reacted strongly to H5N1, but only weakly to the H1N1, H3N2 and B strains.

Next, the reactivity of ostrich IgY was examined in immunocytochemistry. MDCK cells were infected with the Indonesia strain of H5N1 and stained with antibodies. The cytoplasm of infected cells was strongly labeled with IgY from rH5-immune eggs, but not with pre-immune IgY (Fig. 3). In contrast, IgY from eggs immunized with HA from the vaccine strains showed only weak immunoreactivity to the H5N1-infected cells.

These results indicate that high titers of antibodies against the H5N1 virus are collected from eggs immunized with H5 antigens. For this reason, antibodies from the H5-hyperimmune eggs were used for further examinations.

Neutralization assays for H5N1 infection. MDCK cells and embryonated eggs were used for the neutralization assays. The H5N1 virus was reacted with serial dilutions of IgY from preimmune or rH5-immunized eggs, and neutralization was determined by the observation of CPE and embryonic death. IgY with rH5-immunization showed very strong inhibitory

effects on H5N1 infection in either MDCK or chick embryos (Fig. 4 and Table II). In contrast, H5N1 infection was not blocked in either of the analyses, even at the highest concentrations of preimmune IgY, indicating no neutralization activity on the H5N1 infection.

Effect of ostrich IgY on H5N1-infected chicks. The inhibitory activity of IgY from rH5-immunized ostrich was examined in H5N1-infected chicks in order to elucidate the potential therapeutic uses of ostrich IgY.

A high dosage of the H5N1 Indonesian strain ($10^{7.9}$ TCID₅₀) was intra-nasally challenged to 10-day-old chicks. All the birds (100%) died by 5-days post-inoculation and experienced severe clinical symptoms including dehydration and depression. In contrast, the mortality rate of the infected birds was dramatically decreased by the intra-muscular injection with anti-H5 IgY: only 25% of the birds died by viral infection, and the survivors had no symptoms (Fig. 5). All infected birds died when the preimmune IgY was injected instead of the rH5-immune antibody. Accordingly, the anti-H5 IgY had a therapeutic effect on H5N1-infected chickens, even at the low dosage.




Figure 4. Neutralization activities on ostrich IgY for H5N1 infection. (A) MDCK cells infected with H5N1 virus show severe CPE on their monolayer sheet. IgY from H5 immunized ostrich (anti-H5 IgY) inhibit CPE, while preimmune IgY has no effect on the cells. Bar, 50 μm . (B) Neutralization activities of ostrich IgY on H5N1 infection were analyzed using MDCK cells and embryonated chick embryos. Anti-H5 IgY shows inhibitory activities on H5N1 infection in both MDCK and chick embryos, while preimmune IgY has no effect, even at the highest dosage.

Table II. Neutralizing titers of ostrich IgY against H5N1 strain.

Antibodies against indicated HA antigens	Neutralizing titers (50% inhibition)	
	MDCK cells ($\mu\text{g/ml}$)	Embryonic eggs ($\mu\text{g/ml}$)
Preimmune IgY	>384	>768
rH5	6.7	63.4

The neutralization assays were performed using MDCK cells and chick embryos. The neutralizing titers are indicated as the 50% inhibition on CPE or embryonic death.

Discussion

In the present study, large-scale and economical antibodies against the highly pathogenic avian influenza virus H5N1 were generated by employing the female ostrich. To do so, HA antigens were immunized to obtain neutralization antibodies. Since HA binds to the receptors and is essential for viral entrance into the host cells, these antigens are widely used in the vaccine for influenza (17,18). Recently, various attempts have been made to prevent H5N1 pandemic infections (7,19,20). H5 antigens are used to develop vaccines against H5N1, and therapeutic monoclonal antibodies by H5 vaccinations can be evaluated before the emergence of human-adapted H5N1 strains (19,21).




Figure 5. Effects of ostrich IgY on H5N1 infected chicks. The chicks were intra-nasally challenged with high titered H5N1 virus and infected with ostrich IgY. All infected birds died by 5 days, and preimmune IgY had no inhibitory effect on mortality. In contrast, the mortality of infected chicks was remarkably decreased by an injection with IgY from H5-immunized ostrich (anti-H5 IgY).

The HA antigens were also highly antigenic in the ostrich; we could obtain ample IgY from the female birds by immunizing them with the HA antigens of the human influenza vaccine strains of H1N1, H3N2, B and the highly pathogenic avian influenza virus H5N1. Since ostrich IgY are polyclonal, the antibodies generated with the HA of the human vaccine strains are slightly reactive to H5, which are likely to be raised from the homology of epitopes among these antigens.

We used the IgY from the rH5-immunized ostrich for neutralization and therapeutic assays against H5N1 infections,

because this antibody had a much higher titer on the H5 antigen in comparison to other antibodies with human vaccine strains. In addition, rH5-immunized IgY was highly reactive to the H5N1-infected MDCK cells in immunocytochemistry. Therefore, this rH5-immunized IgY might be developed as a diagnostic tool for the H5 virus. In an ELISA against the H5 antigen, the titer of rH5-immunized IgY reached maximum levels at 8 weeks post initial immunization in the serum and yolk. The antibodies were successfully transferred from blood to yolk, which can then be non-invasively obtained by simple collection of the eggs, making it possible to obtain a high amount of anti-H5N1 antibody (200 g in total) from one female ostrich in a year. In neutralization assays, rH5-immunized IgY could inhibit H5N1 infection successfully in either MDCK cells or embryonic eggs. The H5N1 virus is highly virulent to chick embryos and it is difficult to protect them from virus infection using antibodies. However, in the present study ostrich IgY could inhibit H5N1 infection even in the embryos, although the volume of antibodies needed for viral protection was larger in embryos than in MDCK cells.

For the protection of poultry, neutralizing antibodies to HA neuraminidase proteins have been reported to provide primary protection against highly pathogenic avian influenza (22). In addition, the prophylactic and therapeutic efficacy of human monoclonal antibodies was elucidated by the H5N1 infections (21). One particularly interesting and potentially important finding of the present study was the observation that the chicks were protected by a low dosage of H5 immunized IgY from the H5N1 infection. This finding leads to the potential development of ostrich IgY for therapeutic treatment against an H5N1 pandemic. For the protection of poultry, a neutralizing antibody to the haemagglutinin and neuraminidase proteins provides the primary protection against the highly pathogenic avian influenza. It has been reported that chicken IgY has been used for therapeutic purposes: IgY preparations are administered to patients suffering from infectious diseases including rotaviruses, enterotoxigenic *Escherichia coli*, coronavirus, *Salmonella spp.*, *Edwardsiella tarda*, *Yersinia ruckeri*, *Staphylococcus* and *Pseudomonas* (23,24). Therefore, we believe that ostrich IgY is a promising alternative to antibiotics for passive immunotherapy application as well as for chicken antibody.

Because most of the human population has no immunity to H5N1, the most effective method for the reduction of the impact of a pandemic has been reported to be prophylaxis with a safe and effective vaccine (6). In addition, anti-viral materials are recommended for the prevention of an influenza pandemic (8). In this study, a large amount of neutralization antibodies against H5N1 were produced with cost effectiveness, which indicates the potential use of ostrich antibody for industrial purposes, thus including the development of anti-viral materials. It is speculated that the IgY on the filters can capture the viral particles of H5N1 by antigen-antibody reactions, so the virus would be trapped by the filters and not pass through them. In addition, the virus might be neutralized on the filters, because HA is masked with IgY and cannot enter host cells. In the preliminary study, we found that the infectivity of H5N1 was dramatically decreased by the filters adsorbed with ostrich IgY generated by H5 immunizations (data not shown). We are now attempting to apply the ostrich

IgY onto masks of either the N95 or surgical type to protect humans from an influenza pandemic caused by the H5N1 virus. In the event of such a pandemic, where effective vaccine and antiviral medicines may be lacking, disrupting the environmental transmission of the H5N1 virus will be the only viable strategy for the protection of the people. We believe that filters impregnated with ostrich antibodies will be a powerful tool for protection against the influenza pandemic.

Acknowledgements

The authors thank Drs Saeko Morikawa, Satoshi Hiroi, Takahiro Yumisashi and Tetsuo Kase at the Osaka Prefectural Institute of Public Health for their helpful comments. This study was supported in part by Research Fellowships from the Japan Society for the Promotion of Science (Nos. 17688013 and 18658115) and by an innovation grant from the Japan Science and Technology Agency (JST; No. 1804).

References

1. McCauley JW, Pullen LA, Forsyth M, Penn CR and Thomas GP: 4-Guanidino-Neu5Ac2en fails to protect chickens from infection with highly pathogenic avian influenza virus. *Antiviral Res* 27: 179-186, 1995.
2. Alexander DJ and Brown IH: Recent zoonoses caused by influenza A viruses. *Rev Sci Tech* 19: 197-225, 2000.
3. Rimmelzwaan GF and Osterhaus AD: Influenza vaccine: new developments. *Curr Opin Pharmacol* 1: 491-496, 2001.
4. Normile D: North Korea collaborates to fight bird flu. *Science* 308: 175, 2005.
5. Leroux-Roels I, Borkowski A, Vanvolleghem T, Drame M, Clement F, Hons E, Devaster JM and Leroux-Roels G: Antigen sparing and cross-reactive immunity with an adjuvanted rH5N1 prototype influenza vaccine: a randomised controlled trial. *Lancet* 370: 580-589, 2007.
6. Poland GA: Vaccines against avian influenza - a race against time. *N Engl J Med* 354: 1411-1413, 2006.
7. Sharpe M, Lynch D, Topham, Major D, Wood J and Loudon P: Protection of mice from H5N1 influenza challenge by prophylactic DNA vaccination using particle mediated epidermal delivery. *Vaccine* 25: 6392-6398, 2007.
8. CDC: Experiences with influenza-like illness and attitude regarding influenza prevention - United States, 2003-04 influenza season. *MMWR* 53: 1156-1158, 2004.
9. Schade R, Pfister C, Halatsch R and Henklein P: Polyclonal IgY antibodies from chicken egg yolk as an alternative to the production of mammalian IgG type antibodies in rabbits. *ATLA* 19: 403-419, 1991.
10. Schade R, Schniering A and Hlinak A: Polyclonal avian antibodies extracted from egg yolk as an alternative to the production of antibodies in mammals - a review. *ALTEX* 9: 43-56, 1992.
11. Gross M and Speck J: Avian yolk antibodies in diagnosis and research. *Dtsch Tierarztl Wochenschr* 103: 417-422, 1996.
12. Larsson A, Balow RM, Lindahl TL and Forsberg PO: Chicken antibodies: taking advantage of evolution-a review. *Poult Sci* 72: 1807-1812, 1993.
13. Leslie GA and Clem LW: Phylogeny of immunoglobulin structure and function. III. Immunoglobulins of the chicken. *J Exp Med* 130: 1337-1352, 1969.
14. Adachi K, Hagimori K, Kato H, Fukuda K, Kikuta M and Tsukamoto Y: Potential role of SC1, a cell adhesion molecule, in mammary gland tumors. *Mol Med Rep* (In press).
15. Reed LJ and Muench H: A simple method of estimating fifty percent end-points. *Am J Hyg* 27: 493-797, 1938.
16. Lu L, Yu L and Kwang J: Baculovirus surface-displayed hemagglutinin of H5N1 influenza virus sustains its authentic cleavage, hemagglutination activity, and antigenicity. *Biochem Biophys Res Commun* 358: 404-409, 2007.
17. Bosch FX, Orlin M, Klenk HD and Rott R: The structure of the hemagglutinin, a determinant for the pathogenicity of influenza viruses. *Virology* 95: 197-207, 1979.

18. Takada A, Kuboki N, Okazaki K, Ninomiya A, Tanaka H, Ozaki H, Itamura S, Nishimura H, Enami M, Tashiro M, Shortridge KF and Kida H: A virulent avian influenza virus as a vaccine strain against a potential human pandemic. *J Virol* 73: 8303-8307, 1999.
19. Yang ZY, Wei CJ, Kong WP, Wu L, Xu L, Smith DF and Nabel GJ: Immunization by avian H5 influenza hemagglutinin mutants with altered receptor binding specificity. *Science* 317: 825-828, 2007.
20. Kistner O, Howard OK, Spruth M, Wodal W, Bruhl P, Gerencer M, Crowe BA, Savidis-Dacho H, Livey I, Reiter M, Mayerhofer I, Tauer C, Grillberger L, Mundt W, Falkner FG and Barrett PN: Cell culture (Vero) derived whole virus (H5N1) vaccine based on wild-type virus strain induces cross-protective immune responses. *Vaccine* 25: 6028-6036, 2007.
21. Simmons CP, Bernasconi NL, Suguitan JAL, Mills K, Ward JM, Chau NVV, Tran Tinh, Hien TT, Sallusto F, Ha DQ, Farrar J, De-Jong MD, Lanzavecchia A and Subbarao K: Prophylactic and therapeutic efficacy of human monoclonal antibodies against H5N1 influenza. *PLoS Med* 4: 928-936, 2007.
22. Suarez DL and Schultz-Cherry S: Immunology of avian influenza virus: a review. *Dev Comp Immunol* 24: 269-283, 2000.
23. Mine Y and Kovacs-Nolan J: Chicken egg yolk antibodies as therapeutics in enteric infectious disease: a review. *J Med Food* 5: 159-169, 2002.
24. Carlander D, Olesen H, Kollberg H, Johannesson M, Wejaker PE and Larsson A: Oral administration of specific yolk antibodies (IgY) may prevent *Pseudomonas aeruginosa* infections in patients with cystic fibrosis: A phase I feasibility study. *Pediatr Pulmonol* 35: 433-440, 2003.

ORIGINALITY REPORT

%24
SIMILARITY INDEX

%14
INTERNET SOURCES

%19
PUBLICATIONS

%7
STUDENT PAPERS

PRIMARY SOURCES

1 Sharpe, M.. "Protection of mice from H5N1 influenza challenge by prophylactic DNA vaccination using particle mediated epidermal delivery", *Vaccine*, 20070821 **%3**
Publication

2 Kazuhide Adachi, Retno Damajanti Soejoedono, Ekowati Handharyani, Marie Inai, Yasuhiro Tsukamoto. "Therapeutic Trial of an Endothelin Receptor Agonist for the Highly Pathogenic Avian Influenza A/H5N1 Virus Infection in Chicks", *Health*, 2014 **%2**
Publication

3 www.recentmedicalfindings.com **%2**
Internet Source

4 www.oalib.com **%2**
Internet Source

5 M. Keith Howard, Nicolas Sabarth, Helga Savidis-Dacho, Daniel Portsmouth et al. "H5N1 Whole-Virus Vaccine Induces Neutralizing Antibodies in Humans Which Are Protective in a **%1**

Mouse Passive Transfer Model", PLoS ONE, 2011

Publication

6

www.mdpi.com

Internet Source

% 1

7

Yasuhiro Tsukamoto, Osamu Maeda, Genshi Shigekawa, Stuart Greenberg, Barry Hendler. "Ostrich Antibody and Its Application to Skin Diseases, a Review and Case Report", Health, 2018

Publication

% 1

8

crofsblogs.typepad.com

Internet Source

% 1

9

doras.dcu.ie

Internet Source

% 1

10

Li, Xiao-Yu, Li-Ji Jin, Tim A. McAllister, Kim Stanford, Jing-Yi Xu, Ya-Nan Lu, Yu-Hong Zhen, Yong-Xin Sun, and Yong-Ping Xu. "Chitosan–Alginate Microcapsules for Oral Delivery of Egg Yolk Immunoglobulin (IgY)", Journal of Agricultural and Food Chemistry, 2007.

Publication

<% 1

11

www.tandfonline.com

Internet Source

<% 1

12

www.researchsquare.com

Internet Source

<% 1

13

"Oral and Poster Manuscripts", Influenza and Other Respiratory Viruses, 05/2011

Publication

<% 1

14

pdfs.semanticscholar.org

Internet Source

<% 1

15

obihiro.repo.nii.ac.jp

Internet Source

<% 1

16

Kawin Punyokun, Ratchanee Hongprayoon, Prapansak Srisapoome, Theerapol Sirinarumitr. " The production of anti- egg yolk immunoglobulin and evaluation of its stability and neutralisation efficacy ", Food and Agricultural Immunology, 2013

Publication

<% 1

17

Cao, Zhiliang, Jiazi Meng, Xingxing Li, Ruiping Wu, Yanxin Huang, and Yuxian He. "The Epitope and Neutralization Mechanism of AVFluIgG01, a Broad-Reactive Human Monoclonal Antibody against H5N1 Influenza Virus", PLoS ONE, 2012.

Publication

<% 1

18

www.vetsci.org

Internet Source

<% 1

cvi.asm.org

19

Internet Source

<% 1

20

goescholar.uni-goettingen.de

Internet Source

<% 1

21

Xiao-Yu Li, Li-Ji Jin, Tim A. McAllister, Kim Stanford, Jing-Yi Xu, Ya-Nan Lu, Yu-Hong Zhen, Yong-Xin Sun, Yong-Ping Xu.

"Chitosan–Alginate Microcapsules for Oral Delivery of Egg Yolk Immunoglobulin (IgY)",
Journal of Agricultural and Food Chemistry,
2007

Publication

<% 1

22

Submitted to National University of Singapore

Student Paper

<% 1

23

"Vaccines for Pandemic Influenza", Springer
Science and Business Media LLC, 2009

Publication

<% 1

24

YUKA KUBOTA, NAOKI KIRIMURA, HATSUKI SHIBA, KAZUHIDE ADACHI, YASUHIRO TSUKAMOTO. "SC1, an immunoglobulin-superfamily cell adhesion molecule, is involved in the brain metastatic activity of lung cancer cells", Oncology Letters, 2015

Publication

<% 1

25

J.W. McCauley, L.A. Pullen, M. Forsyth, C.R. Penn, G.P. Thomas. "4-Guanidino-Neu5Ac2en

<% 1

fails to protect chickens from infection with highly pathogenic avian influenza virus", *Antiviral Research*, 1995

Publication

26

"SARS- and Other Coronaviruses", Springer Science and Business Media LLC, 2008

Publication

27

academicjournals.org

Internet Source

28

Zhixun Xie, Yao-shan Pang, Jiabo Liu, Xianwen Deng, Xiaofei Tang, Jianhua Sun, Mazhar I. Khan. "A multiplex RT-PCR for detection of type A influenza virus and differentiation of avian H5, H7, and H9 hemagglutinin subtypes", *Molecular and Cellular Probes*, 2006

Publication

29

Yoshiki Matsumoto. "Age-related changes in growth hormone-immunoreactive cells in the anterior pituitary gland of Jcl: Wistar-TgN (ARGHGEN) 1Nts rats (Mini rats)", *Congenital Anomalies*, 12/2006

Publication

30

van Maurik, A.. "Seasonal influenza vaccine elicits heterosubtypic immunity against H5N1 that can be further boosted by H5N1 vaccination", *Vaccine*, 20100217

Publication

<% 1

<% 1

<% 1

<% 1

<% 1

31

Eva Maria Huber. "Structural and Functional Characterization of the Immunoproteasome", Springer Science and Business Media LLC, 2013

Publication

<% 1

32

Huan H. Nguyen. "Prophylactic and Therapeutic Efficacy of Avian Antibodies Against Influenza Virus H5N1 and H1N1 in Mice", PLoS ONE, 04/13/2010

Publication

<% 1

33

repozitorium.omikk.bme.hu

Internet Source

<% 1

34

S Gamo. "Patterns of hobo elements and their effects in natural populations of *Drosophila melanogaster* in Japan", Heredity, 06/2006

Publication

<% 1

35

Eric J Yager, Hansi J Dean, Deborah Heydenburg Fuller. "Prospects for developing an effective particle-mediated DNA vaccine against influenza", Expert Review of Vaccines, 2014

Publication

<% 1

36

www.int-res.com

Internet Source

<% 1

37

iovs.arvojournals.org

Internet Source

<% 1

38

Georgia Sotiropoulou, Georgios Pampalakis, Evangelia Prosnikli, Gregory P Evangelatos, Evangelia Livanou. "Development and immunochemical evaluation of a novel chicken IgY antibody specific for KLK6", Chemistry Central Journal, 2012

Publication

<% 1

39

real-j.mtak.hu

Internet Source

<% 1

40

edepot.wur.nl

Internet Source

<% 1

41

data.epo.org

Internet Source

<% 1

42

Submitted to University of Leicester

Student Paper

<% 1

43

Submitted to University of East London

Student Paper

<% 1

44

Quentin Canelle, Walthère Dewé, Bruce L. Innis, Robbert van der Most. " Evaluation of potential immunogenicity differences between TM and TM ", Human Vaccines & Immunotherapeutics, 2016

Publication

<% 1

45

L. Huang. "Characterizing the Limits of Human Visual Awareness", Science, 08/10/2007

Publication

<% 1

46

O. Odinsen, D. Parker, F. Radebe, M. Guness, D. A Lewis. "Early Detection of Human Immunodeficiency Virus Type 1-Specific B-Lymphocyte-Derived Antibodies in a High-Risk Population", *Clinical and Vaccine Immunology*, 2009

Publication

<% 1

47

Petr O. Ilyinskii, Galini Thoidis, Alexander M. Shneider. "Development of a Vaccine Against Pandemic Influenza Viruses: Current Status and Perspectives", *International Reviews of Immunology*, 2009

Publication

<% 1

48

J Goto. "Prevention of hepatitis C virus infection in a chimpanzee by vaccination and epitope mapping of antiserum directed against hypervariable region 1", *Hepatology Research*, 2001

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES

< 5
WORDS