

# Nutrition, Mental Status and Level of 8-hydroxy- 2- deoxyguanosine (OHdG) Urine as Predictors of Premenstrual Syndrome (PMS) in Adolescent Girls

*by* Elizabet C Jusuf

---

**Submission date:** 15-Jun-2021 08:46PM (UTC+0700)

**Submission ID:** 1606927692

**File name:** Nutrition,\_Mental\_Status\_and\_Level\_of\_8-OHdG,\_IJCRR\_2020.pdf (552.88K)

**Word count:** 3967

**Character count:** 19406



# Nutrition, Mental Status and Level of 8-hydroxy-2-deoxyguanosine (OHdG) Urine as Predictors of Premenstrual Syndrome (PMS) in Adolescent Girls

Julian Purnawati<sup>1</sup>, Andi Wardihan Sinrang<sup>1</sup>, Elizabet Catherine Jusuf<sup>2</sup>, Erlyn Limoa<sup>3</sup>, Mardiana Ahmad<sup>1</sup>, Andi Nilawati Usman<sup>1</sup>

<sup>1</sup>Department of Midwifery, Graduate School, Hasanuddin University, Indonesia; <sup>2</sup>Department of Obstetrics and Gynecology, Faculty of Medicine, Hasanuddin University, Indonesia; <sup>3</sup>Department of Psychiatry, Faculty of Medicine, Hasanuddin University, Indonesia.

## ABSTRACT

**Objective:** This study was aimed to determine the prevalence of PMS, the relationship of nutritional intake, mental status, and levels of 8-OHdG in urine with PMS events and to find a predictor model of PMS events.

**Methods:** Observational analytic study design case-control study and cross-sectional study (hybrid method), samples of 90 students class X and XI at SMAN 21 Makassar was conducted in February-March 2020. Data included 2x24 hour food recall, food frequency questionnaire, stress, depression and anxiety scores, and 8-OHdG levels in girls' urine. Data analysis using chi-square test, receiver operating characteristic curve analysis, and logistic regression.

**Results:** The prevalence of PMS was 28.9%. There was a significant correlation between fat intake, calcium, zinc, stress, depression, and urine 8-OHdG levels with the incidence of PMS ( $p < 0.05$  and  $p < 0.01$ ). Stress and 8-OHdG levels in urine as a predictor of PMS events ( $p < 0.01$ ;  $R^2 = 0.208$ ).

**Conclusion:** Stress and 8-OHdG levels in urine can be a predictor of PMS events.

**Key Words:** Nutrition, Mental Status, 8-hydroxy-2-deoxyguanosine, Premenstrual syndrome

## INTRODUCTION

PMS is a series of symptoms in the form of physical and emotional disturbance or discomfort in women 1-2 weeks before menstruation (luteal phase).<sup>1,2</sup> The prevalence of PMS is 37.5% in Saudi Arabia, 17.5% in Japan, 41% in America, 43.2-52.2% in Brazil, 10-12% in women in Europe and the highest rate of 98% occurs in Asian women.<sup>1</sup> PMS impacts social relationships, work, academic achievement. This also has an impact on the economy, which is increasing absenteeism, reducing academic efficiency and student achievement, incurring additional costs for health care, thereby reducing the quality of life.<sup>1-3</sup> PMS also affects the occurrence of pre-natal depression to postpartum depression.<sup>4,5</sup>

Previous research shows that several factors such as hormonal disorders, genetic, psychological factors, lack of vitamins and minerals, lifestyle (smoking, alcohol consumption,

eating habits, and lack of exercise) as well as oxidative stress are associated with the occurrence of PMS.<sup>1,2,6,7</sup> High carbohydrate, fat and protein intake<sup>1,2,8</sup>, lack of calcium, vitamin, and zinc intake<sup>9-11</sup>, and psychological factors (stress, depression, and anxiety) are thought to play a role in the occurrence of PMS.<sup>1,3,12-15</sup>

Research also shows oxidative stress is higher in women who experience PMS.<sup>7</sup> 8-OHdG levels are the main biomarkers of oxidative stress and show high stability in urine.<sup>16</sup> Adolescent girls aged 15-17 as a transition phase from childhood to adulthood with dense activity inside/outside of school are vulnerable to experiencing PMS. This study was conducted to determine the prevalence of PMS, the relationship of nutritional intake, mental status, and levels of 8-OHdG in urine with the incidence of PMS and determine the predictor models of PMS events in adolescents.

### Corresponding Author:

Andi Wardihan Sinrang, Department of Midwifery, Graduate School, Hasanuddin University, Indonesia.

E-mail: [wardihan@pasca.unhas.ac.id](mailto:wardihan@pasca.unhas.ac.id)

ISSN: 2231-2196 (Print)

ISSN: 0975-5241 (Online)

Received: 18.07.2020

Revised: 23.09.2020

Accepted: 07.11.2020

Published: 07.12.2020

## MATERIALS AND METHODS

### Research site

This research was conducted in February-March 2019 at SMAN 21 Makassar. Ethical permission has been obtained by the ethics committee of Medicine Faculty of Hasanuddin University with Recommendation Number 149/UN.4.6.4.5.31/PP36/2020. All subjects who participated were given information about the study's purpose and aim and asked for informed consent before they participated.

### Data Types and Sources

Data include intake of carbohydrates, protein, fat, calcium, zinc, vitamin D, stress, depression, anxiety, and urine 8-OHdG levels.

### Data collection technique

Using 2x24 hours food recall questionnaire, food frequency questionnaires (FFQ), Depression Anxiety Stress Scale (DASS) 42. Sampling the urine by the researcher, then stored in the refrigerator at -20°C and examined using 8-OHdG ELISA Kit at Hasanuddin University Medical Research Center (HUM-RC) Laboratory.

### Data Analysis

The collected data were reviewed, double-checked for completeness and accuracy, and corrected before being entered

into the spreadsheet. Questionnaires were coded and being analyzed by using STATA 14.2. Descriptive statistics including mean and standard deviation (SD). ROC curve analysis used to determine the cut-off point for 8-OHdG. A Chi-square test was used to examine the relationship among variables. Odds ratio with 95% confidence interval also used to present the strength of association between risk factors and outcomes. Then, variables with p-values (<0.25) were entered into multiple logistic regression analyses to examine their association as exposure with PMS as outcomes, and also to determine the predictor model of PMS. P-values <0.05 were considered statistically significant for all tests.

## RESULTS

Screening 370 students class X and XI with a total of 22 classes to determine the incidence of PMS obtained as many as 107 teenagers experiencing PMS (28.9%). Data shows that the characteristics of nutrition intake and mental status especially anxiety among the control groups and PMS groups were in the same categories that are excess of protein, enough for carbohydrates and fat, lack of calcium, zinc and vitamin D, and moderate anxiety. Adolescents in PMS group having moderate stress, mild depression, and high levels of urine 8-OHdG while in the control group were mild stress, not depressed, and low levels of 8-OHdG (Table 1).

Table 1: Respondent characteristics

Variables	$\bar{X} \pm SD$		
	PMS (N=45)	Not PMS (N=45)	Total (N=90)
Age (years)	16.02 ± 0.72	15.86 ± 0.72	15.94 ± 0.72
Menstrual duration (day)	5.91 ± 1.08	5.62 ± 1.23	5.76 ± 1.16
Menstrual cycle (day)	29.24 ± 1.13	29.33 ± 1.14	29.28 ± 1.13
Carbohydrates (grams)	297.41 ± 93.27	311.04 ± 106.25	304.23 ± 99.64
Protein (grams)	89.77 ± 98.62	75.26 ± 23.60	82.51 ± 71.67
Fat (grams)	74.27 ± 38.77	70.34 ± 26.31	72.30 ± 33.01
Calcium(mg)	383.72 ± 353.82	588.88 ± 426.51	486.30 ± 403.07
Vitamin D (µg)	6.71 ± 4.74	7.60 ± 4.77	7.16 ± 4.75
Zinc (mg)	8.98 ± 3.52	10.09 ± 3.32	9.54 ± 3.45
Stress (skor)	21.42 ± 7.43	16.44 ± 6.49	18.93 ± 7.37
Depression (skor)	12.48 ± 4.59	9.93 ± 5.36	11.21 ± 5.13
Anxiety (skor)	10.40 ± 3.30	10.40 ± 2.51	10.40 ± 2.92
8-OHdG (ng/mL)	17.94 ± 10.72	9.31 ± 4.39	13.63 ± 9.22

ROC (receiver operating characteristic) curve analysis to determine the cut-off point with a yield of 11.49 ng/mL, with sensitivity of 68.89%, and specificity of 71.11% (Figure 1).




Figure 1: 8-OHdG ROC Curve.

Table 2: Relationship of nutrition intake, mental status and 8-OHdG level in urine with PMS

Variables	Group		X <sup>2</sup>	p-value	OR	95% CI
	Case (PMS) (n=45)	Control (Not PMS) (n=45)				
<b>Nutrition Intake</b>						
<b>Carbohydrates n (%)</b>						
More	16 (35.56)	21 (46.67)	1.28	0.525	0.5	0.21-1.54
Less	13 (28.89)	12 (26.67)				
Enough	16 (35.56)	12 (26.67)				
<b>Protein n (%)</b>						
More	24 (53.33)	24 (53.33)	0.42	0.807	1.1	0.45-2.93
Less	8 (17.78)	6 (13.33)				
Enough	13 (28.89)	15 (33.33)				
<b>Fat n (%)</b>						
More	19 (42.22)	14 (31.11)	6.72	0.035	3.4	1.14-10.62
Less	19 (42.22)	13 (28.89)				
Enough	7(15.56)	18 (40.00)				
<b>Calcium n (%)</b>						
Less	39 (86.67)	30 (66.67)	5.03	0.025	3.2	1.12-9.37
Enough	6 (13.33)	15 (33.33)				
<b>Zinc n (%)</b>						
Less	35 (77.78)	25 (55.56)	5.00	0.025	2.8	1.11-7.00
Enough	10 (22.22)	20 (44.44)				
<b>Vitamin D n (%)</b>						
Less	38 (84.44)	34 (75.56)	1.11	0.292	1.7	0.61-5.04
Enough	7 (15.56)	11 (24.44)				

Table 2: (Continued)

Variables	Group		X <sup>2</sup>	p-value	OR	95% CI
	Case (PMS) (n=45)	Control (Not PMS) (n=45)				
<b>Mental Status</b>						
<b>Stress n (%)</b>						
High	19 (42.22)	4 (8.89)	17.01	0.001	10.1	2.50-41.38
Medium	11 (24.44)	8 (17.78)			2.9	0.82-10.58
Low	8 (17.78)	18 (40.00)			0.9	0.28-3.23
Not experience	7 (15.56)	15 (33.33)				
<b>Depression n (%)</b>						
Medium	21 (46.67)	14 (31.11)	8.15	0.017	3.4	1.26-9.41
Low	14 (31.11)	8 (17.78)			4.0	1.28-12.6
Not experience	10 (22.22)	23 (51.11)				
<b>Anxiety n (%)</b>						
Medium	32 (71.11)	30 (66.67)	2.42	0.298	0.6	0.19-2.26
Low	5 (11.11)	10 (22.22)			0.3	0.06-1.47
Not experience	8 (17.78)	5 (11.11)				
<b>8-OHdG n (%)</b>						
High	30 (66.67)	13 (28.89)	12.86	0.000	4.9	2.01-12.03
Low	15 (33.33)	32 (71.11)				

Table 3: Multivariate Analysis of Nutrient Intake, Mental Status and 8-OHdG Urine Levels with PMS

Variables	Models				
	1	2	3	4	5
<b>Fat</b>					
More	2.4 (0.75-8.03)		1.5 (0.43-5.81)	1.4 (0.37-5.36)	1.7 (0.26-4.26)
Less	2.1 (0.63-7.33)		1.3 (0.36-5.22)	1.23 (0.31-4.79)	0.69 (0.14-3.26)
Reference	1		1	1	1
<b>8-OHdG</b>					
High	4.00** (1.56-10.35)	4.1** (1.51-11.49)	3.8* (1.33-11.13)	5.3* (1.50-18.99)	5.2* (1.44-19.00)
Low (Reference)	1	1	1	1	1
<b>Stress</b>					
High		9.0** (2.09-39.05)	8.0** (1.80-35.97)	8.7* (1.63-47.14)	9.2* (1.66-51.9)
Medium		1.8 (0.47-7.45)	1.7 (0.42-7.07)	1.9 (0.45-8.46)	1.9 (0.42-8.44)
Low		1.1 (0.30-4.04)	1.09 (0.29-4.00)	1.1 (0.32-4.45)	1.08 (0.27-9.32)
Not experience (Reference)		1	1	1	1
<b>Depression</b>					
Medium				0.7 (0.19-2.96)	0.7 (0.18-3.08)
Low				2.3 (0.59-9.07)	2.26 (0.54-9.32)
Not experience (Referenc)				1	1
<b>Calcium</b>					
Less					1.4 (0.30-6.91)
Reference					1

Table 3: (Continued)

Variables	Models				
	1	2	3	4	5
Zinc					2.2
Less					(0.58-8.19)
Reference					1
R2%	12.5	20.8	21.2	23.2	25.2
Deviance	2975.7	2437.3	2411.7	2291.5	2175.2
N	90	90	90	90	90

\*= significant, *p*-value <0.05  
 \*\*= significant, *p*-value <0.01

The result of the analysis showed that intake of fat, lack of calcium and zinc, stress, depression, and high levels of urine 8-OHdG had a statistically significant relationship with the incidence of PMS. P-values for fat, calcium, zinc, and depression were <0.05, while stress and urine 8-OHdG were <0.01 (Table 2).

Fat, calcium, zinc, stress, depression, and urine 8-OHdG in model 5 can predict the incidence of PMS by 25.2% (R2= 0.252). From the results of the analysis with several models, researchers tend to choose model 2 as a predictor of PMS events, that is urine 8-OHdG levels and stress as models that are more effective and efficient, statistically and practically meaningful, which affect the incidence of PMS. Urine 8-OHdG levels and stress can predict the incidence of PMS by 20.8% (*p*<0.01; R2= 0.208) (Table 3).

### DISCUSSION

The prevalence of PMS is 28.9%. Intake of fat, calcium, zinc, stress, depression, and urine 8-OHdG levels were significantly associated with the incidence of PMS (*p*<0.05 and *p*<0.01). These results are in line with previous studies where excess fat can cause PMS.<sup>2,8</sup> Carbohydrates, protein, and vitamin D are not associated with the incidence of PMS. Less fat intake is also at high risk for PMS. In theory, fat (cholesterol) is the basic ingredient of the formation of steroid hormones. Progestins, androgens, and estrogens can be synthesized *in situ* in various ovarian tissues with cholesterol as a common steroid precursor.<sup>17</sup> PMS occurs as one of the effects of the imbalance of the hormones estrogen and progesterone. So that excess or lack of fat intake can affect the stability of hormone levels in the body. Fat is also related to adipose tissue which will affect body mass index (BMI). However, this study did not measure BMI in adolescents. BMI is associated with the incidence of PMS, so that despite a lack of fat intake, on the other hand, there are risk factors such as BMI that can contribute to causing PMS.<sup>11,12</sup>

Lack of calcium and zinc intake are also at risk for PMS. The average calcium intake of adolescents is only 486.30 mg from the standard 1.200 mg/day. Calcium plays an important role in the synthesis of neurotransmitters (serotonin) and affects mood symptoms and emotional irregularities in PMS.<sup>9</sup> The recommended standard of zinc requirement for adolescents is 16 mg/day (ages 13-15 years) and 14 mg/day (16-18 years) while the average intake of adolescent zinc in this study is 9.54 mg/day. Zinc deficiency can increase the risk of depression<sup>10</sup> and anxiety which are symptoms of PMS through decreased levels of gamma-aminobutyric acid (GABA).<sup>11</sup> Higher TAC (Total Antioxidant Capacity) and zinc serum are also associated with a reduced risk of experiencing PMS<sup>18</sup>. Although not significantly related, in theory, the amount of vitamin D in the body affects the absorption of calcium where the absorption process will not take place properly without the support of adequate vitamin D. Besides that, other factors influence the incidence of PMS, namely an increase in inflammatory cytokines. Vitamin D in this case can act as an anti-inflammatory metabolite that may affect some of the symptoms of PMS.

Stress is related to PMS events. Adolescents with severe stress are 10.1 times more at risk of experiencing PMS than those without stress. Risk increases with stress levels. This finding is in line with studies in which women with moderate and severe stress were 2.49 and 4.9 times at risk for PMS.<sup>3</sup> Adolescence is a dramatic change in the functioning of the Hypothalamic Pituitary Axis (HPA), the stress response, is also a significant period of sustained nerve maturation, especially in stress-sensitive cortical and limbic areas. Prolonged or repeated stress exposure results in a higher sensitivity to this stressor, which in turn leads to maladaptive neurobehavioral development<sup>19</sup>. Stress causes mood changes through the decreased beta-endorphin brain and increased adrenal cortisol. The effect of cortisol causes hormonal imbalances and creates psychological symptoms and mood changes which are symptoms of PMS.<sup>16-18</sup>

Anxiety does not indicate any association with PMS. As for adolescents who experience depression are also at risk of experiencing PMS. This can be caused by a lack of calcium and zinc intake in the adolescent. Subjects who are depressed show high susceptibility to the sympathetic nerves and this condition in response to stress. Increased ROS (Reactive Oxygen Species) by activation of the sympathetic nervous system causes oxidative damage to DNA. Depression was found to be related to increased oxidative stress and decreased antioxidant status. It is possible that women who are depressed can make a lifestyle that triggers an increase in oxidative stress, the presence of fluctuations from estradiol and FSH (follicle-stimulating hormone) associated with depressive symptoms that can increase oxidative stress, and finally, oxidative stress can reduce serotonin and norepinephrine levels that lead to depressive symptoms.<sup>14,20</sup>

Adolescents with high levels of 8-OHdG in urine are also at risk for PMS. High levels of 8-OHdG can be caused by high fat intake, lack of calcium and zinc, stress, and depression in adolescents. 8-OHdG levels tend to increase during the proliferation phase. On the other hand, 20-50% of women complain of depression, anxiety, and fatigue 7-10 days before menstruation which is one of the symptoms of PMS.<sup>15</sup> Increased oxidative stress and decreased antioxidant capacity can occur in PMS, and an imbalance from the oxidant/antioxidant system can also be a cause or effect of various types of stress symptoms that occur in PMS.<sup>6</sup>

Stress and 8-OHdG urine levels are predictors of PMS events in this study. In line with the findings<sup>13</sup> stress is the strongest predictor of the severity of PMS symptoms, while physical exercise and nutritional intake are not related to PMS. This finding is reinforced by other studies where the intensity of PMS is associated with an increase in psychological predictors namely depression, anxiety, stress, and poor sleep quality.<sup>12,18-20</sup>

Characteristics of respondents who are teenagers and high school students with dense intracurricular and extracurricular activities so that they are vulnerable to stress. Psychological stress ahead of school exams also increases the production of ROS and oxidative stress, thereby causing damage to biomolecules that continue to damage cells and tissues. This condition is characterized by increased levels of 8-OHdG in urine.<sup>15</sup> Increased oxidative stress and decreased antioxidant capacity can occur during PMS. This indicates an antioxidant imbalance before menstruation and can be related to the pathogenesis of PMS.<sup>6,7</sup>

## CONCLUSION

Intake of fat, calcium, zinc, stress, depression, and levels of 8-OHdG in urine can affect the occurrence of PMS in adolescents. Stress and 8-OHdG levels in urine can be a pre-

dictor of PMS. A special counselling group for adolescent reproductive health is needed and education about the importance of antioxidant-rich nutrients in counteracting oxidative stress. Further research on oxidative stress can be carried out to better understand its role in the incidence of Premenstrual Syndrome (PMS).

## ACKNOWLEDGMENT

Authors acknowledge the immense help received from the scholars whose articles are cited and included in references to this manuscript. The authors are also grateful to authors / editors / publishers of all those articles, journals, and books from which the literature for this article has been reviewed and discussed.

**Conflict of Interest:** Nil

**Source of Funding:** Nil

## REFERENCES

1. Mohebbi M, Amir S, Akbari A, Mahmodi Z, Nasiri M. Comparison between The Lifestyle of University Students with and without Premenstrual Syndromes. *Electron Physician*. 2017;9(6):4489-96.
2. Hashim MS, Obaideen AA, Jahrami HA, Radwan H, Hamad HJ, Owais AA, et al. Premenstrual Syndrome is Associated with Dietary and Lifestyle Behaviors among University Students. *Nutrients* 2019;11(1939):1-18.
3. Fernandez MDM, Mendez CR, Takkouche B. Psychological Factors and Premenstrual Syndrome : A Spanish Case-control Study. *PLoS One* 2019;14(3):1-13.
4. Amiel Castro RT, Pataky EA, Ehlert U. Associations between premenstrual syndrome and postpartum depression : A systematic literature review. *Biol Psychol* 2019 Oct;147:107612.
5. Roomruangwong C, Kanchanatawan B, Sirivichayakul S, Mahieu B, Nowak G, Maes M. Lower Serum Zinc and Higher CRP Strongly Predict Prenatal Depression and Physio-somatic Symptoms, Which All Together Predict Postnatal Depressive Symptoms. *Mol Neurobiol* 2017;54:1500-12.
6. Duvan CI, Cumaoglu A, Turhan NO, Karasu C, Kafali H. Oxidant/antioxidant Status in Premenstrual Syndrome Oxidant /antioxidant Status in Premenstrual Syndrome. *Arch Gynecol Obs*. 2011;283:299-304.
7. Al-Gubhory KH, Ismail L, editors. *Nutritional Antioxidant Therapies : Treatments and Perspectives*. Springer International Publishing; 2017.
8. Vani RK, Veena K, Subitha L, Hemanth Kumar V, Bupathy A. Menstrual Abnormalities in School Going Girls – Are They Related to Dietary and Exercise Pattern? *J Clin Diagnostic Res* 2013;7(11):2537-40.
9. Hollins-Martin C, Akker O van den, Martin C, Preedy VR, editors. *Handbook of Diet and Nutrition in The Menstrual Cycle, Periconception and Fertility*. 7th ed. Wagenigen Academic Publishers; 2014. Pg 95.
10. Wang J, Um P, Dickerman BA, Liu J. Zinc , Magnesium , Selenium and Depression : A Review of the Evidence, Potential Mechanisms and Implications. *Nutrients* 2018;10(584):1-19.

11. Russo AJ. Decreased Zinc and Increased Copper in Individuals with Anxiety. *Nutr Metab Insights* 2011;4:1–5.
12. Prabhavathi K, Thilip Kumar G, Hemamalini R V, Poomima K N, Saravanan A Study of psychological predictors and sleep quality in different grades of premenstrual syndrome. *Natl J Physiol Pharm Pharmacol.* 2018;8(3).
13. Walton LM, Machamer L, Asumbrado R, Behrens MA. Physiotherapy & Physical Rehabilitation Relationship between Nutrition (REAP), Exercise (VSAQ), and Stress on Premenstrual Syndrome Severity (PSST): Correlation , Cross-Section, Purposive Sample of 75 Females Ages 18-55. *J Physiother Phys Rehabil.* 2018;3(2):1–6.
14. Iida T, Inoue K, Ito Y, Ishikawa H, Kagiono M, Teradaira R, et al. Comparison of Urinary Levels of 8-Hydroxy-2'-deoxyguanosine between Young Females with and without Depressive Symptoms during Different Menstrual Phases. *Acta Med Okayama* 2015;69(1):45–50.
15. Iida T, Ito Y, Ishikawa H, Kanazashi M, Teradaira R. Effects of Psychological Stress from a National License Examination on the Urine 8-Hydroxy-Deoxyguanosine Levels in Young Female Students , Taking into Account the Menstrual Cycle. *Open J Prev Med* 2018;8:21–31.
16. Zanolin ME, Girardi P, Degan P, Rava M, Olivieri M, Gennaro G Di, et al. Measurement of a Urinary Marker (8-hydroxydeoxyguanosine, 8-OHdG) of DNA Oxidative Stress in Epidemiological Surveys: a Pilot Study. *Int J Biol Markers* 2015;30(3):341–5.
17. Fritz MA, Speroff L. *Clinical Gynecologic Endocrinology and Infertility.* 8th ed. Philadelphia: Lippincott Williams and Wilkins; 2011.
18. Fathizadeh S, Amani R, Haghighizadeh MH, Hormozi R. Comparison of Serum Zinc Concentrations and Body Antioxidant Status between Young Women with Premenstrual Syndrome and Normal Controls : A Case-control Study. *Int J Reprod Biomed* 2016;14(11):699–704.
19. Romeo RD. The Teenage Brain: The Stress Response and the Adolescent Brain. *Curr Dir Psychol Sci* 2014;22(2):140–5.
20. Hirose A, Terauchi M, Akiyoshi M, Owa Y, Kato K, Kubota T. Depressive Symptoms are Associated with Oxidative Stress in Middle-aged Women : a Cross-sectional Study. *Biopsychosoc Med* 2016;10(12):1–6.

# Nutrition, Mental Status and Level of 8-hydroxy- 2-deoxyguanosine (OHdG) Urine as Predictors of Premenstrual Syndrome (PMS) in Adolescent Girls

## ORIGINALITY REPORT

14%

SIMILARITY INDEX

12%

INTERNET SOURCES

11%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://ijcrr.com">ijcrr.com</a> Internet Source	3%
2	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	2%
3	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet Source	2%
4	Submitted to Aspen University Student Paper	1%
5	Sarina Ali, Gemini Alam, Samrichard. "Ajwa date fruit (Phoenix dactylifera L.) in increasing hemoglobin (Hb) level to teenage girl", Enfermería Clínica, 2020 Publication	1%
6	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	1%
7	<a href="http://bpsmedicine.biomedcentral.com">bpsmedicine.biomedcentral.com</a> Internet Source	1%

8	"20th International Congress of Nutrition: Granada, Spain, September 15–20, 2013", Annals of Nutrition and Metabolism, 2013 Publication	<1 %
9	pesquisa.bvsalud.org Internet Source	<1 %
10	journal.frontiersin.org Internet Source	<1 %
11	www.birpublications.org Internet Source	<1 %
12	www.lib.ncsu.edu Internet Source	<1 %
13	Eliane Aragão da Silva, Daniel Alvarez Pires. "Prevalence of premenstrual syndrome and its psychological effects among university students who participate and do not participate in resistance training", Revista Brasileira de Ciências do Esporte, 2021 Publication	<1 %
14	e-journal.unair.ac.id Internet Source	<1 %
15	Fatemeh Abdi, Gity Ozgoli, Fatemeh Sadat Rahnemaie. "A systematic review of the role of vitamin D and calcium in premenstrual syndrome", Obstetrics & Gynecology Science, 2019	<1 %

16

Qingcheng Song, Hongzhi Hu, Xiangtian Deng, Xin Xing, Wei Chen, Yingze Zhang. "The Predictive Role of Platelet-to-Lymphocyte Ratio and Systemic Immune-Inflammation Index in Young and Middle-Aged Patients with Tibial Plateau Fractures", Research Square Platform LLC, 2021

Publication

---

<1 %

17

Rajia Rajia, Muh Nasrum Massi, Mardiana Ahmad, Sharvianty Arifuddin, Upik Anderiani Miskad. "Parity, exposure to cigarette smoke and the presence of bacteria *Stenotrophomonas maltophilia* are related to preterm labor incidence", Enfermería Clínica, 2020

Publication

---

<1 %

18

Shao-Wen Zhang. "Molecularly imprinted monolith in-tube solid-phase microextraction coupled with HPLC/UV detection for determination of 8-hydroxy-2'-deoxyguanosine in urine", Analytical and Bioanalytical Chemistry, 07/23/2009

Publication

---

<1 %

19

Mariani, E.. "Oxidative stress in brain aging, neurodegenerative and vascular diseases: An overview", Journal of Chromatography B, 20051115

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On