

THE EFFECT OF ANTI-DIABETIC  
DRUGS REGIMEN  
COMBINATION WITH  
METFORMIN COMPARED  
WITHOUT METFORMIN TO  
CONVERSION OF ACID-FAST  
BACILLUS SPUTUM SMEAR IN  
NEW CASE OF

---

**Submission date:** 27-Oct-2021 01:25PM (UTC+0700)

**Submission ID:** 168539633

**File name:** dr.\_Fransiskus\_wabia.pdf (334.27K)

**Word count:** 3721

**Character count:** 19177

PULMONARY TUBERCULOSIS PATIENTS AND TYPE

*by* Andi Makbul Aman

## THE EFFECT OF ANTI-DIABETIC DRUGS REGIMEN COMBINATION WITH METFORMIN COMPARED WITHOUT METFORMIN TO CONVERSION OF ACID-FAST BACILLUS SPUTUM SMEAR IN NEW CASE OF PULMONARY TUBERCULOSIS PATIENTS AND TYPE 2 DIABETES MELLITUS

Fransiscus A Wabia<sup>1\*</sup>, Harun Iskandar<sup>1\*</sup>, Husaini Umar<sup>1\*</sup>, Syakib Bakri<sup>1</sup>,  
Andi Makbul Aman<sup>1</sup>, Hasyim Kasim<sup>1</sup>, Haerani Rasyid<sup>1</sup>, Erwin Arif<sup>1</sup>, Agus  
Sudarso<sup>1</sup>, dan Arifin Seweng<sup>2</sup>

7  
<sup>1</sup>Departement of Internal Medicine, Faculty of Medicine, Hasanuddin University, Makassar,  
Indonesia

<sup>2</sup>Biostatistics Department, Faculty of Public Health, Hasanuddin University, Makassar,  
Indonesia

### ABSTRACT

**Background :** *This study aims to determine the effect of a combination regimen of metformin and without metformin on the conversion of sputum smear in new cases of pulmonary TB patients with type 2 DM.*

**Materials and Methods:** *This cross-sectional study was conducted at Wahidin Sudirohusodo*

*Hospital, Makassar Center for Lung Health, Kasi-Kasi, and Sudiang Community Health Center from August 2019. This study involved 83 new cases of pulmonary TB subjects with T2DM. New cases of pulmonary TB patients were obtained by examining AFB sputum in DM patients who had symptoms of pulmonary TB and had no history of suffering or receiving previous anti-TB drug therapy. Pulmonary TB patients receive category 1 anti-TB drugs and DM subjects were divided into 2 groups is metformin and non-metformin. Then, smear sputum is examined in the 2-month intensive phase of anti-TB drug therapy to assess the conversion of AFB sputum.*

**Results:** *From a total of 83 new cases of pulmonary TB and T2DM, 44 subjects received a combination regimen of metformin and 39 subjects non-metformin. In this study, there were 42 (95.5%) subjects who conversion sputum smear in the metformin group more higher than in the non-metformin group but statistically it was not significant ( $p = 0.282$ ).*

**Conclusion:** *There was no significant relationship between the metformin and non-metformin groups on the conversion of AFB sputum smear in new cases of pulmonary TB patients and T2DM.*

**Keywords:** *Tuberculosis, Diabetes Mellitus, Metformin, Sputum Smear Conversion.*

## 5 INTRODUCTION

Diabetes mellitus (DM) is a metabolic disorder characterized by hyperglycemia due to defects in insulin secretion, insulin action or both.<sup>1</sup> DM is a serious long-term condition with a major impact in the lives and well-being of individuals, families and communities around the

world. Aging, changes in lifestyle, socioeconomic factors, and population growth have led to an increase in the prevalence of DM, especially type 2 DM.<sup>2</sup> The International Diabetes Federation (IDF) organization globally estimates that DM sufferers will increase to reach 578 million in 2030 and 700 million in 2045. Prevalence DM in Indonesia is around 10.7 million and is in the 7th position out of 10 countries with the highest number of DM sufferers in the world. In 2012, there was a prevalence of about 15% or 1.04 million people from the DM population who suffered from pulmonary TB.<sup>3</sup> The relationship between DM and TB has long

been known where the first report was documented by Avicenna (980-1027 AD) more than a thousand years ago but studies in the last 10-15 years have shown that DM as a risk factor for TB.<sup>2,3</sup> The increased risk of TB transmission in DM patients is two to three times, especially inpatients with positive acid fast bacillus (AFB).<sup>4</sup> The pathophysiological mechanism of the effect of DM as a predisposing risk factor for TB is chronic hyperglycemia causing cellular immune system depression, alveolar macrophage dysfunction, low levels of IFN- $\gamma$ , pulmonary microangiopathy and micronutrient deficiency.<sup>2</sup>

In DM with TB, the priority of treatment is to treat TB while at the same time keeping blood glucose levels under control.<sup>5,6</sup> From a recent study, it was explained that metformin is the first choice in TB patients with DM because of the low risk of hypoglycemia, its effectiveness, low cost, the effect of minimal clinically relevant interactions with rifampin and

its own efficacy in providing a strong protective effect against TB.<sup>7</sup> It was reported that metformin independently can increase specific immunity in TB patients, reduce inflammation and increase the efficacy of TB treatment.<sup>8</sup> Some of the roles of metformin in TB therapy are controlling the secretion of IL-10 thus altering the host immune response. IL-10 itself works to inhibit the response of CD4 + T cells by inhibiting the function of APC cells infected with microbacteria. IL-10 also plays an important role in metabolic disorders such as diabetes because of its effect on insulin sensitivity by increase peripheral glucose uptake. Metformin therapy can be considered as a new strategy in increasing anti-TB efficacy because of its ability to control IL-10 secretion so as to alter the host immune response to TB infection and its ability to influence insulin sensitivity.<sup>9,10</sup> Metformin also inhibits complex-I function which can also inhibit NDH-I as a potential candidate for antibiotics against TB.

In addition, metformin acts as an activator of Adenosine Monophosphate-Activated Protein Kinase (AMPK) in modulating the immune system and another role for metformin is to inhibit the growth of mycobacteria by inducing the production of mitochondrial reactive oxygen species (mROS).<sup>13,14</sup>

In research by Ye-Jin Lee et al (2018) and Novita et al (2018) found that the metformin combination anti diabetic drugs (ADD) group had AFB sputum smear conversion was higher than the non- metformin combination group.<sup>15,16</sup>

24

## MATERIALS AND METHOD

This cross sectional study was conducted at Wahidin Sudirohusodo Hospital, Makassar Center for Lung Health, Kasi-Kasi, and Sudiang Community Health Center from August 2019. This study involved 83 new cases of pulmonary TB subjects with DM. The research variables were age, sex, BMI, chest radiograph lesions, initial sputum smear load of AFB and conversion of the sputum smear at the end of the 2nd month of intensive therapy where all these variables were subjects who received the metformin and nonmetformin combination regimen.

Subjects included in the inclusion criteria were male and female, aged  $\geq 18$  years, bacteriological diagnosis of new pulmonary TB and laboratory-confirmed DM. Subjects included in the exclusion criteria were tuberculosis patients who had relapsed, failed treatment

or resistant multidrugs (MDR-TB). Diagnosis of pulmonary tuberculosis is defined when the sputum smear examination shows M.tb (*Mycobacterium Tuberculosis*) with bacterial load + 1 / + 2 / + 3. The diagnosis of DM is based on fasting blood glucose levels  $\geq 126$  mg / dL, current

blood glucose  $\geq 200$  mg / dl, HbA1c  $\geq 6.5\%$ . The metformin combination regimen is defined as metformin in combination with other anti-diabetic drugs it is other ADD (glimepiride) and insulin. Total daily dose of metformin 1000-1500 mg / day. Combination regimens without metformin are defined as other anti-diabetic drugs not combined with metformin it is the combination of other ADD (glimepiride) and insulin or insulin alone. Body mass index

(BMI)

$<18.5$  kg / m<sup>2</sup> it is underweight, BMI 18.5-22.9 kg / m<sup>2</sup> it is normoweight, and BMI  $\geq 23.0$  kg

/ m<sup>2</sup> it is overweight. Minimal lesion chest X-ray is defined when the pulmonary tuberculosis process affects a portion of one or two lungs with an area no more than between the 2 front ribs

(the volume of the lung that lies above the chondrosternal junction of the second rib and the spinous processes of the IV thoracic vertebrae or the corpus of the V thoracic vertebrae) and no cavities were found. Extensive chest X-ray lesions is defined when the disease process was

broader than the minimal lesions.

This research provides the ethical requirements of the Human Biomedical Research Commission, Hasanuddin University Medical Faculty No: 607 / UN4.6.4.5.31 / PP36 / 2020. Data analysis was performed using SPSS version 26 (IBM Corp, Armonk, New York, USA). The statistical test used was the Chi Square test and the Fisher Exact test to assess the relationship between variables. The statistical test results were considered significant if the P value was  $<0.05$ .

## RESULTS

In this study, a total of 83 new cases of pulmonary TB subjects with DM were obtained, of which 44 subjects received a combination regimen of metformin and 39 subjects who received a combination regimen without metformin. The sputum smear was checked twice it is before the intensive therapy and the end of the 2nd month of intensive therapy. From a total

of 83 subjects, it was found that the conversion of AFB sputum smear was more than 77 (92.8%) subjects and 6 (7.2%) subjects who did not have conversion (Table 1).

**Table 1. Distribution of Research Variable Categories (n = 83)**

Variabel		n	%
SEX	Male	48	57.8
	Female	35	42.2
AGE	≥ 40 y.o	78	94.0
	< 40 y.o	5	6.0
BMI	Underweight	9	11.7
	Normoweight	49	63.6
	Overweight	19	24.7
Lesion Chest X-ray	Minimal	11	13.3
	Extensive	72	86.7
Initial smear load of AFB	+1	56	67.5
	+2	17	20.5
	+3	10	12.0
Combination Type Anti Diabetic Drugs	Metformin	44	53.0
	Non metformin	39	47.0
Combination Type Anti Diabetic Drugs	Yes	77	92.8
	No	6	7.2

This study found that the metformin group had 42 (95.5%) more conversions of sputum smear compared to 35 (89.7%) in the non-metformin group. However, statistically there was no significant relationship between metformin and non-metformin groups with AFB sputum smear conversion with  $p > 0.05$ . (Table 5.2)

**Table 2 Relationship between Metformin and Without Meetformin with BTA conversion**

Combination Type Anti Diabetic Drugs		Conversion		Total	P value
		Yes	No		
Metformin	N	42	2	44	p= 0,282
	%	95,5%	4,5%	100,0%	
Non metformin	N	35	4	39	
	%	89,7%	10,3%	100,0%	
Total	N	77	6	83	
	%	92,8%	7,2%	100,0%	

Table 3 shows the higher conversion of AFB sputum smears in the metformin combination regimen group on the study variables are gender, age, BMI less and more, chest x-ray lesions and initial grading of AFB sputum smear + 1. Meanwhile, in the combination regimen group

without metformin, the highest conversion of BTA sputum smears occurred in the study variables are normal BMI, initial grading of BTA+2 and BTA+3 sputum smear. However, there was no statistically significant relationship between the combination regimen of metformin and without metformin groups with conversion of AFB sputum smear according to the study variables ( $p>0.05$ ).

**Table 3. The Relationship Between Combination Regimens With Metformin and Without Metformin with Conversion of Sputum Smear According to Research Variables.**

Variables	Combination Regiment		Conversion		Total	P value
			Yes	No		
<b>Sex</b>						
Male	Metformin	n (%)	21 (95.5)	1 (4.5)	22 (100.0)	0.564
	Non Metformin	n (%)	24 (92.3)	2 (7.7)	26 (100.0)	
Female	Metformin	n (%)	21 (95.5)	1 (4.5)	22 (100.0)	0.306
	Non Metformin	n (%)	11 (84.6)	2 (8.6)	13 (100.0)	
<b>Age</b>						
≥ 40 y.o	Metformin	n (%)	41 (95.3)	2 (4.7)	43 (84.6)	0.652
	Non Metformin	n (%)	32 (91.4)	3 (8.6)	35 (100.0)	
< 40 y.o	Metformin	n (%)	1 (100.0)	0 (0.0)	1 (100.0)	1.000
	Non Metformin	n (%)	3 (80.0)	1 (20.0)	4 (100.0)	
<b>BMI</b>						
Underweight	Metformin	n (%)	3 (100.0)	0 (0.0)	3 (100.0)	0.417
	Non Metformin	n (%)	4 (66.7)	2 (33.)	6 (100.0)	
Normoweight	Metformin	n (%)	20 (90.9)	2 (9.1)	22 (100.0)	0.613
	Non Metformin	n (%)	25 (92.6)	2 (7.4)	27 (100.0)	
Overweight	Metformin	n (%)	15(100.0)	0 (0.0)	15 (100.0)	P*
	Non Metformin	n (%)	4 (100.0)	0 (0.0)	4 (100.0)	
<b>Chest X-ray</b>						
Minimal lesions	Metformin	n (%)	8 (100.0)	0 (0.0)	8 (100.0)	0.273
	Non-Metformin	n (%)	2 (66.7)	1 (33.3)	3 (100.0)	
Extensive lesions	Metformin	n (%)	34 (94.4)	2 (5.6)	36 (100.0)	0.500
	Non-Metformin	n (%)	33 (91.7)	3 (8.3)	36 (100.0)	
<b>Initial smear load of AFB</b>						
+1	Metformin	n (%)	32 (97.0)	1 (3.0)	33 (100.0)	0.365
	Non-Metformin	n (%)	21 (91.3)	2 (8.7)	23 (100.0)	
+2	Metformin	n (%)	7 (87.5)	1 (12.5)	8 (100.0)	0.471
	Non-Metformin	n (%)	9 (100.0)	0 (0.0)	9 (100.0)	
+3	Metformin	n (%)	2 (66.7)	1 (33.3)	3 (100.0)	0.533
	Non-Metformin	n (%)	6 (85.7)	1 (14.3)	7 (100.0)	

P\*: can't be tested

## DISCUSSION

The priority of treatment in DM with TB is to treat TB while at the same time keeping blood glucose levels under control.<sup>5,6</sup> The first choice of oral ADD therapy in DM patients with TB is metformin. DM.<sup>15</sup> Thus, the goal of TB treatment with DM is achieved is the conversion of smear sputum AFB after completion of intensive therapy at the end of the 2nd month.<sup>17</sup> In this study, it was found that the age of ≥ 40 years and male gender had the most TB with DM compared to age <40 years and women. It is known that age ≥ 40 years is

the prevalence of the peak age at diagnosis of T2DM which has an average age range between 45-60 years.<sup>18</sup> Workneh et al. (2017) identified that the comorbidity of TB with DM increases with age.<sup>19</sup> The higher prevalence of TB with DM in men is most often due to tobacco smoking

and alcohol consumption so that more access to health centers than women. In addition, normal

BMI is also the most common, almost the same as Tripti et al (2017).<sup>20</sup> Nutritional factors are a risk factor for TB infection and are associated with malnutrition and DM.<sup>20</sup> From the results of the chest radiograph and Initial sputum smear load of AFB, it was found that the most extensive lesions of the chest X-ray and sputum smear of AFB +1. This is the same as Layali et al (2019).<sup>22</sup> But it is different from Barreda et al (2020) who got the most sputum smear of AFB +2 /+3.<sup>23</sup> This occurs because of a decrease in the immune system and a defect in alveolar

macrophages or T lymphocytes resulting in an increase in the number of bacteria in the sputum

and greater expansion of the lesion chest x-ray in TB with DM.<sup>24</sup> There was no statistically significant relationship between the combination regimen with metformin and without metformin with the conversion of sputum smear AFB by sex, age, BMI, chest X-ray lesions and initial grading of sputum smear AFB. This study found no significant relationship between the combination regimen with metformin and without metformin with conversion of sputum smear AFB ( $p > 0.05$ ), it is different from Novita et al (2017) and Al-Shaer et al (2018) who get a significant relationship, but in we study the metformin combination regimen had a higher smear conversion of sputum AFB (95.5%) similar with Novita et al (2017) and Al-Shaer et al (2018).<sup>16,25</sup> As it is known, metformin has an effect on TB infection, which is to suppress the growth of AMPK-dependent intracellular TB germs which facilitate lysosomal phagocyte fusion in cells infected with TB germs such as in a pilot retrospective cohort study by Singhal et al. (2014) in Singapore.<sup>26</sup> According to Lee YJ et al (2017) metformin can be an adjunct to anti-TB drugs to increase conversion of AFB sputum smear after intensive therapy at the end of 2nd month while metformin is widely used as the first choice oral ADD in DM patients.<sup>14,26</sup>

## CONCLUSION

TB with DM subjects who received metformin had more smear sputum smear conversion than subjects without metformin.

## CONFLICT OF INTEREST

There are no Ethical or Financial issues, Conflicts of interests, or animal experiments related to this research.

## IRB INFORMATION

This research provides the ethical requirements of the Human Biomedical Research Commission, Hasanuddin University Medical Faculty No: 607 / UN4.6.4.5.31 / PP36 / 2020.

## REFERENCES

1. ADA. *Classification and Diagnosis of Diabetes Mellitus*.care.diabetesjournals.org. Diabetes Care.2020;43:S14-S31.<https://doi.org/10.2337/dc20-S002>
2. ParvanehBaghaei, Majid Marjani, PedramJavanmard,Payam Tabarsi, Mohammad Reza Masjedi. *Diabetes mellitus and tuberculosis facts and controversies*. Journal of Diabetes & Metabolic Disorders. 2013. 12:58. <http://www.jdmndonline.com/content/12/1/58>
3. Nageswara rao gopathi, Venu Mandava, Sravani Makala. *Tuberculosis and Diabetes : the deadly duo* Int J Adv Med.ijam20150552.. 2015 Aug;2(3):241-245. DOI: <http://dx.doi.org/10.18203/2349-3933>.
4. Stanley S. Schwartz, Solomon Epstein, Barbara E. Corkey, Struan F.A. Grant, James R. Gavin III, Richard B. Aguilar. *The Time is Right for a New Classification System for Diabetes Rationale and Implications of the E-cell-centric Classification Schema*. Diabetes Care,Vol 39.2016; 39(2): 179-186. <https://doi.org/10.2337/dc15-1585>
5. A. D. Harries, A.M.V. Kumar, S. Satyanarayana, Y. Lin, R. Zachariah, K. Lönnroth,et al. *Diabetes mellitus and tuberculosis: programmatic management issues*.TheUnion.Int J Tuberc Lung Dis.2015.19(8):879–886. <http://dx.doi.org/10.5588/ijtld.15.0069>.
6. Jiyan M. R, Vadgama P. K, Pandey A. S, Modh D. A. *Clinical Profile and outcome of Tuberculosis in Patients with Diabetes*. Int J Res Med. 2015; 4(2);36-40.
7. Lin Yan, Anthony D Harries, Ajay.M.V.Kumar, Julia A.Critchley, Reinout van Crevel, Philip Owiti et al. *Management Of Diabetes Mellitus-Tuberculosis*. A Guide to the Essential Practice First Edition. 2019. International Union Against Tuberculosis and Lung Disease. 1 ed. 2019: 33-57. <https://www.theunion.org>
8. SrujithaMarupuru, P.Senapati, S.Pathadka, S.S.Miraj, M.K.Unnikrishnana, M.K.Manu. *Protective effect of metformin against tuberculosis infections in diabetic patients: an observational study of south Indian tertiary healthcare facility*. braz j infect dis 2017;21(3):312–316.<http://dx.doi.org/10.1016/j.bjid.2017.01.001>
9. Beth Kelly, G.M. Tannahill, M.P.Murphy, Luke A.J.O'Neill. *Metformin inhibits the production of reactive oxygen species from NADH:Ubiquinone oxidoreductase to limit induction of IL-1 $\beta$  and boosts IL-10 in lipopolysaccharide-activated macrophages*. JBiolChem. 2015.Vol 290(33):20348–20359. DOI 10.1074/jbc.M115.662114
10. Vashisht R, Brahmachari SK. *Metformin as a potential combination therapy with existing front-line antibiotics for tuberculosis*. J Transl Med. 2015;13 :81-83
11. PacifiqueNdishimye, Fouad Seghrouchni , B.Domokos, O.Soritau, A.Sadak, Danilea.H et al. *Evaluation of interleukin-10 levels in the plasma of patients with various stage of tuberculosis*. Clujul Medical. 2015; 88(2):164-7. DOI: 10.15386/cjmed-459.
12. Blanca I. Restrepo. *Metformin: Candidate host-directed therapy for tuberculosis in diabetes and non-diabetes patients*. Elsevier. 2016:101S:S69-S72. doi: 10.1016/j.tube.2016.09.008.
13. Ming-Chia Lee, Chen-Yuan Chiang, Chih-Hsin Lee, Cheng-Maw Ho, Chia Hao Chang, Jann-Yuan Wang et al. *Metformin use is associated with a low risk of tuberculosis among newly diagnosed diabetes mellitus patients with normal renal function: A nationwide cohort study with validated diagnostic criteria*. PLOS ONE. 2018; 13(10): e0205807. <https://doi.org/10.1371/journal.pone.0205807>

14. Sheng-Wei Pan, Yung-Feng Yen, Yu Ru Kou, Pei-Hung Chuang, Vincent Yi-Fong Su, Jia-Yih Feng et al. *The risk of tb in patients with type 2 diabetes initiating metformin vs sulfonylurea treatment*. 2017.Chest 153(6). DOI: 10.1016/j.chest.2017.11.040
15. Ye-Jin Lee, Sung Koo Han, Ju Hee Park, Jung Kyu Lee, Deog Keom Kim, Hee Soon Chung et al. *The effect of metformin on culture conversion in tuberculosis patients with diabetes mellitus*. Korean J Intern Med 2018;33:933-940. <https://doi.org/10.3904/kjim.2017.249>
16. Bernadette Dian Novita, Mulyohadi Ali, Agung Pranoto, EndangIsbandiatiSoediono, Ni Made Mertaniasih. *Metformin induced autophagy in diabetes mellitus Tuberculosis co-infection patients: A case study*. Indian Journal of Tuberculosis 66 (2019) 64-69. <https://doi.org/10.1016/j.ijtb.2018.04.003>.
17. Ratnawati, Dewi Wijaya, A.M Nazaruddin, FarizNurwidya, Erlina Burhan. *The relationship between hemoglobin A1C levels and sputum conversion time in indonesian patients with new cases of pulmonary tuberculosis*. J Nat Sc Biol Med 2018;9:217-21. DOI: 10.4103/jnsbm.JNSBM\_24\_18.
18. ThanmaiKaleru, Varun K. Vankeshwaram, Ankush Maheshwary, Divya Mohite, Safeera Khan. *Diabetes Mellitus in the Middle-Aged and Elderly Population (>45 Years) and Its Association With Pancreatic Cancer: An Updated Review*.2020. Cureus 12(6): e8884. DOI: 10.7759/cureus.8884.
19. Workneh MH, Bjune GA, Yimer SA. *Prevalence and associated factors of tuberculosis and diabetes mellitus comorbidity: a systematic review*. PloS one. 2017; 12(4):e0175925.
20. TriptiPande, Sophie Huddart, Wilbin Xavier, SrivathsaKulavalli, Tanya Chen, Madhukar Pai et al. *Prevalence of diabetes mellitus amongst hospitalized tuberculosis patients at an Indian tertiary care center: A descriptive analysis*. PLOS ONE. 2018. <https://doi.org/10.1371/journal.pone.0200838>.
21. Wahiduddin,W, Pranoto. A, Sudjarwo S. *KendaliGlikemik pada Pasien Diabetes MelitusTipe 2 dengan dan tanpaTuberkulosisParu*. Media Kesehatan Masyarakat Indonesia, 15(1), pp. 99-109.
22. Dana JauharaLayali, Bintang YM Sinaga, ParluhutanSiagian, Putri C. Eyanoe. *HubunganLesiTuberkulosisParuDengan Diabetes MelitusTerhadap Kadar HbA1c*. Respir Indo. Vol. 39 (3). 2019. <http://www.jurnalrespirologi.org>.
23. Nadia Nilda Barreda, Maria B. Arriaga, Juan.G.Aliaga, Katty Lopez, Oswaldo Martin Sanabria, Thomas A. Carmo et al. *Severe pulmonary radiological manifestations are associated with a distinct biochemical profile in blood of tuberculosis patients with dysglycemia*. BMC Infectious Diseases (2020) 20:139. <https://doi.org/10.1186/s12879-020-4843-0>.
24. ADA. *Standards of Medical Care in Diabetes*. 2020. Diabetes Care 2020;43.
25. Mohammad H. Al-Shaer, Hazem Elewa, YosraAlkabab, Lama H. Nazer, Scott K. Heysell. *Fixed-dose combination associated with faster time to smear conversion compared to separate tablets of anti-tuberculosis drugs in patients with poorly controlled diabetes and pulmonary tuberculosis in Qatar*. BMC. Infectious Diseases (2018) 18:384. <https://doi.org/10.1186/s12879-018-3309-0>

26. Amit Singhal, Liu Jie, Pavanish Kumar, Gan Suay Hong, M.K.ShingLeow , B.Paleja et al.  
*Metformin as adjunct antituberculosis therapy. Science Translational  
Medicine.*2014.19;6(263):263ra159. doi: 10.1126/scitranslmed.3009885.

# THE EFFECT OF ANTI-DIABETIC DRUGS REGIMEN COMBINATION WITH METFORMIN COMPARED WITHOUT METFORMIN TO CONVERSION OF ACID-FAST BACILLUS SPUTUM SMEAR IN NEW CASE OF PULMONARY TUBERCULOSIS PATIENTS AND TYPE

## ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

5%

STUDENT PAPERS

## PRIMARY SOURCES

1	Submitted to Universitas Indonesia Student Paper	4%
2	<a href="http://www.jdmdonline.com">www.jdmdonline.com</a> Internet Source	2%
3	<a href="http://www.theunion.org">www.theunion.org</a> Internet Source	1%
4	<a href="http://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a> Internet Source	1%
5	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet Source	1%
6	Waleed Al Abdulmonem, Zafar Rasheed, Suliman A. Alsagaby, Abdullah S.M. Aljohani, Abdullah Alkhamiss, Ahmed Ali Ahmed. "Impact of ERCC2 Lys751Gln (rs13181), ERCC2 Asp312Asn (rs1799793) and XRCC1 Arg399Gln (rs25487) polymorphisms on the risk of	1%

# prostate cancer among cases in central region of Saudi Arabia", Gene Reports, 2021

Publication

---

7	Andi Irham, Husaini Umar, Fabiola Maureen Shinta Adam, Idar Mappangara, Syakib Bakri, Risna Halim, Alfian Zainuddin. "Concordance between Triglyceride Glucose Index and Admission Insulin Resistance Index in Non-diabetic Subjects of Acute Coronary Syndrome", Open Access Macedonian Journal of Medical Sciences, 2021 Publication	1 %
8	<a href="http://bmcnutr.biomedcentral.com">bmcnutr.biomedcentral.com</a> Internet Source	1 %
9	<a href="http://rjdnmd.org">rjdnmd.org</a> Internet Source	1 %
10	<a href="http://clinicaltrials.gov">clinicaltrials.gov</a> Internet Source	<1 %
11	<a href="http://fieldresearch.msf.org">fieldresearch.msf.org</a> Internet Source	<1 %
12	Noton K. Dutta, Michael L. Pinn, Petros C. Karakousis. "Metformin Adjunctive Therapy Does Not Improve the Sterilizing Activity of the First-Line Antitubercular Regimen in Mice", Antimicrobial Agents and Chemotherapy, 2017 Publication	<1 %

---

13	<a href="http://giapjournals.com">giapjournals.com</a> Internet Source	<1 %
14	<a href="http://jmhg.springeropen.com">jmhg.springeropen.com</a> Internet Source	<1 %
15	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	<1 %
16	A. Badawi, B. Gregg, D. Vasileva. "Systematic analysis for the relationship between obesity and tuberculosis", Public Health, 2020 Publication	<1 %
17	Garrett Teskey, Ruoqiong Cao, Hicret Islamoglu, Albert Medina et al. "The Synergistic Effects of the Glutathione Precursor, NAC and First-Line Antibiotics in the Granulomatous Response Against Mycobacterium tuberculosis", Frontiers in Immunology, 2018 Publication	<1 %
18	Jaswinder Kaur Virk, Ajudhia Nath Kalia, Vinod Kumar Gauttam, Minky Mukhija, Gautam Rath. "Development and characterization of spheroidal antidiabetic polyherbal formulation from fresh vegetable juice: A novel approach", Journal of Food Biochemistry, 2020 Publication	<1 %
19	Y. Ma, Y. Pang, W. Shu, Y.-H. Liu, Q.-P. Ge, J. Du, L. Li, W.-W. Gao. "Metformin reduces the	<1 %

relapse rate of tuberculosis patients with diabetes mellitus: experiences from 3-year follow-up", European Journal of Clinical Microbiology & Infectious Diseases, 2018

Publication

---

20	<a href="https://discovery.ucl.ac.uk">discovery.ucl.ac.uk</a> Internet Source	<1 %
21	<a href="http://ejournal.upnvj.ac.id">ejournal.upnvj.ac.id</a> Internet Source	<1 %
22	<a href="http://res.mdpi.com">res.mdpi.com</a> Internet Source	<1 %
23	<a href="http://tbassnindia.org">tbassnindia.org</a> Internet Source	<1 %
24	<a href="http://www.scielo.br">www.scielo.br</a> Internet Source	<1 %
25	"Advances in Host-Directed Therapies Against Tuberculosis", Springer Science and Business Media LLC, 2021 Publication	<1 %

---

Exclude quotes  On

Exclude matches  < 5 words

Exclude bibliography  On