

## Importance of philopatry and hydrodynamics in the recruitment of bioeroding sponges on Indonesian coral reefs

Joseph Marlow<sup>A,C</sup>, Abdul Haris<sup>B</sup> and James Bell<sup>A</sup>

<sup>A</sup>Victoria University of Wellington, School of Biological Sciences, Kelburn Parade, Wellington, 6140, New Zealand.

<sup>B</sup>Research and Development Centre on Marine, Coastal and Small Islands, Hasanuddin University, Gedung Pusat Kegiatan Penelitian Lantai 5, Tamalanrea, Makassar, Indonesia.

<sup>C</sup>Corresponding author. Email: joe.marlow@vuw.ac.nz

**Abstract.** Increasingly frequent large-scale coral mortality events are occurring across the globe, leading to a rise in available reef substrata and promoting an increase in the abundance of other benthic taxa. One such group are bioeroding sponges, which can benefit considerably from reef degradation. However, the occupation of new substrates is partially reliant upon larval recruitment, and currently little is known about the factors affecting bioeroding sponge recruitment. In this study we investigated the potential drivers of bioeroding sponge recruitment using a 2-year deployment of experimental calcareous substrates across seven reefs in the Wakatobi region of Indonesia. Recruitment was observed for five bioeroding sponge species, namely *Cliona orientalis*, *Clathrospira cf. aurivillii*, *Clathrospira hancocki* and two presently unidentified brown clionoids, *Cliona* aff. *viridis* sp. A and *Cliona* aff. *viridis* sp. B. Recruits were present on 69% of the experimental substrates but had a low mean ( $\pm$ s.e.m.) spatial coverage of just  $0.42 \pm 0.13\%$ . Total recruitment and species assemblage structure were correlated with local adult abundance, water flow and substrate cues. Our results suggest that any proliferation of bioeroding sponges on newly available substrate following coral mortality is likely to be conditional on local adult abundance and hydrodynamics.

**Additional keywords:** benthos, bioerosion, larvae.

Received 25 June 2018, accepted 6 November 2018, published online 18 January 2019

### Introduction

Coral reefs are in a state of global decline as sources of anthropogenic stress increase (Bellwood *et al.* 2004; Bruno and Selig 2007; De'ath *et al.* 2012; Jackson *et al.* 2014). In recent years, these declines have largely been attributed to man-made climate change, mass coral bleaching associated with thermal anomalies (Hughes *et al.* 2003, 2017; Eakin *et al.* 2010; DeCarlo *et al.* 2017). Historically, corals have dominated coral reef substrate through competitive interactions with less-competitive taxa. However, large-scale coral die-offs, associated with bleaching events (Glynn 1984; Normile 2016) and other stressors (Aronson and Precht 2001; Kayal *et al.* 2012), are increasingly exposing a greater proportion of bare calcareous substrate on coral reefs. On degraded and stressed reefs, where coral recruitment can be low (Kuffner *et al.* 2006; Hoey *et al.* 2011) and spatial competitive ability is reduced (Aerts 2000; Vermeij *et al.* 2010), other taxa can colonise and competitively maintain this substrate, preventing the re-establishment of hard coral dominance. For example, large-scale coral cover declines in the Caribbean have coincided with regime shifts to macroalgal dominance (Lessios *et al.* 1984; Aronson and Precht 2001; Gardner *et al.* 2003; Mumby 2009; Eakin *et al.* 2010;

Jackson *et al.* 2014). The situation appears to be different in the Indo-Pacific, where, despite some declines in coral cover (Bruno and Selig 2007), macroalgal-dominated reefs are rare (Bruno *et al.* 2009). This has led some authors (e.g. Norström *et al.* 2009; Bell *et al.* 2013) to question what other benthic taxa may occupy the space left by coral mortality and potentially dominate degraded reefs. Recent bleaching events (and subsequent coral mortality) have been associated with large increases in bioeroding sponge abundance on reefs across the globe (Sheppard *et al.* 2002; Schönberg and Ortiz 2009; Carballo *et al.* 2013). These increases and the resilience demonstrated by some bioeroding sponges to elevated water temperature and reduced pH have led several authors to propose bioeroding sponges as possible 'winners' on degraded coral reefs (e.g. Schönberg and Ortiz 2009; Chaves-Fonnegra *et al.* 2018).

The ability of bioeroding sponges to occupy newly available substrate following coral mortality is likely to occur through lateral encroachment, fragmentation, larval recruitment or a combination of these processes (Lopez-Victoria and Zea 2004, 2005; Zilberberg *et al.* 2006; Chaves-Fonnegra and Zea 2011). Larval recruitment is vital for sponge population maintenance (Pineda *et al.* 2009, 2010; McMurray *et al.* 2010; Knapp *et al.*

2016), but may also be important for regulating adult distribution (Uriz *et al.* 1998) and supporting the expansion of a species into novel habitats. Recruitment rates in sponges are generally considered to be low (Zea 1993) and appear to be particularly low from the few studies that indirectly assessed recruitment in bioeroding sponges (Kiene and Hutchings 1994; Pari *et al.* 1998, 2002). Recruitment of internal bioeroders has been studied in French Polynesia and the Great Barrier Reef (GBR), and these studies have revealed a successional pattern from initial colonisation by polychaetes and sipunculans to an increasing abundance of boring molluscs and sponges with time (Kiene and Hutchings 1994; Pari *et al.* 1998, 2002; Tribollet *et al.* 2002).

Although sponges are capable of asexual reproduction and dispersal (e.g. Lopez-Victoria and Zea 2004), the prevailing reproductive mode is sexual and dispersal relies on a free-living larval stage (Uriz *et al.* 2008). As such, recruitment is thought to be limited by larval production (Mariani *et al.* 2001; Piscitelli *et al.* 2011), survival (e.g. Uriz *et al.* 1996), dispersal (Maldonado 2006), habitat selection and settlement (Maldonado and Uriz 1998) and post-settlement survival (Maldonado and Young 1996; de Caralt *et al.* 2007). The presettlement lifespan of sponge larvae is generally short, with larvae normally only remaining in the water column for minutes to days and usually for less than 2 weeks (Maldonado 2006). Once larvae contact the seabed, substratum exploration occurs at a scale of centimetres to metres, and can last longer than the planktonic phase (Maldonado 2006). Little is known about the mechanisms that influence final settlement location selection, but chemical signals produced by conspecifics, competitors or substrates are thought to be involved, as is the case for other invertebrate larva (Pawlik 1992). Evidence for these cues in sponges (e.g. microbial biofilms) are only beginning to emerge (Whalan *et al.* 2008, 2012; Abdul Wahab *et al.* 2011; Whalan and Webster 2014).

Most of what is currently known about reproduction and recruitment in bioeroding sponges comes from studies on *Cliona celata*, *C. delix*, *C. tenuis* and *C. viridix*. From these studies, it appears that oviparous embryonic development is the most widespread in bioeroding sponges (although potentially with internal fertilisation; Rosell 1993; Mariani *et al.* 2000; Maldonado and Riesgo 2008; González-Rivero *et al.* 2013). Gamete development occurs on an annual cycle and release is usually a highly synchronised event that has been related to water temperature (Mariani *et al.* 2000; Piscitelli *et al.* 2011; González-Rivero *et al.* 2013). Clonidae larvae are clavabastula (Maldonado 2006; Maldonado and Riesgo 2008) and particularly weak swimmers, often exhibiting predominantly crawling rather than swimming behaviour (Mariani *et al.* 2000, 2001, 2006). Consequently, dispersal is low and philopatric (Mariani *et al.* 2006). There are exceptions to these generalisations; for example, the Caribbean sponge *C. delix* exhibits asynchronous and multiple spawning events, and has wide larval dispersal (<350 km; Chaves-Fonnegra *et al.* 2015, 2016).

Despite some studies indirectly investigating bioeroding sponge recruitment within the Indo-Pacific (Kiene and Hutchings 1994; Tribollet *et al.* 2002), our knowledge of the drivers of recruitment is still very limited. Understanding these factors is important given the increase in potential habitat availability due to regional decreases in coral cover (Bruno

and Selig 2007). Declines in coral cover may not only promote habitat availability for bioeroding sponges, but also the abundance of other benthic taxa that may act as cues for sponge recruitment. This study focused on the recruitment of bioeroding sponges on Indonesian coral reefs. Using a 2-year deployment of experimental calcareous substrates, recruitment was analysed in relation to water flow, substrate composition and local adult sponge abundance.

## Materials and methods

### Study area

This study was conducted within the Wakatobi UNESCO Biosphere Reserve in south-east Sulawesi, Indonesia. Bioeroding sponge recruitment was studied at seven reef sites fringing the islands of Hoga and Kaledupa (Fig. 1; Table 1; Marlow *et al.* 2018a). The sites (Buoy 1, Buoy 3, Kaledupa 1, Kaledupa Double Spur, Pak Kasim's, Ridge 1 and Sampela 1; abbreviated as B1, B3, K1, KDS, PK, R1 and S1) were chosen to represent a variety of reef types, environmental conditions and levels of reef degradation.

This research was conducted under a permit (182/SIP/FRP/E5/Dit.KI/VI/2016) from the Indonesian Ministry of Research and Technology (RISTEK).

### Bioeroding sponge recruitment

Bioeroding sponge recruitment was studied using a 2-year deployment of experimental calcareous substrates. The cubic calcareous recruitment blocks (10 × 10 × 10 cm) were made of cemented reef rock (limestone) obtained from a mine on the local coralline island of Wanci. The blocks were sourced in this way because the local availability of suitable dead coral was insufficient. Furthermore, previous work (Marlow *et al.* 2018b) has established the ecological relevance of these blocks, because bioeroding sponges erode into them at the same rates as into *Porites lutea*, a locally abundant coral species. Prior to deployment, recruitment blocks were cleaned with a wire brush (but not chemically cleaned) and cured in seawater for 24 h. Deployment occurred in July and August 2014; at each of the seven sites and at depths of 10 and 5 m, five blocks were attached 15 m apart to suitable patches of dead substrate using marine epoxy ( $n = 70$ ). Each block was then photographed in reference to notable reef features to aid future recovery.

In July and August 2016, the recruitment blocks were recovered from the reef sites. Upon locating each block, each exposed side of the block was examined for any bioeroding sponges, noting species presence and location and the presence of any taxa that may be mistaken for sponges during image analysis. Blocks were then photographed *in situ* (every exposed side) and removed from the reef. In the laboratory, each block was subsequently split open using a hammer and chisel to either confirm in-water observations or locate previously unobserved sponges.

Recruitment was analysed using the *in situ* images and imaging software VidAna (ver. 1.0.1, J. Hedley, The University of Queensland, St Lucia, Qld, Australia, see <http://www.marinespatialecologylab.org/resources/vidana/>). From the images of each side of the block, recruits were located (with the aid of field notes) and marked (Fig. 2). Recruitment was




Fig. 1. Map of sponge recruitment study sites within the Wakatobi UNESCO Biosphere Reserve in south-east Sulawesi. B1, Buoy 1; B3, Buoy 3; K1, Kaledupa I; KDS, Kaledupa Double Spur; PK, Pak Kasim's; R1, Ridge I; S1, Sampela I.

Table 1. Reef site descriptions

Coral cover data sourced from Marlow *et al.* (2018a), stressor information sourced from Clifton *et al.* (2010)

Reef	Reef type	Live coral (%)	Main localised stressors
Buoy 1	Steep wall	19	Fisheries
Buoy 3	Steep wall	34	None
Kaledupa 1	Sloping reef	11	Fisheries
Kaledupa Double Spur	Sloping reef	15	Fisheries
Pak Kasim's	Sloping reef	24	None
Ridge 1	Steep wall	28	Fisheries
Sampela 1	Sloping reef	12	Fisheries, sedimentation, coral mining

measured in 61 ms of percentage cover of sponge recruits (per species) in relation to the total area of the recruitment block side. Each side of the recruitment block was considered a subsample and data was averaged 43 for all five sides of the block.

Previous research (J. Marlow, J. J. Bell, A. Haris, M. Shaffer, and C. H. L. Schönberg, unpubl. data) has used macromorphological differences and spicule and phylogenetic analyses to identify the common bioeroding sponge species of the Wakatobi region. That research identified eight bioeroding sponge species, namely *Cliona orientalis*, *Cliona* cf. *schmidtii*, *Clathrospira* cf. *aurivillii*, *Clathrospira hancocki*, *Spherospongia* cf. *vagabunda*, *Zyzygia criceta* and two unknown clionoids, *Cliona* aff. *viridis* sp. A and B. These species are recognisably differentiable at the macromorphological level and so, in the present study,

bioeroding sponges species were identified by the naked eye. In the rare occurrences of identification uncertainty, species identity was confirmed by spicule analysis of bleach-dissolved tissue using light microscopy.

#### Biotic and abiotic predictor variables

Data were collected to determine the extent to which bioeroding sponge recruitment was correlated with the local adult population, substrate cues or flow.

The local adult bioeroding sponge population and substrate composition adjacent to each recruitment block were assessed using photographic quadrats, with images taken during block deployment in 2014. Each 1-m<sup>2</sup> quadrat was divided into 16 subdivisions and centrally placed around every recruitment




Fig. 2. Example screen shots of the VidAna software (ver. 1.0.1, J. Hedley, see <http://www.marinespatialecologylab.org/resources/vidana/>). (a) Image depicting one side of the recruitment block before measurement; (b) the same block after the recruits had been located and tagged.

block. Photographs were then taken of each subdivision, and the presence of any bioeroding sponges within the subdivisions was noted. The images of each photographic quadrat subdivision (and in combination with field notes) were subsequently examined using VidAna software; bioeroding sponge presence and cover was measured in terms of percentage cover of each species per subdivision. Benthic composition (other than bioeroding sponge) of the substrate adjacent to each block was measured using Coral Point Count (CPCc) software (<https://cnso.nova.edu/cpee/index.html>, accessed 1 December 2018; Kohler and Gill 2006). Fifteen points were randomly allocated to each subdivision image ( $n = 240$  per photographic quadrat) and the substrate beneath each point was then allocated to one of the following categories: macroalgae, coralline algae (CCA), hard coral, sponge, soft coral, dead coral + algae, rubble, dead platform, recently dead coral (bleached), sand, silt, sediment on rock or 'other'. Bioeroding sponge cover and substrate composition data from each subdivision were averaged to create mean values for the entire square metre.

Analysis of the composition of the other benthic colonists of the recruitment blocks in 2016 was used to determine whether substrate cues at finer spatial scale were affecting bioeroding sponge recruitment. Previous studies have found that the recruitment of bioeroding sponges can take between 2 and 4 years (Kiene and Hutchings 1994; Piri *et al.* 1998, 2002; Tribollet *et al.* 2002). Therefore it was assumed that the majority of the other benthic types were already present on the blocks at the time of bioeroding sponge recruitment. The *in situ* images of recruitment blocks that were taken in 2016 for the purposes of sponge recruitment analysis were used for the analysis of the benthic composition of recruitment blocks. Twenty points were randomly allocated to each block side image ( $n = 100$  per block) and the benthic taxa beneath each point identified as one of the following categories: ascidians, encrusting bryozoans, macroalgae, coralline algae, hard coral, soft coral, sponges, dead carbonate, settled sediment and 'other'. Data from each side of the block was taken as a subsample and averaged to create mean block benthic composition.

Flow rates were measured using mean proportional flow for each site by means of gypsum 'clods' (Doty 1971; Jokiel and Morrissey 1993). Clods were deployed for 24 h at depths of 10 and 5 m at each site on three occasions in June–August 2015, and proportional differences in water flow were determined from differences in percentage dry weight loss (see Marlow *et al.* 2018b).

#### Statistical analysis

Univariate statistical analyses were performed in SPSS (ver. 23, IBM Corp., Armonk, NY, USA) to test the hypothesis that total recruitment differed among sites and between depths or was affected by the local abundance of adult bioeroding sponges, substrate cues or flow. To test the hypothesis that these same factors would also affect the assemblage composition of bioeroding sponge recruits, multivariate analyses were performed using the PRIMER-E (ver. 6, Plymouth Marine Laboratory, Plymouth, UK) in the permutational multivariate analysis of variance (PERMANOVA) add on.

#### Bioeroding sponge recruitment analysis

Differences in total recruit abundance (species combined) among sites and between depths was tested using a general linear model (GLM) and Tukey's *post hoc* tests using fourth root-transformed data. Correlations between recruit abundance and all biotic and abiotic predictor variables (localised substrate, block benthos, flow rates and adult bioeroding sponge abundance) were assessed using Spearman's rank correlation coefficient.

Sponge recruitment was highly variable: recruit assemblage data (average percentage cover per species per recruitment block) ranged from the very small (e.g. a single papillae) to quite large (e.g. many merging papillae) and contained many zeros. Therefore, the multivariate data was square root transformed and resemblance matrices were constructed using Bray–Curtis similarity coefficients. Because of the absence of bioeroding sponge recruits on many recruitment blocks, a dummy species with an abundance value of 1.0 was added to each recruitment block

**Table 2. Average percentage cover of bioeroding sponge recruits for individual species per site**

Data are given as the mean  $\pm$  s.e.m. Also shown is total mean recruitment (all species combined) per site and mean for species across all recruitment blocks. Not shown are Species that did not recruit are not shown. B1, Buoy 1; B3, Buoy 3; K1, Kaledupa 1; KDS, Kaledupa Double Spur; PK, Pak Kasim's; R1, Ridge 1; S1, Sampela 1

Site	<i>Clothosa cf. aurivillii</i>	<i>Clothosa hancocki</i>	<i>Cliona orientalis</i>	<i>Cliona aff. viridis</i> sp. A	<i>Cliona aff. viridis</i> sp. B	Total
B1	0	0	0	0	0.100 $\pm$ 0.060	0.100 $\pm$ 0.060
B3	0.004 $\pm$ 0.003	0.008 $\pm$ 0.008	0	0	0.092 $\pm$ 0.030	0.104 $\pm$ 0.025
K1	0	0	0	0.213 $\pm$ 0.140	0.058 $\pm$ 0.016	0.270 $\pm$ 0.194
KDS	0.036 $\pm$ 0.017	0	1.297 $\pm$ 0.500	0	0	1.333 $\pm$ 0.778
PK	0.218 $\pm$ 0.211	0	0	0.040 $\pm$ 0.040	0.026 $\pm$ 0.024	0.284 $\pm$ 0.220
R1	0.040 $\pm$ 0.040	0	0.364 $\pm$ 0.342	0.012 $\pm$ 0.012	0.290 $\pm$ 0.240	0.706 $\pm$ 0.402
S1	0.056 $\pm$ 0.030	0	0.007 $\pm$ 0.005	0	0.058 $\pm$ 0.013	0.120 $\pm$ 0.049
Mean	0.052 $\pm$ 0.033	0.001 $\pm$ 0.001	0.237 $\pm$ 0.125	0.034 $\pm$ 0.027	0.092 $\pm$ 0.038	0.417 $\pm$ 0.134

(Clarke *et al.* 2006). Differences in the composition of recruit assemblages among sites and between depths (and any interactions) were assessed using a two-factor PERMANOVA, and results were displayed graphically using principal coordinate analysis (PCO). The species of recruits that contributed most towards any site or depth differences were further investigated using similarity percentage analysis (SIMPER).

Differences in bioeroding sponge recruit assemblages were also analysed in relation to differences in biotic and abiotic explanatory variables using a distance-based multiple linear regression model (DistLM). Explanatory variables consisted of: (1) local adult bioeroding sponge abundance; and (2) benthic composition of the adjacent substrate and recruitment blocks (percentage cover), and flow rates. In order to find the most parsimonious model, Akaike's information criterion (AIC), a step-wise procedure, adjusted for small sample sizes (AICc) was used (Burnham and Anderson 2004). DistLM outputs were represented graphically with a distance-based redundancy ordination (dbRDA) in two-dimensional (2D) bubble plots of the highest recruiting species (species that recruited  $\geq 10$  recruitment blocks).

#### Benthic characterisation and flow analysis

The significance of differences in the benthic composition of the recruitment blocks and adjacent substrate among sites and between depths was analysed using PERMANOVAs based on Bray-Curtis resemblance matrices constructed using square root-transformed data. These were characterised graphically using PCO with overlaying Pearson's correlation vectors. The same methodology was repeated to characterise the differences in assemblage composition of the adult bioeroding sponges adjacent to the recruitment blocks.

The significance of proportional differences in water flow rates (cloud dissolution rates) among sites was tested using a GLM and Tukey's *post hoc* tests.

Where appropriate, data are given as the mean  $\pm$  s.e.m.

## Results

### Recruitment

After 2 years of deployment, five species of bioeroding sponges had recruited to most of the deployed recruitment blocks, and bioeroding sponge recruits were present at every site and depth category.




Fig. 3. Two-dimensional representation (principal coordinate analysis; PCO) of site similarities with regard to recruited bioeroding sponge species assemblage. B1, Buoy 1; B3, Buoy 3; K1, Kaledupa 1; KDS, Kaledupa Double Spur; PK, Pak Kasim's; R1, Ridge 1; S1, Sampela 1.

Of the 70 recruitment blocks deployed across the 7 reef sites, 65 were recovered and 45 of these (69.2%) had at least one sponge recruit. However, mean cover ( $0.42 \pm 0.13\%$ ) and species richness ( $1.2 \pm 0.1$  species per block) were generally low. Average recruit cover ranged from  $0.10 \pm 0.06$  at B1 to  $1.3 \pm 0.8\%$  at KDS and was not significantly different among sites or between depths.

Five bioeroding sponge species were found to have recruited to the experimental substrate blocks. These species were *C. aff. viridis* sp. B (28 blocks), *Cl. cf. aurivillii* (11 blocks), *C. orientalis* (11 blocks), *C. aff. viridis* sp. A (3 blocks) and *Cl. hancocki* (1 block). Despite recruiting to the most number of blocks, *C. aff. viridis* sp. B was not as abundant overall as *C. orientalis*, predominantly due to large recruitment events for the latter species at KDS and R1 (see Table 2).

The assemblage composition of the recruits differed significantly among sites (PERMANOVA, pseudo  $F = 3.876$ ,  $P = 0.001$ ; Fig. 3), but not between depths; in addition, there was no interaction between sites and depths. SIMPER showed that the biggest differences occurred between KDS and both K1

and B1 (both 100% dissimilarity), driven by the dominance of *C. orientalis* in the KDS assemblage, which was absent at K1 and B1. The smallest differences were between B3 and S1 (63.98% dissimilarity), K1 and S1 (72.4% dissimilarity) and B3 and K1 (70.52% dissimilarity); at all these sites, *C. aff. viridis* sp. B was abundant, but the relative importance of other species, such as *C. aff. viridis* sp. A and *Cl. cf. aurivillii*, differed (Table 2).

**Biotic and abiotic variables**

Local substrate composition adjacent to each block in 2014 differed significantly among sites (Fig. 4a) and between depths (Fig. 4b), and there was a significant interaction between depth and site (Table 3). The benthic community on the recruitment blocks in 2016 was also significantly different among sites, but not between depths and there was no significant interaction between site and depth (Table 3; Fig. 4c).

Significant differences in water flow (using the proxy clod dissolution) were found between the sites (GLM,  $F_{(6,76)} = 16.906, P < 0.001$ ). *Post hoc* Tukey's tests showed that flow at KDS was significantly ( $P < 0.05$ ) higher than all sites except K1, and that flow rates at B1 and B3 were significantly lower than at KDS, K1 and R1.

All eight regionally common species of bioeroding sponges (Marlow et al. 2018a) were found to be present in the 1 m<sup>2</sup> adjacent to the recruitment blocks. The species composition of these adjacent bioeroding sponge assemblages differed significantly among sites and between depths, but there was no significant interaction between site and depth (Table 3). Most sites were associated with assemblages that were dominated by *Cl. cf. aurivillii*, but KDS was characterised by a high abundance of *C. orientalis*, and several blocks at K1, KDS and R1 were adjacent to assemblages characterised by the presence of *C. aff. viridis* sp. A (Fig. 5)

**Effects of biotic and abiotic variables on recruitment**


Total recruit percentage cover was positively correlated with local adult bioeroding sponge abundance ( $r_s = 0.338, P = 0.006$ ; Fig. 6a) and flow ( $r_s = 0.333, P = 0.007$ ; Fig. 6b), and was

negatively correlated with the abundance of algae-covered dead coral in the adjacent substrate ( $r_s = -0.318, P = 0.010$ ). None of the other measured biotic or abiotic variables was significantly correlated with total recruitment.

Variation in the local abundance of adult sponges explained 26.6% (Table 4) of the variation of the recruit assemblage composition, with the DistLM selecting three variables: *C. orientalis* (14.8%), *C. aff. viridis* sp. A (6.6%) and *Cl. cf. aurivillii* (5.1%). Species-specific dbRDA bubble plots (Fig. 7) indicate that this is driven primarily by higher recruitment of *C. orientalis* at locations where adult populations were already more established. Recruitment of *C. aff. viridis* sp.


**Table 3. Results of permutational multivariate analysis of variance (PERMANOVA) of the significance of differences in the composition of the local substrate, recruitment block benthos and local adult bioeroding community**

Factor	d.f.	SS	MS	Pseudo F	P <sub>perm</sub>
<b>Local substrate</b>					
Site	6	255.09	42.514	5.6117	0.001
Depth	1	29.118	29.118	3.8435	0.001
Site × depth	6	89.538	14.923	1.9698	0.001
Residual	51	386.38	7.576		
Total	64	768			
<b>Recruitment block benthos</b>					
Site	1	8.4527	8.4527	0.91856	0.529
Depth	6	97.837	16.306	1.772	0.002
Site × depth	6	66.141	11.023	1.1979	0.172
Residual	51	469.31	9.2021		
Total	64	640			
<b>Local adult bioeroding sponges</b>					
Site	1	1185.4	1185.4	2.2846	0.047
Depth	6	10289	1714.9	3.3053	0.001
Site × depth	6	4500	750	1.4455	0.072
Residual	51	26461	518.84		
Total	64	42602			




**Fig. 4.** Two-dimensional representation (principal coordinate analysis; PCO) of benthic community composition of the substrate adjacent to (a, b) and on the recruitment blocks (c). (a, b) The same PCO is shown, but points are either designated by site (a) or depth (b). Overlaid vectors represent substrate components that have a Pearson's correlation coefficient >0.4 with either of the PCO axes. B1, Buoy 1; B3, Buoy 3; K1, Kaledupa 1; KDS, Kaledupa Double Spur; PK, Pak Kasim's; R1, Ridge 1; S1, Sampela 1.

B was also associated with locations where the adult population of *Cl. cf. aurivillii* was high. Variation in flow and substrate composition (both adjacent to and on the recruitment blocks) explained 31.3% of variation in the recruit assemblage (Table 4); the best DistLM model selected flow (14.5%), macroalgae (3.6%) and bare substrate (3.2%) on the recruitment blocks, and algae-covered dead coral (4.3%), recently dead coral (2.9%) and rubble (2.8%) adjacent to the recruitment blocks. Species-specific dbRDA (Fig. 8) bubble plots indicate that recruitment of both *Cl. cf. aurivillii* and *C. aff. viridis* sp. B was associated with low-flow sites and a higher cover of recently dead cover in adjacent benthos. The recruitment of *C. orientalis* was more associated with higher-flow sites with low rubble and macroalgae cover in the adjacent benthos.



**Fig. 5.** Two-dimensional representation (principal coordinate analysis; PCO) of recruitment block similarities with regard to their adjacent bioeroding sponge assemblage composition. Overlaid vectors represent Pearson's correlation of the same data with either of the PCO axes. Only *Clathrospira* (*Cl.*) *cf. aurivillii*, *Clathrospira* (*C.*) *orientalis* and *C. aff. viridis* sp. A have correlation coefficients  $>0.4$ . B1, Buoy 1; B3, Buoy 3; K1, Kaledupa 1; KDS, Kaledupa Double Spur; PK, Pak Kasim's; R1, Ridge 1; S1, Sampela 1.



**Fig. 6.** Total bioeroding sponge recruitment (percentage cover) in relation to the (a) local abundance of adult bioeroding sponges (percentage cover) and (b) average water flow. Note data in (a) are represented on log axes; error bars in (b) represent the s.e.m.

## Discussion


Over the course of this 2-year study, five species of bioeroding sponge were observed to have recruited to most of the deployed experimental blocks, across all sites and depths. Species composition of recruit assemblages was distinctly different among sites, apparently due to species-specific differences in the effects of philopatry, hydrodynamic regime and substrate cues. Although spatial cover of the recruits was generally low compared with the adult cover of these sponges in the region (Marlow *et al.* 2018a), compared with other studies (Kiene and Hutchings 1994; Pari *et al.* 1998, 2002) the proportion of blocks that exhibited bioeroding sponge recruitment was very high for this length of study.

Total bioeroding sponge recruitment and assemblage composition was associated with the local abundance of adult bioeroding sponges, suggesting limited larval dispersal (or a preference for larval settlement among conspecifics) for some species. Although it is impossible to be conclusive about dispersal without the use of genetic markers, *ex situ* larval manipulation (*in situ* larval sampling (e.g. Mariani *et al.* 2006; Whalan *et al.* 2008; Chaves-Fonnegra *et al.* 2015), the results of this study suggest limited larval dispersal for *C. orientalis* and a wider dispersal for *C. aff. viridis* sp. B and *Cl. cf. aurivillii*. Recruitment of *C. orientalis* at sites where this species was already well established is consistent with the limited larval dispersal of the Mediterranean bioeroding sponge *C. viridis*, which is highly correlated with the abundance of adults (Mariani *et al.* 2000). The correlation is attributed to clusters of eggs adhering to the substratum adjacent to 'mother' sponges and the predominantly crawling behaviour of larvae (Mariani *et al.* 2000, 2001, 2006). Heightened larval settlement in the presence of conspecific larvae (and to a lesser extent adults) is also known to occur in sponges (Ettinger-Epstein *et al.* 2008). Limited dispersal and philopatric aggregated recruitment is thought to facilitate settlement to favourable habitats (Pawlik 1992), and is particularly important for zooxanthellate bioeroding sponges (such as *C. orientalis* and *C. viridis*) that require a well-lit calcareous habitat (López-Victoria and Zea 2005). The alternative strategy of a wider dispersal by *C. aff. viridis* sp. B


**Table 4.** Table of sequential test results of biotic and abiotic predictor variables selected as the best solution for bioeroding sponge recruitment by distance-based linear model (DistLM) analyses

AICc, Akaike's information criterion corrected for small sample size; SS, sum of squares; proportion of variation; cumulative proportion of variation

	55 AICc	SS (trace)	Pseudo-F	P	Prop.	Cumul.
Species (adult bioeroding sponges)						
<i>Cliona orientalis</i>	422.3	7036.3	10.976	0	0.1484	0.1484
<i>Cliona</i> aff. <i>viridis</i> sp. A	419.2	3121.2	5.1926	0.01	0.0658	0.2142
<i>Clathrospira cf. aurivillii</i>	417.1	2438.9	4.2717	0.01	0.0514	0.2656
Factor (local substrate, block benthos and flow)						
Flow	422.5	6871.6	10.675	0	0.1449	0.1449
Algae-covered dead coral (adjacent substrate)	421.3	2060	3.3181	0.02	0.0434	0.1883
Macroalgae (block)	420.7	1697.9	2.8148	0.04	0.0358	0.2241
Bare substrate (block)	420.3	1499.3	2.5488	0.05	0.0316	0.2558
Recently dead coral (adjacent substrate)	420.2	1376.5	2.3943	0.09	0.029	0.2848
Rubble (adjacent substrate)	420.1	1345.1	2.3951	0.04	0.0284	0.3131



**Fig. 7.** Distance based redundancy analysis (dBRDA) ordinations of fitted models for recruit assemblage composition in relation to adult population. Overlaid two-dimensional bubbles represent recruit abundance of (a) *Clathrospira cf. aurivillii*, (b) *Cliona orientalis* and (c) *Cliona* aff. *viridis* sp. B. Overlaid vectors represent adult abundance components that were selected by the distance-based linear model (DistLM) analysis.



**Fig. 8.** Distance based redundancy analysis (dBRDA) ordinations of fitted models for recruit assemblage composition in relation to flow and substrate composition (both adjacent to and on the recruitment blocks). Overlaid two-dimensional bubbles represent the recruit abundance of (a) *Clathrospira cf. aurivillii*, (b) *Cliona orientalis* and (c) *Cliona* aff. *viridis* sp. B. Overlaid vectors represent components that were selected by the distance-based linear model (DistLM) analysis.

and *Cl. cf. aurivillii* is likely to incur greater presettlement mortality, but also has a greater chance of avoiding localised extinction (Uriz et al. 1998; Maldonado 2006).

Dispersal distance is likely to be a consequence of not just larval mobility, but also local hydrodynamics, which was

found to affect recruit abundance and assemblage composition. Overall recruitment was greatest at the two sites with the highest degree of water movement (KDS and R1) and least at the sites with lowest water movement (B1 and B3). Research on the tropical non-bioeroding sponge *Cartierospongia*

*foliascens* (Abdul Wahab *et al.* 2014) has shown that hydrodynamics can have a positive effect on fecundity, with consequentially higher recruitment. This may explain the generally lower recruitment levels on reefs with a low degree of water movement where *C. aff. viridis* sp. B was the only species to recruit in considerable numbers. Interestingly, the high overall recruitment at the high-flow sites was largely influenced by the recruitment of *C. orientalis* on these reefs. It may be assumed that a high degree of water flow would increase larval dispersal and not the philopatry observed in this species. However, increased flow may also promote fecundity in the local adult sponge population, as has been shown in *C. foliascens* (Abdul Wahab *et al.* 2014). Furthermore, crawling larvae may be decoupled from the effects of high flow rates at the reef level because they are able to manoeuvre within the boundary layer, where they are subject to much less flow speed and turbulence than midwater larvae (Maldonado 2006).

Evidence for the effects of substrate cues on recruitment was not as consistent as that of flow or localised adult population, but there were some associations that warrant discussion. The most consistent was the presence of algae-covered dead coral in substrate adjacent to the recruitment blocks, which was negatively correlated with overall recruitment and was the strongest substrate indicator in the DistLM analysis. Algae, particularly macroalgae, have consistently been shown to be a strong spatial competitor with bioeroding sponges (López-Victoria *et al.* 2006; Chaves-Fonnegra and Zea 2011; González-Rivero *et al.* 2012, 2016) and are known to deter the settlement of invertebrate larvae, such as corals (Arnold *et al.* 2010). Moreover, a recent assessment of bioeroding sponge cover on the GBR (Ramsby *et al.* 2017) found that increases in the abundance of *C. orientalis* were most likely to occur on reefs with low macroalgal cover. The results of the present study suggest that this could be due to enhanced recruitment at these sites. To a lesser degree, the DistLM analysis also showed that recruitment (particularly *Cl. cf. aurivillii* and *C. aff. viridis* sp. B) was associated with higher cover of recently dead coral and coral rubble in the adjacent substrate. This positive relationship strengthens the suggestion that recruitment on uncolonised calcareous substrate is favoured among presettlement bioeroding sponge larvae (López-Victoria and Zea 2005), potentially due to unique microbial biofilms, which are increasingly recognised as important for facilitating settlement in sponges and other marine invertebrates (Whalan *et al.* 2008; Abdul Wahab *et al.* 2011; Whalan and Webster 2014).

Three locally extant species, namely *C. cf. schmidtii*, *S. cf. vagabunda* and *Z. criceta*, did not recruit to the experimental substrates during the 2-year deployment. Interspecies differences in recruitment rates could be due to interspecies differences in fecundity or due to number of pre- and post-settlement factors. Although many bioeroding sponge species have single annual reproductive events (Mariani *et al.* 2000; Rosell and Uriz 2002; Piscitelli *et al.* 2011; González-Rivero *et al.* 2013), recruitment in some species of sponges can often vary inter-annually, with 'pulses' of high recruitment occurring in some years and not in others (e.g. McMurray *et al.* 2010). For these 'pulse' species (e.g. *S. vagabunda*; Kelly 1986), the time scale of this study may have been insufficient to detect recruitment. The

sides of the recruitment blocks were also all equally exposed, which could possibly deter settlement by sponges with a preference for cryptic habitats. Numerous studies have found large differences in sponge abundance between cryptic and exposed surfaces (Maldonado and Young 1996; Stabler *et al.* 2016), which is thought to be a function of negative phototaxis in settlement, reduced exposure to predators or settled sediment (Maldonado and Young 1996). Species such as *C. cf. schmidtii* that are more common in cryptic habitats (J. Marlow, pers. obs.) may require more cryptic substrate for recruitment than offered by the recruitment blocks.

The recruitment of bioeroding sponges to the majority of experimental substrates after only 2 years of deployment in the Wakatobi is considerably more rapid than in other comparable studies. A deployment of experimental substrates by Tribollet *et al.* (2002) in the GBR found no bioeroding sponge presence after 12 months of deployment. A similar absence of bioeroding sponges was found after 15 months of deployment on Palmyra Atoll in the central Pacific (Elmer 2016) and after 24 months in French Polynesia, where only 5% of experimental substrates were colonised by bioeroding sponges (Pari *et al.* 1998). Kiene and Hutchings (1994) found a similar trend around Lizard Island in the GBR, and suggested that succession to bioeroding sponge occupation may take  $\geq 4$  years to appear after substrate becomes available. The disparity between the studies could be due to differences in the timing and investment of reproductive output between the respective species assemblages or differences in adult population densities (Kelly 1986; McMurray *et al.* 2010).

Our observations of rapid bioeroding sponge recruitment indicate that coral mortality in the region could benefit these sponges. Successful species are likely to have reproductive modes that allocate considerable amounts of energy to egg production (e.g. *C. viridis*; Mariani *et al.* 2001; Piscitelli *et al.* 2011), have multiple intra-annual spawning events (e.g. *C. delirix*; Chaves-Fonnegra *et al.* 2016) or exhibit constant recruitment (e.g. *C. tenuis*; González-Rivero *et al.* 2013) or chance-related recruitment in the cases of those that recruit at more irregular intervals (e.g. *S. vagabunda*; Kelly 1986). In addition, although wider larval dispersal (e.g. *Cl. cf. aurivillii* and *C. aff. viridis* sp. B) is generally considered more opportunistic (Mariani *et al.* 2006), species that have more restricted dispersal (e.g. *C. orientalis*) are potentially able to proliferate on more local scales (Uriz *et al.* 1998).

In conclusion, coral mortality may increase the availability of calcareous habitat for bioeroding sponges but recruitment to these novel substrates could also be reliant upon the proximity of adult populations and local hydrodynamics. Although species with wider larval dispersal may benefit from more regional scales of coral mortality (given the right settlement cues), more limited dispersal strategies could result in localised proliferation and even dominance on some degraded reefs.

#### Conflicts of interest

The authors declare that they have no conflicts of interest.

#### Declaration of funding

This study was funded by a Victoria University of Wellington doctoral scholarship awarded to Joseph Marlow and facilitated

by Operation Wallacea, which provided travel funds and access to research facilities on Hoga, Indonesia.

#### Acknowledgments

The authors thank Christine Schönberg for providing invaluable support in the identification of the Wakatobi bioeroding sponge species. The authors also thank all the volunteer and Indonesian staff at Operation Wallacea who helped facilitate this study.

#### References

- Abdul Wahab, M. A., de Nys, R., and Whalan, S. (2011). Larval behaviour and settlement cues of a brooding coral reef sponge. *Coral Reefs* **30**, 451–460. doi:10.1007/s00338-011-0727-5
- Abdul Wahab, M. A., de Nys, R., Abdo, D., Webster, N., and Whalan, S. (2014). The influence of habitat on post-settlement processes, larval production and recruitment in a common coral reef sponge. *Journal of Experimental Marine Biology and Ecology* **461**, 162–172. doi:10.1016/j.jembe.2014.08.006
- Aerts, L. A. (2000). Dynamics behind standoff interactions in three reef sponge species and the coral *Montastraea cavernosa*. *Marine Ecology* **21**, 191–204. doi:10.1046/j.1439-0485.2000.00685.x
- Arnold, S. N., Steneck, R. S., and Mumby, P. J. (2010). Running the gauntlet: inhibitory effects of algal turfs on the processes of coral recruitment. *Marine Ecology Progress Series* **414**, 91–105. doi:10.3354/meps08724
- Aronson, R. B., and Precht, W. F. (2001). White-band disease and the changing face of Caribbean coral reefs. *Hydrobiologia* **460**, 25–38. doi:10.1023/A:1013103928980
- Bell, J. J., Davy, S. K., Jones, T., Taylor, M. W., and Webster, N. S. (2013). Could some coral reefs become sponge reefs as our climate changes? *Global Change Biology* **19**, 2613–2624. doi:10.1111/gcb.12212
- Bellwood, D. R., Hughes, T. P., Folke, C., and Nyström, M. (2004). Confronting the coral reef crisis. *Nature* **429**, 827–833. doi:10.1038/NATURE02691
- Bruno, J. F., and Selig, E. R. (2007). Regional decline of coral cover in the Indo-Pacific: timing, extent, and subregional comparisons. *PLoS One* **2**, e711. doi:10.1371/JOURNAL.PONE.0000711
- Bruno, J. F., Sweatman, H., Precht, W. F., Selig, E. R., and Schutte, V. G. (2009). Assessing evidence of phase shifts from coral to macroalgal dominance on coral reefs. *Ecology* **90**, 1478–1484. doi:10.1890/08-1781.1
- Burnham, K. P., and Anderson, D. R. (2004). Multimodel inference understanding AIC and BIC in model selection. *Sociological Methods & Research* **33**, 261–304. doi:10.1177/0049124104268644
- Carballo, J. L., Bautista, E., Nava, H., Cruz-Barraza, J. A., and Chávez, J. A. (2013). Boring sponges, an increasing threat for coral reefs affected by bleaching events. *Ecology and Evolution* **3**, 872–886. doi:10.1002/ece3.452
- Chaves-Fonnegra, A., and Zea, S. (2011). Coral colonization by the encrusting excavating Caribbean sponge *Cliona delitrix*. *Marine Ecology* **32**, 162–173. doi:10.1111/j.1439-0485.2010.00416.x
- Chaves-Fonnegra, A., Feldheim, K. A., Secord, J., and Lopez, J. V. (2015). Population structure and dispersal of the coral-excavating sponge *Cliona delitrix*. *Molecular Ecology* **24**, 1447–1466. doi:10.1111/mec.13134
- Chaves-Fonnegra, A., Maldonado, M., Blackwelder, P., and Lopez, J. V. (2016). Asynchronous reproduction and multi-spawning in the coral-excavating sponge *Cliona delitrix*. *Journal of the Marine Biological Association of the United Kingdom* **96**, 515–528. doi:10.1017/S0025315415000636
- Chaves-Fonnegra, A., Riegl, B., Zea, S., Lopez, J. V., Smith, T., Brandt, M., and Gilliam, D. S. (2018). Bleaching events regulate shifts from corals to excavating sponges in algae-dominated reefs. *Global Change Biology* **24**, 773–785. doi:10.1111/gcb.13962
- Clarke, K. R., Somerfield, P. J., and Chapman, M. G. (2006). On resemblance measures for ecological studies, including taxonomic dissimilarities and a zero-adjusted Bray–Curtis coefficient for denuded assemblages. *Journal of Experimental Marine Biology and Ecology* **330**, 55–80. doi:10.1016/j.jembe.2005.12.017
- Clifton, J., Unsworth, R. K., and Smith, D. J. (2010). 'Marine Research and Conservation in the Coral Triangle: The Wakatobi National Park.' (Nova Science Publishers: New York, NY, USA.)
- de Caralt, S., Ojens, H., Uriz, M. J., and Wijffels, R. H. (2007). Cultivation of sponge larvae: settlement, survival, and growth of juveniles. *Marine Biotechnology* **9**, 592–605. doi:10.1007/s10126-007-9013-5
- De'ath, G., Fabricius, K. E., Sweatman, H., and Puotinen, M. (2012). The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 17995–17999. doi:10.1073/pnas.1208909109
- DeCarlo, T. M., Cohen, A. L., Wong, G. T., Davis, K. A., Lohmann, P., and Soong, K. (2017). Mass coral mortality under local amplification of 2°C ocean warming. *Scientific Reports* **7**, 44586. doi:10.1038/SREP44586
- Doty, M. S. (1971). Measurement of water movement in reference to benthic algal growth. *Botanica Marina* **14**, 32–35. doi:10.1515/BOTM.1971.14.1.32
- Eakin, C. M., Morgan, J. A., Heron, S. F., Smith, T. B., Liu, G., Alvarez-Filip, L., Baca, B., Bartels, E., Bastidas, C., and Bouchon, C. (2010). Caribbean corals in crisis: record thermal stress, bleaching, and mortality in 2005. *PLoS One* **5**, e13969. doi:10.1371/JOURNAL.PONE.0013969
- Elmer, F. (2016). Factors affecting coral recruitment and calcium carbonate accretion rates on a Central Pacific coral reef. Ph.D. Thesis, Victoria University of Wellington, Wellington, New Zealand.
- Ettinger-Epstein, P., Whalan, S., Battershill, C. N., and de Nys, R. (2008). A hierarchy of settlement cues influences larval behaviour in a coral reef sponge. *Marine Ecology Progress Series* **365**, 103–113. doi:10.3354/meps07503
- Gardner, T. A., Cote, I. M., Gill, J. A., Grant, A., and Watkinson, A. R. (2003). Long-term region-wide declines in Caribbean corals. *Science* **301**, 958–960. doi:10.1126/SCIENCE.1086050
- Glynn, P. W. (1984). Widespread coral mortality and the 1982–83 El Niño warming event. *Environmental Conservation* **11**, 133–146. doi:10.1017/S0376892900013825
- González-Rivero, M., Ferrari, R., Schönberg, C. H. L., and Mumby, P. J. (2012). Impacts of macroalgal competition and parrotfish predation on the growth of a common bioeroding sponge. *Marine Ecology Progress Series* **444**, 133–142. doi:10.3354/meps09424
- González-Rivero, M., Ereskovsky, A. V., Schönberg, C. H. L., Ferrari, R., Fromont, J., and Mumby, P. J. (2013). Life-history traits of a common Caribbean coral-excavating sponge, *Cliona tenuis* (Porifera: Hadromerida). *Journal of Natural History* **47**, 2815–2834. doi:10.1080/00222933.2013.802042
- González-Rivero, M., Bozec, Y. M., Chollett, J., Ferrari, R., Schönberg, C. H. L., and Mumby, P. J. (2016). Asymmetric competition prevents the outbreak of an opportunistic species after coral reef degradation. *Oecologia* **181**, 161–173. doi:10.1007/s00442-015-3541-x
- Hoey, A. S., Pratchett, M. S., and Cvitanovic, C. (2011). High macroalgal cover and low coral recruitment undermines the potential resilience of the world's southernmost coral reef assemblages. *PLoS One* **6**, e25824. doi:10.1371/JOURNAL.PONE.0025824
- Hughes, T. P., Baird, A. H., Bellwood, D. R., Card, M., Connolly, S. R., Folke, C., Grosberg, R., Hoegh-Guldberg, O., Jackson, J. B., Kleypas, J., Lough, J. M., Marshall, P., Nyström, M., Palumbi, S. R., Pandolfi, S. R., Rosen, B., and Roughgarden, J. (2003). Climate change, human impacts, and the resilience of coral reefs. *Science* **301**, 929–933. doi:10.1126/SCIENCE.1085046
- Hughes, T. P., Kerry, J. T., Álvarez-Norega, M., Álvarez-Romero, J. G., Anderson, K. D., Baird, A. H., Babcock, R. C., Beger, M., Bellwood, D. R., and Berkelmans, R. (2017). Global warming and recurrent mass bleaching of corals. *Nature* **543**, 373–377. doi:10.1038/NATURE21707

- Jackson, J. B. C., Donovan, M., Cramer, K., and Lam, V. (Eds) (2014). Status and trends of Caribbean coral reefs 1970–2012. Global Coral Reef Monitoring Network, International Union for the Conservation of Nature Global Marine and Polar Program, Washington, DC, USA.
- Jokiel, P. L., and Morrissey, J. I. (1993). Water motion on coral reefs—evaluation of the clod card technique. *Marine Ecology Progress Series* **93**, 175–181. doi:10.3354/MEPS093175
- Kayal, M., Vercelloni, J., De Loma, T. L., Bosserelle, P., Chancerelle, Y., Geoffroy, S., Stievenart, C., Michonneau, F., Penn, L., and Planes, S. (2012). Predator crown-of-thorns starfish (*Acanthaster planci*) outbreak, mass mortality of corals, and cascading effects on reef fish and benthic communities. *PLoS One* **7**, e47363. doi:10.1371/JOURNAL.PONE.0047363
- Kelly, M. (1986). Systematics and ecology of the sponges of Motupore Island, Papua New Guinea. M.Sc. Thesis, University of Auckland, Auckland, New Zealand.
- Kiene, W., and Hutchings, P. (1994). Bioerosion experiments at Lizard Island, Great Barrier Reef. *Coral Reefs* **13**, 91–98. doi:10.1007/BF00300767
- Knapp, I. S., Williams, G. J., and Bell, J. J. (2016). Temporal dynamics and persistence of sponge assemblages in a central Pacific atoll lagoon. *Marine Ecology* **37**, 1147–1153. doi:10.1111/MAEC.12303
- Kohler, K. E., and Gill, S. M. (2006). Coral Point Count with Excel extensions (CPCe): a Visual Basic program for the determination of coral and substrate coverage using random point count methodology. *Computers & Geosciences* **32**, 1259–1269. doi:10.1016/J.CAGEO.2005.11.009
- Kuffner, I. B., Walters, L. J., Becerro, M. A., Paul, V. J., Ritson-Williams, R., and Beach, K. S. (2006). Inhibition of coral recruitment by macroalgal and cyanobacteria. *Marine Ecology Progress Series* **323**, 107–117. doi:10.3354/MEPS323107
- Lessios, H. A., Robertson, D. R., and Cubit, J. D. (1984). Spread of *Diadema* mass mortality through the Caribbean. *Science* **226**, 335–337. doi:10.1126/SCIENCE.226.4672.335
- López-Victoria, M., and Zea, S. (2004). Storm-mediated coral colonization by an excavating Caribbean sponge. *Climate Research* **26**, 251–256. doi:10.3354/CR026251
- López-Victoria, M., and Zea, S. (2005). Current trends of space occupation by encrusting excavating sponges on Colombian coral reefs. *Marine Ecology* **26**, 33–41. doi:10.1111/J.1439-0485.2005.00036.X
- López-Victoria, M., Zea, S., and Weil, E. (2006). Competition for space between encrusting excavating Caribbean sponges and other coral reef organisms. *Marine Ecology Progress Series* **312**, 113–121. doi:10.3354/MEPS312113
- Maldonado, M. (2006). The ecology of the sponge larva. *Canadian Journal of Zoology* **84**, 175–194. doi:10.1139/Z05-177
- Maldonado, M., and Riesgo, A. (2008). Reproduction in the Phylum Porifera: a synoptic overview. *Treballs de la Societat Catalana de Biologia* **59**, 29–49.
- Maldonado, M., and Uriz, M. J. (1998). Microrefuge exploitation by subtidal encrusting sponges: patterns of settlement and post-settlement survival. *Marine Ecology Progress Series* **174**, 141–150. doi:10.3354/MEPS174141
- Maldonado, M., and Young, C. M. (1996). Effects of physical factors on larval behavior, settlement and recruitment of four tropical demosponges. *Marine Ecology Progress Series* **138**, 169–180. doi:10.3354/MEPS138169
- Mariani, S., Uriz, M., and Turon, X. (2000). Larval bloom of the oviparous sponge *Cliona viridis*: coupling of larval abundance and adult distribution. *Marine Biology* **137**, 783–790. doi:10.1007/S002270000400
- Mariani, S., Piscitelli, M., and Uriz, M. (2001). Temporal and spatial co-occurrence in spawning and larval release of *Cliona viridis* (Porifera: Hadromerida). *Journal of the Marine Biological Association of the United Kingdom* **81**, 565–567. doi:10.1017/S0025315401004246
- Mariani, S., Uriz, M., Turon, X., and Alcoverro, T. (2006). Dispersal strategies in sponge larvae: Integrating the life history of larvae and the hydrologic component. *Oecologia* **149**, 174–184. doi:10.1007/S00442-006-0429-9
- Marlow, J., Schönberg, C. H. L., Davy, S. K., Haris, A., Jompa, J., and Bell, J. J. (2018a). Bioeroding sponge assemblages: the importance of substrate availability and sediment. *Journal of the Marine Biological Association of the United Kingdom*. [Published online early 22 March 2018]. doi:10.1017/S0025315418000164
- Marlow, J., Smith, D., Werorilangi, S., and Bell, J. J. (2018b). Sedimentation limits the erosion rate of a bioeroding sponge. *Marine Ecology* **39**, e12483. doi:10.1111/MAEC.12483
- McMurray, S. E., Henkel, T. P., and Pawlik, J. R. (2010). Demographics of increasing populations of the giant barrel sponge *Aestospongia muta* in the Florida Keys. *Ecology* **91**, 560–570. doi:10.1890/08-2060.1
- Mumby, P. J. (2009). Phase shifts and the stability of macroalgal communities on Caribbean coral reefs. *Coral Reefs* **28**, 761–773. doi:10.1007/S00338-009-0506-8
- Normile, D. (2016). El Niño's warmth devastating reefs worldwide. *Science* **352**, 15–16. doi:10.1126/SCIENCE.352.6281.15
- Norström, A. V., Nyström, M., Lokrantz, J., and Folke, C. (2009). Alternative states on coral reefs: beyond coral-macroalgal phase shifts. *Marine Ecology Progress Series* **376**, 295–306. doi:10.3354/MEPS07815
- Pari, N., Peyrot-Clausade, M., Le Campion-Alsumard, T., Hutchings, P., Chazottes, V., Golubic, S., Le Campion, J., and Fontaine, M. (1998). Bioerosion of experimental substrates on high islands and on atoll lagoons (French Polynesia) after two years of exposure. *Marine Ecology Progress Series* **166**, 119–130. doi:10.3354/MEPS166119
- Pari, N., Peyrot-Clausade, M., and Hutchings, P. A. (2002). Bioerosion of experimental substrates on high islands and atoll lagoons (French Polynesia) during 5 years of exposure. *Journal of Experimental Marine Biology and Ecology* **276**, 109–127. doi:10.1016/S0022-0981(02)00243-5
- Pawlik, J. R. (1992). Chemical ecology of the settlement of benthic marine invertebrates. *Oceanography and Marine Biology* **30**, 273–335.
- Pineda, J., Reyns, N. B., and Starczak, V. R. (2009). Complexity and simplification in understanding recruitment in benthic populations. *Population Ecology* **51**, 17–32. doi:10.1007/S10144-008-0118-0
- Pineda, J., Porn, F., Starczak, V., and Blythe, J. (2010). Causes of decoupling between larval supply and settlement and consequences for understanding recruitment and population connectivity. *Journal of Experimental Marine Biology and Ecology* **392**, 9–21. doi:10.1016/J.JEMBE.2010.04.008
- Piscitelli, M., Corriero, G., Gaiino, E., and Uriz, M. (2011). Reproductive cycles of the sympatric excavating sponges *Cliona celata* and *Cliona viridis* in the Mediterranean Sea. *Invertebrate Biology* **130**, 1–10. doi:10.1111/J.1744-7410.2010.00216.X
- Ramsby, B. D., Hoogenboom, M. O., Whalan, S., Webster, N. S., and Thompson, A. (2017). A decadal analysis of bioeroding sponge cover on the inshore Great Barrier Reef. *Scientific Reports* **7**, 2706. doi:10.1038/S41598-017-02196-Z
- Rosell, D. (1993). Effects of reproduction in *Cliona viridis* (Hadromerida) on zooxanthellae. *Scientia Marina* **57**, 405–413.
- Rosell, D., and Uriz, M. (2002). Excavating and endolithic sponge species (Porifera) from the Mediterranean: species descriptions and identification key. *Organisms, Diversity & Evolution* **2**, 55–86. doi:10.1078/1439-6092-00033
- Schönberg, C. H. L., and Ortiz, J. (2009). Is sponge bioerosion increasing? In 'Proceedings of the 11th International Coral Reef Symposium', 7–11 July 2008, Fort Lauderdale, FL, USA. (Eds B. M. Riegl and R. E. Dodge.) Vol. 1, pp. 520–523. (National Coral Reef Institute: Fort Lauderdale, FL, USA.)
- Sheppard, C. R. C., Spalding, M., Bradshaw, C., and Wilson, S. (2002). Erosion vs recovery of coral reefs after 1998 El Niño: Chagos reefs, Indian Ocean. *Ambio* **31**, 40–48. doi:10.1579/0044-7447-31.1.40

- Stubler, A. D., Stevens, A. K., and Peterson, B. J. (2016). Using community-wide recruitment and succession patterns to assess sediment stress on Jamaican coral reefs. *Journal of Experimental Marine Biology and Ecology* **474**, 29–38. doi:10.1016/j.jembe.2015.09.018
- Tribollet, A., Decherf, G., Hutchings, P., and Peyrot-Clausade, M. (2002). Large-scale spatial variability in bioerosion of experimental coral substrates on the great barrier reef (Australia): importance of microborers. *Coral Reefs* **21**, 424–432.
- Uriz, M., Turon, X., Becerro, M., and Galera, J. (1996). Feeding deterrence in sponges. The role of toxicity, physical defenses, energetic contents, and life-history stage. *Journal of Experimental Marine Biology and Ecology* **205**, 187–204. doi:10.1016/S0022-0981(96)02609-3
- Uriz, M. J., Maldonado, M., Turon, X., and Martí, R. (1998). How do reproductive output, larval behaviour and recruitment contribute to adult spatial patterns in Mediterranean encrusting sponges? *Marine Ecology Progress Series* **167**, 137–148. doi:10.3354/MEPS167137
- Uriz, M. J., Turon, X., and Mariani, S. (2008). Ultrastructure and dispersal potential of sponge larvae: tufted versus evenly ciliated parenchymellae. *Marine Ecology* **29**, 280–297. doi:10.1111/j.1439-0485.2008.00229.x
- Vermeij, M. J., Van Moorselaar, I., Engelhard, S., Hömlein, C., Vonk, S. M., and Visser, P. M. (2010). The effects of nutrient enrichment and herbivore abundance on the ability of turf algae to overgrow coral in the Caribbean. *PLoS One* **5**, e14312. doi:10.1371/JOURNAL.PONE.0014312
- Whalan, S., and Webster, N. S. (2014). Sponge larval settlement cues: the role of microbial biofilms in a warming ocean. *Scientific Reports* **4**, 4072. doi:10.1038/SREP04072
- Whalan, S., Ettinger-Epstein, P., Battershill, C., and de Nys, R. (2008). Larval vertical migration and hierarchical selectivity of settlement in a brooding marine sponge. *Marine Ecology Progress Series* **368**, 145–154. doi:10.3354/MEPS07573
- Whalan, S., Webster, N. S., and Negri, A. P. (2012). Crustose coralline algae and a cnidarian neuropeptide trigger larval settlement in two coral reef sponges. *PLoS One* **7**, e30386. doi:10.1371/JOURNAL.PONE.0030386
- Zea, S. (1993). Recruitment of demosponges (Porifera, Demospongiae) in rocky and coral reef habitats of Santa Marta, Colombian Caribbean. *Marine Ecology* **14**, 1–21. doi:10.1111/j.1439-0485.1993.tb00361.x
- Zilberberg, C., Maldonado, M., and Solé-Cava, A. (2006). Assessment of the relative contribution of asexual propagation in a population of the coral-excavating sponge *Cliona delitrix* from the Bahamas. *Coral Reefs* **25**, 297–301. doi:10.1007/S00338-006-0094-9

Handling Editor: Jacob Johansen

# Bioeroding sponges

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

10%

INTERNET SOURCES

11%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1%
2	<a href="http://researchonline.jcu.edu.au">researchonline.jcu.edu.au</a> Internet Source	1%
3	Submitted to Longwood College Student Paper	1%
4	Edward Castañeda-Moya. "Patterns of Root Dynamics in Mangrove Forests Along Environmental Gradients in the Florida Coastal Everglades, USA", <i>Ecosystems</i> , 09/07/2011 Publication	1%
5	<a href="http://www.publish.csiro.au">www.publish.csiro.au</a> Internet Source	<1%
6	Darren J. Coker, Joseph D. DiBattista, Tane H. Sinclair-Taylor, Michael L. Berumen. "Spatial patterns of cryptobenthic coral-reef fishes in the Red Sea", <i>Coral Reefs</i> , 2017 Publication	<1%
7	<a href="http://hal.archives-ouvertes.fr">hal.archives-ouvertes.fr</a>	

Internet Source

<1%

8

Ahmed A. Issa, Awatief F. Hifney, Khayria M. Abdel-Gawad, Mohamed Gomaa. "Spatio temporal and environmental factors influencing macroalgal  $\beta$  diversity in the Red Sea, Egypt", *Botanica Marina*, 2014

Publication

<1%

9

Muhammad Azmi Abdul Wahab, Ben Radford, Mike Cappo, Jamie Colquhoun et al. "Biodiversity and spatial patterns of benthic habitat and associated demersal fish communities at two tropical submerged reef ecosystems", *Coral Reefs*, 2018

Publication

<1%

10

[www.cambridge.org](http://www.cambridge.org)

Internet Source

<1%

11

Mancini, PL, KA Hobson, and L Bugoni. "Role of body size in shaping the trophic structure of tropical seabird communities", *Marine Ecology Progress Series*, 2014.

Publication

<1%

12

*Coral Reefs of the World*, 2016.

Publication

<1%

13

[repository.essex.ac.uk](http://repository.essex.ac.uk)

Internet Source

<1%

14	Submitted to University of Wales, Bangor Student Paper	<1%
15	Encyclopedia of Earth Sciences Series, 2011. Publication	<1%
16	Emily Howe, Charles A. Simenstad, Andrea Ogston. "Detrital shadows: estuarine food web connectivity depends on fluvial influence and consumer feeding mode", Ecological Applications, 2017 Publication	<1%
17	elib.suub.uni-bremen.de Internet Source	<1%
18	epubs.scu.edu.au Internet Source	<1%
19	www.oncotarget.com Internet Source	<1%
20	espace.library.uq.edu.au Internet Source	<1%
21	Roff, G., T. R. Clark, C. E. Reymond, J.-x. Zhao, Y. Feng, L. J. McCook, T. J. Done, and J. M. Pandolfi. "Palaeoecological evidence of a historical collapse of corals at Pelorus Island, inshore Great Barrier Reef, following European settlement", Proceedings of The Royal Society B Biological Sciences, 2012. Publication	<1%

22	H.T Yap, R.M Alvarez, H.M Custodio, R.M Dizon. "Physiological and ecological aspects of coral transplantation", Journal of Experimental Marine Biology and Ecology, 1998 Publication	<1%
23	<a href="http://www.ou.edu">www.ou.edu</a> Internet Source	<1%
24	<a href="http://www.trialsjournal.com">www.trialsjournal.com</a> Internet Source	<1%
25	Peter Deines. "Methane-derived carbon flows through methane-oxidizing bacteria to higher trophic levels in aquatic systems", Environmental Microbiology, 5/2007 Publication	<1%
26	Submitted to University of Queensland Student Paper	<1%
27	<a href="http://hal.upmc.fr">hal.upmc.fr</a> Internet Source	<1%
28	Submitted to CSU, Long Beach Student Paper	<1%
29	<a href="http://peerj.com">peerj.com</a> Internet Source	<1%
30	Ellis, Lauren M., Heather L. Hunt, and Rémy Rochette. "Effect of background substrate on recruitment of benthic marine invertebrates to	<1%

subtidal cobble-filled collectors", *Marine Biology*, 2015.

Publication

- 
- 31 Edmunds, P. J., R. Steneck, R. Albright, R. C. Carpenter, A. P. Y. Chui, T.-Y. Fan, S. Harii, H. Kitano, H. Kurihara, L. Legendre, S. Mitarai, S. Muko, Y. Nozawa, J. Padilla-Gamino, N. N. Price, K. Sakai, G. Suzuki, M. J. H. van Oppen, A. Yarid, and R. D. Gates. "Geographic variation in long-term trajectories of change in coral recruitment: a global-to-local perspective", *Marine and Freshwater Research*, 2015. <1%
- 32 [diposit.ub.edu](http://diposit.ub.edu) <1%  
Internet Source
- 
- 33 [www.nature.com](http://www.nature.com) <1%  
Internet Source
- 
- 34 [www.amjbot.org](http://www.amjbot.org) <1%  
Internet Source
- 
- 35 [ediss.uni-goettingen.de](http://ediss.uni-goettingen.de) <1%  
Internet Source
- 
- 36 J.J. Bell, J. Jompa, A. Haris, S. Werorilangi, M. Shaffer, C. Mortimer. "Domination of mesophotic ecosystems in the Wakatobi Marine National Park (Indonesia) by sponges, soft corals and other non-hard coral species",

Journal of the Marine Biological Association of  
the United Kingdom, 2018

Publication

- 
- |    |   |     |
|----|---|-----|
| 37 | A. Tribollet. "Cross-shelf differences in the pattern and pace of bioerosion of experimental carbonate substrates exposed for 3 years on the northern Great Barrier Reef, Australia", <i>Coral Reefs</i> , 11/2005<br>Publication | <1% |
| 38 | <a href="http://www.swiofp.net">www.swiofp.net</a><br>Internet Source   | <1% |
| 39 | <a href="http://www.ceab.csic.es">www.ceab.csic.es</a><br>Internet Source   | <1% |
| 40 | <a href="http://www.com.univ-mrs.fr">www.com.univ-mrs.fr</a><br>Internet Source   | <1% |
| 41 | <a href="http://citeseerx.ist.psu.edu">citeseerx.ist.psu.edu</a><br>Internet Source   | <1% |
| 42 | <a href="http://ir.canterbury.ac.nz">ir.canterbury.ac.nz</a><br>Internet Source   | <1% |
| 43 | Submitted to University of Plymouth<br>Student Paper  | <1% |
| 44 | Edlin Guerra-Castro. "Spatial variability and human disturbance of sponge assemblages associated with mangrove roots in the southern Caribbean", <i>Marine and Freshwater Research</i> ,  | <1% |
-

2011

Publication

- 
- 45** Montano, Simone, Giovanni Strona, Davide Seveso, Davide Maggioni, and Paolo Galli. "Widespread occurrence of coral diseases in the central Maldives", *Marine and Freshwater Research*, 2015. <1%  
Publication
- 
- 46** Holly M. Bennett, Christine Altenrath, Lisa Woods, Simon K. Davy, Nicole S. Webster, James J. Bell. " Interactive effects of temperature and CO on sponges: from the cradle to the grave ", *Global Change Biology*, 2017 <1%  
Publication
- 
- 47** [cmsdata.iucn.org](http://cmsdata.iucn.org) <1%  
Internet Source
- 
- 48** [onlinelibrary.wiley.com](http://onlinelibrary.wiley.com) <1%  
Internet Source
- 
- 49** K. Koop, D. Booth, A. Broadbent, J. Brodie et al. "ENCORE: The Effect of Nutrient Enrichment on Coral Reefs. Synthesis of Results and Conclusions", *Marine Pollution Bulletin*, 2001 <1%  
Publication
- 
- 50** Feitosa, Caroline Vieira, Simone Marques, Maria Elisabeth de Araujo, and Beatrice Padovani Ferreira. "Reproduction of French <1%
-

angelfish *Pomacanthus paru* (Teleostei: Pomacanthidae) and implications for management of the ornamental fish trade in Brazil", *Marine and Freshwater Research*, 2015.

Publication


---

51	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	<1%
52	<a href="http://biopublisher.ca">biopublisher.ca</a> Internet Source	<1%
53	<a href="http://www.un.org">www.un.org</a> Internet Source	<1%
54	<a href="http://www.rsmas.miami.edu">www.rsmas.miami.edu</a> Internet Source	<1%
55	<a href="http://luvia.dihma.upv.es">luvia.dihma.upv.es</a> Internet Source	<1%
56	<a href="http://boris.unibe.ch">boris.unibe.ch</a> Internet Source	<1%
57	Patrick, P., and N. A. Strydom. "Larval fish variability in response to oceanographic features in a nearshore nursery area : larval fish nearshore assemblages", <i>Journal of Fish Biology</i> , 2014. Publication	<1%
58	J. A. Scealy. "Aquatic macroinvertebrate communities on wood in an Australian lowland	<1%

---

river: experimental assessment of the interactions of habitat, substrate complexity and retained organic matter", Marine and Freshwater Research, 2007

Publication

- 
- |    |   |     |
|----|---|-----|
| 59 | <a href="http://www.arochoa.org">www.arochoa.org</a><br>Internet Source | <1% |
|----|---|-----|
- 
- |    |   |     |
|----|---|-----|
| 60 | <a href="http://researcharchive.vuw.ac.nz">researcharchive.vuw.ac.nz</a><br>Internet Source | <1% |
|----|---|-----|
- 
- |    |  |     |
|----|--|-----|
| 61 | Submitted to The School for Field Studies<br>Student Paper | <1% |
|----|--|-----|
- 
- |    |   |     |
|----|---|-----|
| 62 | <a href="http://journals.plos.org">journals.plos.org</a><br>Internet Source | <1% |
|----|---|-----|
- 
- |    |   |     |
|----|---|-----|
| 63 | Haapkyll  , J., J. Melbourne-Thomas, and M. Flavell. "The association between coral communities and disease assemblages in the Wakatobi Marine National Park, south-eastern Sulawesi, Indonesia", Marine and Freshwater Research, 2015.<br>Publication | <1% |
|----|---|-----|
- 
- |    |   |     |
|----|---|-----|
| 64 | "Coral Reefs of the Eastern Tropical Pacific", Springer Nature, 2017<br>Publication | <1% |
|----|---|-----|
- 
- |    |   |     |
|----|---|-----|
| 65 | Alexander V. Ereskovsky. "The Comparative Embryology of Sponges", Springer Nature, 2010 | <1% |
|----|---|-----|
-

66	<a href="http://lrd.yahooapis.com">lrd.yahooapis.com</a> Internet Source	<1%
67	<a href="http://www.science.gov">www.science.gov</a> Internet Source	<1%
68	"Climate Change, Ocean Acidification and Sponges", Springer Nature, 2017 Publication	<1%
69	<a href="http://darchive.mblwhoilibrary.org">darchive.mblwhoilibrary.org</a> Internet Source	<1%
70	Ettinger-Epstein, P, S Whalan, CN Battershill, and R de Nys. "A hierarchy of settlement cues influences larval behaviour in a coral reef sponge", Marine Ecology Progress Series, 2008. Publication	<1%
71	"Complete Issue pdf 74-4", ICES Journal of Marine Science, 2017 Publication	<1%
72	D. E. P. Bryant, A. Rodriguez-Ramirez, S. Phinn, M. González-Rivero, K. T. Brown, B. P. Neal, O. Hoegh-Guldberg, S. Dove. "Comparison of two photographic methodologies for collecting and analyzing the condition of coral reef ecosystems", Ecosphere, 2017	<1%

73

Héctor Nava, Carlos Alberto Emmanuel García-Madrigal, José Luis Carballo. "Relationships between boring sponge assemblages and the availability of dead coral substrate on Mexican Pacific coral reefs", Journal of the Marine Biological Association of the United Kingdom, 2018

Publication

<1%

74

Powell, Abigail, David J. Smith, Leanne J. Hepburn, Timothy Jones, Jade Berman, Jamaluddin Jompa, and James J. Bell. "Reduced Diversity and High Sponge Abundance on a Sedimented Indo-Pacific Reef System: Implications for Future Changes in Environmental Quality", PLoS ONE, 2014.

Publication

<1%

75

DA Smale, T Wernberg, MA Vanderklift. "Regional-scale variability in the response of benthic macroinvertebrate assemblages to a marine heatwave", Marine Ecology Progress Series, 2017

Publication

<1%

Exclude quotes  On

Exclude matches  < 1 words

Exclude bibliography  On

