



**The 3rd FSSAT International Conference Attendance List Room 3, Moderator Dr. Hari Iswoyo, SP.,M.Sc.**

No.	Kode ABS	Name	Name of Institution	Date of attendance at international conferences	Title	Status of attendance
1	ABS-258	Fajri Ansari	Environment and Forestry Research and Development Institute of Makassar	January 8th 2021	Development of Bulu Sipong Park (PT Semen Tonasa) to support the preservation of the diversity of ecosystem karst	Presenter
2	ABS-259	Andi Siady Hamzah	Faculty of Forestry, Hasanuddin University	January 8th 2021	Bird diversity and structure in different landcover in mining concession	Presenter
3	ABS-8	Burhanuddin Rasyid	Department of Soil Science, Faculty of Agriculture, Universitas Hasanuddin	January 8th 2021	Different water content and soil amendment effect on water use efficiency in maize growth	Presenter
4	ABS-286	Wardiman Dg. Sipato	Management of Environment, Postgraduate, Hasanuddin University	January 8th 2021	Policy analysis of the adaptation of makassar city's government for climate change and global warming	Presenter
5	ABS-35	Eka Wisdawi	The Agricultural Science, Graduate School, Hasanuddin University	January 8th 2021	Screening and identification cellulolytic fungi at rhizosfer of safira taro plant	Presenter
6	ABS-291	Musfira M	Program Study of Agrotechnology, Faculty of Agriculture, Universitas Hasanuddin	January 8th 2021	Application of humic substances and bottom ash compounds in improving soil quality in lime stone post-mining land	Presenter
7	ABS-38	Aqshan Shadikin Nurdin	Department of Forestry, Faculty of Agriculture, University of Khairun, Ternate, North Moluccas, Indonesia	January 8th 2021	Composition and nutritional content of honey trigona sp in the tikep forest management unit (kph) north moluccas	Presenter
8	ABS-294	Mukrimin	Department of Forestry, Faculty of Forestry, Hasanuddin University	January 8th 2021	Dinamic of soil properties under an exotic pine (Pinus sp.) species growing in Bantaeng Regency	Presenter
9	ABS-45	Intan Giliang Cempaka	BPTP Jawa Tengah, Semarang, Indonesia	January 8th 2021	Characterization of mango (Mangifera indica L.) blungko from Rembang, Central Java	Presenter
10	ABS-55	Inwanto	Faculty of Agriculture, Pattimura University, Ambon, Indonesia	January 8th 2021	Estimated carbon stock of various mangrove zonation in Marsegu Island, West Seram, Maluku	Presenter
11	ABS-57	Risma Riyandani	Faculty of Agriculture, Universitas Hasanuddin	January 8th 2021	Utilization of liquid organic fertilizers for banana stems and coconut husk to increase potassium (K) nutrients in alfisols and corn	Presenter
12	ABS-60	Arini Putri Hanifa, Rinaldi Sjahri	School of Agriculture and Environment of Massey University, Laboratory of Plant Bioscience and Reproduction Biotechnology, Universitas Hasanuddin	January 8th 2021	Agro-morphological variation of pigmented rice landraces from South Sulawesi grown in a temperate glasshouse of New Zealand	Presenter
13	ABS-61	Arinana	Forest Product Department, Faculty of Forestry and Environmental, IPB University, Bogor	January 8th 2021	The daytime indoor and outdoor temperatures of subterranean termite <i>Coptotermes curvignathus holmgren</i> (Isoptera: Rhinotermitidae) tunnel	Presenter
14	ABS-91	A. Mushawwir	Animal Science Faculty, University of Padjadjaran	January 8th 2021	Molecular marker for thermoregulation of pasundan cattle in different ambient temperature	Presenter


  
 Makassar, January 8th 2021  
 Dr. Ir. Asmita Ahmad, ST., M.Si  
 Chair of IC-FSSAT

**The 3rd FSSAT International Conference Attendance List Room 3, Moderator Dr. Hari Iswoyo, SP.,M.Sc.**

No.	Kode ABS	Name	Name of Institution	Date of attendance at international conferences	Title	Status of attendance
1	ABS-258	Fajri Ansari	Environment and Forestry Research and Development Institute of Makassar	January 8th 2021	Development of Bulu Sipong Park (PT Semen Tonasa) to support the preservation of the diversity of ecosystem karst	Presenter
2	ABS-259	Andi Siady Hamzah	Faculty of Forestry, Hasanuddin University	January 8th 2021	Bird diversity and structure in different landcover in mining concession	Presenter
3	ABS-8	Burhanuddin Rasyid	Department of Soil Science, Faculty of Agriculture, Universitas Hasanuddin	January 8th 2021	Different water content and soil amendment effect on water use efficiency in maize growth	Presenter
4	ABS-286	Wardiman Dg. Sipato	Management of Environment, Postgraduate, Hasanuddin University	January 8th 2021	Policy analysis of the adaptation of makassar city's government for climate change and global warming	Presenter
5	ABS-35	Eka Wisdawati	The Agricultural Science, Graduate School, Hasanuddin University	January 8th 2021	Screening and identification cellulolytic fungi at rhizosfer of safira taro plant	Presenter
6	ABS-291	Musfira M	Program Study of Agrotechnology, Faculty of Agriculture, Universitas Hasanuddin	January 8th 2021	Application of humic substances and bottom ash compounds in improving soil quality in lime stone post-mining land	Presenter
7	ABS-38	Aqshan Shadikin Nurdin	Department of Forestry, Faculty of Agriculture, University of Khairun, Ternate, North Moluccas, Indonesia	January 8th 2021	Composition and nutritional content of honey trigona sp in the tikep forest management unit (kph) north moluccas	Presenter
8	ABS-294	Mukrimin	Department of Forestry, Faculty of Forestry, Hasanuddin University	January 8th 2021	Dinamic of soil properties under an exotic pine (Pinus sp.) species growing in Bantaeng Regency	Presenter
9	ABS-45	Intan Giliang Cempaka	BPTP Jawa Tengah, Semarang, Indonesia	January 8th 2021	Characterization of mango (Mangifera indica L.) blungko from Rembang, Central Java	Presenter
10	ABS-55	Inwanto	Faculty of Agriculture, Pattimura University, Ambon, Indonesia	January 8th 2021	Estimated carbon stock of various mangrove zonation in Marsegu Island, West Seram, Maluku	Presenter
11	ABS-57	Risma Riyandani	Faculty of Agriculture, Universitas Hasanuddin	January 8th 2021	Utilization of liquid organic fertilizers for banana stems and coconut husk to increase potassium (K) nutrients in alfisols and corn	Presenter
12	ABS-60	Arini Putri Hanifa, Rinaldi Sjahri	School of Agriculture and Environment of Massey University, Laboratory of Plant Bioscience and Reproduction Biotechnology, Universitas Hasanuddin	January 8th 2021	Agro-morphological variation of pigmented rice landraces from South Sulawesi grown in a temperate glasshouse of New Zealand	Presenter
13	ABS-61	Arinana	Forest Product Department, Faculty of Forestry and Environmental, IPB University, Bogor	January 8th 2021	The daytime indoor and outdoor temperatures of subterranean termite <i>Coptotermes curvignathus holmgren</i> (Isoptera: Rhinotermitidae) tunnel	Presenter
14	ABS-91	A. Mushawwir	Animal Science Faculty, University of Padjadjaran	January 8th 2021	Molecular marker for thermoregulation of pasundan cattle in different ambient temperature	Presenter


  
 Makassar, January 8th 2021  
 Dr. Ir. Asmita Ahmad, ST., M.Si  
 Chair of IC-FSSAT

PAPER • OPEN ACCESS

## Peer review declaration

To cite this article: 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **807** 011002

View the [article online](#) for updates and enhancements.

You may also like

- [Peer review declaration](#)
- [Peer Review Declaration](#)
- [Peer review declaration](#)

## Peer review declaration

All papers published in this volume of IOP Conference Series: Earth and Environmental Science have been peer reviewed through processes administered by the Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

- **Type of peer review:**  
Single-blind
- **Conference submission management system:**  
We conduct paper submission through submission management system (<http://konfrenzi.com/>). The manuscript from author has submitted before the conference through the conference website (<https://fssat3.unhas.ac.id/>).
- **Number of submissions received:**  
Total received abstract = 364 papers
- **Number of submissions sent for review:**  
355 full papers (19 papers rejected)
- **Number of submissions accepted:**  
291 full papers
- **Acceptance Rate (Number of Submissions Accepted / Number of Submissions Received X 100):**  
79.9 %
- **Average number of reviews per paper:**  
6 papers per reviewer
- **Total number of reviewers involved:**  
47 reviewers
- **Any additional info on review process:**  
There are some steps on review process
  1. Abstract review and acceptance  
At the first stage, abstracts submitted to the committee through the online system available in the website were reviewed by a Scientific Advisory Board/Reviewers for the suitability of the paper's topic to the theme of the conference. Abstracts that suit the theme were accepted and author were notified of the acceptance through the system. Abstract submission and review were conducted simultaneously until abstract submission deadline on October 31, 2020. As many as 364 abstracts submitted through system were reviewed. 1 abstract was rejected as not comply with the theme of the conference. Accepted abstracts were allowed to be presented in an oral or live presentation.
  2. Submission of Full Paper  
At the second stage, authors with abstracts declared accepted by the Scientific Advisory Board/Reviewer and will be presented in the conference were requested to submit a complete manuscript according to the IOP Publishing format. The deadline for receipt of a complete manuscript (full paper) was on December 15, 2020.
  3. Revision of Full Paper  
At the third stage, following the conference that took place on January 8-9, 2021, Authors were requested to revise the full paper according to the input and



suggestions from the conference audiences and re-submit the manuscript through the system until January 31, 2021.

4. Peer review of Full Paper

At the Fourth stage, full paper previously submitted through system were reviewed following a single blind peer review process. The papers were sent to the Scientific Advisory Boards that consisted of 47 reviewers according to topic of the papers and reviewer's competency. The reviewers are from international scientists and from the Faculty of Agriculture, and Faculty of Animal Husbandry, Hasanuddin University. Review was conducted on content and format of the paper based on IOP Conference Series. The review process was carried out starting from February to March 7, 2021. At this stage, Reviewer submit a decision for the acceptance of the papers for publication in IOP Conference Series.

5. Revision of Reviewed Full Paper

At the Fifth Stage, Review results were returned to the authors for revision and re-submission online for those accepted for publication (deadline on April 15, 2021). Following the submission of revised manuscript. The submitted papers were then subjected to proofread stage and check for similarity using Turnitin. Papers with similarity check more than 25% were returned to the authors for further revision. Subsequently, papers were prepared for submission to IOP Conference Series.

- **Contact person for queries:**

The chief of the conference: Dr. Asmita Ahmad, ST., M.Si.

Mobile phone: +6281342542792

E-mail: [asmita.ahmad@agri.unhas.ac.id](mailto:asmita.ahmad@agri.unhas.ac.id)

PAPER • OPEN ACCESS

## Agro-morphological variation of pigmented rice landraces from South Sulawesi grown in a temperate glasshouse of New Zealand

To cite this article: A P Hanifa *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **807** 022026

View the [article online](#) for updates and enhancements.



**ECS** **240th ECS Meeting**  
Digital Meeting, Oct 10-14, 2021  
**We are going fully digital!**  
Attendees register for free!  
**REGISTER NOW**

# Agro-morphological variation of pigmented rice landraces from South Sulawesi grown in a temperate glasshouse of New Zealand

A P Hanifa<sup>1,2</sup>, J Millner<sup>1</sup>, C McGill<sup>1</sup> and R Sjahril<sup>3</sup>

<sup>1</sup> School of Agriculture and Environment of Massey University, Palmerston North  
New Zealand

<sup>2</sup> AIAT of South Sulawesi, Jl. Perintis Kemerdekaan KM 17.5 Makassar, Indonesia

<sup>3</sup> Laboratory of Plant Bioscience and Reproduction Biotechnology, Universitas  
Hasanuddin, Makassar, South Sulawesi 90245, Indonesia

Email : rinaldi.sjahril@gmail.com

**Abstract.** The present study is aimed to compare the agro-morphological traits of 15 pigmented rice landraces and two modern rice varieties grown in the same environment. The selected rice were cultivated in a greenhouse of Massey University, New Zealand using a randomised complete block design with five replications. Data collection included agro-morphological observation and measurement. Quantitative data were analysed with analysis of variance, followed with Tukey HSD. Result showed considerable variability in seedling vigour, lodging, plant height, panicle exertion, panicle thresh ability, spikelet fertility, and maturity among the studied pigmented rice landraces. Panicle exertion and spikelet fertility indicate plant cope toward environmental change, which was performed better by many landraces than modern varieties. Variability in morphological characteristics was also recorded among tested varieties. Utilisation of the trait diversity can be useful for future breeding programs, which can lead to this crops' improvement, including pest and disease resistance and grain quality research.

## 1. Introduction

Several research on the diversity of local pigmented rice across Indonesia have been carried out [1,2,3]. In particular, the physicochemical properties of pigmented rice from South Sulawesi have been revealed [4,5], confirming the potential of such varieties for nutraceutical products. Therefore, pigmented rice is considered as speciality rice, which deserves a better premium price in both the domestic and export market.

The pigmented rice landraces possess the immense potential of most valuable genes which can be utilised in the current breeding program to evolve varieties that have not only high yield potential and grain quality of nutritional contents but also resistant to biotic and abiotic stresses. Systematic study and characterisation of such germplasm is not only crucial for utilising the appropriate attribute-based donors but also essential for protecting the exotic rice.

Morphological rice characteristics are less documented in Indonesia [3]; hence the diversity of rice germplasms remain untapped. Plant morphologies are the primary basis for the classification of plants or an interpretation of plant adaptability. Moreover, the link between the morphologies of plants and



productivity has resulted in the concept of a desirable plant type (idiotype) [6]. Hence, there is a necessity to do the process of collection, exploitation and evaluation of untapped germplasms [7].

The study was aimed to compare the variability of 15 pigmented rice landraces and two modern varieties using agro-morphological characters and to provide useful information to facilitate the choice of genitors for further crop improvement.

## 2. Materials and methods

### 2.1 Seed materials

Seeds of 16 traditional pigmented rice varieties were collected from four regencies in South Sulawesi of Indonesia, namely: North Toraja, Tana Toraja, Enrekang, and Jeneponto (table 1). The seed collection is a part of local variety exploration program organised by a team from the Plant Biosciences & Reproduction Biotechnology Laboratory, Faculty of Agriculture, Hasanuddin University). Seed samples were collected during the timeframe of June - November. Local varieties are named according to farmers' information. In this research, two modern variety were used as comparison, *i.e.* Inpari 24 and Ciherang.

**Table 1.** List of pigmented rice landraces used in the present study and their source of origin.

Variety	Grain colour and type	Sub-district/Regency	Altitude (m asl)
Ko'bo	Black	Balusu, North Toraja	900
Lotong	Black	Nanggala, North Toraja	780
Ambo	Black	Balusu, North Toraja	903
Barri Rarang	Red	Bengkelekila, North Toraja	836
Lalodo	Black	Balusu North Toraja	905
Lea	Red	Sangalla, Tana Toraja	791
Lotong Tanduk	Black	Nanggala, North Toraja	908
Maminyak	Red	Baraka, Enrekang	727
Mandoti	Red	Salukanan, Enrekang	789
Balan	Red	Baraka, Enrekang	536
Kamida	Red	Curio, Enrekang	847
Jambu	Red	Banua, Enrekang	607
Le'leng	Black	Turatea, Jeneponto	115
Bakka Eja	Red	Turatea, Jeneponto	115
Punu Eja	Red	Turatea, Jeneponto	153
Inpari24*	Red	Purchased from Indonesian Rice Research Center, West Java	
Ciherang*	White	Purchased from Regional Crop Seed Center of Maros, South Sulawesi	

\* Modern variety (nationally released)

### 2.2 Timing and methods.

The study was conducted in Plant Growth Unit at Massey University, Turitea campus (40°22'40"S 175°36'48"E, elevation: 35 m above sea level) from December 2017 (summer) to July 2018 (winter). Rice plants for the preliminary research were cultivated in a controlled environment. The experiment was set up as a Randomised Block Design with five replicates. Potting mix media with slow released fertiliser was used. Drip irrigation was established in week 4, where each pot was placed in 9.6 L spout

bucket to achieve flooded/saturated condition. Fertilisation was done when leaf nitrogen status was assessed as low based on a Leaf Color Chart (score  $\leq 3$ ) (Rice Knowledge Bank, 2018). Peters® Professional Allrounder (N-P-K : 20-20-20+TE) at 1g/L was used as fertiliser, incorporating with drip irrigation (5 L per day).

### 2.3 Data collection and analysis.

The accessions were characterised according to Standard Evaluation System for Rice [8]. A total of seven agronomic traits data (table 2) and twelve of morphological characteristics (table 3) were collected. Simple statistical parameters *i.e.* mean and standard error were determined for the quantitative traits using statistical package SPSS 15.0. The data after compiling was statistically analysed using SPSS 15.0 and Tukey post hoc test (Honest Significance Difference) test was applied to test the significance of varietal differences. Pearson correlation was also used to investigate any association between parameters.

**Table 2.** Descriptions and scales/codes of agronomic traits to evaluate 15 pigmented rice landraces from South Sulawesi [8].

Descriptors	Observed phenotypic classes	Evaluation phase
Seedling vigour (Vg) /vegetative	1-extra weak;9-very weak;3-vigorous;5-normal;7-vigorous	Seedling vigour:2 Vegetative vigour:3
Tillering ability (Ti)	1-very high (>25);3-good(20-25);5-medium(10-19);7-low(5-9);9-very low (<5)	At growth stage 5
Plant height	1-semidwarf (lowland:<110cm;upland:<90cm);2-intermediate (lowland:110-130cm;upland(90-125cm);9-tall(lowland>130cm;upland:>125cm)	At growth stage 7-9
Panicle exertion	1-enclosed;3-partly exerted (panicle base is slightly beneath the collar of flag leaf blade);5-just exerted (panicle base coincides with the collar of the flag leaf blade);7-moderately well exerted (panicle base is above the collar of the flag leaf blade);9-well exerted(panicle base appears well above the collar of the flag leaf blade)	At growth stage 7-9 (near maturity)
Panicle threshability	1-difficult (less than 1%);3-moderately difficult (1-5%);intermediate (6-15%);7-loose (26-50%);9-easy(51-100%)	At growth stage 9
Spikelet fertility	1-highly fertile(>90%);3-fertile(75-89%);partly sterile(50-74%);highly sterile(<50% to trace);9-0%	At growth stage 9
Maturity	Number of days from seeding to grain ripening (85% of grains on panicle are mature)	At growth stage 9

**Table 3.** Morphological character descriptors used to evaluate 15 pigmented rice landraces from South Sulawesi.

Descriptions	Observed phenotypic classes	Evaluation phase
Leaf length	Actual measurement in cm of the widest portion of the leaf blade just below the flag leaf	At growth stage 6
Leaf blade colour	1-light green;2-green;3-dark green;4-purple tips;5-purple margins;6:purple blotch;7:purple	At growth stage 4-6
Ligule length	Enter actual measurement of ligules measured in mm from the base of collar to the tip. Sample size 5	At growth stage 4-5
Ligule colour	0-absent;1:white;2:purple lines;3:purple	At growth stage 4-5
Ligule shape	0-absent;1-acute to acuminate;2-cleft;3-truncate	At growth stage 3-4
Auricle colour	0-absent;1-light green;2-purple	At growth stage 4-5
Culm internode colour	1-green;2-light gold;3-purple lines;4-purple	At growth stage 7-9
Panicle length	Enter actual measurement in cm from panicle base to tip	At growth stage 8
Awning	0-absent;1-short and partly awned;5-short and fully awned;7:long and partly awned;9-long and fully awned	At growth stage: 7-9
Awn colour	0-awnless;1-straw;2-gold;3-brown;4-red;5-purple;6-black	At growth stage: 6
Grain width	Enter the actual measurement of width in mm as the distance across the fertile lemma and the palea in the widest point	At growth stage:9
Grain length/width ratio	Ratio of length in mm as the distance from the base of the lowermost sterile lemma to the tip (apiculus) of the fertile lemma/palea per width	At growth stage:9

### 3. Results and discussion

#### 3.1 Agronomic traits

The assessed pigmented rice landraces showed considerable variability for the studied agromorphological characteristics. Seven agronomic traits comparison among studied varieties are presented in table 4.

*3.1.1 Maturity.* Number of days to achieve the maturity of rice seeds were in the range from 157 cm in Jambu to 215 cm in each variety of Lallodo and Mandoti. Modern varieties reached their maturity longer than expected, indicating a response to the environmental difference from their original habitat. Based on observation, plants with early maturity relatively have intermediate to semi-dwarf height, such as Balan, Bakka Eja, Inpari 24 and Ciherang. However, Jambu has early maturity but fall into the tall group.

Generally, the maturity days reached in this trial were longer than reported maturity in references [9,10] which under five months if grown in Toraja regions for some varieties like

Lallodo, Ambo, Mandoti and Lotong. Environmental factors such as light reception and temperature are some of the key factors.

*3.1.2 Plant height.* Plant height can be seen in three classes, namely, intermediate (Balan), semi-dwarf (Bakka Eja, Inpari, Ciherang) and tall type for the remaining tested varieties. 5.8% intermediate height, 17.6% semi-dwarf, 76.5% tall in height. Traditional rice varieties typically have tall stature whereas modern varieties/early maturing types that tend to be semi-dwarf. In general, most tested varieties grew higher than their common height in their origin of habitat. On the other hand, decreased height, earlier maturity, and longer panicles were recorded from upland rice of South Sulawesi under lowland cultivation [11]. Semi-dwarf rice plant is a primary target for improving the lodging resistance [12], but the drawback is, the canopy for photosynthesis and number of biomass are limited, and then decrease the grain yield [13]

*3.1.3 Panicle exertion.* There are four groups of panicle exertion, which are dominated by well exerted (58.8%), whereas partly exerted was shown in two modern varieties. Most traditional varieties such as Kobo, Lotong, Ambo, Ra'rang, Lotong Tanduk, Maminyak, Kamida, Jambu, Leleng and Punu Eja showed well-exserted panicles. In contrast, modern varieties have partly exerted panicles.

Panicle exertion is one of the important agronomic traits since it can be an indication of genetic defect caused by environmental or pest -disease attack [14]. Incomplete panicle exertion indicates cold injury at the reproductive stage, which lead to damaged grain yield as disease incidence raises [15]. The long and well-exserted panicles are the typical characters of upland rice panicles, as supported by [16], that upland traditional rice are adapted to harvesting and storage by panicles which are long and well exerted.

*3.1.4 Threshability.* Six traditional varieties and two modern varieties were observed to be in the easy category of threshability. Grain characteristics of the ease threshability group are awnless. In contrast, awned grains have intermediate to moderately difficult of threshability.

The result confirmed the threshability of awned genotypes is more difficult than awnless. Some farmers in Toraja prefer to have moderate-intermediate threshability of panicles to minimise losses during post-harvest, as they transport their stacks of panicles manually and store the panicles with the stalks, without shattering [17].

*3.1.5 Seedling vigour.* Seedling vegetative vigour fell into three groups: weak (Kobo, Lotong, Lotong Tanduk), normal (Lea, Kamida) and the remainder as vigorous (table 4). In general, most tested varieties, including modern varieties had acceptable seedling vigour.

*3.1.6 Spikelet fertility.* Our result only have two groups of spikelet fertility: partly sterile and highly sterile. Some traditional varieties perform better than the modern ones. Unexpected sterility of panicles may resulted from microclimate discrepancy in pot trial compared to the origin. Moreover there was disruption from the heat fan which caused dried panicle during anthesis.

The insufficient spikelet fertility in this research might also resulted from cold stress and low solar radiation during reproductive phase. According to [18], during reproductive period, low temperature-induced spikelet sterility, could be associated with the physiological response mechanisms in the pollen, anther, and stigma.

**Table 4.** Agronomic traits of pigmented rice landrace of South Sulawesi and two modern rice varieties for comparison

Variety	Maturity (days)	Plant height (cm)	Panicle exertion	Threshability	Seedling vegetative vigour	Spikelet fertility	Tillering ability
Kobo	195	Tall	Well exerted	Moderately difficult	Weak	Partly sterile	Medium
Lotong	202	Tall	Well exerted	Moderately difficult	Weak	Highly sterile	Medium Very high
Ambo	190	Tall	Well exerted	Intermediate	Vigorous	Partly sterile	Good Very high
Ra'rang	195	Tall	Well exerted	Easy	Vigorous	Highly sterile	Good Very high
Lalodo	215	Tall	Moderately well	Intermediate	Vigorous	Highly sterile	Good Very high
Lea	202	Tall	Moderately well	Easy	Normal	Highly sterile	Good Very high
L. Tanduk	195	Tall	Well exerted	Intermediate	Weak	Highly sterile	Good Very high
Maminyak	190	Tall	Well exerted	Moderately difficult	Vigorous	Partly sterile	Good Very high
Mandoti	215	Tall	Moderately well	Moderately difficult	Vigorous	Highly sterile	Very high
Balan	170	Intermediate	Just exerted	Easy	Vigorous	Highly sterile	Very high
Kamida	198	Tall	Well exerted	Easy	Normal	Highly sterile	Very high
Jambu	157	Tall	Well exerted	Intermediate	Vigorous	Highly sterile	Very high
Le'leng	198	Tall	Well exerted	Intermediate	Vigorous	Partly sterile	Very high
Bakka Eja	160	Semidwarf	Just exerted	Easy	Vigorous	Highly sterile	Very high
Punu Eja	198	Tall	Well exerted	Easy	Vigorous	Partly sterile	Very high
Inpari 24	160	Semidwarf	Partly exerted	Easy	Vigorous	Highly sterile	Very high
Ciherang	160	Semidwarf	Partly exerted	Easy	Vigorous	Highly sterile	Very high

*3.1.7 Tillering ability.* The tillering ability of all varieties in this study appears to be medium (10-19 tillers per plant) to very high (more than 25 tillers/plant). The ideal planting space is 25 cm to allow the appropriate growth of tillers and minimise disease in humid condition. Most of the varieties showed a very high tillering ability while few varieties namely Maminyak, Lea and Ra'rang had good tillering ability and only Kobo as well as Lotong had medium tillering ability.

Similarly [10] reported significant difference in traits such as maturity, height, tiller number, seed weight, and productivity among 29 rice landraces from Toraja region. A few of them have coloured grain, such as Pare Lea, Pare Lallodo, and Pare Ambo. However, the tiller number reported by [9] was 3-4 times higher (17-30) than tiller number reported in [10]. These two different studies were undertaken in different locations but in the Toraja region hence variability in result is noted.

### 3.2 Morphological characteristics

*3.2.1 Quantitative measurements.* Ra'rang has the longest ligule length among the landraces, which similar to modern Inpari 24. Jambu has the shortest ligule. correlation coefficient of ligule length and grain length/width ratio was 0.115 Which interpreted as little if any correlation [19]. Conversely, a little negative correlations were found between ligule length and leaf length, panicle length, and grain width.

The most fundamental factor in the diversity of grain colours is the genetic factor that expresses the grain colour of the variety. Result of bioactive content assessment of pigmented rice landrace of South Sulawesi also confirmed the high anthocyanin content of black rice, and considerable amount of phenolic content in red rice [5].

Maminyak has the longest leaf (78.80 cm), whilst Ciherang, Balan, and Inpari have the shortest leaf length. has the shortest leaf length (45 cm;48 cm and 51 cm respectively). Moderate positive correlation ( $r=0.577$   $N=80$   $p<0.01$ ) was found between leaf length and plant height.

The longest panicle length was found in Lotong (36.5 cm), which significantly differ from Rarang, Lallodo, Lea, Lotong Tanduk, Mandoti, Balan, Kamida, Leleng, Bakka Eja, Punu Eja and two modern varieties (Ciherang and Inpari 24) (Table 5). As expected, the shortest panicle was found in Ciherang. Panicle length has low positive correlation with leaf length ( $r=0.488$   $N=80$   $p<0.001$ ) (Table 5). Multiple genes controlled panicle length, a typical quantitative trait, and may be affected by environmental conditions significantly. In addition, breeding varieties with large panicle and the ideal plant architecture traits are significant for increasing the grain yield of rice [20].

**Table 5.** Measurement of morphological characteristics of pigmented rice landrace of South Sulawesi and two modern rice varieties for comparison.

Variety	Ligule length	Leaf length	Panicle length	Grain's length/width ratio	Grain width
Kobo	14.20±0.66edc	63.00±2.19gfedcb	32.00±2.30ba	2.78±0.07gf	2.90±0.13cba
Lotong	17.20±1.24cba	76.10±3.26cba	36.50±0.87a	3.14±0.08fedc	3.00±0.05ba
Ambo	13.80±1.24edc	63.40±2.71fedcb	29.60±0.53cdba	2.64±0.04g	3.00±0.06ba
Ra'rang	20.40±1.63a	68.40±3.16edcba	21.80±2.89hgfed	2.92±0.12gfed	2.84±0.08cba
Lallodo	14.80±0.50edcb	72.60±1.89dcba	21.80±1.98hgfed	3.32±0.11dc	2.72±0.09dcba
Lea	13.00±1.38edc	70.40±2.25dcba	23.40±1.78hgfedc	2.84±0.05gfe	2.86±0.07cba

L. Tanduk	11.50±0.50ed	63.00±2.00gfedcb	23.00±2.00hgfedc	2.77±0.05	3.25±0.05
Maminyak	13.60±0.93edc	78.80±2.96a	29.20±1.59edcba	2.89±0.15gfed	2.76±0.08dcba
Mandoti	12.00±0.71edc	76.00±3.41cba	25.40±0.98gfedcb	3.11±0.09fedc	2.86±0.02cba
Balan	15.60±0.51dcba	48.20±2.44hg	20.20±1.36hgf	3.19±0.05fedc	2.66±0.05dc
Kamida	11.00±0.45ed	59.20±0.86hgfed	27.80±1.66fedcb	3.09±0.10gfedc	2.70±0.03dcb
Jambu	10.00±0.55e	77.40±2.58ba	30.04±1.16cba	3.25±0.06edc	2.82±0.04cba
Le'leng	12.60±0.81edc	59.40±2.01hgfed	25.20±0.73gfedcb	2.91±0.02gfed	3.04±0.06a
Bakka Eja	12.60±0.81edc	53.90±1.35hgfe	21.20±1.01hgfe	3.85±0.08ba	2.48±0.06ed
Punu Eja	12.00±0.95edc	62.00±4.12gfedc	22.40±1.69hgfedc	2.99±0.12gfed	3.00±0.10ba
Inpari 24	20.20±1.20ba	51.00±1.87hgf	18.93±1.22hg	4.13±0.08a	2.24±0.05e
Ciherang	14.00±1.10edc	45.40±5.57h	15.40±0.51h	3.52±0.10cb	2.18±0.04e

Values followed the same letter are not different ( $P < 0.05$ ).

**3.2.2 Qualitative observations.** Morphological characteristics of tested varieties observed under glass house conditions are presented in table 6. An important observed feature is the extent of pigmentation found in various plant parts including auricles, ligules, awns, culm internodes, and leaf blades among different varieties. While light green was the typical colour of auricles in most varieties tested in this experiment, only Ambo and Lotong Tanduk have the purple colour in the auricles. Awned seeds found in 52.9% of tested varieties. One of Wild rice species' distinct feature in morphology is a long awn [21]. Shorter awn or awnless grain in domesticated wheats facilitate harvesting, handling and storage [22].

The awn color of awned seed types ranged from straw (Kobo, Lotong, Lallodo, Maminyak, Mandoti, Jambu, Le'leng) to red (Ambo and Lotong Tanduk). Each variety of Ambo as well as Lotong Tanduk also shared similarity in culm internode color (purple), ligule color (purple), and leaf blade colour (purple in the margin), along with kernel colour of black. The culm internode colour of Ambo, L. Tanduk, Kamida were purple, while Punu Eja has purple line for their culm internode colour. Most varieties (88.2%) have white ligule colour while the ligule shape was cleft in all varieties. Plants with purple ligule colour (Ambo and Lotong Tanduk) also have purple margin leaf blade colour, and the remaining varieties are in green colour of leaf blade (table 6). Compared with black rice found in Central Java, namely Matesih, Klaten and Cempo, all of them have white ligule in cleft shape, green leaf blade color and white auricle [23].

**Table 6.** Morphological characteristics observation of pigmented rice landrace of South Sulawesi and two modern rice varieties for comparison

Variety	Auricle color	Awn color	Awning	Culm internode color	Ligule color	Ligule shape	Leaf blade color
Kobo	Light green	Straw	Long and Well	Green	White	cleft	Green
Lotong	Light green	Straw	Long and Well	Green	White	cleft	Green
Ambo	Purple	Red	Long and Well	Purple	Purple	cleft	Purple margin
Rarang	Light green	-	Awnless	Green	White	cleft	Green
Lallodo	Light green	Straw	Short and partly	Green	White	cleft	Green
Lea	Light green	-	Awnless	Green	White	cleft	Green
Lotong Tanduk	Purple	Red	Short and partly	Purple	Purple	cleft	Purple margin
Maminyak	Light green	Straw	Long and Well	Green	White	cleft	Green
Mandoti	Light green	Straw	Long and Well	Green	White	cleft	Green
Balan	Light green	-	Awnless	Green	White	cleft	Green
Kamida	Light green	-	Awnless	Purple	White	cleft	Green
Jambu	Light green	Straw	Long and Well	Green	White	cleft	Green
Le'leng	Light green	Straw	Long and Well	Green	White	cleft	Green
Bakka Eja	Light green	-	Awnless	Green	White	cleft	Green
Punu Eja	Light green	-	Awnless	Purple line	White	cleft	Green
Inpari 24	Light green	-	Awnless	Green	White	cleft	Green
Ciherang	Light green	-	Awnless	Green	White	cleft	Green

#### 4. Conclusions

The discovery of genetic differences plays a key role in the development of varietal conditions that may contribute to food security. In this study, variability is seen in some agronomic traits and morphological characteristics of pigmented rice landraces.

Some pigmented rice landraces performed better than modern varieties under glasshouse environment of temperate climate. The better performance of pigmented rice landrace indicate the capability in coping towards different environment from their origin of habitat.

Pigmented rice plant idotype can be engineered from selecting desired agronomic traits, ideal architecture, as well as superior physicochemical properties. Farmers perspective and their requirement should be taken into consideration as well, since farmers are the key factor for preserving the germplasm in situ by retaining the varieties intergeneration.

#### References

- [1] Anggraini T, Novelina, Limber U, and Amalia R 2015 Antioxidant activities of some red, black and white rice cultivar from West Sumatra, Indonesia. *Pakistan Journal of Nutrition* **14** 112-7.
- [2] Kristantini T, Basunanda P, Murti R H, & Supriyanta S 2012 Morphological of genetic relationships among black rice landraces from Yogyakarta and surrounding areas. *ARPN Journal of Agricultural and Biological Science* **7** 982-9.
- [3] Shinta S, Indriyani S, and Arisoesilaningsih E 2014 Morphological variation of six pigmented rice local varieties grown in organic rice field at Sengguruh Village, Kepanjen Subdistrict, Malang District. *Journal of Tropical Life Science*, **4** 149-160. Retrieved from <https://jtolis.uib.ac.id/index.php/jtolis/article/view/171>
- [4] Murdifin M, Pakki E, Rahim A, Syaiful S A, Ismail, Evary, Y M, and Bahar M A 2015 Physicochemical properties of Indonesian pigmented rice (*Oryza sativa* Linn.) varieties from South Sulawesi. *Asian Journal of Plant Sciences* **14** 59-65. 10.3923/ajps.2015.59.65
- [5] Hanifa A, Millner J, McGill C, and Sjahril R 2020 Total anthocyanin, flavonoid and phenolic content of pigmented rice landraces from South Sulawesi. *IOP Conference Series: Earth and Environmental Science*,
- [6] Loomis R S and Williams W A 1969 Productivity and the morphology of crop stands patterns with leaves. In *Physiological Aspects of Crop Yield*. ASA-CSSA. Wisconsin.
- [7] Parikh M, Motiramani N K, Rastogi N K and Sharma B 2012 Agro-morphological characterization and assessment of variability in aromatic rice germplasm *Bangladesh Journal of Agricultural Research* **37** 1-8.
- [8] International Rice Research Institute 2013 *Standard Evaluation System for Rice* 5<sup>th</sup> Edition. IRRI. Manila, Philippines.
- [9] Limbongan Y and Djufry F 2015 Characterization and Observation of Five Local Rice Accessions of Toraja Plateau, South Sulawesi *Bul. Plasma Nutrafah* **21** 61-70
- [10] Sahardi, Herniwati, and Djufry F 2015 Keragaman karakter morfologis plasma nutrafah lokal Toraja. Paper presented at the *Prosiding Seminar Nasional Sumber Daya Genetik Pertanian*, Bogor.
- [11] Sahardi and Hanifa, AP 2015 Performance of upland rice under lowland growing condition in Gowa Regency, South Sulawesi. *Prosiding Seminar Nasional Pengembangan Sumber Daya Genetik Pertanian: Potensi Sumber Daya Genetik Spesifik Lokasi Mendukung Ketahanan dan Swasembada Pangan Nasional*. Bogor: 27<sup>th</sup> May 2015. Balai Besar Pengkajian dan Pengembangan Teknologi Pertanian, Badan Penelitian dan Pengembangan Pertanian,
- [12] Khush G S 1999 *Green revolution: Preparing for the 21st century Genome* **42** 646-655 PMID: 10464789
- [13] Kuroda E, Ookawa T, Ishihara K 1989 Analysis on difference of dry matter production between rice cultivars with plant height in relation to gas diffusion inside stands *Jap J Crop Sci* **58** 374-382 DOI:10.1626/JCS.58.374
- [14] International Rice Research Institute 2002 *Standard Evaluation System for Rice* IRRI <http://www.knowledgebank.irri.org/images/docs/rice-standard-evaluation-system.pdf>

- [15] Pereira da Cruz R, S C Milach and L C Federizzi 2008 Inheritance of panicle exertion in rice *Sci. agric. (Piracicaba, Braz.)* **65** 502-507.
- [16] Chandler Jr R F. 1979 *Rice in the tropics*. A guide to the development of national programs. Westview Press Boulder, Colorado.
- [17] Hanifa, A P 2018 Evaluation of pigmented rice landraces from Toraja, Sulawesi. Confirmation report. School of Agriculture and Environment. Massey University. Unpublished
- [18] Zeng Y, Zhang Y, Xiang J, Uphof N T, Pan X, and Zhu D 2017 Effects of low temperature stress on spikelet-related parameters during anthesis in Indica–Japonica hybrid rice *Frontiers in Plant Science*, **8** 10.3389/fpls.2017.01350
- [19] Asuero, G A Sayago & A G González 2006 *The correlation coefficient: An overview, critical reviews in analytical chemistry* **36** 41-59, DOI: 10.1080/10408340500526766
- [20] Yao X Y, Li Q, Liu J, Jiang S -K, Yang S -L, Wang, J -Y (2015). Dissection of QTLs for plant height and panicle length traits in rice under different environment. *Sci. Agr. Sin.* **48** 407–414 DOI: 10.3864/j.issn.05781752.2015.03.01
- [21] Furuta T, Komeda N, Asano K, Uehara K, Gamuyao R, Angeles-Shim R B, Ashikari M 2015 Convergent loss of awn in two cultivated rice species *Oryza sativa* and *Oryza glaberrima* is caused by mutations in different loci. *G3 (Bethesda, Md.)* **5** 2267-2274. 10.1534/g3.115.020834
- [22] Mach J 2015 Domesticated versus wild rice? Bring it awn! *The Plant cell* **27** 1818. <https://doi.org/10.1105/tpc.15.00504>
- [23] Purwanto E and Meidini A 2018 *Morphology, production, and chemical content performance of black rice Matesih accession with several comparisons*. Paper presented at the IOP Conference Series: Earth and Environmental Science.

PAPER • OPEN ACCESS

## Preface

To cite this article: 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **807** 011001

View the [article online](#) for updates and enhancements.

## You may also like

- [Smartphone-based GNSS Method for Covid-19 Information on Android \(Case Study: Surabaya City\)](#)  
A H Akbar and M N Cahyadi
- [The Effect of Climate on the Outbreak of Covid-19: A Review](#)  
Candra Kirana, Samsul, Hesty Yariska Hapsir et al.
- [Impacts of the Covid-19 pandemic on traditional oil mining at Wonocolo Village Kedewan Sub-District Bojonegoro Regency East Java](#)  
L A Rahmawati, N Afiati and T T Putranto

## PREFACE

It is a great honour for us to be the host of the 3rd International Conference on Food Security and Sustainable Agriculture in the Tropics (FSSAT) on January 8th – 9th, 2021 at Agriculture Faculty of Hasanuddin University, Makassar City of South Sulawesi Province, Indonesia. This conference is a series of international scientific seminars held by the Faculty of Agriculture, Universitas Hasanuddin, Makassar, Indonesia, which has been held annually since 2017.

This conference was held as a forum for exchanging information on productivist paradigm in Indonesian food policies to feed 260 million people targets the self-sufficiency of rice, sago, corn, soybean, and meat production, including the infrastructure development such as dams, irrigation channels building, and the expansion of paddy and sago fields in the Eastern Indonesia. The unintended outcomes of these efforts are the marginalization of the local food system and dispossession of the local communities from their land and food culture. As a consequence, the human-environmental relationship has changed dramatically posing calamitous disaster and slowly disappearing foodways through local wisdom in everyday life of the local community. It also brings health issues due to the excessive use of chemical agricultural inputs in production, and the increase of degenerative disease and obesity due the unhealthy practices in the consumption level.

The International Conference on Food Security and Sustainable Agriculture in the Tropics will be held every two years, the first in 2017, the second in 2019, and the third in 2021. The covid-19 conditions that have hit the whole world have resulted in restrictions on activities to reduce the spread of Covid-19 sufferers so that the conference model from the form of a live conference has turned into a virtual conference.

The condition of the Covid-19 pandemic has raised concerns about food security throughout the world, including Indonesia, through restrictions on community activities. Several things have implications for the existence of Covid-19, including the occurrence of a very significant supply chain disruption caused by a reduction in capacity to produce food, closing transportation routes so that it will slow down the distribution of food from producers to consumers. The inability of people to consume enough healthy and nutritious food will reduce the immune system and increase disease risk. Thus ensuring an affordable food supply chain is essential during the Covid-19 outbreak. To solve this problem globally, cooperation is needed that involves scientists, academics, practitioners, professionals, governments, entrepreneurs, and communities across countries. To build a network between stakeholders, an international conference held by the Hasanuddin University Faculty of Agriculture in 2021 is important as a forum for building connectivity between researchers and/or research institutions. At this conference, a multi-disciplinary approach based on the knowledge and experience of scientists, researchers, practitioners, and policymakers will be brought together.

If the meeting cannot be held virtually, then solutions in dealing with the food crisis during the Covid-19 pandemic will not find a way out and add to the length of the crisis period chain.

This conference presents keynote speakers from 5 countries, namely Dr. Ir. Bayu Krisnamurthi, M.Si. from IPB University, Indonesia, Prof. Dr. Ir. Sumbangan Baja, M.Phil from Universitas Hasanuddin, Indonesia, Prof. Takuya Sugahara from Ehime University, Japan, Dr. Ravindra C Joshi from CABI South-East Asia, Dr. Lau Wei Hong from University

Putra Malaysia, Mr. Moh, Guo-Jhong from ICDF Taiwan. First plenary session and discussion with 2 speakers for 1 hour and the second plenary and discussion with 3 speakers for 2 hours. For participant presentations (oral presentation and video presentation) it was held in 2 sessions, each session was divided into 10 rooms. The number of participants per room was 11-14 participants with an allocation of presentations for 7 minutes/participants conducted in parallel as many as 7 parallel sessions. Discussions session were held in parallel in each parallel session with an allocation of 3 minutes per participant. Virtual FSSAT 3 implementation using the zoom meeting application by the Faculty of Agriculture, Universitas Hasanuddin has which is obtained by subscription.

The seminar participants were attended by lecturers and students from various universities as well as researchers from various research institutions both from the ministry of agriculture and from Agricultural Technology Research Center. The institution of the participants came from Agricultural Technology Research Center, some polytechnic, institute and University, Indonesian Center for Estate Crops Research and Development, International Coconut Community, Center for Agricultural Technology Research and Development, PT Vale Indonesia Tbk, Sorowako. Some from overseas universities: University of New England, Armidale, Australia, and School of Agriculture and Environment of Massey University, Palmerston North New Zealand. Overall participant numbers around 1.034 participants.

There were no significant technical obstacles, speakers and participants from overseas also presented their papers fluently. Apart from sharing zoom rooms, we also created a social media group (WhatsApp) for each class to facilitate communication between participants and organizers.

All published articles are subject to a rigorous selection process and are peer-reviewed by international and national reviewers. On this occasion, we would like to thank all the authors and the team who have actively participated in the 3<sup>rd</sup> FSSAT conference, expert reviewers who are very responsive to the technical program committee, experienced senior publication chair, Publication Management Center (PMC) of Universitas Hasanuddin, authors and scholars who are enthusiastic in research. Sincerely thank you for your warm support and look forward to working with you in the next future.

Warm Regards,

Dr. Asmita Ahmad, ST., MSi.

Conference Chairman

Faculty of Agriculture, Universitas Hasanuddin

**COMMITTEES**

Steering Committee	Dean, Faculty of Agricultural, Hasanuddin University Dean, Faculty of Forestry, Hasanuddin University Dean, Faculty of Animal Science, Hasanuddin University Dean, Faculty of Marine Science and Fishery, Hasanuddin University
Organizing Committee	Chair: Dr. Asmita Ahmad, ST., MSi. Vice Chair: Andi Dirpan, STP., MSi., PhD. Secretary: Dr. Suleha Thamrin, SP., MSi. Treasurer: St. Hasnah A. Rahman, SP.
Scientific Advisory Board	Prof. Dr.sc.agr.Ir. Baharuddin (Indonesia) Prof. Takuya Sugahara (Japan) Dr. Ravindra C Joshi (Philippines) Dr. Lau Wei Hong (Malaysia) Mr. Guo-Jhong Moh (Taiwan) Dr. Ir. Bayu Krisnamurthi, M.Si (Indonesia) Prof. Ir. Sumbangan Baja, M.Phil, PhD(Indonesia)  Dr. Muh. Hatta Jamil, SP., M.Si. (Indonesia) Dr.rer.Nat. Zainal, STP. M.Food.Tech (Indonesia) Dr. Ir. Novaty E. Dunga, MP (Indonesia) Prof. Dr. Ir. Nurriaty Agus, MS. (Indonesia) Dr. Ir. Rahmadani, M.Si. (Indonesia) Prof. Dr. Ir. Tutik Kuswinanti, M.Sc. (Indonesia) Prof. Dr. Ir. Meta Mahendrasatta, MS. (Indonesia) Dr. Ir. Amir Yassi, MS. (Indonesia) Dr. Rismaneswati, SP., MP. (Indonesia) Dr. Nixia Tenriawaru, SP., M.Si. (Indonesia) Dr. Iqbal, STP. M.Si. (Indonesia) Ir. Rinaldi Sjahril, M.Agr., PhD. (Indonesia) Dr. Ir. Rusnadi Padjung, M.Sc. (Indonesia) Muhammad Arsyad, SP., M.Si., PhD. (Indonesia) Dr. Ir. Andi Nasruddin, M.Sc. (Indonesia) Prof. Dr. Ir. Sikstus Gusli, M.Sc. (Indonesia) Prof. Ade Rosmana, M.Sc. (Indonesia) Prof. Dr. Ir. Christianto Lopulisa, M.Sc. (Indonesia) Prof, Dr. Ir. Salengke, M.Sc. (Indonesia) Prof. Dr. Ir. Yunus Musa, M.Sc. (Indonesia) Prof. Dr. Ir. M. Saleh Ali (Indonesia) Prof. Dr. Ir. Musrizal Muin M,Sc(Indonesia)
Secretariat	Sartika Laban, SP., MP., Ph.D Dr. Abdul Aziz, S.TP., M.Si. Nirmala Juita, SP., M.Sc. Hamdayanti, SP., M.Si. Ahmad Fauzan Adzima, SP., M.Sc. Magfirah, SP. Nilarisa Meganita, S.TP

Paper, Poster, and Prosiding	Muhammad Junaid, SP., MP, Ph.D Dr. Ir. Asmiaty Sahur, MP. Rasyida Bakri, SP., M.Sc Muspilah Djalal, STP., M.Sc. Dhini Amriani Mutia Mey Nindi Zulkifly
Publicity and Publishing	Ifayanti, SP., MP., Ph.D Rusli M. Rukka, SP., M.Si. Rahmansyah Dermawan, SP. M.Si Andi Rahmayanti, S.TP. M.Si Iswar Fajar Sultan, SP. Andi Junila, SP. Rizza Nurul Aprillia.
Website	Muh. Tahir Sapsal, STP., MSi. Samsuar, STP., M.Si. Dr. Adiansyah, S.TP, M.Si Yudiono, SE
Registration and Certificate	Dr. Muh. Asfar, S.TP., M.Si Dr. Ir. Katriani Mantja, MP Husnul Mubarak, S.TP., M.Si Ir. Nurlina Kasim, M.Si Cri Wahyuni Brahmiyanti, SP., M.Si Tigin Dariarti, SP
Finance and Corporate Partnership	Dr. Letty Fudjaja, SP., M.Si Dyah Yumeina R. Datu, S.TP., M.Agri.,PhD. Pipi Diansari, SE., M.Si., PhD. Dr. Ir. Heliawaty Ch. Lakare, M.Si Dr. Ir. Abd. Haris Bahrin, M.Si Ikmalul Akmal, S.Sos
Keynote/Invited Speaker team	Hari Iswoyo, SP., MA., PhD Dr. Olly Sunny Hutabarat, S.TP. Dr. Ir. Vien Sartika Dewi, M.Si Dr. Andi Nur Faidah, R, S.TP., M.Si NiMadeViantikaSuliandri,SP.,M.Ag
Event, Exhibition and Technical Program	Februardi Bastian, STP., M.Si., Ph.D. Asman, SP., MP. Abd. Mollah, SP., M.Si. Achmad Amiruddin, SP.,M.Si Nuniek Widiyani, SP., MP Abd. Ishak, S.Sos Kurniati Latief

**EDITORS & REVIEWERS**

Ir. Muhammad Arsyad, S.P., M.Si, Ph.D

Prof. Dr. Ir. Didi Rukmana, M.S

Prof. Dr. Ir. Muslim Salam, M.Ec

Dr. A. Nixia Tenriawaru, S.P., M.Si.

Achmad Amiruddin, S.P., M.Si

Muhammad Khaliqi, S.P., M.Si

Andi Azrarul Amri, S.P., M.Si

Dwi Ahrisa Putri, S.P., M.Si

Andi Tenri Darhyati, SP, M.Si

Ikawati, S.T.P., M.Si

Ayu Wulandary, S.P., M.Si

Ayu Anisa Amir, S.P., M.Si

Ary Bakhtiar., SP., M.Si

Dian Utami Zainuddin, S.Si., M.Si.

Departemen Ilmu Tanah

Dr. Rismaneswati, S.P., M.P

Dr. Asmita Ahmad, S.T., M.Si

Sartika Laban, S.P., M.P., Ph.D

James Nana Ofori, B.Sc., M.Sc (Ghana)

Supriadi, S.P., M.Si (Japan)

Nirmala Juita, S.P., M.Si

Indryani Bali, S.P., M.Sc

Prof. Dr. Ir. Yunus Musa, M.S

Ir. Rinaldi Sjahril, M.Agr., Ph.D

Ir. Hari Iswoyo, S.P., M.A., Ph.D

Dr. Ir. Muh. Riadi, M.P

Dr. Ir. Rusnadi Padjung, M.Sc

Ifayanti Ridwan Saleh, S.P., M.P., Ph.D

Dr. Ir. Asmiaty Sahur, M.P

Rahmansyah Dermawan, S.P., M.Si

Nuniek Widiyani, S.P., M.Si

Prof. Dr. Ir. Ade Rosmana, M.Sc

Prof. Dr. Ir. Andi Nasruddin, M.Sc

Prof. Dr. Ir. Tutik Kuswinanti, M.Sc  
Muhammad Junaid. S.P.,M.P.,Ph.D  
Dr. Sulaeha Thamrin, S.P.,M.Si  
Hamdayanty S.P.,M.Si  
Prof. Dr. Ir. Meta Mahendradatta  
Andi Dirpan, S.TP.,M.Si.,Ph.D  
Dr. Andi Nur Faidah Rahman., STP., MSi  
JamyangTashi Wangdi, M.Sc (Bhutan)  
Nur Alim Bahmid, S.TP.,M.Si (Netherland)  
Muspilah Djalal, S.TP.,M.Sc  
Dr. Iqbal, S.TP.,M.Si  
Dr. Haerani, S.TP.,M.Eng.Sc  
Dr.rer.nat. Olly S. Hutabarat, S.TP.,M.Si  
Samsuar, S.TP.,M.Si  
Husnul Mubarak, S.TP.,M.Si  
Dr. Kasmiyati Kasim, SPt, MP

PAPER • OPEN ACCESS

## Agro-morphological variation of pigmented rice landraces from South Sulawesi grown in a temperate glasshouse of New Zealand

To cite this article: A P Hanifa *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **807** 022026

View the [article online](#) for updates and enhancements.



**ECS** **240th ECS Meeting**  
Digital Meeting, Oct 10-14, 2021  
**We are going fully digital!**  
Attendees register for free!  
**REGISTER NOW**

# Agro-morphological variation of pigmented rice landraces from South Sulawesi grown in a temperate glasshouse of New Zealand

A P Hanifa<sup>1,2</sup>, J Millner<sup>1</sup>, C McGill<sup>1</sup> and R Sjahril<sup>3</sup>

<sup>1</sup> School of Agriculture and Environment of Massey University, Palmerston North New Zealand

<sup>2</sup> AIAT of South Sulawesi, Jl. Perintis Kemerdekaan KM 17.5 Makassar, Indonesia

<sup>3</sup> Laboratory of Plant Bioscience and Reproduction Biotechnology, Universitas Hasanuddin, Makassar, South Sulawesi 90245, Indonesia

Email : rinaldi.sjahril@gmail.com

**Abstract.** The present study is aimed to compare the agro-morphological traits of 15 pigmented rice landraces and two modern rice varieties grown in the same environment. The selected rice were cultivated in a greenhouse of Massey University, New Zealand using a randomised complete block design with five replications. Data collection included agro-morphological observation and measurement. Quantitative data were analysed with analysis of variance, followed with Tukey HSD. Result showed considerable variability in seedling vigour, lodging, plant height, panicle exertion, panicle thresh ability, spikelet fertility, and maturity among the studied pigmented rice landraces. Panicle exertion and spikelet fertility indicate plant cope toward environmental change, which was performed better by many landraces than modern varieties. Variability in morphological characteristics was also recorded among tested varieties. Utilisation of the trait diversity can be useful for future breeding programs, which can lead to this crops' improvement, including pest and disease resistance and grain quality research.

## 1. Introduction

Several research on the diversity of local pigmented rice across Indonesia have been carried out [1,2,3]. In particular, the physicochemical properties of pigmented rice from South Sulawesi have been revealed [4,5], confirming the potential of such varieties for nutraceutical products. Therefore, pigmented rice is considered as speciality rice, which deserves a better premium price in both the domestic and export market.

The pigmented rice landraces possess the immense potential of most valuable genes which can be utilised in the current breeding program to evolve varieties that have not only high yield potential and grain quality of nutritional contents but also resistant to biotic and abiotic stresses. Systematic study and characterisation of such germplasm is not only crucial for utilising the appropriate attribute-based donors but also essential for protecting the exotic rice.

Morphological rice characteristics are less documented in Indonesia [3]; hence the diversity of rice germplasms remain untapped. Plant morphologies are the primary basis for the classification of plants or an interpretation of plant adaptability. Moreover, the link between the morphologies of plants and



productivity has resulted in the concept of a desirable plant type (idiotype) [6]. Hence, there is a necessity to do the process of collection, exploitation and evaluation of untapped germplasms [7].

The study was aimed to compare the variability of 15 pigmented rice landraces and two modern varieties using agro-morphological characters and to provide useful information to facilitate the choice of genitors for further crop improvement.

## 2. Materials and methods

### 2.1 Seed materials

Seeds of 16 traditional pigmented rice varieties were collected from four regencies in South Sulawesi of Indonesia, namely: North Toraja, Tana Toraja, Enrekang, and Jeneponto (table 1). The seed collection is a part of local variety exploration program organised by a team from the Plant Biosciences & Reproduction Biotechnology Laboratory, Faculty of Agriculture, Hasanuddin University). Seed samples were collected during the timeframe of June - November. Local varieties are named according to farmers' information. In this research, two modern variety were used as comparison, *i.e.* Inpari 24 and Ciherang.

**Table 1.** List of pigmented rice landraces used in the present study and their source of origin.

Variety	Grain colour and type	Sub-district/Regency	Altitude (m asl)
Ko'bo	Black	Balusu, North Toraja	900
Lotong	Black	Nanggala, North Toraja	780
Ambo	Black	Balusu, North Toraja	903
Barri Rarang	Red	Bengkelekila, North Toraja	836
Lalodo	Black	Balusu North Toraja	905
Lea	Red	Sangalla, Tana Toraja	791
Lotong Tanduk	Black	Nanggala, North Toraja	908
Maminyak	Red	Baraka, Enrekang	727
Mandoti	Red	Salukanan, Enrekang	789
Balan	Red	Baraka, Enrekang	536
Kamida	Red	Curio, Enrekang	847
Jambu	Red	Banua, Enrekang	607
Le'leng	Black	Turatea, Jeneponto	115
Bakka Eja	Red	Turatea, Jeneponto	115
Punu Eja	Red	Turatea, Jeneponto	153
Inpari24*	Red	Purchased from Indonesian Rice Research Center, West Java	
Ciherang*	White	Purchased from Regional Crop Seed Center of Maros, South Sulawesi	

\* Modern variety (nationally released)

### 2.2 Timing and methods.

The study was conducted in Plant Growth Unit at Massey University, Turitea campus (40°22'40"S 175°36'48"E, elevation: 35 m above sea level) from December 2017 (summer) to July 2018 (winter). Rice plants for the preliminary research were cultivated in a controlled environment. The experiment was set up as a Randomised Block Design with five replicates. Potting mix media with slow released fertiliser was used. Drip irrigation was established in week 4, where each pot was placed in 9.6 L spout

bucket to achieve flooded/saturated condition. Fertilisation was done when leaf nitrogen status was assessed as low based on a Leaf Color Chart (score  $\leq 3$ ) (Rice Knowledge Bank, 2018). Peters® Professional Allrounder (N-P-K : 20-20-20+TE) at 1g/L was used as fertiliser, incorporating with drip irrigation (5 L per day).

### 2.3 Data collection and analysis.

The accessions were characterised according to Standard Evaluation System for Rice [8]. A total of seven agronomic traits data (table 2) and twelve of morphological characteristics (table 3) were collected. Simple statistical parameters *i.e.* mean and standard error were determined for the quantitative traits using statistical package SPSS 15.0. The data after compiling was statistically analysed using SPSS 15.0 and Tukey post hoc test (Honest Significance Difference) test was applied to test the significance of varietal differences. Pearson correlation was also used to investigate any association between parameters.

**Table 2.** Descriptions and scales/codes of agronomic traits to evaluate 15 pigmented rice landraces from South Sulawesi [8].

Descriptors	Observed phenotypic classes	Evaluation phase
Seedling vigour (Vg) /vegetative	1-extra weak;9-very weak;3-vigorous;5-normal;7-vigorous	Seedling vigour:2 Vegetative vigour:3
Tillering ability (Ti)	1-very high (>25);3-good(20-25);5-medium(10-19);7-low(5-9);9-very low (<5)	At growth stage 5
Plant height	1-semidwarf (lowland:<110cm;upland:<90cm);2-intermediate (lowland:110-130cm;upland(90-125cm);9-tall(lowland>130cm;upland:>125cm)	At growth stage 7-9
Panicle exertion	1-enclosed;3-partly exerted (panicle base is slightly beneath the collar of flag leaf blade);5-just exerted (panicle base coincides with the collar of the flag leaf blade);7-moderately well exerted (panicle base is above the collar of the flag leaf blade);9-well exerted(panicle base appears well above the collar of the flag leaf blade)	At growth stage 7-9 (near maturity)
Panicle threshability	1-difficult (less than 1%);3-moderately difficult (1-5%);intermediate (6-15%);7-loose (26-50%);9-easy(51-100%)	At growth stage 9
Spikelet fertility	1-highly fertile(>90%);3-fertile(75-89%);partly sterile(50-74%);highly sterile(<50% to trace);9-0%	At growth stage 9
Maturity	Number of days from seeding to grain ripening (85% of grains on panicle are mature)	At growth stage 9

**Table 3.** Morphological character descriptors used to evaluate 15 pigmented rice landraces from South Sulawesi.

Descriptions	Observed phenotypic classes	Evaluation phase
Leaf length	Actual measurement in cm of the widest portion of the leaf blade just below the flag leaf	At growth stage 6
Leaf blade colour	1-light green;2-green;3-dark green;4-purple tips;5-purple margins;6:purple blotch;7:purple	At growth stage 4-6
Ligule length	Enter actual measurement of ligules measured in mm from the base of collar to the tip. Sample size 5	At growth stage 4-5
Ligule colour	0-absent;1:white;2:purple lines;3:purple	At growth stage 4-5
Ligule shape	0-absent;1-acute to acuminate;2-cleft;3-truncate	At growth stage 3-4
Auricle colour	0-absent;1-light green;2-purple	At growth stage 4-5
Culm internode colour	1-green;2-light gold;3-purple lines;4-purple	At growth stage 7-9
Panicle length	Enter actual measurement in cm from panicle base to tip	At growth stage 8
Awning	0-absent;1-short and partly awned;5-short and fully awned;7:long and partly awned;9-long and fully awned	At growth stage: 7-9
Awn colour	0-awnless;1-straw;2-gold;3-brown;4-red;5-purple;6-black	At growth stage: 6
Grain width	Enter the actual measurement of width in mm as the distance across the fertile lemma and the palea in the widest point	At growth stage:9
Grain length/width ratio	Ratio of length in mm as the distance from the base of the lowermost sterile lemma to the tip (apiculus) of the fertile lemma/palea per width	At growth stage:9

### 3. Results and discussion

#### 3.1 Agronomic traits

The assessed pigmented rice landraces showed considerable variability for the studied agromorphological characteristics. Seven agronomic traits comparison among studied varieties are presented in table 4.

*3.1.1 Maturity.* Number of days to achieve the maturity of rice seeds were in the range from 157 cm in Jambu to 215 cm in each variety of Lallodo and Mandoti. Modern varieties reached their maturity longer than expected, indicating a response to the environmental difference from their original habitat. Based on observation, plants with early maturity relatively have intermediate to semi-dwarf height, such as Balan, Bakka Eja, Inpari 24 and Ciherang. However, Jambu has early maturity but fall into the tall group.

Generally, the maturity days reached in this trial were longer than reported maturity in references [9,10] which under five months if grown in Toraja regions for some varieties like

Lallodo, Ambo, Mandoti and Lotong. Environmental factors such as light reception and temperature are some of the key factors.

*3.1.2 Plant height.* Plant height can be seen in three classes, namely, intermediate (Balan), semi-dwarf (Bakka Eja, Inpari, Ciherang) and tall type for the remaining tested varieties. 5.8% intermediate height, 17.6% semi-dwarf, 76.5% tall in height. Traditional rice varieties typically have tall stature whereas modern varieties/early maturing types that tend to be semi-dwarf. In general, most tested varieties grew higher than their common height in their origin of habitat. On the other hand, decreased height, earlier maturity, and longer panicles were recorded from upland rice of South Sulawesi under lowland cultivation [11]. Semi-dwarf rice plant is a primary target for improving the lodging resistance [12], but the drawback is, the canopy for photosynthesis and number of biomass are limited, and then decrease the grain yield [13]

*3.1.3 Panicle exertion.* There are four groups of panicle exertion, which are dominated by well exerted (58.8%), whereas partly exerted was shown in two modern varieties. Most traditional varieties such as Kobo, Lotong, Ambo, Ra'rang, Lotong Tanduk, Maminyak, Kamida, Jambu, Leleng and Punu Eja showed well-exserted panicles. In contrast, modern varieties have partly exerted panicles.

Panicle exertion is one of the important agronomic traits since it can be an indication of genetic defect caused by environmental or pest -disease attack [14]. Incomplete panicle exertion indicates cold injury at the reproductive stage, which lead to damaged grain yield as disease incidence raises [15]. The long and well-exserted panicles are the typical characters of upland rice panicles, as supported by [16], that upland traditional rice are adapted to harvesting and storage by panicles which are long and well exerted.

*3.1.4 Threshability.* Six traditional varieties and two modern varieties were observed to be in the easy category of threshability. Grain characteristics of the ease threshability group are awnless. In contrast, awned grains have intermediate to moderately difficult of threshability.

The result confirmed the threshability of awned genotypes is more difficult than awnless. Some farmers in Toraja prefer to have moderate-intermediate threshability of panicles to minimise losses during post-harvest, as they transport their stacks of panicles manually and store the panicles with the stalks, without shattering [17].

*3.1.5 Seedling vigour.* Seedling vegetative vigour fell into three groups: weak (Kobo, Lotong, Lotong Tanduk), normal (Lea, Kamida) and the remainder as vigorous (table 4). In general, most tested varieties, including modern varieties had acceptable seedling vigour.

*3.1.6 Spikelet fertility.* Our result only have two groups of spikelet fertility: partly sterile and highly sterile. Some traditional varieties perform better than the modern ones. Unexpected sterility of panicles may resulted from microclimate discrepancy in pot trial compared to the origin. Moreover there was disruption from the heat fan which caused dried panicle during anthesis.

The insufficient spikelet fertility in this research might also resulted from cold stress and low solar radiation during reproductive phase. According to [18], during reproductive period, low temperature-induced spikelet sterility, could be associated with the physiological response mechanisms in the pollen, anther, and stigma.

**Table 4.** Agronomic traits of pigmented rice landrace of South Sulawesi and two modern rice varieties for comparison

Variety	Maturity (days)	Plant height (cm)	Panicle exertion	Threshability	Seedling vegetative vigour	Spikelet fertility	Tillering ability
Kobo	195	Tall	Well exerted	Moderately difficult	Weak	Partly sterile	Medium
Lotong	202	Tall	Well exerted	Moderately difficult	Weak	Highly sterile	Medium Very high
Ambo	190	Tall	Well exerted	Intermediate	Vigorous	Partly sterile	Good Very high
Ra'rang	195	Tall	Well exerted	Easy	Vigorous	Highly sterile	Good Very high
Lalodo	215	Tall	Moderately well	Intermediate	Vigorous	Highly sterile	Good Very high
Lea	202	Tall	Moderately well	Easy	Normal	Highly sterile	Good Very high
L. Tanduk	195	Tall	Well exerted	Intermediate	Weak	Highly sterile	Good Very high
Maminyak	190	Tall	Well exerted	Moderately difficult	Vigorous	Partly sterile	Good Very high
Mandoti	215	Tall	Moderately well	Moderately difficult	Vigorous	Highly sterile	Very high
Balan	170	Intermediate	Just exerted	Easy	Vigorous	Highly sterile	Very high
Kamida	198	Tall	Well exerted	Easy	Normal	Highly sterile	Very high
Jambu	157	Tall	Well exerted	Intermediate	Vigorous	Highly sterile	Very high
Le'leng	198	Tall	Well exerted	Intermediate	Vigorous	Partly sterile	Very high
Bakka Eja	160	Semidwarf	Just exerted	Easy	Vigorous	Highly sterile	Very high
Punu Eja	198	Tall	Well exerted	Easy	Vigorous	Partly sterile	Very high
Inpari 24	160	Semidwarf	Partly exerted	Easy	Vigorous	Highly sterile	Very high
Ciherang	160	Semidwarf	Partly exerted	Easy	Vigorous	Highly sterile	Very high

*3.1.7 Tillering ability.* The tillering ability of all varieties in this study appears to be medium (10-19 tillers per plant) to very high (more than 25 tillers/plant). The ideal planting space is 25 cm to allow the appropriate growth of tillers and minimise disease in humid condition. Most of the varieties showed a very high tillering ability while few varieties namely Maminyak, Lea and Ra'rang had good tillering ability and only Kobo as well as Lotong had medium tillering ability.

Similarly [10] reported significant difference in traits such as maturity, height, tiller number, seed weight, and productivity among 29 rice landraces from Toraja region. A few of them have coloured grain, such as Pare Lea, Pare Lallodo, and Pare Ambo. However, the tiller number reported by [9] was 3-4 times higher (17-30) than tiller number reported in [10]. These two different studies were undertaken in different locations but in the Toraja region hence variability in result is noted.

### 3.2 Morphological characteristics

*3.2.1 Quantitative measurements.* Ra'rang has the longest ligule length among the landraces, which similar to modern Inpari 24. Jambu has the shortest ligule. correlation coefficient of ligule length and grain length/width ratio was 0.115 Which interpreted as little if any correlation [19]. Conversely, a little negative correlations were found between ligule length and leaf length, panicle length, and grain width.

The most fundamental factor in the diversity of grain colours is the genetic factor that expresses the grain colour of the variety. Result of bioactive content assessment of pigmented rice landrace of South Sulawesi also confirmed the high anthocyanin content of black rice, and considerable amount of phenolic content in red rice [5].

Maminyak has the longest leaf (78.80 cm), whilst Ciherang, Balan, and Inpari have the shortest leaf length. has the shortest leaf length (45 cm;48 cm and 51 cm respectively). Moderate positive correlation ( $r=0.577$   $N=80$   $p<0.01$ ) was found between leaf length and plant height.

The longest panicle length was found in Lotong (36.5 cm), which significantly differ from Rarang, Lallodo, Lea, Lotong Tanduk, Mandoti, Balan, Kamida, Leleng, Bakka Eja, Punu Eja and two modern varieties (Ciherang and Inpari 24) (Table 5). As expected, the shortest panicle was found in Ciherang. Panicle length has low positive correlation with leaf length ( $r=0.488$   $N=80$   $p<0.001$ ) (Table 5). Multiple genes controlled panicle length, a typical quantitative trait, and may be affected by environmental conditions significantly. In addition, breeding varieties with large panicle and the ideal plant architecture traits are significant for increasing the grain yield of rice [20].

**Table 5.** Measurement of morphological characteristics of pigmented rice landrace of South Sulawesi and two modern rice varieties for comparison.

Variety	Ligule length	Leaf length	Panicle length	Grain's length/width ratio	Grain width
Kobo	14.20±0.66edc	63.00±2.19gfedcb	32.00±2.30ba	2.78±0.07gf	2.90±0.13cba
Lotong	17.20±1.24cba	76.10±3.26cba	36.50±0.87a	3.14±0.08fedc	3.00±0.05ba
Ambo	13.80±1.24edc	63.40±2.71fedcb	29.60±0.53cdba	2.64±0.04g	3.00±0.06ba
Ra'rang	20.40±1.63a	68.40±3.16edcba	21.80±2.89hgfed	2.92±0.12gfed	2.84±0.08cba
Lallodo	14.80±0.50edcb	72.60±1.89dcba	21.80±1.98hgfed	3.32±0.11dc	2.72±0.09dcba
Lea	13.00±1.38edc	70.40±2.25dcba	23.40±1.78hgfedc	2.84±0.05gfe	2.86±0.07cba

L. Tanduk	11.50±0.50ed	63.00±2.00gfedcb	23.00±2.00hgfedc	2.77±0.05	3.25±0.05
Maminyak	13.60±0.93edc	78.80±2.96a	29.20±1.59edcba	2.89±0.15gfed	2.76±0.08dcba
Mandoti	12.00±0.71edc	76.00±3.41cba	25.40±0.98gfedcb	3.11±0.09fedc	2.86±0.02cba
Balan	15.60±0.51dcba	48.20±2.44hg	20.20±1.36hgf	3.19±0.05fedc	2.66±0.05dc
Kamida	11.00±0.45ed	59.20±0.86hgfed	27.80±1.66fedcb	3.09±0.10gfedc	2.70±0.03dcb
Jambu	10.00±0.55e	77.40±2.58ba	30.04±1.16cba	3.25±0.06edc	2.82±0.04cba
Le'leng	12.60±0.81edc	59.40±2.01hgfed	25.20±0.73gfedcb	2.91±0.02gfed	3.04±0.06a
Bakka Eja	12.60±0.81edc	53.90±1.35hgfe	21.20±1.01hgfe	3.85±0.08ba	2.48±0.06ed
Punu Eja	12.00±0.95edc	62.00±4.12gfedc	22.40±1.69hgfedc	2.99±0.12gfed	3.00±0.10ba
Inpari 24	20.20±1.20ba	51.00±1.87hgf	18.93±1.22hg	4.13±0.08a	2.24±0.05e
Ciherang	14.00±1.10edc	45.40±5.57h	15.40±0.51h	3.52±0.10cb	2.18±0.04e

Values followed the same letter are not different ( $P < 0.05$ ).

**3.2.2 Qualitative observations.** Morphological characteristics of tested varieties observed under glass house conditions are presented in table 6. An important observed feature is the extent of pigmentation found in various plant parts including auricles, ligules, awns, culm internodes, and leaf blades among different varieties. While light green was the typical colour of auricles in most varieties tested in this experiment, only Ambo and Lotong Tanduk have the purple colour in the auricles. Awned seeds found in 52.9% of tested varieties. One of Wild rice species' distinct feature in morphology is a long awn [21]. Shorter awn or awnless grain in domesticated wheats facilitate harvesting, handling and storage [22].

The awn color of awned seed types ranged from straw (Kobo, Lotong, Lallodo, Maminyak, Mandoti, Jambu, Le'leng) to red (Ambo and Lotong Tanduk). Each variety of Ambo as well as Lotong Tanduk also shared similarity in culm internode color (purple), ligule color (purple), and leaf blade colour (purple in the margin), along with kernel colour of black. The culm internode colour of Ambo, L. Tanduk, Kamida were purple, while Punu Eja has purple line for their culm internode colour. Most varieties (88.2%) have white ligule colour while the ligule shape was cleft in all varieties. Plants with purple ligule colour (Ambo and Lotong Tanduk) also have purple margin leaf blade colour, and the remaining varieties are in green colour of leaf blade (table 6). Compared with black rice found in Central Java, namely Matesih, Klaten and Cempo, all of them have white ligule in cleft shape, green leaf blade color and white auricle [23].

**Table 6.** Morphological characteristics observation of pigmented rice landrace of South Sulawesi and two modern rice varieties for comparison

Variety	Auricle color	Awn color	Awning	Culm internode color	Ligule color	Ligule shape	Leaf blade color
Kobo	Light green	Straw	Long and Well	Green	White	cleft	Green
Lotong	Light green	Straw	Long and Well	Green	White	cleft	Green
Ambo	Purple	Red	Long and Well	Purple	Purple	cleft	Purple margin
Rarang	Light green	-	Awnless	Green	White	cleft	Green
Lallodo	Light green	Straw	Short and partly	Green	White	cleft	Green
Lea	Light green	-	Awnless	Green	White	cleft	Green
Lotong Tanduk	Purple	Red	Short and partly	Purple	Purple	cleft	Purple margin
Maminyak	Light green	Straw	Long and Well	Green	White	cleft	Green
Mandoti	Light green	Straw	Long and Well	Green	White	cleft	Green
Balan	Light green	-	Awnless	Green	White	cleft	Green
Kamida	Light green	-	Awnless	Purple	White	cleft	Green
Jambu	Light green	Straw	Long and Well	Green	White	cleft	Green
Le'leng	Light green	Straw	Long and Well	Green	White	cleft	Green
Bakka Eja	Light green	-	Awnless	Green	White	cleft	Green
Punu Eja	Light green	-	Awnless	Purple line	White	cleft	Green
Inpari 24	Light green	-	Awnless	Green	White	cleft	Green
Ciherang	Light green	-	Awnless	Green	White	cleft	Green

#### 4. Conclusions

The discovery of genetic differences plays a key role in the development of varietal conditions that may contribute to food security. In this study, variability is seen in some agronomic traits and morphological characteristics of pigmented rice landraces.

Some pigmented rice landraces performed better than modern varieties under glasshouse environment of temperate climate. The better performance of pigmented rice landrace indicate the capability in coping towards different environment from their origin of habitat.

Pigmented rice plant idotype can be engineered from selecting desired agronomic traits, ideal architecture, as well as superior physicochemical properties. Farmers perspective and their requirement should be taken into consideration as well, since farmers are the key factor for preserving the germplasm in situ by retaining the varieties intergeneration.

#### References

- [1] Anggraini T, Novelina, Limber U, and Amalia R 2015 Antioxidant activities of some red, black and white rice cultivar from West Sumatra, Indonesia. *Pakistan Journal of Nutrition* **14** 112-7.
- [2] Kristantini T, Basunanda P, Murti R H, & Supriyanta S 2012 Morphological of genetic relationships among black rice landraces from Yogyakarta and surrounding areas. *ARPN Journal of Agricultural and Biological Science* **7** 982-9.
- [3] Shinta S, Indriyani S, and Arisoesilaningsih E 2014 Morphological variation of six pigmented rice local varieties grown in organic rice field at Sengguruh Village, Kepanjen Subdistrict, Malang District. *Journal of Tropical Life Science*, **4** 149-160. Retrieved from <https://jtolis.uib.ac.id/index.php/jtolis/article/view/171>
- [4] Murdifin M, Pakki E, Rahim A, Syaiful S A, Ismail, Evary, Y M, and Bahar M A 2015 Physicochemical properties of Indonesian pigmented rice (*Oryza sativa* Linn.) varieties from South Sulawesi. *Asian Journal of Plant Sciences* **14** 59-65. 10.3923/ajps.2015.59.65
- [5] Hanifa A, Millner J, McGill C, and Sjahril R 2020 Total anthocyanin, flavonoid and phenolic content of pigmented rice landraces from South Sulawesi. *IOP Conference Series: Earth and Environmental Science*,
- [6] Loomis R S and Williams W A 1969 Productivity and the morphology of crop stands patterns with leaves. In *Physiological Aspects of Crop Yield*. ASA-CSSA. Wisconsin.
- [7] Parikh M, Motiramani N K, Rastogi N K and Sharma B 2012 Agro-morphological characterization and assessment of variability in aromatic rice germplasm *Bangladesh Journal of Agricultural Research* **37** 1-8.
- [8] International Rice Research Institute 2013 *Standard Evaluation System for Rice* 5<sup>th</sup> Edition. IRRI. Manila, Philippines.
- [9] Limbongan Y and Djufry F 2015 Characterization and Observation of Five Local Rice Accessions of Toraja Plateau, South Sulawesi *Bul. Plasma Nutrafah* **21** 61-70
- [10] Sahardi, Herniwati, and Djufry F 2015 Keragaman karakter morfologis plasma nutrafah lokal Toraja. Paper presented at the *Prosiding Seminar Nasional Sumber Daya Genetik Pertanian*, Bogor.
- [11] Sahardi and Hanifa, AP 2015 Performance of upland rice under lowland growing condition in Gowa Regency, South Sulawesi. *Prosiding Seminar Nasional Pengembangan Sumber Daya Genetik Pertanian: Potensi Sumber Daya Genetik Spesifik Lokasi Mendukung Ketahanan dan Swasembada Pangan Nasional*. Bogor: 27<sup>th</sup> May 2015. Balai Besar Pengkajian dan Pengembangan Teknologi Pertanian, Badan Penelitian dan Pengembangan Pertanian,
- [12] Khush G S 1999 *Green revolution: Preparing for the 21st century Genome* **42** 646-655 PMID: 10464789
- [13] Kuroda E, Ookawa T, Ishihara K 1989 Analysis on difference of dry matter production between rice cultivars with plant height in relation to gas diffusion inside stands *Jap J Crop Sci* **58** 374-382 DOI:10.1626/JCS.58.374
- [14] International Rice Research Institute 2002 *Standard Evaluation System for Rice* IRRI <http://www.knowledgebank.irri.org/images/docs/rice-standard-evaluation-system.pdf>

- [15] Pereira da Cruz R, S C Milach and L C Federizzi 2008 Inheritance of panicle exertion in rice *Sci. agric. (Piracicaba, Braz.)* **65** 502-507.
- [16] Chandler Jr R F. 1979 *Rice in the tropics*. A guide to the development of national programs. Westview Press Boulder, Colorado.
- [17] Hanifa, A P 2018 Evaluation of pigmented rice landraces from Toraja, Sulawesi. Confirmation report. School of Agriculture and Environment. Massey University. Unpublished
- [18] Zeng Y, Zhang Y, Xiang J, Uphof N T, Pan X, and Zhu D 2017 Effects of low temperature stress on spikelet-related parameters during anthesis in Indica–Japonica hybrid rice *Frontiers in Plant Science*, **8** 10.3389/fpls.2017.01350
- [19] Asuero, G A Sayago & A G González 2006 *The correlation coefficient: An overview, critical reviews in analytical chemistry* **36** 41-59, DOI: 10.1080/10408340500526766
- [20] Yao X Y, Li Q, Liu J, Jiang S -K, Yang S -L, Wang, J -Y (2015). Dissection of QTLs for plant height and panicle length traits in rice under different environment. *Sci. Agr. Sin.* **48** 407–414 DOI: 10.3864/j.issn.05781752.2015.03.01
- [21] Furuta T, Komeda N, Asano K, Uehara K, Gamuyao R, Angeles-Shim R B, Ashikari M 2015 Convergent loss of awn in two cultivated rice species *Oryza sativa* and *Oryza glaberrima* is caused by mutations in different loci. *G3 (Bethesda, Md.)* **5** 2267-2274. 10.1534/g3.115.020834
- [22] Mach J 2015 Domesticated versus wild rice? Bring it awn! *The Plant cell* **27** 1818. <https://doi.org/10.1105/tpc.15.00504>
- [23] Purwanto E and Meidini A 2018 *Morphology, production, and chemical content performance of black rice Matesih accession with several comparisons*. Paper presented at the IOP Conference Series: Earth and Environmental Science.

PAPER • OPEN ACCESS

## Preface

To cite this article: 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **807** 011001

View the [article online](#) for updates and enhancements.

## You may also like

- [Smartphone-based GNSS Method for Covid-19 Information on Android \(Case Study: Surabaya City\)](#)  
A H Akbar and M N Cahyadi
- [The Effect of Climate on the Outbreak of Covid-19: A Review](#)  
Candra Kirana, Samsul, Hesty Yariska Hapsir et al.
- [Impacts of the Covid-19 pandemic on traditional oil mining at Wonocolo Village Kedewan Sub-District Bojonegoro Regency East Java](#)  
L A Rahmawati, N Afiati and T T Putranto

## PREFACE

It is a great honour for us to be the host of the 3rd International Conference on Food Security and Sustainable Agriculture in the Tropics (FSSAT) on January 8th – 9th, 2021 at Agriculture Faculty of Hasanuddin University, Makassar City of South Sulawesi Province, Indonesia. This conference is a series of international scientific seminars held by the Faculty of Agriculture, Universitas Hasanuddin, Makassar, Indonesia, which has been held annually since 2017.

This conference was held as a forum for exchanging information on productivist paradigm in Indonesian food policies to feed 260 million people targets the self-sufficiency of rice, sago, corn, soybean, and meat production, including the infrastructure development such as dams, irrigation channels building, and the expansion of paddy and sago fields in the Eastern Indonesia. The unintended outcomes of these efforts are the marginalization of the local food system and dispossession of the local communities from their land and food culture. As a consequence, the human-environmental relationship has changed dramatically posing calamitous disaster and slowly disappearing foodways through local wisdom in everyday life of the local community. It also brings health issues due to the excessive use of chemical agricultural inputs in production, and the increase of degenerative disease and obesity due the unhealthy practices in the consumption level.

The International Conference on Food Security and Sustainable Agriculture in the Tropics will be held every two years, the first in 2017, the second in 2019, and the third in 2021. The covid-19 conditions that have hit the whole world have resulted in restrictions on activities to reduce the spread of Covid-19 sufferers so that the conference model from the form of a live conference has turned into a virtual conference.

The condition of the Covid-19 pandemic has raised concerns about food security throughout the world, including Indonesia, through restrictions on community activities. Several things have implications for the existence of Covid-19, including the occurrence of a very significant supply chain disruption caused by a reduction in capacity to produce food, closing transportation routes so that it will slow down the distribution of food from producers to consumers. The inability of people to consume enough healthy and nutritious food will reduce the immune system and increase disease risk. Thus ensuring an affordable food supply chain is essential during the Covid-19 outbreak. To solve this problem globally, cooperation is needed that involves scientists, academics, practitioners, professionals, governments, entrepreneurs, and communities across countries. To build a network between stakeholders, an international conference held by the Hasanuddin University Faculty of Agriculture in 2021 is important as a forum for building connectivity between researchers and/or research institutions. At this conference, a multi-disciplinary approach based on the knowledge and experience of scientists, researchers, practitioners, and policymakers will be brought together.

If the meeting cannot be held virtually, then solutions in dealing with the food crisis during the Covid-19 pandemic will not find a way out and add to the length of the crisis period chain.

This conference presents keynote speakers from 5 countries, namely Dr. Ir. Bayu Krisnamurthi, M.Si. from IPB University, Indonesia, Prof. Dr. Ir. Sumbangan Baja, M.Phil from Universitas Hasanuddin, Indonesia, Prof. Takuya Sugahara from Ehime University, Japan, Dr. Ravindra C Joshi from CABI South-East Asia, Dr. Lau Wei Hong from University

Putra Malaysia, Mr. Moh, Guo-Jhong from ICDF Taiwan. First plenary session and discussion with 2 speakers for 1 hour and the second plenary and discussion with 3 speakers for 2 hours. For participant presentations (oral presentation and video presentation) it was held in 2 sessions, each session was divided into 10 rooms. The number of participants per room was 11-14 participants with an allocation of presentations for 7 minutes/participants conducted in parallel as many as 7 parallel sessions. Discussions session were held in parallel in each parallel session with an allocation of 3 minutes per participant. Virtual FSSAT 3 implementation using the zoom meeting application by the Faculty of Agriculture, Universitas Hasanuddin has which is obtained by subscription.

The seminar participants were attended by lecturers and students from various universities as well as researchers from various research institutions both from the ministry of agriculture and from Agricultural Technology Research Center. The institution of the participants came from Agricultural Technology Research Center, some polytechnic, institute and University, Indonesian Center for Estate Crops Research and Development, International Coconut Community, Center for Agricultural Technology Research and Development, PT Vale Indonesia Tbk, Sorowako. Some from overseas universities: University of New England, Armidale, Australia, and School of Agriculture and Environment of Massey University, Palmerston North New Zealand. Overall participant numbers around 1.034 participants.

There were no significant technical obstacles, speakers and participants from overseas also presented their papers fluently. Apart from sharing zoom rooms, we also created a social media group (WhatsApp) for each class to facilitate communication between participants and organizers.

All published articles are subject to a rigorous selection process and are peer-reviewed by international and national reviewers. On this occasion, we would like to thank all the authors and the team who have actively participated in the 3<sup>rd</sup> FSSAT conference, expert reviewers who are very responsive to the technical program committee, experienced senior publication chair, Publication Management Center (PMC) of Universitas Hasanuddin, authors and scholars who are enthusiastic in research. Sincerely thank you for your warm support and look forward to working with you in the next future.

Warm Regards,

Dr. Asmita Ahmad, ST., MSi.

Conference Chairman

Faculty of Agriculture, Universitas Hasanuddin

**COMMITTEES**

Steering Committee	Dean, Faculty of Agricultural, Hasanuddin University Dean, Faculty of Forestry, Hasanuddin University Dean, Faculty of Animal Science, Hasanuddin University Dean, Faculty of Marine Science and Fishery, Hasanuddin University
Organizing Committee	Chair: Dr. Asmita Ahmad, ST., MSi. Vice Chair: Andi Dirpan, STP., MSi., PhD. Secretary: Dr. Suleha Thamrin, SP., MSi. Treasurer: St. Hasnah A. Rahman, SP.
Scientific Advisory Board	Prof. Dr.sc.agr.Ir. Baharuddin (Indonesia) Prof. Takuya Sugahara (Japan) Dr. Ravindra C Joshi (Philippines) Dr. Lau Wei Hong (Malaysia) Mr. Guo-Jhong Moh (Taiwan) Dr. Ir. Bayu Krisnamurthi, M.Si (Indonesia) Prof. Ir. Sumbangan Baja, M.Phil, PhD(Indonesia)  Dr. Muh. Hatta Jamil, SP., M.Si. (Indonesia) Dr.rer.Nat. Zainal, STP. M.Food.Tech (Indonesia) Dr. Ir. Novaty E. Dunga, MP (Indonesia) Prof. Dr. Ir. Nurriaty Agus, MS. (Indonesia) Dr. Ir. Rahmadani, M.Si. (Indonesia) Prof. Dr. Ir. Tutik Kuswinanti, M.Sc. (Indonesia) Prof. Dr. Ir. Meta Mahendrasatta, MS. (Indonesia) Dr. Ir. Amir Yassi, MS. (Indonesia) Dr. Rismaneswati, SP., MP. (Indonesia) Dr. Nixia Tenriawaru, SP., M.Si. (Indonesia) Dr. Iqbal, STP. M.Si. (Indonesia) Ir. Rinaldi Sjahril, M.Agr., PhD. (Indonesia) Dr. Ir. Rusnadi Padjung, M.Sc. (Indonesia) Muhammad Arsyad, SP., M.Si., PhD. (Indonesia) Dr. Ir. Andi Nasruddin, M.Sc. (Indonesia) Prof. Dr. Ir. Sikstus Gusli, M.Sc. (Indonesia) Prof. Ade Rosmana, M.Sc. (Indonesia) Prof. Dr. Ir. Christianto Lopulisa, M.Sc. (Indonesia) Prof, Dr. Ir. Salengke, M.Sc. (Indonesia) Prof. Dr. Ir. Yunus Musa, M.Sc. (Indonesia) Prof. Dr. Ir. M. Saleh Ali (Indonesia) Prof. Dr. Ir. Musrizal Muin M,Sc(Indonesia)
Secretariat	Sartika Laban, SP., MP., Ph.D Dr. Abdul Aziz, S.TP., M.Si. Nirmala Juita, SP., M.Sc. Hamdayanti, SP., M.Si. Ahmad Fauzan Adzima, SP., M.Sc. Magfirah, SP. Nilarisa Meganita, S.TP

Paper, Poster, and Prosiding	Muhammad Junaid, SP., MP, Ph.D Dr. Ir. Asmiaty Sahur, MP. Rasyida Bakri, SP., M.Sc Muspilah Djalal, STP., M.Sc. Dhini Amriani Mutia Mey Nindi Zulkifly
Publicity and Publishing	Ifayanti, SP., MP., Ph.D Rusli M. Rukka, SP., M.Si. Rahmansyah Dermawan, SP. M.Si Andi Rahmayanti, S.TP. M.Si Iswar Fajar Sultan, SP. Andi Junila, SP. Rizza Nurul Aprillia.
Website	Muh. Tahir Sapsal, STP., MSi. Samsuar, STP., M.Si. Dr. Adiansyah, S.TP, M.Si Yudiono, SE
Registration and Certificate	Dr. Muh. Asfar, S.TP., M.Si Dr. Ir. Katriani Mantja, MP Husnul Mubarak, S.TP., M.Si Ir. Nurlina Kasim, M.Si Cri Wahyuni Brahmiyanti, SP., M.Si Tigin Dariarti, SP
Finance and Corporate Partnership	Dr. Letty Fudjaja, SP., M.Si Dyah Yumeina R. Datu, S.TP., M.Agri.,PhD. Pipi Diansari, SE., M.Si., PhD. Dr. Ir. Heliawaty Ch. Lakare, M.Si Dr. Ir. Abd. Haris Bahrin, M.Si Ikmalul Akmal, S.Sos
Keynote/Invited Speaker team	Hari Iswoyo, SP., MA., PhD Dr. Olly Sunny Hutabarat, S.TP. Dr. Ir. Vien Sartika Dewi, M.Si Dr. Andi Nur Faidah, R, S.TP., M.Si NiMadeViantikaSuliandri,SP.,M.Ag
Event, Exhibition and Technical Program	Februardi Bastian, STP., M.Si., Ph.D. Asman, SP., MP. Abd. Mollah, SP., M.Si. Achmad Amiruddin, SP.,M.Si Nuniek Widiyani, SP., MP Abd. Ishak, S.Sos Kurniati Latief

**EDITORS & REVIEWERS**

Ir. Muhammad Arsyad, S.P., M.Si, Ph.D

Prof. Dr. Ir. Didi Rukmana, M.S

Prof. Dr. Ir. Muslim Salam, M.Ec

Dr. A. Nixia Tenriawaru, S.P., M.Si.

Achmad Amiruddin, S.P., M.Si

Muhammad Khaliqi, S.P., M.Si

Andi Azrarul Amri, S.P., M.Si

Dwi Ahrisa Putri, S.P., M.Si

Andi Tenri Darhyati, SP, M.Si

Ikawati, S.T.P., M.Si

Ayu Wulandary, S.P., M.Si

Ayu Anisa Amir, S.P., M.Si

Ary Bakhtiar., SP., M.Si

Dian Utami Zainuddin, S.Si., M.Si.

Departemen Ilmu Tanah

Dr. Rismaneswati, S.P., M.P

Dr. Asmita Ahmad, S.T., M.Si

Sartika Laban, S.P., M.P., Ph.D

James Nana Ofori, B.Sc., M.Sc (Ghana)

Supriadi, S.P., M.Si (Japan)

Nirmala Juita, S.P., M.Si

Indryani Bali, S.P., M.Sc

Prof. Dr. Ir. Yunus Musa, M.S

Ir. Rinaldi Sjahril, M.Agr., Ph.D

Ir. Hari Iswoyo, S.P., M.A., Ph.D

Dr. Ir. Muh. Riadi, M.P

Dr. Ir. Rusnadi Padjung, M.Sc

Ifayanti Ridwan Saleh, S.P., M.P., Ph.D

Dr. Ir. Asmiaty Sahur, M.P

Rahmansyah Dermawan, S.P., M.Si

Nuniek Widiyani, S.P., M.Si

Prof. Dr. Ir. Ade Rosmana, M.Sc

Prof. Dr. Ir. Andi Nasruddin, M.Sc

Prof. Dr. Ir. Tutik Kuswinanti, M.Sc  
Muhammad Junaid. S.P.,M.P.,Ph.D  
Dr. Sulaeha Thamrin, S.P.,M.Si  
Hamdayanty S.P.,M.Si  
Prof. Dr. Ir. Meta Mahendradatta  
Andi Dirpan, S.TP.,M.Si.,Ph.D  
Dr. Andi Nur Faidah Rahman., STP., MSi  
JamyangTashi Wangdi, M.Sc (Bhutan)  
Nur Alim Bahmid, S.TP.,M.Si (Netherland)  
Muspilah Djalal, S.TP.,M.Sc  
Dr. Iqbal, S.TP.,M.Si  
Dr. Haerani, S.TP.,M.Eng.Sc  
Dr.rer.nat. Olly S. Hutabarat, S.TP.,M.Si  
Samsuar, S.TP.,M.Si  
Husnul Mubarak, S.TP.,M.Si  
Dr. Kasmiyati Kasim, SPt, MP

PAPER • OPEN ACCESS

## Peer review declaration

To cite this article: 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **807** 011002

View the [article online](#) for updates and enhancements.

You may also like

- [Peer review declaration](#)
- [Peer Review Declaration](#)
- [Peer review declaration](#)

## Peer review declaration

All papers published in this volume of IOP Conference Series: Earth and Environmental Science have been peer reviewed through processes administered by the Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

- **Type of peer review:**  
Single-blind
- **Conference submission management system:**  
We conduct paper submission through submission management system (<http://konfrenzi.com/>). The manuscript from author has submitted before the conference through the conference website (<https://fssat3.unhas.ac.id/>).
- **Number of submissions received:**  
Total received abstract = 364 papers
- **Number of submissions sent for review:**  
355 full papers (19 papers rejected)
- **Number of submissions accepted:**  
291 full papers
- **Acceptance Rate (Number of Submissions Accepted / Number of Submissions Received X 100):**  
79.9 %
- **Average number of reviews per paper:**  
6 papers per reviewer
- **Total number of reviewers involved:**  
47 reviewers
- **Any additional info on review process:**  
There are some steps on review process
  1. Abstract review and acceptance  
At the first stage, abstracts submitted to the committee through the online system available in the website were reviewed by a Scientific Advisory Board/Reviewers for the suitability of the paper's topic to the theme of the conference. Abstracts that suit the theme were accepted and author were notified of the acceptance through the system. Abstract submission and review were conducted simultaneously until abstract submission deadline on October 31, 2020. As many as 364 abstracts submitted through system were reviewed. 1 abstract was rejected as not comply with the theme of the conference. Accepted abstracts were allowed to be presented in an oral or live presentation.
  2. Submission of Full Paper  
At the second stage, authors with abstracts declared accepted by the Scientific Advisory Board/Reviewer and will be presented in the conference were requested to submit a complete manuscript according to the IOP Publishing format. The deadline for receipt of a complete manuscript (full paper) was on December 15, 2020.
  3. Revision of Full Paper  
At the third stage, following the conference that took place on January 8-9, 2021, Authors were requested to revise the full paper according to the input and



suggestions from the conference audiences and re-submit the manuscript through the system until January 31, 2021.

4. Peer review of Full Paper

At the Fourth stage, full paper previously submitted through system were reviewed following a single blind peer review process. The papers were sent to the Scientific Advisory Boards that consisted of 47 reviewers according to topic of the papers and reviewer's competency. The reviewers are from international scientists and from the Faculty of Agriculture, and Faculty of Animal Husbandry, Hasanuddin University. Review was conducted on content and format of the paper based on IOP Conference Series. The review process was carried out starting from February to March 7, 2021. At this stage, Reviewer submit a decision for the acceptance of the papers for publication in IOP Conference Series.

5. Revision of Reviewed Full Paper

At the Fifth Stage, Review results were returned to the authors for revision and re-submission online for those accepted for publication (deadline on April 15, 2021). Following the submission of revised manuscript. The submitted papers were then subjected to proofread stage and check for similarity using Turnitin. Papers with similarity check more than 25% were returned to the authors for further revision. Subsequently, papers were prepared for submission to IOP Conference Series.

- **Contact person for queries:**

The chief of the conference: Dr. Asmita Ahmad, ST., M.Si.

Mobile phone: +6281342542792

E-mail: [asmita.ahmad@agri.unhas.ac.id](mailto:asmita.ahmad@agri.unhas.ac.id)