

Herbal Plants as a Treatment for Halitosis in Children: A Systematic Review

by

Submission date: 17-Apr-2021 09:35AM (UTC+0700)

Submission ID: 1561530835

File name: Plants_as_a_Treatment_for_Halitosis_in_Children_A_Systematic.pdf (441.63K)

Word count: 7038

Character count: 38739

Herbal Plants as a Treatment for Halitosis in Children: A Systematic Review

Harun Achmad¹, Irene Edith Rieuwpassa², ArniIrawaty Djais³, Eriska Riyanti⁴, RistiSaptarini Primarti⁴, Fathimah⁵

39

¹Department of Pediatric Dentistry, Faculty of Dentistry, Hasanuddin University, Indonesia

²Department of Oral Biology, Faculty of Dentistry, Hasanuddin University, Indonesia

³Department of Periodontology, Faculty of Dentistry, Hasanuddin University, Indonesia

⁴Department of Pediatric Dentistry, Faculty of Dentistry, Padjadjaran University, Indonesia

⁵Clinical Dental Student, Faculty of Dentistry, Hasanuddin University, Indonesia

Corresponding Author: harunachmader@gmail.com

ABSTRACT

Introduction: Halitosis is an important multifactorial health problem that affects a person's psychological, social life and quality of life. Halitosis is very common in children. It should be noted that the main causes of halitosis in children include dental caries, poor oral and dental hygiene. Common treatments for halitosis include brushing, flossing using dental floss, chewing gum, and using both chemical and herbal mouthwashes. Long-term use of this chemical has side effects in the form of discoloration of the teeth, triggering tartar formation, allergic reactions, as well as desquamation (flaking) of the oral mucosa. Herbal plants are plants that produce active substances and can be used as an alternative to treat several diseases, including halitosis. **Aim:** This systematic review reviews some of the literature on the potential of herbal plants for treating halitosis in children. **Methods:** In this systematic review, article searches were conducted on Google Search and Pubmed. Studies published from 2010 to 2020. As much 84 articles were rated, including 84 articles from electronic databases, 0 from manual searches. 49 records were screened, 35 records excluded, 37 full-text articles were assessed for eligibility and 12 full-text articles were included. **Result:** There are 12 articles about medicinal plants that can be used to treat halitosis in children. **Conclusion:** There are 12 articles on various active substances in herbal plants that can treat halitosis in children.

Keywords: Herbal plants, Halitosis, Children

INTRODUCTION

Human breath consists of very complex substances with various odors which can give rise to unpleasant situations such as halitosis. Halitosis is an important multifactorial health problem that affects a person's psychological, social life and quality of life.¹ Halitosis comes from the Latin word halitus which means breath and osis means a pathological change used to describe an unpleasant or unpleasant odor that comes from the breath in a person's mouth. Halitosis, also known as *fetor ex ore*, *fetor oris* or *oral malodor*, is a general term that denotes bad breath to describe and characterize halitosis.²

This undesirable condition is a common complaint for both sexes and for all age groups. According to the American Dental Association (ADA), halitosis is the main problem after caries and periodontal disease that is complained of by people in the world. Halitosis is

very common, with more than 50% of the general population suffering from halitosis.³The prevalence of halitosis sufferers in each country varies. In China, patients with halitosis reached 27.5% with a sample of 2500 people. Research conducted in America shows that the prevalence of halitosis sufferers is quite high, reaching 50% of the total population in America. Also in other literature states, the prevalence of halitosis in children is quite high, ranging from 5% to 75%.^{4,5}

Halitosis is very common in children. It should be noted that the main causes of halitosis in children include dental caries, poor oral and dental hygiene. The results of Basic Health Research in 2018 stated that, the total proportion of dental and oral problems was 57.6%; the largest proportion were tooth decay/cavity/pain (45.3%). The prevalence of caries or cavities in children and adults is also quite high, including the prevalence of cavities in early childhood and is still very high, namely 93%. Tooth and mouth disease is caused due to the presence of bacteria that colonize the oral cavity to form a biofilm/plaque layer. This will cause various diseases in the oral cavity, one of which is halitosis.^{6,7}

Halitosis is basically caused by two things, namely physiological and pathological. The physiological source of this halitosis comes from conditions on the surface of the tongue. The bacteria found on the tongue surface of a healthy patient are different from that of a patient with halitosis.⁸The pathological source involves the severity of pocket gums which is known as periodontal disease. The main cause of halitosis is the release of Volatile Sulfur Compound (VSCs) which is caused by the decay activity of gram negative microorganisms. The cause of most halitosis comes from the oral cavity, which is around 80-90%, such as poor oral hygiene, deep caries, periodontal disease, oral infection, xerostomia, ulceration, pericoronitis, leftover food in the mouth and tongue coating, while halitosis originates from from outside the oral cavity occurs only slightly, namely about 10-20% such as systemic diseases, diseases of the respiratory or gastrointestinal system, liver organ failure and metabolic disorders.⁹

A healthy oral cavity allows a person to communicate effectively, enjoy various types of food, improve quality of life, self-confidence, and have a better social life. The existence of halitosis has a negative impact and can even trigger stress. Halitosis can cause harm not only to sufferers but also to others and can affect a person's social life such as shame, avoidance of social interactions and decreased self-confidence in children.^{4,5}

Common treatments for halitosis include brushing, flossing using dental floss, chewing gum, and using mouthwash both chemicals and herbs that usually help freshen breath. Long-term use of these chemicals has side effects in the form of discoloration of teeth, triggering formation of tartar, allergic reactions, as well as desquamation (peeling) of the oral mucosa.^{5,10}Therefore, one of the alternative ingredients that can be used to reduce halitosis is herbal plants. Herbal plants are plants that produce active substances and are used as alternatives to treat several diseases, including oral diseases. Currently, approximately 40% of drugs available use natural sources, either directly or indirectly, and 25% of them are obtained from plants. Research on herbal plants in dentistry has been used as analgesic, anti-inflammatory, antimicrobial, curative and regenerative, with minimal toxicity, better biocompatibility and lower cost. The World Health Organization (WHO) also recommends and encourages the use of plants as alternative therapies for oral hygiene.^{11,12}Therefore, this

systematic study aims to examine more deeply the potential of herbal plants to treat halitosis in children.

SEARCH METHODS

Data Source

The data was collected by searching the literature on article search sites, namely Google search and Pubmed which were published from 2010 to 2020, the search was carried out in December 2020. The data search was carried out systematically using keywords *herbal plant, halitosis, children*

Research Criteria

A. Inclusion criteria

1. Articles published from the years 2010-2020
2. Articles in English
3. Scientific articles that have been published and available online
4. Articles that examine herbal plants can be used to treat halitosis

B. Exclusion Criteria

1. Articles included in systematic reviews, literature review, interview and editorials
2. Articles that cannot be accessed for free

Data Collection

The data that will be used in this research are secondary data. The data is obtained from articles that are searched for in the article database which will then be reviewed according to the research criteria set by the researcher.




Figure 1. A diagram showing the selection of articles for review

The literature search was carried out on the online database, Pubmed, using keywords, namely *herbal plants*, *halitosis*, *children* found 84 articles. After eliminating duplicate articles, the titles and abstracts of each article were analyzed on 84 articles excluding 69 articles. The full-text articles in the remaining 12 articles were analyzed and excluded as many as 34 articles for the following reasons: articles that use chemicals in treating halitosis (n=25) and systematic review articles (n=12).

Table 1. Article about herbal plants as a treatment for halitosis in children

No.	Author	Year	Title	Conclusion
1	Minmin Tian, Anthony Bryan Hanley, Michael W.J. Dodds, Bds, & Ken Yaegaki ¹³	2013	Chewing gum containing allyl isothiocyanate from mustard seed extract is effective in reducing volatile sulfur compounds responsible for oral malodor	Research shows that allyl isothiocyanate from mustard seed extract can effectively reduce the concentration of volatile sulfur compounds in the mouth.
2	Nazliniwaty Nazliniwaty, Lia Laila ¹⁴	2019	Formulation and Antibacterial Activity of Plectranthus amboinicus (Lour.) Spreng Leaves Ethanolic Extract as Herbal Mouthwash Against Halitosis Caused Bacteria	The ethanol extract of the leaves of plectranthus amboinicus (Lour.) Spreng can be formulated as a mouthwash and has antibacterial activity against Staphylococcus aureus and Streptococcus mutans bacteria that cause halitosis.
3	Muneco Tanaka, Masahiro Toe, Hideki Nagata, Miki Ojima, Masae Kuboniwa, Katsumasa Shimizu ¹⁵	2010	Effect of Eucalyptus-Extract Chewing Gum on Oral Malodor: A Double-Masked, Randomized Trial	Chewing eucalyptus extract gum had long-term effects on olganolectic scores, vscs levels, and tongue-coating scores. These findings suggest that eucalyptus extract gum can reduce bad breath by reducing tongue-coating accumulation.
4	Marie-Pierre Morin, Telma Blanca Lombardo Bedran, Jade Fournier-Larente, Bruno Haas, Jabrane Azelmat ¹⁶	2015	Green tea extract and its major constituent epigallocatechin-3-gallate inhibit growth and halitosis-related properties of Solobacterium moorei	Our research carries supporting evidence that the polyphenol in green tea (<i>Camellia sinensis</i>) has a number of properties that may contribute to reducing bacterial halitosis.

5	Abdolhossein Moghbel, Ahmad Farjzadeh, Nasrin Aghel., Homaun Agheli, Nafiseh Raisi ¹⁷	2015	26 Evaluation of the Effect of Green Tea Extract on Mouth Bacterial Activity in the Presence of Propylene Glycol ³⁰	Using green tea extract as a herbal mouthwash is safe and harmless, especially for children. These results suggest that green tea may be able to prevent plaque formation on the teeth so that it can treat halitosis in children.
6	Supanee Rassameemasuang, Pakkarada Phusudsawang, Vanida Sangalungkam ¹⁸	2013	Effect of Green Tea Mouthwash on Oral Malodor	Green tea mouthwash can significantly reduce VSC levels after gargling for 4 weeks.
7	Q c zeng, a z wu and j pika ¹⁹	2010	19 The effect of green tea extract on the removal of sulfur-containing oral malodor volatiles in vitro and its potential application in chewing gum	17 Increasing the pH solution from 7.5 to 8.0 was found to significantly increase the effectiveness of green tea extract for methanethiol removal in vitro. Green tea extract was also found to remove hydrogen sulfide and effectively increase pH in the oral cavity to an alkaline condition.
8	Pier Francesco Porciani, Simone Grandini ²⁰	2012	21 The Effect of Zinc Acetate and Magnolia Bark Extract Added to Chewing Gum on Volatile Sulfur-Containing Compounds in the Oral Cavity	16 Chewing gum containing zinc acetate and magnolia bark extract can significantly reduce oral VSC levels.
9	Kiyoko watanabe, Hiroko H, Toshizo T, Nobuhiro H ²¹	2017	36 Effects of French pine bark extract chewing gum on oral malodor and salivary bacteria	The results showed ²⁴ that French pine bark extract gum was effective in reducing bad breath by reducing the accumulation of tongue-coating and the number of hydrogen sulfide-producing bacteria in saliva.

10	PratibhaMamgain, Abhishek Kandwal, Ravindra K. ²²	2016	<p>11 Comparative Evaluation of Triphala and ElaDecoction With 0.2% Chlorhexidine as Mouthwash in the Treatment of Plaque-Induced Gingivitis and Halitosis:A Randomized Controlled Clinical Trial</p>	Organic triphala and ela decoctions are easy to prepare economically and are just as effective when compared to chlorhexidine mouthwash.
11	Kriti Sharma, Shashidhar Acharya, EshanVerma, Deepak Singhal, NishuSingla ²³	22 2019	Efficacy of chlorhexidine, hydrogen peroxide and tulsi extract mouthwash in reducing halitosis using spectrophotometric analysis: A randomized controlled trial	Tulsi extract may not have the same properties as chlorhexidine and hydrogen peroxide mouthwash. However, tulsi is effective in reducing halitosis, plaque and gingivitis. Apart from that, this Tulsi Extract can be used to treat halitosis due to the lack of side effects, economics and cost effectiveness. Spectrophotometric techniques appear to be promising Methods for evaluation of halitosis
12	AbirMajbauddin, Isamu Kodani and Kazuo Ryoke ²⁴	13 2015	The Effect of Bamboo Leaf Extract Solution and Sodium Copper ChlorophyllinSolution on Growth and Volatile Sulfur Compounds Production of Oral Malodor Associated Some Anaerobic Periodontal Bacteria	High concentrations of both bamboo leaf extract solutions (0.16%) and sodium copper chlorophyllin (0.25%) showed high inhibitory ability in the four anaerobic periodontal bacteria (<i>porphyromonasgingivalis</i> (pg), <i>prevotellainterimidai</i> (pi), <i>fusobacteriumnucleatum</i> (fn) ²⁷ and <i>prevotellanigresce</i> (pn) and produce <i>sulfur volatile compounds</i> that contribute to bad breath.These compounds may have beneficial effects on oral health care.

DISCUSSION

The use of medicinal plants is one of the solutions in solving health problems that are often faced by the community, apart from using chemical drugs in the preventive and medicinal stage.²⁹ The use of medicinal plants has a major impact on the sustainability and biodiversity of plants. According to the World Health Organization (WHO), medicinal plants will be the best source for getting various drugs in a disease. Various formulas are found such as mouthwash, toothpaste and gum derived from plant extracts that are able to treat halitosis. Therefore, these plants must be investigated to determine their properties, safety and efficacy as a treatment for halitosis.^{11,25}

Based on several studies, there are many plants that can be used as medicine to treat halitosis. Research conducted by Minmin Tian, et al, showed that the use of mustard seed extract in the form of chewing gum can effectively reduce the concentration of volatile sulfur compounds in the mouth that play a role in halitosis. This is because the main constituent of mustard seed extract is allyl isothiocyanate (AITC) which has antibacterial properties which will be formed and released through enzymatic reactions of glucosinolates and myrosinase in aqueous solutions. Apart from that, AITCs also exhibit various health benefits such as anti-inflammatory, anticancer and anti-tumor properties.¹³

Hydrogen sulfide, methyl mercaptan and dimethyl sulfide are the main components of halitosis. Where the presence of AITC content in mustard seeds can effectively reduce the VSC concentration in the mouth for a longer time. The reaction of AITC with VSCs occurs through a nucleophilic addition mechanism where the carbon center lacks electrons in the AITC molecule so that it reacts with nucleophilic substances such as hydrogen sulfide to form a ditiocarbamate complex. However, in this study, it was shown that the use of AITC in chewing gum did not affect the taste of the chewing gum and could reduce VSC effectively. In addition, chewing gum with 0.01% AITC and 0.1% zinc lactate can significantly reduce total VSC levels by up to 180 minutes. The effect will be seen when AITC is combined with zinc lactate without affecting sensory stimulation in reducing halitosis.^{13,26}




Figure 1 Release of AITC from chewing gum after 12 and 20 minutes of chewing.

Source: Tian M, Hanley AB, Dodds MWJ. Allyl isothiocyanate from mustard seed is effective in reducing the levels of volatile sulfur compounds responsible for intrinsic oral malodor. *J Breath Res.* 2013;7(2).¹³

Muneo Tanaka et al also mentioned that the use of eucalyptus herbal plant extracts in the gum formula has a long-term effect on the organoleptic score, VSCs levels, and tongue-coating scores that cause halitosis. Eucalyptus globulus is a plant that has many properties to treat various diseases, namely cold medicine, overcoming pulmonary tuberculosis, diabetes,

asthma, disinfectant, antiseptic, antibacterial, antiviral, insect repellent or repellent, and antifungal. The chemical compounds that play a role are 1,8-cineol, α -terpineol, quinate, luteolin, and proanthocyanidin. These findings suggest that eucalyptus extract gum can reduce bad breath by reducing tongue-coating accumulation. So that Eucalyptus has the potential to be developed into a drug derived from natural ingredients.^{15,27}

Research conducted by Pier Francesco et al using chewing gum containing zinc acetate and Magnolia bark extract (MBE) significantly reduced oral VSC levels. Magnolia bark extract (MBE) is a traditional medicine isolated from the bark of *Magnolia officinalis*. This extract is extracted by boiling steam and hot water or by extraction of organic solvents. MBE is usually used by Chinese people for the treatment of fever, headache, pain relief, antibacterial and stress reduction.²⁸

The mechanism for zinc and MBE to reduce oral VSC production is through different routes. Where Zinc acetate when in contact with saliva will release Zn^{2+} ions, which can form stable mercaptides with VSC and its precursors with sulfur, while Magnolia bark extract (MBE) has antibacterial and anti-plaque activity when given in chewing gum or candy. In addition, neolignans from magnolia can also damage bacterial membranes. These considerations suggest a possible synergistic effect of MBE and Zinc acetate to enhance the antihalitotic effect of the active ingredients. Additionally, chewing gum itself may have a mechanical role in reducing plaque by increasing salivary flow. The results of this trial indicated that chewing gum containing zinc acetate and MBE significantly reduced the oral VSC levels at the time of chewing after one hour compared to control chewing gum which did not have this effect.²⁰

Generally, saliva plays an important role in eliminating bad odors in the mouth. Chewing stimulates salivary flow, along with cleansing the oral cavity and reducing bad odors. This latest study shows that chewing gum of various flavors stimulates salivary flow and that chewing for a long time is effective at maintaining salivary flow. In addition, regular use of chewing gum could result in a significant reduction in plaque scores. The microorganisms that cause bad breath are gram-negative, anaerobic bacteria similar to the bacteria that cause periodontitis. In this study, chewing PYC (French pine bark extract) gum for 4 weeks resulted in a significant reduction in oral hydrogen sulfide-producing organisms in saliva compared to baseline. PYC has bacteriostatic activity against various microorganisms including cariogenic and periodontopathic bacteria. Therefore, PYC gum contains an effective concentration to reduce the number of microorganisms that cause halitosis.^{21,29}

In contrast to the research conducted by Nazliniwaty et al, using the leaf extract of *Plectranthusamboinicus* (Lour.) Spreng (EEPL), it can be formulated in the form of a mouthwash that is proven to have antibacterial activity against *Staphylococcus aureus* and *Streptococcus mutans* which causes halitosis. In this study using ethanol as a solvent produced most of the non-volatile compounds. Non-volatile compounds from *Plectranthusamboinicus* (Lour.) Spreng leaves have been reported to have antibacterial activity against gram-positive bacteria such as *Staphylococcus aureus*. In accordance with the results reported by Valgas et al and Gurgelet al, leaf extract of *Plectranthusamboinicus* (Lour.) spreng showed very strong results as an antibacterial because it contains flavonoid compounds in the extract against *Staphylococcus aureus* which causes halitosis.^{14,30}

Several studies have also identified chemical compounds contained in the leaves of *Plectranthusamboinicus* (Lour.) Spreng that play a role in antibacterial activity, namely essential oils such as p-cymene, thymol, β -caryophyllene and γ -terpinene in the leaves of *Plectranthusamboinicus* (Lour.) Spreng which has antibacterial properties against *Staphylococcus aureus*.^{14,31}

Apart from *Staphylococcus aureus* and streptococcus mutans which causes halitosis, *Solobacteriummoorei* is also anaerobic gram-positive bacteria considered as the main causative agent of halitosis. Marie-Pierre Morin et al. Stated that the use of green tea extract containing polyphenols can reduce halitosis associated with *Solobacteriummoorei* bacteria. In this study, researchers found the active substance associated with green tea polyphenols, namely the presence of EGCG (constituent epigallocatechin-3-gallate) which has bactericidal properties to control halitosis. This suggests that green tea extract contains additional bioactive components. Green tea extract and EGCG can absorb cell membranes resulting in leakage of calcein-AM. Previous studies have also reported that green tea also contains theaflavins and catechins which can play a role in permanently damaging the cytoplasmic membrane of bacteria.^{16,20,32}




Figure 2. Transmission electron microscopy analysis of *S. moorei*. Panel A: Control untreated bacteria. Panel B: Bacteria treated (4 h) with green tea extract (500 µg/ml). Panels C and D: Bacteria treated (4 h) with EGCG (250 µg/ml)

Source : (Morin MP, Bedran TBL, Fournier-Larente J, Haas B, Azelma¹⁸, Grenier D. Green tea extract and its major constituent epigallocatechin-3-gallate inhibit growth and halitosis-related properties of *Solobacterium moorei*. BMC Complement Altern Med [Internet]. 2015;15(1):1–11)¹⁶

The formation of biofilms by *S. moorei* is the main step in the occurrence of halitosis. Although the polyphenols in green tea do not show specific anti-biofilm effects in *S. moorei*, they can affect biofilms through two mechanisms. First, the biofilm formation by *S. moorei* on EGCG treated surfaces was particularly affected. Second, green tea extract and EGCG can reduce the previously formed *S. moorei* biofilm. The mechanism involved in biofilm

desorption may be related to ⁶ the ability of green tea polyphenols to alter the integrity of *S. moorei* cells.^{16,33,34}

In previous studies, it has also been shown that *S. moorei* produces high VSC levels when the presence of cysteine and β -galactosidase are associated with VSC production from mucin. This enzyme can deglycosylate salivary glycoproteins, which can then be degraded into peptides and amino acids by host proteolytic enzymes and bacteria before being converted into VSC. So that β -galactosidase activity in saliva has been associated with halitosis. In this study, showed that green tea extract and EGCG can inhibit β -galactosidase activity in *S. moorei* which can contribute to reducing halitosis.^{32,35,36,37}

This study is also in line with the research by Abdolhosseinmoghbel et al using green tea extract to treat halitosis in children. Green tea contains flavonoids, tannins, vitamins, fluorides and other mineral salts. Some of the antioxidants and antimicrobial agents of green tea can increase strength in teeth. Tannins are biosynthetic materials that have a strong antibacterial effect. In this study using propylene glycol (PG) as a solvent rather than ethyl alcohol. In Yamamoto et al's research on green tea in the United States, the use of green tea containing 0.5% tannins with 2% ethyl alcohol solvent reduced various types of staphylococcus by only about 15%. But in Iranian the use of PG 10%, green tea mouthwash 1% and free tannins can decrease the number of bacteria by about 48%. In addition, using PG as a substitute for hazardous ethyl alcohol is not only useful as a co-solvent but also as a chemical stabilizer and causes microbial death and is safe for use for children. This proves that propylene glycol is better used than using ethyl alcohol.^{17,38,39,40}

Studies show that toothpaste and mouthwash containing green tea can fight bacteria and prevent plaque formation. Plaque is the cause of bad breath. Green tea can prevent bad breath by consuming mouthwash before and after brushing your teeth, or mixing it with toothpaste products. A herbal mouthwash formulation from Iranian green tea extract containing 1% tannin with 10% propylene glycol can reduce the number of aerobic oral bacteria by 45-64% and can also prevent plaque formation on teeth that results in halitosis.^{17,41,42}

Supanee et al. also mentioned the ability of green tea mouthwash to reduce bad breath caused by the antimicrobial activity of green teacatechins, especially in *Porphyromonasgingivalis*, as well as the ability of green tea catechins to convert VSCs into odorless compounds. It has been shown that active catechins and their derivatives can inhibit tissue damage through *P. gingivalis* proteinase activity, thus, catechins and their derivatives can reduce the source of microbial spoilage. According to this study, green tea mouthwash showed a comparable percentage of VSC reduction to mouthwashes containing 0.05% chlorhexidine, 0.05% cetylpyridinium chloride, and 0.14% zinc-lactate.^{43,44,45,46}




Figure 3: Level of volatile sulfur compounds before and after mouthwash usage: * significantly different compared to baseline ($P < 0.05$) and † significantly different compared to placebo ($P < 0.05$).

Source : (Porciani PF, Grandini S. Effect of green tea-added tablets on volatile sulfur-containing compounds in the oral cavity. J Clin Dent. 2016;27(4):110–3).²⁰

In the study of Pratibhamamgain et al. using triphala and organic ela decoction was effective when compared with chlorhexidine mouthwash in inhibiting halitosis. Triphala has antimicrobial properties and inhibits the growth of activity against bacterial isolates, besides that it is also shown to have antioxidant and free radical reduction properties. Ela churna in combination with Triphala decoction to treat halitosis has an immunomodulatory, and antimicrobial action against the organisms that cause dental caries. Triphala is efficient in inhibiting plaque biofilm and effective in treating biofilm infections present in teeth. Similar results were seen in this study where the plaque index increased for the Triphala and Ela groups at baseline 14 and 21 days. Triphala concentration was 0.6% compared to chlorhexidine concentration of 0.1% and both were efficient in controlling gingivitis and plaque biofilm formation. Also, it appears that Triphala and chlorhexidine show the same inhibitory effect on microbial counts, except for *Lactobacillus*, where Triphala has shown better results than chlorhexidine.^{22,47,48}

In addition, *Ocimum sanctum* (Tulsi) pharmacological activity has spread in the field of dentistry. In this study, tulsi mouthwash was formulated and had an effect on bad odor assessed after 15 days. A study has analyzed the effect of various concentrations of *Ocimum sanctum* extract ranging from 0.5 to 10%, and it was observed that a 4% extract concentration was optimal as an antibacterial agent against the normal flora of the oral cavity; thus, in this study using a concentration of 4%. *Ocimum sanctum* has been tested against various microorganisms such as *Candida albicans*, *Staphylococcus aureus*, enteric pathogens, *Klebsiella*, *Escherichia coli* and *Proteus*.^{49,50}

In this study, *Ocimum sanctum* extract mouthwash prevented plaque formation. In addition, the stems and leaves of *Ocimum sanctum* also contain various kinds of active ingredients that have antibacterial properties, including saponins, flavonoids, triterpenoids and tannins which form high molecular complexes, lyse bacteria on the tooth surface and saliva. This study demonstrated a significant reduction in VSCs, organoleptic, plaque and gingival scores in the tulsi extract mouthwash group, which could be attributed to the compounds isolated from the tulsi extract. However, after 15 days of intervention, the mean

percentage reduction in VSC and organoleptic scores in the chlorhexidine and hydrogen peroxide groups was significantly greater than in the tulsi mouthwash group. Therefore, it can be concluded that tulsi mouthwash, although effective in reducing VSC which is responsible for halitosis, but does not have the same efficacy against chlorhexidine and hydrogen peroxide mouthwash.^{23,51,52,53,54,55}

In contrast to the case of bamboo leaf extract which has components such as beta carotene, alpha tocopherol, gamma tocopherol which has strong anti-bacterial activity properties. Apart from that, the phenolic compounds possessed by bamboo leaves themselves are also proven to have a strong bactericidal action. The highest concentration of BLES (Bamboo leaf extract solution) in combination with SCCS (sodium copper chlorophyllin solution) was able to inhibit the growth and proliferation of four bacteria in a concentration-dependent test. BLES and SCCS certainly have an inhibitory effect on the growth and proliferation of anaerobic bacteria against high concentrations of bacteria showing superior susceptibility. In this context, high concentrations of both bamboo leaf extract solution (0.16%) and sodium copper chlorophyllin (0.25%) showed high inhibitory ability in the four anaerobic periodontal bacteria (*porphyromonasgingivalis* (PG), *prevotellainterimidai* (PI), *fusobacteriumnucleatum* (FN) and *prevotellanigracence* (PN) and produce sulfur volatile compounds that contribute to bad breath. These compounds may have a beneficial effect on oral health care. New antibacterial agents fight the bacteria that cause halitosis and related oral diseases.^{24,56,57}




Figure 4. Growth rate of four strains after 1 day incubation, followed by additional 3 days incubation with 0.16% BLES or 0.25% SCCS. The values **P < 0.001 and *P < 0.01 are shown with 1 day incubation.

Source : (Majbauddin A, Kodani I, Ryoke K. The effect of bamboo leaf extract solution and sodium copper chlorophyllin solution on growth and volatile sulfur compounds production of oral malodor associated some anaerobic periodontal bacteria. *Yonago Acta Med.* 2015;58(3):129–36)²⁴

CONCLUSION

Recent literature shows that herbal plants that have been described have various kinds of active substances that can be the right solution in treating halitosis. This herbal plant can be formulated in preparations as mouthwash, toothpaste, chewing gum or even other products

that can reduce halitosis in children, so this systematic review can be used as a reference in the development of herbal plants in the future, especially dentistry.

REFERENCES

1. Choudhary A, Aggarwal A, Kishore M. Halitosis- A Review REVIEW ARTICLE. 2012;(March).1-5
2. Irianti R, Pandelaki K, Mintjelungan C. Description of Knowledge About Halitosis in Workers at the Port of Manado. 2015;3(1).
3. Yoga IGKM, Giri P ratna K, Sarjana K. BA L I D E N T A L JOUR NAL Acrylic release to the elderly in Penatahan Village. Bali Dent J [Internet]. 2018;2(1):45–52. Available from: <http://jkg-udayana.org/ojs/index.php/bdj/article/view/135>
4. Senjaya AA. TREATMENT OF HALITOSIS Asep Arifin Senjaya 1. Scale Husada. 2011;8(2):126–31.
5. Aylikci B, Çolak H. Halitosis: From diagnosis to management. J Nat Sci Biol Med. 2013;4(1):14–23.
6. Sri Hernawati, Balqis Salsabila Setya Aldianah, Pujiana Endah AI. The Effectiveness of Red Pomegranate (*Punica granatum* Linn) Extract Mouthwash Against the Number of Oral Bacteria Colony. Malaysian J Med Heal. 2020;16(4):2636-9346,.
7. Astrid Wylleman 1, Feie Vuylsteke 2, Christel Dekeyser 3, Wim Teughels 4, Marc Quirynen 4 IL. Alternative therapies in controlling oral malodour: a systematic review. J Breath Res [Internet]. 2020; Available from: <https://pubmed.ncbi.nlm.nih.gov/33227726/>
8. Care PD, Karimi M. The Causes of Holitosis in Children. Pediatr Dent Care. 2017;2(1):2–4.
9. Yulimatussa AP, Blambangan BGPB, Dewi JC, Herdianto RS, Mumtaza I, Nafii MM, et al. Knowledge of Handling Halitosis in Oral Health Problems. J Farm Community. 2016;3(2):28–32.
10. Torsten M, Gómez-Moreno G, Aguilar-Salvatierra A. Drug-related oral malodour (halitosis): a literature review. Eur Rev Med Pharmacol Sci. 2017;21(21):4930–4.
11. Khan DMA. Introduction and Importance of Medicinal Plants and Herbs. In 2016. Available from: https://www.nhp.gov.in/introduction-and-importance-of-medicinal-plants-and-herbs_mtl
12. Irfan I, Rochmah YS, Yusuf M, Aditya G. The Effectiveness of Gambir Leaves (*Uncaria Gambir* Roxb) To Reduce Halitosis Caused by Study Plaques at the Zuhriyah Orphanage and Islamic Boarding School, Sleman, Yogyakarta. ODONTO Dent J. 2015;2(1):52.
13. Tian M, Hanley AB, Dodds MWJ. Allyl isothiocyanate from mustard seed is effective in reducing the levels of volatile sulfur compounds responsible for intrinsic oral malodor. J Breath Res. 2013;7(2).
14. Nazliniwaty N, Laila L. Formulation and antibacterial activity of plectranthus amboinicus (Lour.) spreng leaves ethanolic extract as herbal mouthwash against halitosis caused bacteria. Open Access Maced J Med Sci. 2019;7(22):3900–3.
15. Tanaka M, Toe M, Nagata H, Ojima M, Kuboniwa M, Shimizu K, et al. Effect of Eucalyptus-Extract Chewing Gum on Oral Malodor: A Double-Masked, Randomized

- Trial. *J Periodontol.* 2010;81(11):1564–71.
16. Morin MP, Bedran TBL, Fournier-Larente J, Haas B, Azelmat J, Grenier D. Green tea extract and its major constituent epigallocatechin-3-gallate inhibit growth and halitosis-related properties of *Solobacterium moorei*. *BMC Complement Altern Med [Internet]*. 2015;15(1):1–11.
 17. Széliga ME, Nacucchio MC. Pharmaceutical products. *Pharm Policy Law.* 2015;17(1–2):81–9.
 18. Rassameemasuang S, Phusudsawang P, Sangalungkarn V. Effect of Green Tea Mouthwash on Oral Malodor. *ISRN Prev Med.* 2013;2013:1–6.
 19. Zeng QC, Wu AZ, Pika J. The effect of green tea extract on the removal of sulfur-containing oral malodor volatiles in vitro and its potential application in chewing gum. *J Breath Res.* 2010;4(3).
 20. Porciani PF, Grandini S. Effect of green tea-added tablets on volatile sulfur-containing compounds in the oral cavity. *J Clin Dent.* 2016;27(4):110–3.
 21. Watanabe K, Hiramine H, Toyama T, Hamada N. Effects of French pine bark extract chewing gum on oral malodor and salivary bacteria. *J Nutr Sci Vitaminol (Tokyo).* 2018;64(3):185–91.
 22. Mamgain P, Kandwal A, Mamgain RK. Comparative Evaluation of Triphala and Ela Decoction With 0.2% Chlorhexidine as Mouthwash in the Treatment of Plaque-Induced Gingivitis and Halitosis: A Randomized Controlled Clinical Trial. *J Evidence-Based Complement Altern Med.* 2017;22(3):468–72.
 23. Sharma K, Acharya S, Verma E, Singhal D, Singla N. Efficacy of chlorhexidine, hydrogen peroxide and tulsi extract mouthwash in reducing halitosis using spectrophotometric analysis: A randomized controlled trial. *J Clin Exp Dent.* 2019;11(5):e457–63.
 24. Majbauddin A, Kodani I, Ryoke K. The effect of bamboo leaf extract solution and sodium copper chlorophyllin solution on growth and volatile sulfur compounds production of oral malodor associated some anaerobic periodontal bacteria. *Yonago Acta Med.* 2015;58(3):129–36.
 25. Ranjan Malhotra 1, Vishakha Grover, Anoop Kapoor DS. Comparison of the effectiveness of a commercially available herbal mouthrinse with chlorhexidine gluconate at the clinical and patient level. *J Indian Soc Periodontol .* 2011;15(4):349–52.
 26. Francisco Wilker Mustafa Gomes Muniz1, 1, Stephanie Anagnostopoulos Friedrich1 CFS and CKR. The impact of chewing gum on halitosis parameters: a systematic review. *J Breath Res [Internet]*. 11(1):1–5. Available from: <https://iopscience.iop.org/article/10.1088/1752-7163/aa5cc2/meta>
 27. Koswandy LF, Ramadhania ZM. Chemical Compounds Content and Bioactivity of *Eucalyptus globulus* Labill. 2017;24(2):63–78.
 28. Greenberg M, Urnezis P, Tian M. Compressed mints and chewing gum containing magnolia bark extract are effective against bacteria responsible for oral malodor. *J Agric Food Chem.* 2007;55(23):9465–9.
 29. Rohdewald P. A review of the French Maritime Pine Bark Extract (Pycnogenol), a herbal medication with a diverse clinical pharmacology. *Int J Clin Pharmacol Ther.*

- 2002;40(4):158–62.
30. Silalahi M. *Plectranthus Amboinicus* (Lour.) Spreng as a Food and Drug Material and its Bioactivity. *J Din Educator*. 2018;11(2):123.
 31. Patel R, Alam G, M N, PS M, S A, S S. *Plectranthus Amboinicus* (Lour) Spreng : an Overview. *Pharma Res*. 2010;4(December 2010):1–15.
 32. Widyaningrum N, Sumarwati T, Sukryana WS. Effect of vitamin C and vitamin E on the stability of EGCG (epigallocatechin gallate) in the ethyl acetate fraction of green tea leaf extract (*Camellia sinensis* L.). *Journal of Pharmaceutical Sciences*. 2019;15(2):80–5.
 33. Talebi Ardakani MR, Golmohammadi S, Ayremlou S, Taheri S, Daneshvar S, Meimandi M. Antibacterial effect of Iranian green-tea-containing mouthrinse vs chlorhexidine 0.2%: An in vitro study. *Oral Health Prev Dent*. 2014;12(2):157–62.
 34. Shaik JA, Reddy RK. Review Article Prevention and Treatment of White Spot Lesions in Orthodontic Patients. *Contemp Clin Dent*. 2017;8(September):11–9.
 35. Ahmad Muhamud NAS, Amran S. Screening of Epigallocatechin-3-Gallate (EGCG) Content in *Camellia sinensis* Products. *Nat Prod Chem Res*. 2018;06(05).
 36. Priya BM, Anitha V, Shanmugam M, Ashwath B, Sylva SD, Vigneshwari SK. Efficacy of chlorhexidine and green tea mouthwashes in the management of dental plaque-induced gingivitis: A comparative clinical study. *Contemp Clin Dent*. 2015;6(4):505–9.
 37. Vilela MM, Salvador SL, Teixeira IGL, Del Arco MCG, De Rossi A. Efficacy of green tea and its extract, epigallocatechin-3-gallate, in the reduction of cariogenic microbiota in children: a randomized clinical trial. *Arch Oral Biol* [Internet]. 2020;114(April):104727. Available from: <https://doi.org/10.1016/j.archoralbio.2020.104727>
 38. Khurshid Z, Zafar MS, Zohaib S, Najeeb S, Naseem M. Green Tea (*Camellia Sinensis*): Chemistry and Oral Health. *Open Dent J*. 2016;10(1):166–73.
 39. Fajriani F, Andriani JN. Reduction of Salivary *Streptococcus mutans* Colonies in Children After Rinsing with 2.5% Green Tea Solution. *J Dent Indones*. 2015;21(3):79–84.
 40. Alqutami J, Elger W, Grafe N, Hiemisch A, Kiess W, Hirsch C. Dental health, halitosis and mouth breathing in 10-to-15 year old children: A potential connection. *Eur J Paediatr Dent*. 2019;20(4):274–9.
 41. Shashibhushan Kukkalli Kamalaksharappa, Ramya Rai, Prashant Babaji MP. Efficacy of probiotic and green tea mouthrinse on salivary pH. *J indian Soc Pedod Prev Dent*. 2018;36(3):279–82.
 42. Chaitali U. Hambire, Rashmi Jawade, [...] and PBN. Comparing the antiplaque efficacy of 0.5% *Camellia sinensis* extract, 0.05% sodium fluoride, and 0.2% chlorhexidine gluconate mouthwash in children. *J Int Soc Prev Community Dent*. 2015;5(3):218–26.
 43. Simaremare RT, Hasny, Lusian Y. Effectiveness of Mouthwash with Steeping Green Tea and Listerine Solution Against Ohi-S in Students / I Class Viii Bsmpt Smart Private Nation Jl. Titi Kuning Namorambe. *PANNMED*. 2014;9(1):32–5.
 44. L. SS, Kawengian SES, Mariati NW. The Effectiveness of Gargling With Green Tea

- Brewed Water In Reducing Plaque Accumulation. 2015;3(2).
45. Armidin RP, Yanti GN. Effectiveness of rinsing black tea compared to green tea in decreasing streptococcus mutans. *Open Access Maced J Med Sci.* 2019;7(22):3799–802.
 46. Jazaeri M, Pakdek F, Rezaei-Soufi L, Abdolsamadi H, Rafieian N. Cariostatic Effect of Green Tea in Comparison with Common Anticariogenic Agents: An in Vitro Study. *J Dent Res Dent Clin Dent Prospects.* 2015;9(1):44–8.
 47. Srinagesh J. Antibacterial efficacy of triphala against oral streptococci: An in vivo study. *Indian J Dent Res Off Publ Indian Soc Dent Res.* 2012;23(5):696–9.
 48. Salama MT, Alsughier ZA. Effect of Green Tea Extract Mouthwash on Salivary Streptococcus mutans Counts in a Group of Preschool Children: An In vivo Study. *Int J Clin Pediatr Dent.* 2019;12(2):133–8.
 49. M. S, KR S, B. S, G. V, S. R, K. S, et al. Ocimum sanctum: a review on the pharmacological properties. *Int J Basic Clin Pharmacol.* 2016;(January):558–65.
 50. Kaur H, Jain S, Kaur A. Comparative evaluation of the antiplaque effectiveness of green tea catechin mouthwash with chlorhexidine gluconate. *J Indian Soc Periodontol.* 2014;18(2):178–82.
 51. Pandey G, Madhuri S. Pharmacological activities of Ocimum sanctum (Tulsi): A review. *Int J Pharm Sci Rev Res.* 2010;5(1):61–6.
 52. Ningrum WA, Waznah U. Mouthwash Formulation Ethanol Extract of Basil Leaves (*Ocimum basilicum L.*). *Journal Pharmacy.* 2018;2(2):159–66.
 53. Achmad H, Thahir H, Rieuwpassa I, Mardiana AA, Oktawati S, Samad R, Djais AI, Gani A, Singgih MF, Madjid F, Admy SC. The Effectiveness of *Channa striata* Extract Antimicrobial Effect on Periopathogen Bacteria (*Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*). *Systematic Reviews in Pharmacy.* 2020; 11(4): 319-323. doi: [10.31838/srp.2020.4.46](https://doi.org/10.31838/srp.2020.4.46)
 54. Achmad H, Adam AM, Asalui TR, Huldani, Sukmana BI, Putra AP. Use of Sea Cucumber Extract As An Alternative Treatment of Inflammation by Chronic Periodontitis. *International Journal of Pharmaceutical Research.* October-December, 2020. 12(4):3567-3575.
 55. Achmad H, Atjo NA, Arnianti, Lindriani, Ermawati, Mahmud Y, Arniyanti A, Hasanuddin A. The Use of Children's Herbal Mouthwash in Pediatric Dentistry: A Systematic Review. *International Journal of Phamaceutical Research.* July-December, 2020. Volume 12 Supplementary Issue 2: 4508-4515.
 56. Achmad H, Djais AI, Syahrir S, Fitri A, Ramadhany YF. A literature us regarding the use of herbal medicines in pediatric dentistry. *International Journal of Pharmaceutical Research.* 2020. 12,PP. 881-897.
 57. Achmad H, Djais AI, Syahrir S, Fitri A, Ramadhany YF. Impact Covid-19 in pediatric dentistry: A literature review. *International Journal of Pharmaceutical Research,* 2020. 12,p.830-840.

Herbal Plants as a Treatment for Halitosis in Children: A Systematic Review

ORIGINALITY REPORT

20%
SIMILARITY INDEX

15%
INTERNET SOURCES

14%
PUBLICATIONS

10%
STUDENT PAPERS

PRIMARY SOURCES

1 ijpsr.com 1%
Internet Source

2 Submitted to Institute of Graduate Studies,
UiTM 1%
Student Paper

3 Minmin Tian, A Bryan Hanley, Michael W J
Dodds. "Allyl isothiocyanate from mustard
seed is effective in reducing the levels of
volatile sulfur compounds responsible for
intrinsic oral malodor", Journal of Breath
Research, 2013 1%
Publication

4 motorrijder.info 1%
Internet Source

5 Michael Greenberg, Philip Urnezis, Minmin
Tian. "Compressed Mints and Chewing Gum
Containing Magnolia Bark Extract Are
Effective against Bacteria Responsible for Oral
Malodor", Journal of Agricultural and Food
Chemistry, 2007 1%

6

www.science.gov

Internet Source

1 %

7

Marcela Leticia Leal Gonçalves, Ana Carolina Costa da Mota, Alessandro Melo Deana, Lisyanne Araújo de Souza Cavalcante et al. "Antimicrobial photodynamic therapy with Bixa orellana extract and blue LED in the reduction of halitosis—A randomized, controlled clinical trial", *Photodiagnosis and Photodynamic Therapy*, 2020

Publication

1 %

8

Duaa Alsaffar, Hamad Alzoman. "Efficacy of antioxidant mouthwash in the reduction of halitosis: A randomized, double blind, controlled crossover clinical trial", *Journal of Dental Sciences*, 2020

Publication

1 %

9

Submitted to Universitas Hasanuddin

Student Paper

1 %

10

Submitted to Universitas Siliwangi

Student Paper

1 %

11

www.mysciencework.com

Internet Source

1 %

12

ir.amu.ac.in

Internet Source

1 %

13	www.longdom.org Internet Source	1 %
14	www.dovepress.com Internet Source	1 %
15	www.pubfacts.com Internet Source	1 %
16	lib.bioinfo.pl Internet Source	<1 %
17	Submitted to University of Oklahoma Health Science Center Student Paper	<1 %
18	Stefanie J. Gartenmann, Severine L. Steppacher, Yves von Weydlich, Christian Heumann, Thomas Attin, Patrick R. Schmidlin. "The Effect of Green Tea on plaque and gingival inflammation: A systematic review", Journal of Herbal Medicine, 2020 Publication	<1 %
19	Mireille Andriamihaja, Annaïg Lan, Martin Beaumont, Marta Grauso et al. "Proanthocyanidin-containing polyphenol extracts from fruits prevent the inhibitory effect of hydrogen sulfide on human colonocyte oxygen consumption", Amino Acids, 2018 Publication	<1 %

20 N Fujiwara, K Murakami, M Nakao, M Toguchi, H Yumoto, T Amoh, K Hirota, T Matsuo, S Sano, K Ozaki, Y Miyake. "Novel reuterin-related compounds suppress odour by periodontopathic bacteria", Oral Diseases, 2017
Publication

21 184.106.70.54
Internet Source

22 eprints.manipal.edu
Internet Source

23 benthamopen.com
Internet Source

24 www.jstage.jst.go.jp
Internet Source

25 Shin-ichi Tanabe, Daniel Grenier. "Characterization of volatile sulfur compound production by Solobacterium moorei", Archives of Oral Biology, 2012
Publication

26 dentjods.sums.ac.ir
Internet Source

27 www.frontiersin.org
Internet Source

28 academic.oup.com
Internet Source

29

www.ncbi.nlm.nih.gov

Internet Source

<1 %

30

blog.englishteastore.com

Internet Source

<1 %

31

Alessandro Di Cerbo, Federica Pezzuto, Sergio Canello, Gianandrea Guidetti, Beniamino Palmieri. "Therapeutic Effectiveness of a Dietary Supplement for Management of Halitosis in Dogs", *Journal of Visualized Experiments*, 2015

Publication

<1 %

32

Koduru Sravani, A. Suchetha, Darshan B. Mundinamane, Divya Bhat, Nanditha Chandran, H. R. Rajeshwari. "Plant Products in Dental and Periodontal Disease: An overview", *International Journal of Medical and Dental Sciences*, 2015

Publication

<1 %

33

[Submitted to RMIT University](#)

Student Paper

<1 %

34

Yuan Li, Xiaoge Jiang, Jianqi Hao, Yifei Zhang, Ruijie Huang. "Tea polyphenols: application in the control of oral microorganism infectious diseases", *Archives of Oral Biology*, 2019

Publication

<1 %

35

doaj.org

Internet Source

<1 %

36	pubmed.ncbi.nlm.nih.gov Internet Source	<1 %
37	www.amjdent.com Internet Source	<1 %
38	"Chewing gum does not increase salivary flow in older adults", Springer Healthcare News, 2012 Publication	<1 %
39	FDI World Dental Federation. "FREE COMMUNICATIONS SESSIONS 33-46 Friday, August 30, 2013", International Dental Journal, 2013 Publication	<1 %
40	Marc Quirynten, Hong Zhao, Daniel van Steenberghe. "Review of the treatment strategies for oral malodour", Clinical Oral Investigations, 2002 Publication	<1 %
41	journals.plos.org Internet Source	<1 %
42	www.degruyter.com Internet Source	<1 %
43	zombiedoc.com Internet Source	<1 %
44	Isabelle Laleman, Wim Teughels. "Novel natural product - based oral topical rinses	<1 %

and toothpastes to prevent periodontal diseases", Periodontology 2000, 2020

Publication

45

María José Cires, Paola Navarrete, Edgar Pastene, Catalina Carrasco-Pozo et al. "Effect of a proanthocyanidin-rich polyphenol extract from avocado on the production of amino acid-derived bacterial metabolites and the microbiota composition in rats fed a high-protein diet", Food & Function, 2019

Publication

<1 %

46

Nao Suzuki, Masahiro Yoneda, Kazuto Haruna, Yosuke Masuo et al. "Effects of S-PRG eluate on oral biofilm and oral malodor", Archives of Oral Biology, 2014

Publication

<1 %

47

ijrhas.com

Internet Source

<1 %

48

www.mdpi.com

Internet Source

<1 %

49

www.scielo.br

Internet Source

<1 %

50

Damla Aksit Bicak. "A Current Approach to Halitosis and Oral Malodor- A Mini Review", The Open Dentistry Journal, 2018

Publication

<1 %

51

Lianet Monzote, Alexander M. Scherbakov, Ramón Scull, Yamilet I. Gutiérrez et al. "Pharmacological Assessment of the Carvacrol Chemotype Essential Oil From Growing in Cuba ", Natural Product Communications, 2020

Publication

<1 %

52

Radafshar, Golpar, Mahshid Ghotbizadeh, Farzad Saadat, and Nastaran Mirfarhadi. "Effects of green tea (Camellia sinensis) mouthwash containing 1% tannin on dental plaque and chronic gingivitis: a double-blinded, randomized, controlled trial", Journal of Investigative and Clinical Dentistry, 2015.

Publication

<1 %

53

annalsofrscb.ro

Internet Source

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On