

# Differences of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of Moringa Leaf Extract (Moringa Oliefera L.) on Bacteria Aggregatibacter Actinomycetemcomitans and P

**Submission date:** 04-Jun-2021 07:16AM (UTC+0700) *by*

**Submission ID:** 1599972128

**File name:** DIFFER\_1.PDF (1.67M)

**Word count:** 2278

**Character count:** 13697

# Differences of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of Moringa Leaf Extract (*Moringa Oleifera L.*) on Bacteria *Aggregatibacter Actinomycetemcomitans* and *Porphyromonas Gingivalis*

Arni Irawaty Djais<sup>1</sup>, Hasanuddin Tahir<sup>1</sup>, Mochammad Hatta<sup>2</sup>, Harun Achmad<sup>3</sup>, Ainul Wahyuni<sup>4</sup>

<sup>1</sup>Department of Periodontology, Hasanuddin University, Faculty of Dentistry, Hasanuddin University, Indonesia,

<sup>2</sup>Department of Microbiology, Molecular Biology and Immunology Laboratory, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia, <sup>3</sup>Department of Pedodontics, Hasanuddin University, Faculty of Dentistry,

Hasanuddin University, Indonesia, <sup>4</sup>Dental Hospital, Hasanuddin University, Faculty of Dentistry, Hasanuddin University, Indonesia

## Abstract

**Background:** Moringa leaves (*Moringa oleifera L.*) has many nutrients that contain bioactive components such as tannins, flavonoids, and saponins, as antimicrobials. The growth of *Aggregatibacter actinomycetemcomitans* and *Porphyromonas Gingivalis* may be inhibited so that they do not become pathogens and cause periodontitis. **Objective:** The general purpose of this study was to determine the Minimum Inhibitory Concentration (MIC) and Minimal Bactericidal Concentration (MBC) and inhibitory power of moringa leaf extract (*Moringa oleifera*) against bacteria *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*. **Method:** The type of research used was laboratory experimental research. The design of this study is *post test only control group design* using the Kirby Bauer dilution method. With treatment extract concentration of 15%, 20%, 25%, 30%, 35% 100%, and positive control (Metronidazole). The measuring instrument in this study uses a caliper with millimeters (mm). **Results:** The results of the study of bacteria *Porphyromonas gingivalis* obtained from MIC by 20%, MBC 25% and bacteria *Aggregatibacter actinomycetemcomitans* obtained by MIC by 30% and MBC 35%. Kruskal Wallis test results showed that the value of  $p < 0.05$  so it can be concluded that there are significant differences in the inhibition zone of *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*. **Conclusion:** The extract of leaves of Moringa (*Moringa oleifera L.*) were able to inhibit bacterial growth *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*.

**Keywords:** Moringa leaves extract (*Moringa oleifera L.*), inhibitory power, bacteria *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*.

## Introduction

Periodontal disease is one of the most common dental and oral health problems in the community. Based on Indonesia's health profile in 2011, in 2010 there were 92,979 people who visited public hospitals belonging to the health ministry and local governments due to

periodontal disease. Periodontal disease is an infection in the oral cavity that affects periodontal tissues. The main cause of this disease is microorganisms that colonize the surface of the tooth (bacterial plaque). Microorganism culture (bacteria) found in plaque shows the presence of certain gram negative bacteria in periodontitis.<sup>1,2,3</sup>

Periodontitis is an inflammation that affects the supporting tissues of a tooth, caused by microorganisms and can cause progressive damage to the periodontal ligament, alveolar bone and is accompanied by pocket formation.<sup>4</sup> Periodontitis causes permanent tissue

### Corresponding author :

Harun Achmad,

Department of Pedodontic, Hasanuddin University,  
Faculty of Dentistry, Hasanuddin University, Indonesia

destruction characterized by chronic inflammation, migration of the fused epithelium to the apical, loss of connective tissue and loss of alveolar bone.<sup>5</sup>

The main causes of periodontal disease are the presence of microorganisms that colonize the dental plaque. Dental plaque is a structured, soft, yellow substance, which is attached to the tooth surface. The content of dental plaque is various types of microorganisms, especially the remaining bacteria are fungi, protozoa and viruses. Plaque containing pathogenic microorganisms plays an important role in causing and exacerbating periodontal infections.<sup>6</sup> An increasing number of gram-negative organisms in subgingival plaques such as *Porphyromonas gingivalis*, *Actinobacillus actinomycetemcomitans*, *Tannerella forsythia* and *Treponema denticola* infect periodontal infection.<sup>7</sup> *Porphyromonas gingivalis* is a gram-negative anaerobic bacteria involved in the pathogenesis of periodontitis and other inflammatory diseases that destroy dental support tissue. These bacteria can invade the periodontal tissues locally and survive the host defense mechanism by utilizing a panel of virulence factors that cause deregulation of innate immune and inflammatory responses.<sup>8</sup> *Aggregatibacter actinomycetemcomitans* is a gram negative bacterium found in the oral cavity and one of the etiologies of aggressive periodontitis. This bacterium has the ability to produce leukotoxins which can cause damage to the periodontal tissues and one type of bacteria that is considered a periodontal pathogen.<sup>9</sup> Inappropriate and excessive use of antibiotics can result in bacteria being *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis* resistant to antibiotic drugs that have been given. The resistance of these bacteria to antibiotic drugs allows the use of herbal medicines from natural ingredients to be one of the other alternatives in the treatment of periodontitis. The use of herbal medicines from natural ingredients is generally considered safer than the use of modern medicine, because herbal medicines as traditional medicines have relatively fewer side effects than modern drugs.<sup>1</sup>

Natural ingredients used as herbal medicines one of which is Moringa leaves or known as the Latin name *Moringa Oleifera*. Moringa plants are efficacious as anti-cancer, anti-bacterial, hypotensive, inhibiting the activity of bacteria and fungi. This is related to the chemical content contained in it, which is rich in vitamin A and vitamin C, gluconic compounds and isothiocinates.<sup>10</sup>

Kelor leaves contain phytochemicals that make this plant capable of carrying out self-defense mechanisms. Phytochemicals contained include catechol tannins, galia tannins, steroids, triterpenoids, flavonoids, saponins, anthraquinones, alkaloids, and reducing sugars.<sup>11,12</sup> The compound has the ability as a drug, its benefits are as antibiotics, skin care, anti-inflammatory, blood pressure, diabetes, and anemia.

Based on the description above, researchers are interested in conducting research to find out "The effect of Moringa leaf extract on bacterial growth, *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis* cause periodontitis.

## Materials and Method

Type of research used in this study was a laboratory experimental study. The design of this study is *post test only control group design* using methods agar dilution and diffusion. This dilution method is used to determine the MIC (Minimum Inhibition Concentration) value, the lowest concentration that can inhibit the growth of test microbes. The dilution method is done by mixing the sample, the microbial test, and the inoculation media with several variations of dilution. The diffusion method to be used to determine the activity of antimicrobial agents or often also called the inhibitory test. This method uses disk paper that has contained Moringa leaf extract and then put it into a culture medium.

The Kirby Bauer method was carried out in observing certain inhibitory zone diameters and producing a good batch-to-batch, resulting in satisfactory growth of the most pathogenic bacteria. The treatment was carried out 4 times with a concentration of 15%, 20%, 25%, 30%, 35% 100%. The positive control used is Metronidazole. The research tools used are petri dishes, round oases, autoclaves, bunsen, erlenmeyer flasks, suction pipettes, filter devices, rotary evaporators, filter paper, sterile cotton swabs, stationery.

The materials used in the research were (*Moringa oleifera L.*), test bacteria *Aggregatibacter actinomycetemcomitans* and *Porphyromonas gingivalis*. Obtained from the Laboratory of Hasanuddin University Faculty of Medicine, sterile distilled water, 96% ethanol, Metranidazole, *Mueller Hinton Agar* (MHA), label paper and aluminum foil.<sup>9</sup> Samples of (*Moringa oleifera L.*) leaves were cleaned from the remaining dirt. After cleaning the moringa leaves are dried by air. After

drying, the sample is kept in a closed glass container. Oyster mushroom extract is obtained by maceration. The samples were in a closed container and then soaked with 96% ethanol solution and left for 5 days. After 5 days, the soaked sample is filtered using filter paper. The results of the filter are then evaporated using a rotary evaporator, so that a thick extract from the moringa leaf is obtained. The resulting thick extract is inserted into the vaporized container until all ethanol solvents evaporate.

#### Process of Testing Antimicrobial Effects of Diffusion Method

- 7 sterile test tubes were provided and Metranidazole. Dilution of the extract with sterile Aquadest was obtained to obtain concentrations of 15%, 20%, 25%, 30% and 35%, respectively as much as 5 mL.
- Each test tube is filled with the following conditions:
  - Tube 1: 2.5 mL 15% extract + 2.5 mL suspension *Porphyromonas gingivalis*
  - Tube 2: 2.5 mL 20% extract + 2.5 mL suspension *Porphyromonas gingivalis*
  - Tube 3: 2.5 mL extract 25% + 2.5 mL suspension *Porphyromonas gingivalis*
  - Tube 4: 2.5 mL 30% extract + 2.5 mL suspension *Porphyromonas gingivalis*
  - Tube 5: 2.5 mL 35% extract + 2.5 mL suspension *Porphyromonas gingivalis*
  - Tube 6: 2.5 mL extract 100% + 2.5 mL suspension *Porphyromonas gingivalis*



Figure 1. Dilution results of the tube *aggregatibacter actinomycetemcomitans*

Tube 7: 500 mg Metronidazole + suspension *Porphyromonas gingivalis* (positive control)

- Each test tube is filled with the following conditions:

Tube 1: 2.5 mL 15% extract + 2.5 mL suspension *Aggregatibacter actinomycetemcomitans*

Tube 2: 2.5 mL of 20% extract + 2.5 mL suspension *Aggregatibacter actinomycetemcomitans*

Tube 3: 2.5 mL of 25% extract + 2.5 mL suspension *Aggregatibacter actinomycetemcomitans*

Tube 4: 2.5 mL 30% extract + 2.5 mL suspension *Aggregatibacter actinomycetemcomitans*

Tube 5: 2.5 mL of 35% extract + 2.5 mL suspension *Aggregatibacter actinomycetemcomitans*

Tube 6: 2.5 mL 100% extract + 2.5 mL suspension *Aggregatibacter actinomycetemcomitans*

Tube 7: 500 mg Metronidazole + suspension *Aggregatibacter actinomycetemcomitans* (positive control)

#### Research Result

Minimum inhibitory effect of leaf extract Moringa (*Moringa oleifera L.*) against bacteria *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitan*.

The result of the tube dilution test is to see the turbidity level to determine the minimum inhibitory level. Tube dilution test results can be observed in Figures 1 and 2.



Figure 2. Dilution results of tubes *porphyromonas gingivalis*

From the results of tube dilution test observations in Figure 1 the Minimum Inhibitory Level (MIC) can be determined in 30% tubes starting to look clear and Figure 2 Minimal Inhibitory Levels (MIC) can be determined on the tube 20% starts to look clear.



Figure 3 bacterial growth *Aggregatibacter actinomycetemcomitans*




Figure 4 growth of bacteria *Porphyromonas gingivalis*

Minimal Bactericidal Concentration of leaf extract *Moringa oleifera L.* against *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*. From the results of the diffusion test observations in Figure 3, the Minimal Bactericidal Concentration (MBC) seen at a concentration of 35% shows that there is no growth of bacteria *Aggregatibacter actinomycetemcomitans* and in Figure 4 the Minimal Bactericidal Concentration (MBC) can be determined at a concentration of 25%.

### Conclusion

The conclusions of this study are:

Extract (*Moringa oleifera L.*) can inhibit the bacteria *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*.

Minimal Inhibitory Concentration (MIC) of leaf extract (*Moringa oleifera L.*) *Moringa* which can inhibit bacteria *Porphyromonas gingivalis* is 20% with an average inhibition zone formed is 12.9 mm and for bacteria *Aggregatibacter actinomycetemcomitans* is 30% with an average zone inhibition of 10.2 mm.

The Minimal Bactericidal Concentration (MBC)

of leaf extract killing (*Moringa oleifera L.*) which can inhibit bacteria *Porphyromonas gingivalis* is 25% with the average inhibition zone formed is 13.3 mm and for bacteria *Aggregatibacter actinomycetemcomitans* is 35% with zone average inhibition of 11.02 mm.

### Suggestions

Need to be carried out further research on leaf extract (*Moringa oleifera L.*) with different methods to inhibit the growth of *Porphyromonas gingivalis* and *Aggregatibacter actinomycetemcomitans*.

**Conflict of Interest:** There is no conflict of interest in this study.

**Source of Funding:** Domestic government

**Ethical Clearance:** This study obtained a label of ethics escaped by the number: 0082/PL09/KEPKFKG - RSGMUNHAS/2018 and register number UH 17120068 on Oktober 9, 2018.

### References

1. Thresia U, Sapara, OW, Juliatri. Antibacterial Effectiveness of Pacar Air Leaf Extract (Impatiens

- Balsamina L.) Against the Growth of Porphyromonas gingivalis. Pharmacy Scientific Journal - UNSRAT. 2016; 5(4): 10-7.
2. Mardiana Adam, Achmad H. The Relationship of Mineral Fluor Exposure in Water with The Presence of Gingivitis (Study Case in Subdistrict of Tempe, Sengkang City, Wajo District). Journal of International Dental and Medical Research. 2018; 11(2): 470-476
  3. Adam, A.M, Achmad, H. et. al. Efficacy of Mouthwash From Aloe Vera Juice After Scaling Treatment on Patient With Gingivitis: A Clinical Study. PBOCI journal. 2018; 18(1): e3959
  4. Quamilla N. Stress and Periodontitis. Journal of Syiah Kuala dentistry Society. 2016; 1(2): 161-3
  5. Mohamed AH, Elnima El AI, Ghandour IA. Inhibitory Effect of Garlic Extract on Porphyromonas Gingivalis Isolated from Sudanese Aggressive Periodontitis Patients Compared to Chlorhexidine Di- gluconate 0.2%. Journal of Chemical and Pharmaceutical research. 2017; 9(4): 259-63
  6. Beautiful Y, Mattulada IK. Antibacterial effect of crown god fruit extract (Phaleria macrocarpa (Scheff) Boerl) on Porphyromonas Gingivalis as an alternative root canal medicinal material. Makassar Dent J. 2015; 4 (2): 50-3
  7. Susanto C, Ervina I, Agusnar H. In vitro evaluation of antimicrobial effectiveness chitosan based tetracycline gel on some pathogenic periodontal bacteria. International Journal of Applied Dental Sciences. 2017; 3(2): 71-6.
  8. Dima L, Fatimawali, Lolo WA. Antibacterial Activity Test of Moringa oleifera L. Leaf Extract on Escherichia coli and Staphylococcus aureus. Pharmaceutical scientific journal - UNSRAT. 2016; 5 (2): 283-9
  9. Perumal Siddhuraju and Klaus Becker. Antioxidant Properties of Various Solvent Extracts of Total Phenolic Constituents from Three Different Agroclimatic Origins of Drumstick Tree (Moringa Oleifera Lam.). J. Agric. Food Chem. 2003; 51(8): 2144-5
  10. Thakmani, Fatma, Sokindra Kumar, and Shah Alam Han. Estimation of Total Phenolic Content, in-vitro Antioxidant and Anti-Inflammatory Activity of Flowers of Moringa Oleifera. Asian Pacific. J. Tropical Biomed. 2013; 3(8): 623-27
  11. AWD, Samsumaharto RA. Antibacterial Activity Test of Maseration of Moringa Oleifera Leaves (Moringa oleifera, Lamk) Against Staphylococcus aureus Bacteria. 2016; 6 (2): 14-9.
  12. Yudistira FA, Murwani S, Trisunuwati P. Potential of Antimicrobial Moringa oleifera Leaf Extract on Salmonella enteritidis (SP-1-PKH) In Vitro. Malang: Veterinary Medicine Program, Universitas Brawijaya; 2013.

# Differences of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC) of Moringa Leaf Extract (*Moringa Oleifera* L.) on Bacteria *Aggregatibacter Actinomycetemcomitans* and P

## ORIGINALITY REPORT

18%

SIMILARITY INDEX

12%

INTERNET SOURCES

10%

PUBLICATIONS

12%

STUDENT PAPERS

## PRIMARY SOURCES

1	Submitted to Universitas Airlangga Student Paper	3%
2	Submitted to Universitas Katolik Indonesia Atma Jaya Student Paper	2%
3	<a href="http://eprints.ners.unair.ac.id">eprints.ners.unair.ac.id</a> Internet Source	1%
4	Submitted to Universitas Muhammadiyah Yogyakarta Student Paper	1%
5	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1%
6	<a href="http://revista.uepb.edu.br">revista.uepb.edu.br</a> Internet Source	1%
7	Kollakondan Muthaiya Maria John, Muniappan Ayyanar, Thangavel Arumugam, Gansukh Enkhtaivan, Kijoun Jin, Doo Hwan Kim.	1%

"Phytochemical screening and antioxidant activity of different solvent extracts from *Strychnos minor* Dennst leaves", Asian Pacific Journal of Tropical Disease, 2015

Publication

---

8

Mochammad Hatta. "Multi-Locus Variable-Number Tandem Repeat Profiling of *Salmonella enterica* Serovar Typhi Isolates from Blood Cultures and Gallbladder Specimens from Makassar, South-Sulawesi, Indonesia", PLoS ONE, 09/15/2011

Publication

---

9

[garuda.ristekbrin.go.id](http://garuda.ristekbrin.go.id)

Internet Source

---

10

Submitted to Institute of International Studies

Student Paper

---

11

[www.researchgate.net](http://www.researchgate.net)

Internet Source

---

12

Submitted to Canadian International School, Singapore

Student Paper

---

13

Submitted to Harrisburg University of Science and Technology

Student Paper

---

14

[core.ac.uk](http://core.ac.uk)

Internet Source

---

1 %

1 %

1 %

1 %

1 %

1 %

1 %

15	<a href="http://www.academicjournals.org">www.academicjournals.org</a> Internet Source	1 %
16	<a href="http://mafiadoc.com">mafiadoc.com</a> Internet Source	1 %
17	<a href="http://www.omicsonline.org">www.omicsonline.org</a> Internet Source	<1 %
18	Submitted to Universitas Hasanuddin Student Paper	<1 %
19	<a href="http://www.jove.com">www.jove.com</a> Internet Source	<1 %
20	Listyaning Kartiko Kinasih, Idamawati Nababan, Suci Erawati, Rouli Natasia M Simanjuntak. "Effectivity of Jatropha multifida L. Leaves Extract as Antibacterial on Streptococcus mutans using In Vitro Testing Methods", Biomedical Journal of Indonesia, 2021 Publication	<1 %

Exclude quotes  On  
Exclude bibliography  On

Exclude matches  < 5 words