

The Comparative and Usefulness of Platelet Distribution Width in Acute Coronary Syndrome

by Idar Mappangara

Submission date: 25-Apr-2021 01:56PM (UTC+0700)

Submission ID: 1568950770

File name: ative_and_Usefulness_of_Platelet_Distribution_Width_In_Acute.pdf (772.54K)

Word count: 4079

Character count: 19174

RESEARCH ARTICLE

The Comparative and Usefulness of Platelet Distribution Width in Acute Coronary Syndrome

Idar Mappangara, Ali Aspar Mappahya, Sheila Witjaksono*

Department of Cardiology and Vascular Medicine, Faculty of Medicine, Hasanuddin University, Jl. Perintis Kemerdekaan Km.10, Makassar, Indonesia

*Corresponding author. E-mail: sheilawitjaksono@yahoo.com

Received date: Jul 17, 2016; Revised date: Aug 26, 2016; Accepted date: Sep 21, 2016

Abstract

BACKGROUND: Identifying risk factors for acute coronary syndrome (ACS) is important for both diagnostic and prognostic purposes. Abnormal platelet parameters, mainly platelet count (PC), mean platelet volume (MPV) and platelet distribution width (PDW) are thought to be among these risk factors because platelet activation plays a pivotal role in ACS. PDW is a more specific marker of platelet activation, since it does not increase during simple platelet swelling. In this study, the comparison of risk factors along with the PDW between Non ST-Elevation ACS (NSTEMACS) and ST-Elevation ACS (STEMACS) were investigated.

METHODS: This is a cross-sectional study, patients with ACS that admitted at dr.Wahidin Sudirohusodo General Hospital Makassar since December 2014 to March 2015 who undergo coronary angiography were enrolled. All data variables were taken according to the medical record. PDW was taken on first blood examination at hospital

admission. Data was significant if p -value <0.05 and was processed with SPSS version 16.

RESULTS: A total of 63 patients were enrolled. There were significantly higher PDW ($p=0.047$) in STEACS group (13.7 ± 2.0 fL) than in NSTEMACS group (12.6 ± 2.3 fL). There were a significant positive correlation between PDW value and SYNTAX scores (Pearson, $r=0.552$, $p=0.000$). Cardiovascular risk factors such as history of ACS before ($p=0.049$ & OR=10.216) and dyslipidemia ($p=0.049$ & OR=3.915) along with PDW ($p=0.003$ & OR=1.733) were significant predictor of the incidence of multivessel disease in patient with ACS.

CONCLUSION: An increased PDW may predicting more severe disease in ACS based on incidence of STEACS and multivessel disease more often and higher SYNTAX score.

KEYWORDS: platelet distribution width, PDW, acute coronary syndrome, ACS, multivessel disease, SYNTAX

Indones Biomed J. 2016; 8(3): 173-8

Introduction

Cardiovascular diseases (CVD) account for approximately 12 million deaths annually and are the commonest cause of death globally.(1) Ischemic heart disease (IHD) is one of CVD that defined as myocardial impairment due to imbalance between coronary blood flow (supply) and myocardial requirement (demand). It is divided into stable angina secondary to coronary artery disease (CAD) and

acute coronary syndromes (ACS). ACS is composed of patients with acute myocardial infarction with ST-segment elevation on their presenting electrocardiogram, ST-Elevation ACS (STEMACS), and those with unstable angina and non ST-segment elevation myocardial infarction, or non ST-Elevation ACS (NSTEMACS).(2) Conventional risk factors for atherosclerosis include smoking, diabetes mellitus, hypertension, dyslipidemia, obesity and stress which either acting singly or in combination increase the chances of developing coronary atherosclerosis. However,

they only explain part of the cases and other relevant risk factors need to be identified for an accurate calculation of an individual's risk for myocardial infarction.

Platelet indices such as platelet count (PC), mean platelet volume (MPV), and platelet distribution width (PDW) have been well utilized for certain conditions like idiopathic thrombocytopenic purpura (ITP), aplastic anemia and other haematological and myeloproliferative disorders to assess the prognosis but are underutilized for cardiovascular disorders.(3) Platelets played a role in the pathogenesis of cardiovascular disorders including atherosclerosis and its complications such as ACS and sudden cardiac death. In ACS there was hyperactivity and activation of platelet. Platelet activity is reflected by platelet which is indirectly measured by the some parameters. Larger platelets are metabolically and enzymatically more active than small platelets.(4) Platelet indices correlate with functional status of platelet and is an emerging risk marker for atherothrombosis.(5) Platelets with increased number and size of pseudopodia differ in size, possibly affecting PDW. PDW is a more specific marker of platelet activation, since it does not increase during simple platelet swelling.(6)

The most sensitive and specific biomarkers of myocardial damage are troponin I and troponin T, levels of both begin to rise at 2 to 4 hours and peak at 48 hours. Creatine Kinase enzymes begins to rise within 2 to 4 hours of the onset of myocardial infarction, peaks at about 24 hours and returns to normal within approximately 72 hours.(7) Platelet parameters can be detected earlier than markers of myocardial infarction. Platelet indices are easily recorded by automated cell counter and are routinely available in most clinical laboratories. There is an aim to make better use of the platelet parameters generated, as patients with larger platelets can easily be identified during routine haematological analysis and could possibly benefit from timely treatment.(4) The study was undertaken to compare the PDW in STEACS and NSTEMACS.

Methods

This is a cross-sectional study, which took place at Cardiology and Vascular Department, Faculty of Medicine, Hasanuddin University/dr.Wahidin Sudirohusodo General Hospital Makassar. All data variables based on medical record of patients with ACS who admitted at dr.Wahidin Sudirohusodo General Hospital Makassar since December 2014 to March 2015. The inclusion criteria were patients more than 18 years who undergo coronary angiography.

The exclusion criteria were patient with hematology disease and bleeding diathesis, patient with end stage renal disease, patient with carcinoma, patient with active or chronic inflammation, patient with autoimmune disease, patient with active infection, patient with blood transfusion, patient with cardiomyopathy and patient who didn't undergo cardiac catheterization in any cause. Data of blood examination was taken on first blood examination at emergency department when admitted to the hospital.

PDW was taken on first blood examination at hospital admission. Patients were divided into two groups: patients with STEACS and with NSTEMACS. Results were expressed as mean±standard deviation. Independent T tests were used to compare numerical variables. Chi square tests were used to compare categorical variables. Presence of multivessel disease and SYNTAX score based on coronary angiography. Correlation between continuous variables and SYNTAX score were determined by Pearson's correlation test. Predictor of multivessel disease were determined by logistic regression test. $D_{0.05}$ was significant if the p -value <0.05 and processed with SPSS version 16.

Results

A total of 63 cases were studied and were divided further into two groups, patients with STEACS and with NSTEMACS. The mean age of patients was 53.46±8.62 years old, with range between 35-72 years old. The mean body weight was 66.62±12.39 kg, with range between 40-96 kg. The mean body height was 163.19±5.76 cm, with range between 143-175 cm. The mean body mass index was 24.93±4.05 kg/m², with range between 16-36.6 kg/m². The mean waist circumference was 92.04±11.39 cm, with range of 61-119 cm. The mean systolic blood pressure was 135.71±32.27 mmHg, with range of 60-230 mmHg. The mean diastolic blood pressure was 80.48±21.73 mmHg, with range of 0-130 mmHg. The mean random blood glucose was 151.87±78.19 mg/dL, with range of 73-442 mg/dL. The mean leukocyte was 11,970±4,891 / μ L, with range of 5,000-27,000 / μ L. The mean PC was 272,000±74,495 / μ L, with range of 95,000-460,000 / μ L. The mean platelecrit was 0.220±0.06% with range of 0.08-0.36%. The mean MPV was 7.88±1.03 fL, with range of 6.3-11 fL. The mean PDW was 13.25±2.18 fL, with range of 7.8-17.5 fL. The mean total cholesterol was 193.40±47.54 mg/dL, with range of 108-328 mg/dL. The mean high density lipoprotein (HDL) cholesterol was 11.20±33.54 mg/dL, with a range of 9-62 mg/dL. The mean low density lipoprotein (LDL) cholesterol

was 129.97 ± 45.28 mg/dL, with range of 62-247 mg/dL. The mean triglyceride was 168.73 ± 97.49 mg/dL, with range of 61-643 mg/dL. The mean length of stay was 7.9 ± 4.67 days, with range of 1-27 days.

Total number of males was 56 (88.9%) and number of females was 7 (11.1%). The number of males in the STEACS group were 35 (62.5%) compared to NSTEMACS group 21 (37.5%). The numbers of females in the STEACS group were 4 (42.85%) compared to NSTEMACS group 4 (57.14%). In our present study, risk factors were evaluated. Significantly, more number of patients who had history of ACS before and older age were present in NSTEMACS than STEACS. Other risk factors did not differ significantly among both of them. There were no difference in the incidence of multivessel disease, SYNTAX score group and death in hospital between STEACS and NSTEMACS. There was significantly more number of patients who had heart failure in NSTEMACS than STEACS (Table 1).

There was statistically significant different in mean ages between STEACS group (51.45 years old) and NSTEMACS group (56.52 years old) with $p=0.021$. There was statistically significant different in mean leukocyte between STEACS group (14.08 years old) and NSTEMACS group (8.76 years old) with $p=0.000$. There was statistically significant different in mean PDW between STEACS group (13.69 years old) and NSTEMACS group (12.58 years old) with $p=0.047$. There were not statistically significant different in body weight, body height, body mass index, waist circumference, systolic blood pressure, diastolic blood pressure, random blood glucose, trombocyte, plateletcrit, MPV, total cholesterol, LDL cholesterol, HDL cholesterol, triglycerides, uric acid, number of vessel with significant lesion, SYNTAX scores, and length of stay in hospital between STEACS group and NSTEMACS group (Table 2).

In this study, we also performed Pearson correlation test to see platelet indices correlation with SYNTAX score. The results of the analysis with Pearson correlation between the PDW and SYNTAX scores showed a very significant positive correlation (Pearson, $r=0.552$, $p=0.000$); while the indices of other platelets and SYNTAX score as platelet count (Pearson, $r=-0.159$, $p=0.214$), plateletcrit (Pearson, $r=0.065$, $p=0.612$) and MPV (Pearson, $r=0.185$, $p=0.147$) did not show a significant correlation with SYNTAX score.

We analyse risk factor along with platelet indices and other variables with logistic regression to search factor or parameter that can predict the incidence of multivessel disease. The mean PDW in the group with multivessel disease was 13.9 ± 1.9 fL, while in the group without multivessel

Table 1. Comparison of categorical variables in STEACS and NSTEMACS.

Risk Factor	Total	ACS		P
		STEACS	NSTEMACS	
Sex				
Male	56	35	21	0.317
Female	7	3	4	
Family History of CAD				
No	51	29	22	0.248
Yes	12	9	3	
History of ACS				
No	48	35	13	0.000
Yes	15	3	12	
Hypertension				
No	26	18	8	0.225
Yes	37	20	17	
Diabetes Mellitus				
No	52	31	21	0.804
Yes	11	7	4	
Dyslipidemia				
No	31	19	12	0.877
Yes	32	19	13	
History of Stroke				
No	63	38	25	-
Yes	0	0	0	
Smoking				
No	15	6	9	0.065
Yes	48	32	16	
Obesity (BMI > 25 kg/m²)				
No	35	21	14	0.954
Yes	28	17	11	
Central Obesity				
No	28	19	9	0.274
Yes	35	19	16	
Multivessel Disease				
No	22	14	8	0.693
Yes	41	24	17	
SYNTAX score group				
Low (<22)	40	26	14	0.316
Moderate-High (>22)	23	12	11	
Presence of Heart Failure				
No	30	14	16	0.035
Yes	33	24	9	
Death in Hospital				
No	60	37	23	0.328
Yes	3	1	2	

*Data was significant if the p -value < 0.05

disease was 11.9 ± 1.9 fL. Statistical analysis showed that PDW mean significantly different with $p=0.000$. Other platelet indices such as platelet count ($p=0.649$), plateletcrit ($p=0.446$) and MPV ($p=0.591$) do not have statistically significant association with multivessel disease. Other risk factors such as history of ACS ($p=0.009$), dyslipidemia ($p=0.027$) and age ($p=0.024$) also had significant association with multivessel disease (Table 3).

Table 4 shows that history of ACS, dyslipidemia, age and PDW were analyzed by logistic regression. Statistically

Table 2. Comparison the mean of numerical variables in STEACS and NSTEACS.

	SKA Group	n	mean±SD	p
Age (years)	NSTEACS	25	56.52±8.88	0.021
	STEACS	38	51.45±7.93	
	Total	63	53.46±5.62	
Body Weight (kg)	NSTEACS	25	66.36±13.00	0.894
	STEACS	38	66.79±12.14	
	Total	63	66.62±12.39	
Body Height (cm)	NSTEACS	25	162.48±4.74	0.431
	STEACS	38	163.66±6.36	
	Total	63	163.19±5.76	
BMI (kg/m ²)	NSTEACS	25	25.09±4.56	0.801
	STEACS	38	24.83±3.74	
	Total	63	24.93±4.05	
Waist Circumference (cm)	NSTEACS	25	91.84±11.52	0.911
	STEACS	38	92.17±11.45	
	Total	63	92.04±11.39	
Systolic Blood Pressure (mmHg)	NSTEACS	25	141.6±39.34	0.243
	STEACS	38	131.84±26.49	
	Total	63	135.71±32.27	
Diastolic Blood Pressure (mmHg)	NSTEACS	25	81.2±25.55	0.832
	STEACS	38	80±19.17	
	Total	63	80.48±21.73	
Random Blood Glucose (mg/dL)	NSTEACS	25	133.92±45.19	0.141
	STEACS	38	163.68±92.52	
	Total	63	151.87±78.20	
Leukocyte (10 ³ /μL)	NSTEACS	25	8.76±2.21	0.000
	STEACS	38	14.08±5.04	
	Total	63	11.97±4.89	
Trombocyte (10 ³ /μL)	NSTEACS	25	269.48±65.75	0.796
	STEACS	38	274.5±80.53	
	Total	63	272.51±74.50	
Plateletcrit (%)	NSTEACS	25	0.22±0.59	0.739
	STEACS	38	0.21±0.06	
	Total	63	0.22±0.06	
MPV (fL)	NSTEACS	25	8.03±1.17	0.351
	STEACS	38	7.78±0.93	
	Total	63	7.88±1.03	
PDW (fL)	NSTEACS	25	12.58±2.25	0.047
	STEACS	38	13.69±2.05	
	Total	63	13.25±2.18	
Total Cholesterol (mg/dL)	NSTEACS	25	200.8±48.91	0.320
	STEACS	38	188.53±46.63	
	Total	63	193.4±47.54	
HDL Cholesterol (mg/dL)	NSTEACS	25	36.32±9.49	0.111
	STEACS	38	31.71±11.97	
	Total	63	33.54±11.20	
LDL Cholesterol (mg/dL)	NSTEACS	25	138.92±48.80	0.206
	STEACS	38	124.08±42.44	
	Total	63	129.97±45.28	
Triglycerides (mg/dL)	NSTEACS	25	164.36±94.32	0.776
	STEACS	38	171.61±100.67	
	Total	63	168.73±97.49	
Uric Acid (mg/dL)	NSTEACS	25	7.31±2.20	0.212
	STEACS	38	6.57±2.33	
	Total	63	6.86±2.29	
Number of Vessel with Significant Lesion	NSTEACS	25	1.92±1	0.997
	STEACS	38	1.92±0.88	
	Total	63	1.92±0.92	
SYNTAX Score	NSTEACS	25	16.26±10.43	0.829
	STEACS	38	15.67±10.60	
	Total	63	15.91±10.45	
Length of Stay in Hospital (days)	NSTEACS	25	7.44±3.32	0.526
	STEACS	38	8.21±5.4	
	Total	63	7.9±4.67	

*Data was significant if the p-value <0.05

Table 3. Comparison between multivessel disease group.

Risk Factor	Multivessel Disease		p
	Yes (n=41)	No (n=22)	
Sex			
Male	36	20	0.709
Female	5	2	
Family History of CAD			
Yes	8	4	0.898
No	33	18	
History of ACS			
Yes	14	1	0.009
No	27	21	
Hypertension			
Yes	25	12	0.621
No	16	10	
Diabetes Mellitus			
Yes	9	2	0.200
No	32	20	
Dyslipidemia			
Yes	25	7	0.027
No	16	15	
History of Stroke			
Yes	0	0	-
No	41	22	
Smoking			
Yes	31	17	0.883
No	10	5	
Obesity (BMI >25 kg/m²)			
Yes	17	11	0.516
No	24	11	
Central Obesity			
Yes	24	11	0.516
No	17	11	
Presence of Heart Failure			
Yes	23	10	0.420
No	18	12	
Death in hospital			
Yes	3	0	0.194
No	38	22	
Age (years)	55.2±8.7	50.1±7.6	0.024
Body Weight (kg)	65.9±13.5	67.8±10.1	0.563
Body Height (cm)	162.9±6.4	163.6±4.5	0.689
BMI (kg/m²)	24.7±4.3	25.3±3.5	0.561
Waist Circumferences (cm)	92.1±12.4	91.8±9.5	0.920
Systolic Blood Pressure (mmHg)	134.2±29.1	138.6±38.1	0.603
Diastolic Blood Pressure (mmHg)	80.0±20.6	81.4±24.1	0.815
Random Blood Glucose (mg/dL)	153.6±77.7	148.6±80.9	0.812
Leukocyte (10³/μL)	11.2±4.4	13.3±6.1	0.104
PC (10³/μL)	269.3±74.8	278.4±75.3	0.649
Plateletcrit (%)	0.219±0.06	0.278±0.06	0.446
MPV (fL)	7.9±0.8	7.8±1.4	0.591
PDW (fL)	13.9±1.9	11.9±1.9	0.000
Total Cholesterol (mg/dL)	197.3±52.3	186.1±37.0	0.376
HDL Cholesterol (mg/dL)	32.7±12.5	35.1±8.3	0.412
LDL Cholesterol (mg/dL)	137.5±51.6	116±25.7	0.073
Triglycerides (mg/dL)	161.1±104.6	182.9±83.1	0.402
Uric Acid (mg/dL)	7.2±2.6	6.2±1.3	0.082
Length of stay in Hospital (days)	7.3±3.5	9.1±6.3	0.157

*Data was significant if the p-value <0.05

analyse showed that age is no longer a predictor of the incidence of multivessel disease with $p=0.352$; whereas history of ACS ($p=0.049$ & $OR=10.216$), dyslipidemia ($p=0.049$ & $OR=3.915$) and PDW ($p=0.003$ & $OR=1.733$) are a significant predictor of the incidence of multivessel disease.

Table 4. Logistic regression analysis of history SKA, dyslipidemia, age, and PDW with incidence of multivessel disease.

	B	SE	p	OR
History of ACS	2.324	1.181	0.049	10.216
Dyslipidemia	1.365	0.693	0.049	3.915
Age	0.041	0.044	0.352	1.042
PDW	0.550	0.186	0.003	1.733
Constant	-9.643	3.223	0.003	0.000

*Logistic regression analyze was significant if the p-value <0.05

Discussion

This study showed that mean PDW value was higher in STEACS than NSTEMACS. It has been known that STEACS was more severe than NSTEMACS. Pathophysiologically, NSTEMACS somewhat different from STEACS. NSTEMACS caused by complete occlusion of a minor coronary artery or a partial occlusion of a major coronary artery and its causes a partial damage of heart muscle. STEACS caused by a complete occlusion of a major coronary artery and its causes a full damage of heart muscle. Both of them can make some complications. But some complications like cardiogenic shock, left ventricular failure (like in this study), severe mitral regurgitation due to papillary muscle rupture, cardiac tamponade due to ventricular wall rupture are more in STEACS (due to full thickness heart muscle damage) than NSTEMACS.

There were significantly higher leukocytes in STEACS than NSTEMACS. This result is similar with previous study that conclude STEACS patients have significantly higher peak CRP levels compared to NSTEMACS patients and suggest that inflammatory processes play an independent role in the pathogenesis of myocardial infarction.(8) There were significantly higher PDW in STEACS than NSTEMACS. In ACS, rupture of unstable plaque activated the thrombogenic cascade. However, platelet reactivity plays a vital role in the

formation and propagation of intracoronary thrombus.(9) Other platelet indices such as platelet count, platelecrit, and MPV did not significantly different between STEACS and NSTEACS. This result similar with study by Vagdatli, *et al.*, which conclude that PDW is a more specific marker of platelet activation.(6)

PDW has a positive correlation with SYNTAX scores. This result is similar with study by Bekler, *et al.*, that conclude PDW is associated with severity of CAD based on Gensini score (10) and study by Akin, *et al.*, that conclude PDW is a cheap and easy laboratory test which independently associated with high SYNTAX score (11). We also divided patients into multivessel disease group and no multivessel group, so we may predict the incidence of multivessel disease because it associated with poor prognosis.(12) Analyzing the data with logistic regression, we found that PDW among with history of ACS before and dyslipidemia are statistically significant to predicting multivessel disease. We all have known that history of ACS before and dyslipidemia are well known cardiovascular risk factors that had been proven influence the incidence of multivessel disease.

Limitations of the study include the small sample size and only done in one place. The study did not follow up the patient to see that PDW value can be used as a marker of future cardiovascular events.

Conclusion

Platelet distribution width in STEACS group is higher than in NSTEACS group. The higher PDW, SYNTAX score will also be higher. Platelet distribution width is also higher in multivessel group. In this study, PDW may be useful to predict more severe disease in ACS based on incidence of multivessel disease and SYNTAX score.

References

1. Shah SN, Anand MP, Billimoria AR, Kamath SA, Karnad DR. API textbook of Medicine. 8th edition. New Delhi: Jaypee Brothers Medical Publishers; 2008.
2. Fauci AS, Kasper DL, Longo DL, Braunwald E, Hauser S. Harrison's Principles of Internal Medicine. 17th edition. New Delhi: Mc Graw Hill Medical; 2008.
3. Kaito K, Otsubo H, Usui N, Yoshida M, Tanno J, Kurihara E, *et al.* Platelet size deviation width, platelet large cell ratio, and mean platelet volume have sufficient sensitivity and specificity in the diagnosis of immune thrombocytopenia. *Br J Haematol.* 2005; 128: 698-702.
4. Khandekar MM, Khurana AS, Deshmukh SD, Ka-krani AL, Katdare AD, Inamdar AK. Platelet volume indices in patients with coronary artery disease and acute myocardial infarction. *J Clin Pathol.* 2006; 59: 146-9.
5. Lippi G, Filippozzi L, Salvagno GL, Montagnana M, Franchini M, Guidi GC, *et al.* Increased mean platelet volume with acute coronary syndromes. *Arch Pathol Lab Med.* 2009; 133: 1441-3.
6. Vagdatli E, Gounari E, Lazaridou E, Katsibourlia E, Tsikopoulou F, Labrianou I. Platelet distribution width: a simple, practical and specific marker of activation of coagulation. *Hippokratia.* 2010; 14: 28-32.
7. Kumar V, Abbas AK, Fausto N, Aster JC. Robbins and Cotran Pathologic Basis of Disease. 8th edition. New Delhi: Elsevier; 2010.
8. Habib SS, Kurdi MI, Al Aseri Z, Suriya MO. CRP levels are higher in patients with ST elevation than non-ST elevation acute coronary syndrome. *Arq Bras Cardiol.* 2011; 96: 13-7.
9. Martin JF, Bath PM, Burr ML. Influence of platelet size on outcome after MI. *Lancet.* 1991; 338: 1409-11.
10. Bekler A, Ozkan MT, Tenekecioglu E, Gazi E, Yener AU, Temiz A, *et al.* Increased platelet distribution width is associated with severity of coronary artery disease in patients with acute coronary syndrome. *Angiology.* 2015; 66: 638-43.
11. Akin F, Ayca B, Kose N, Altun I, Avsar M, Celik O, *et al.* Relation of platelet indices to severity of coronary artery disease in patients undergoing primary percutaneous coronary intervention. *Perfusion.* 2016; 31: 216-22.
12. Sorajja P, Gersh BJ, Cox DA, McLaughlin MG, Zimetbaum P, Costantini C, *et al.* Impact of multivessel disease on reperfusion success and clinical outcomes in patients undergoing primary PCI for AMI. *Eur Heart J.* 2007; 28: 1709-16.

The Comparative and Usefulness of Platelet Distribution Width in Acute Coronary Syndrome

ORIGINALITY REPORT

21 %
SIMILARITY INDEX

14 %
INTERNET SOURCES

17 %
PUBLICATIONS

10 %
STUDENT PAPERS

PRIMARY SOURCES

- 1** Siti Nurul Mubarakah, I Dewa Agung Susilawati, Sumarno Sumarno, I Ketut Gedhe Muliarta, Djanggan Sargowo. "Porphyromonas gingivalis Induced Fragmentation of Type IV Collagen Through Macrophage-Activated MMP-9: (In Vitro Study of Collagenolytic Mechanism in Pathogenesis of Atherosclerotic Plaque Rupture)", The Indonesian Biomedical Journal, 2009
Publication **2** %
 - 2** www.jove.com
Internet Source **1** %
 - 3** Abhey Chawla, Mahesh H Karigoudar, Anil K Reddy, . Parul, Mahmood Nawaz Khan. "The Study of Correlation between Different Complete Blood Count Parameters in Acute Coronary Syndromes", NATIONAL JOURNAL OF LABORATORY MEDICINE, 2019
Publication **1** %
-

4	M M Khandekar. "Platelet volume indices in patients with coronary artery disease and acute myocardial infarction: an Indian scenario", Journal of Clinical Pathology, 2/1/2006 Publication	1 %
5	Submitted to University of Strathclyde Student Paper	1 %
6	online-medical-help.blogspot.com Internet Source	1 %
7	www.jcnonweb.com Internet Source	1 %
8	Georg Endler. "Mean platelet volume is an independent risk factor for myocardial infarction but not for coronary artery disease", British Journal of Haematology, 5/2002 Publication	1 %
9	Submitted to Higher Education Commission Pakistan Student Paper	1 %
10	www.scielo.br Internet Source	1 %
11	tuberktoraks.org Internet Source	1 %

- | | | |
|----|---|------|
| 12 | Abdollahi, Bita, Mehran Mesgari Abbasi, Parvin Zakeri Milani, Ashraf Sadat Nourdadgar, Seyyed Mehdi Banan Khojasteh, and Vahid Nejati. "Hydro-Methanolic Extract of Cornus Mas L. and Blood Glucose, Lipid Profile and Hematological Parameters of Male Rats", Iranian Red Crescent Medical Journal, 2014.
Publication | 1 % |
| 13 | Jaya Manchanda, Ajay Malik. "Study of platelet indices in pregnancy-induced hypertension", Medical Journal Armed Forces India, 2020
Publication | 1 % |
| 14 | journals.plos.org
Internet Source | 1 % |
| 15 | statperson.com
Internet Source | <1 % |
| 16 | upslide.site
Internet Source | <1 % |
| 17 | jcp.bmj.com
Internet Source | <1 % |
| 18 | www.dovepress.com
Internet Source | <1 % |
| 19 | Pervin, Silpi, Md Saiful Islam, Sheuly Ferdoushi, Mosharaf Hossain, Tuhin Sultana, Md Harisul Hoque, Chowdhury Meshkat | <1 % |

Ahmed, and AN Nashimuddin Ahmed.
"Platelet Distribution width is an Early
Indicator of Acute Coronary Syndrome",
University Heart Journal, 2014.

Publication

20

journals.sagepub.com

Internet Source

<1 %

21

www.sdiarticle2.org

Internet Source

<1 %

22

Neelam Bharihoke, Vaibhavi Subhedar,
Praveen Singh, Poonam Gupta. "MEAN
PLATELET VOLUME (MPV) & OTHER PLATELET
INDICES IN ACUTE MYOCARDIAL INFARCTION
(AMI) & STABLE CORONARY ARTERY DISEASES
(SCAD)", Journal of Evidence Based Medicine
and Healthcare, 2014

Publication

<1 %

23

www.pacificjournals.com

Internet Source

<1 %

24

Submitted to National postgraduate Medical
College of Nigeria

Student Paper

<1 %

25

Robert H. Knopp, Pathmaja Paramsothy.
"Clinical trials report", Current Atherosclerosis
Reports, 2006

Publication

<1 %

www.guideline.gov

26

Internet Source

<1 %

27

Kumari Shilpi, R. M. Potekar. "A Study of Platelet Indices in Type 2 Diabetes Mellitus Patients", Indian Journal of Hematology and Blood Transfusion, 2017

Publication

<1 %

28

dent.unhas.ac.id

Internet Source

<1 %

29

www.hcplive.com

Internet Source

<1 %

30

Habib, Syed Shahid, and Abeer Al Masri. "Relationship of high sensitivity C-reactive protein with presence and severity in angiographically defined coronary artery disease patients", Pakistan Journal of Medical Sciences, 2013.

Publication

<1 %

31

bmrcbd.org

Internet Source

<1 %

32

ir.ymlib.yonsei.ac.kr

Internet Source

<1 %

33

jthc.tums.ac.ir

Internet Source

<1 %

34

m.scirp.org

Internet Source

<1 %

35	med.mahidol.ac.th Internet Source	<1 %
36	"Angiography and interventional cardiology", Journal of the American College of Cardiology, 20050201 Publication	<1 %
37	Bekler, A., M. T. A. Ozkan, E. Tenekecioglu, E. Gazi, A. U. Yener, A. Temiz, B. Altun, A. Barutcu, G. Erbag, and E. Binnetoglu. "Increased Platelet Distribution Width Is Associated With Severity of Coronary Artery Disease in Patients With Acute Coronary Syndrome", Angiology, 2014. Publication	<1 %
38	G A Large. "Contemporary management of acute coronary syndrome", Postgraduate Medical Journal, 2005 Publication	<1 %
39	Ponpan Matangkasombut. "Direct activation of natural killer T cells induces airway hyperreactivity in a non-human primate model of asthma", World Allergy Organization Journal, 11/2007 Publication	<1 %

Exclude bibliography On