

Plasma Levels of Cytokines (IL-10) in Multidrug Resistant Tuberculosis– A literature review

by Irawaty Djaharuddin

Submission date: 31-May-2021 08:20AM (UTC+0700)

Submission ID: 1597419864

File name: 10_in_Multidrug_Resistant_Tuberculosis-_A_Literature_review.pdf (394.96K)

Word count: 4000

Character count: 22122

Review Article

Plasma Levels of Cytokines (IL-10) in Multidrug Resistant Tuberculosis– A literature review

IRAWATY DJAHARUDDIN¹, MUZAKKIR AMIR², SRI RAMADANI³, ETIEN ANDRIYANI⁴, DICKY WAHYUDI⁵

Received: 30.08.20, Revised: 28.09.20, Accepted: 16.10.20

ABSTRACT

Objective: Interleukin-10 (IL-10), interferon gamma (IFN- γ) and tumour necrosis factor (TNF- α) play very important roles in anti-TB cell-mediated immunity, in both MDR-TB and drug-susceptible TB (DS-TB) patients.

Method: The databases used were PubMed and ScienceDirect. The keywords for the search were "Interleukin-10" AND "MDR-TB". The inclusion criteria was review articles and research articles published between January 2015 until August 2020 in free full text English.

Results: IL-10 plays an important role in suppressing macrophage and dendritic cell (DC) function, which helps control and initiate the immune responses. IL-10, which is a suppressor T cell or T-regulatory cytokine, is known to play a critical role during chronic and latent stages of pulmonary TB [53]. The IL-10 production is said to be elevated during the infection, promoting reactivation of TB [54]. The excessive production of this cytokine usually results in failure to control the infection, and this could account for why IL-10 was elevated in the MDR-TB participants.

Conclusions: IL-10 production in MDR-TB might also indicate suppression of the immune response, leading to an inadequate balance of pro- and anti-inflammatory cytokines.

Keywords: IL-10, MDR-TB

30

INTRODUCTION

Tuberculosis (TB) is an infectious disease caused by infection with *Mycobacterium tuberculosis* (M.tb). It is estimated that about one third of the world's population is currently infected by the M.tb. Pulmonary tuberculosis is still an important health problem in the world because this disease affects millions of people and is the second leading cause of death among infectious diseases, after Human Immunodeficiency Virus (HIV). Globally there were 9 million new cases in 2011 and 4 million deaths due to TB. WHO states that out of 22 countries with a high burden of TB (high burden country), Indonesia is in fourth place (after India, China and South Africa) with a TB prevalence of 289 per 100,000 population, a TB incidence of 189 per 100,000 population and a 27 per 100,000 population. Multi-drug resistant tuberculosis (MDR-TB) is defined as in vitro resistance to isoniazid (H) and rifampin (R).

In Indonesia it is defined as resistance to, at least H and R simultaneously, with or without resistance to other first-line anti-tuberculosis (OAT) drugs.³ Data from WHO shows the prevalence is increasing from year to year. Worldwide it is estimated that there are 3.7% new MDR-TB cases and 20% MDR-TB with a history of previous OAT treatment.

METHOD

The databases used were PubMed and ScienceDirect. The keywords for the search were "Interleukin-10" AND "MDR-TB". The inclusion criteria was review articles and research articles published between January 2015 until August 2020 in free full text English. Studies that were published as abstract, editorial, and case report were excluded. The same articles that found on the two database also excluded.




Fig.1: Article extraction flow chart. Authors' compilation.

RESULTS

Tuberculosis (TB) remains one of the world's deadliest communicable diseases.¹ Despite the availability of drugs to treat and control the disease, certain factors including patients' non-compliance, poor nutrition, and co-infection with HIV/AIDS has resulted in increased burden of TB infection, especially in resource-limited countries in sub-Saharan Africa and Asia.² In 2017, the World Health Organization (WHO) reported that 490,000 million cases of multidrug-resistant TB (MDR-TB) emerged in 2016 and an additional 110,000 cases were susceptible to isoniazid but resistant to rifampicin (RR-TB), the most effective first-line anti-TB drug.³ The advent of multidrug-resistant cases pose a disproportionate threat to the global prospects of TB control, and frustrates diagnosis and treatment.^{4,5} The regions most affected by MDR-TB cases are the former Soviet Union (Eastern European countries) and the Russian Federation, where at least one-third of TB cases presenting for treatment have MDR-TB.^{6,7} In sub-Saharan Africa, South Africa is reported to have the highest number of MDR-TB cases.⁸ Population rates of MDR-TB cases in a number of countries across the world have increased over time.^{9,10} In Ghana, MDR-TB (resistance to at least isoniazid and rifampin, plus resistance to several drugs, excluding combined resistance to isoniazid and rifampin) among TB patients has been found to be 8.7%.¹¹ Cytokines have been identified to play multiple roles in the immune and pathological

responses in TB. Interleukin-10 (IL-10), interferon gamma (IFN-γ) and tumour necrosis factor (TNF-α) play very important roles in anti-TB cell-mediated immunity, in both MDR-TB and drug-susceptible TB (DS-TB) patients.^{12,13,14} TNF-α has been identified to play various roles in the immune and pathological responses in TB by preventing the reactivation of persistent tuberculosis, modulating the pulmonary expression of specific immunologic factors and limiting the pathological response of the host.¹⁴ IFN-γ boosts antigen presentation, leading to the recruitment of CD4+ T-lymphocytes and toxic T-lymphocytes,¹² while IL-10 in response to *Mycobacterium tuberculosis* (MTB), may down-regulate the immune response and limit tissue injury. However, overexpression of these cytokines may have a negative impact on the capacity to control infection.¹³ Effective disease management strategies must therefore consider both drug treatment and host immunity. This strategic approach can be successful if studies are conducted on the immune markers involved and devise ways of enhancing their function to maximize effective response, especially in MDR-TB. This study therefore determined the levels of cytokines (IL-10, IFN-γ and TNF-α) that are linked with the ability of MTB to evade the host's immune system and mediate long-term infections in the lungs, leading to chronic tuberculosis.

Table 1: Overview of systematic review and meta-analysis Plasma Levels of Cytokines (IL-10) in Multidrug Resistant Tuberculosis

Author	Objectives	Result	Grade of Recommendation (Joanna Briggs Institute)

<p>Anthony Basingnaa,1,2 Samuel Antwi-Baffour, Dinah Obenewaa Nkansah, Emmanuel Afutu, and Enid Owusu</p>	<p>determined the differences in plasma concentrations of pro-inflammatory (IFN-γ and TNF-α) and anti-inflammatory (IL-10) cytokines in MDR-TB and drug-susceptible (DS) TB patients, in addition to some socio-economic factors</p>	<p>There were statistically significant associations between MDR-TB and factors such as education level ($X^2 = 9.895$, $p = 0.043$), employment status ($X^2 = 19.404$, $p = 0.001$) and alcoholism ($X^2 = 3.971$, $p = 0.046$)</p>	<p>4B</p>
<p>Seok-Yong Eum, Bo-Young Jeon, Jin-Hong Min, Seung-Cheol Kim, Sungae Cho, Seung-Kyu Park, Sang-Nae Cho</p>	<p>examined the relationship between cytokine levels and clinical parameters indicating the state of disease in active pulmonary TB patients.</p>	<p>No significant difference was found in IFN-gamma production between non-MDR-TB and MDR-TB patients, but there was a marked reduction in TNF-alpha production in MDR-TB patients accompanied by a moderate increase in IL-10 levels. In contrast, the presence of cavity was associated with a significant increase in IFN-gamma, whereas no difference in TNF-alpha between the cavity and non-cavity group was observed. Those who have TB lesions in the left lung showed lower levels of IFN-gamma and TNF-alpha and higher IL-10 levels than the patients with lesions on the right side. IFN-gamma levels were significantly increased in those with moderate or advanced lesions compared with patients with mild lesions. TNF-alpha and IL-10 levels did not change with disease severity. The number of <i>M. tuberculosis</i> bacilli in sputum was closely associated with TNF-alpha levels. The patient group with high value (+++) of sputum culture acid-fast bacilli produced significantly reduced levels of TNF-alpha compared with medium (++) and low (+) values</p>	<p>3B</p>

DISCUSSION

Chronic immune activation, which might be because of exposure to a high load of environmental antigens, has mostly characterized the immune profile of people living in sub-Saharan Africa^{14,15,16}. Such exposure has been observed to impair the host's immune response to *M. tuberculosis* and HIV¹⁷, which are widespread in sub-Saharan Africa¹⁸. Infection with intracellular parasites such as *Mycobacterium tuberculosis* is known to induce Th1 immune response¹⁸. The protective immunity against the TB pathogen is said to be mediated by cytokines such as IFN- γ , TNF- α , IL-12, IL-6 and IL-18 during the initial stage of infection¹⁹. IFN- γ has been identified to be significant for the function and maturation of multiple immune cells²⁰. It stimulates macrophages to produce TNF- α , which is an essential component of the innate defence mechanism of the host against pathogenic challenge²¹. IL-10 plays an important role in suppressing macrophage and dendritic cell (DC) function, which helps control and initiate the immune responses²². TNF- α contributes to the pathogenesis of tuberculosis due to its role in the formation and maintenance of granulomas²³.

TNF- α is considered a necessity in the removal of bacteria in inflammatory lesions, and hence it has been found to be one of the key cytokines in controlling MTB infection^{24,25}. Therefore, it was not surprising that elevated plasma levels of TNF- α were found in MDR-TB participants compared to the DS-TB participants. This correlated with a Chinese study that showed elevated levels of TNF- α in TB patients compared to controls²⁶. Several studies have reported increased levels of TNF- α in the serum of TB patients^{27,28,29}. TNF- α is a factor both in the protection against tuberculosis and in immunopathology. The high levels of TNF- α in MDR-TB patients in the current study may be due to a marked tissue necrosis during the disease occurrence that led to progressive TB and eventually MDR. This might have resulted in the release of TNF- α into circulation and eventually contributed to systemic indicators of TB, such as fever and cachexia. The role of IFN- γ as the main macrophage-activator (Th1 cytokine) has been clearly established in animal models infected with *M. tuberculosis*³⁰. Intracellular mycobacteria are destroyed due to IFN- γ action on macrophages

^{31,32}. IFN- γ has been shown to stimulate macrophages, ³ leading to the following: production of IFN- α , oxygen free radicals and nitric oxide; increase in the surface display ³⁵ MHC antigens and Fc receptors; increase in the intracellular concentration of some antibiotics; and decrease in lysosomal pH ^{33,34,35}. The significantly higher plasma levels of IFN- γ in MDR-TB compared to DS-TB observed in this study may be because more IFN- γ action on macrophages will be needed to destroy intracellular mycobacteria that show multi-resistance to anti-TB drugs.

IL-10, which is a suppressor T cell or T-regulatory cytokine, is known to play a critical role during chronic and latent stages of pulmonary TB³⁶. The IL-10 production is said to be elevated during the ⁴ infection, promoting reactivation of TB ^{37,38,39}. The excessive production of this cytokine usually results in failure to control the infection, and this could account for why IL-10 was elevated in the MDR-TB participants. Increased production of IL-10 in patients with active disease including MDR-TB has been reported in Turkey ^{37,40,41,42}. This high IL-10 production in MDR-TB might also indicate suppression of the ²³ immune response, leading to an inadequate balance of pro- and anti-inflammatory cytokines.

CONCLUSION

⁹ The levels of both pro- (IFN- γ , TNF- α) and anti-inflammatory (IL-10) cytokines ³³ are observed to be significantly higher in MDR-TB patients compare DS-TB patients. The findings of this study emphasize the potential roles of IFN- γ , TNF- α and IL-10 in the host response to MTB during drug resistance development. This study also showed a statistically significant association between MDR-TB and factors such as education level, employment status and alcohol intake. Alcohol intake can be considered as a very important risk factor due to the alcoholism attitude of most Ghanaians, in both urban and rural areas..

Financing: There are no public, commercial, and private sources of funding.

¹⁰ **Conflict of Interests:** The authors report no conflict of interests. The authors alone are responsible for the content and writing of the paper.

Acknowledgements: None.

REFERENCES

1. Lönnroth K., Migliori G.B., Abubakar I., D'Ambrosio L., de Vries G., Diel R., Douglas P., Falzon D., Gaudreau M.A., Goletti D., et al.

Towards tuberculosis elimination: An action framework for low-incidence countries. *Eur. Respir. J.* 2015;45:928–952. doi: 10.1183/09031936.00214014. [PMC free article] [PubMed] [CrossRef] [Google Scholar]

- Jassal M.S., Bishai W.R. Epidemiology and challenges to the elimination of global tuberculosis. *Clin. Infect. Dis.* 2010;50(Suppl. 3):S156–234. doi: 10.1086/651486. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- World Health Organization .WHO Global Tuberculosis Report 2017. (WHO/HTM/TB/2017.04) WHO; Geneva, Switzerland: 2017. [(accessed on 13 December 2018)]. Available online: https://www.who.int/tb/publications/global_report/gtbr2017_main_text.pdf. [Google Scholar]
- Kurz S.G., Furin J.J., Bark C.M. Drug-Resistant Tuberculosis: Challenges and Progress. *Infect. Dis. Clin. N. Am.* 2016;30:509–522. doi: 10.1016/j.idc.2016.02.010. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Diel R., Vandeputte J., de Vries G., Stillo J., Wanlin M., Nienhaus A. Costs of tuberculosis disease in the European Union: A systematic analysis and cost calculation. *Eur. Respir. J.* 2014;43:554–565. doi: 10.1183/09031936.00079413. [PubMed] [CrossRef] [Google Scholar]
- Skrachina A., Hurevich H., Zalutskaya A., Sahalchik E., Astrauko A., Hoffner S., Rusovich V., Dadu A., de Colombani P., Dara M., et al. Multidrug-resistant tuberculosis in Belarus: The size of the problem and associated risk factors. *Bull. World Health Organ.* 2013;91:36–45. doi: 10.2471/BLT.12.104588. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Ulmasova D.J., Uzakova G., Tillyashayhov M.N., Turaev L., van Gemert W., Hoffmann H., Zignol M., Kremer K., Gombogaram T., Gadoev J., et al. Multidrug-resistant tuberculosis in Uzbekistan: Results of a nationwide survey, 2010 to 2011. *Eur. Surveill.* 2013;17:20609. doi: 10.2807/1560-7917.ES2013.18.42.20609. [PubMed] [CrossRef] [Google Scholar]
- Berhan A., Berhan Y., Yizengaw D. A Meta-Analysis of Drug Resistant Tuberculosis in Sub-Saharan Africa: How Strongly Associated with Previous Treatment and HIV Co-Infection? *Ethiop. J. Health Sci.* 2013;23:271–282. doi: 10.4314/ejhs.v23i3.10. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
- Zignol M., van Gemert W., Falzon D., Sismanidis C., Glaziou P., Floyd K., Raviglione M. Surveillance of anti-tuberculosis drug resistance in the world: An updated analysis, 2007–2010. *Bull. World Health Organ.* 2012;90:111D–119D. doi:

- 10.2471/BLT.11.092585. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
10. Abubakar I., Zignol M., Falzon D., Raviglione M., Ditiu L., Masham S., Adetifa I., Ford N., Cox H., Lawn S.D., et al. Drug-resistant tuberculosis: Time for visionary political leadership. *Lancet Infect. Dis.* 2013;13:529–539. doi: 10.1016/S1473-3099(13)70030-6. [PubMed] [CrossRef] [Google Scholar]
 11. Owusu-Dabo E., Adjei O., Meyer C.G., Horstmann R.D., Enimil A., Kruppa T.F., Bonsu F., Browne E.N.L., Chinbuah M.A., Osei I., et al. *Mycobacterium tuberculosis* drug resistance, Ghana. *Emerg. Infect. Dis.* 2006;12:1170–1172. doi: 10.3201/eid1207.051028. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 12. Domingo-Gonzalez R., Prince O., Cooper A., Khader S.A. Cytokines and Chemokines in *Mycobacterium tuberculosis* Infection. *Microbiol. Spectr.* 2016;4:1–58. [PMC free article] [PubMed] [Google Scholar]
 13. Adrian T.B.R., Montiel J.L., Fernandez G., Valecillo A. Role of Cytokines and other Factors Involved in the *Mycobacterium Tuberculosis* Infection. *World J. Immunol.* 2015;5:16–50. doi: 10.5411/wji.v5.i1.16. [CrossRef] [Google Scholar]
 14. Olsen A., Chen Y., Ji Q., Zhu G., De Silva A.D., Vilcheze C., Weisbrod T., Li W., Xu J., Larsen M., et al. Targeting *Mycobacterium tuberculosis* Tumor Necrosis Factor Alpha-Downregulating Genes for the Development of Antituberculous Vaccines. *MBio.* 2016;7:e01023-15. doi: 10.1128/mBio.01023-15. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 15. Ferreira-Borges C., Parry C.D., Babor T.F. Harmful Use of Alcohol: A Shadow over Sub-Saharan Africa in Need of Workable Solutions. *Int. J. Environ. Res. Public Health.* 2017;14:346. doi: 10.3390/ijerph14040346. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 16. O'Connell R., Chishinga N., Kinyanda E., Patel V., Ayles H., Weiss H.A., Seedat S. Prevalence and correlates of alcohol dependence disorder among TB and HIV infected patients in Zambia. *PLoS ONE.* 2013;8:e74406. [PMC free article] [PubMed] [Google Scholar]
 17. Imtiaz S., Shield K.D., Roerecke M., Samokhvalov A.V., Leboth K., Rehm J. Alcohol consumption as a risk factor for tuberculosis: Meta-analyses and burden of disease. *Eur. Respir. J.* 2017;50:1700216. doi: 10.1183/13993003.00216-2017. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 18. Macintyre K., Bloss E. Alcohol brewing and the African tuberculosis epidemic. *Med Anthropol.* 2011;30:126–135. doi: 10.1080/01459740.2011.552457. [PubMed] [CrossRef] [Google Scholar]
 19. Walker N.F., Meintjes G., Wilkinson R.J. HIV-1 and the immune response to TB. *Future Virol.* 2013;8:57–80. doi: 10.2217/fvl.12.123. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 20. Roider J.M., Muenchhoff M., Goulder P.J. Immune activation and paediatric HIV-1 disease outcome. *Curr. Opin. HIV AIDS.* 2016;11:146–155. doi: 10.1097/COH.000000000000231. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 21. Mohammed H., Assefa N., Mengistie B. Prevalence of extrapulmonary tuberculosis among people living with HIV/AIDS in sub-Saharan Africa: A systemic review and meta-analysis. *HIV/AIDS.* 2018;10:225–237. doi: 10.2147/HIV.S176587. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 22. Amelio P., Portevin D., Reither K., Mhimbira F., Mpina M., Tumbo A., Nickel B., Marti H., Knopp S., Ding S., et al. Mixed Th1 and Th2 *Mycobacterium tuberculosis*-specific CD4 T cell responses in patients with active pulmonary tuberculosis from Tanzania. *PLoS Negl. Trop. Dis.* 2017;11:e0005817. doi: 10.1371/journal.pntd.0005817. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 23. Gupta A., Kaul A., Tsolaki A.G., Kishore U., Bhakta S. *Mycobacterium tuberculosis*: Immune evasion, latency and reactivation. *Immunobiology.* 2012;217:363–374. doi: 10.1016/j.imbio.2011.07.008. [PubMed] [CrossRef] [Google Scholar]
 24. Ellison M.A., Gearheart C.M., Porter C.C., Ambruso D.R. IFN- γ alters the expression of diverse immunity related genes in a cell culture model designed to represent maturing neutrophils. *PLoS ONE.* 2017;12:e0185956. doi: 10.1371/journal.pone.0185956. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
 25. Amanda M., Elena S., Mauricio A., Rajko R. TNF- α in Tuberculosis: A Cytokine with a Split Personality. *Inflamm Allergy Drug Targets.* 2009;8:53-62. [PubMed] [Google Scholar]
 26. Redford P.S., Murray P.J., O'Garra A. The role of IL-10 in immune regulation during *M. tuberculosis* infection. *Mucosal Immunol.* 2011;4:261–270. doi: 10.1038/mi.2011.7. [PubMed] [CrossRef] [Google Scholar]
 27. Flynn J.L., Chan J. Immunology of tuberculosis. *Annu. Rev. Immunol.* 2001;19:93–129. doi: 10.1146/annurev.immunol.19.1.93. [PubMed] [CrossRef] [Google Scholar]
 28. Mattos A.M., Almeida Cde S., Franken K.L. Increased IgG1, IFN-gamma, TNF-alpha and IL-6

- responses to *Mycobacterium tuberculosis* antigens in patients with tuberculosis are lower after chemotherapy. *Int. Immunol.* 2010;22:775–782. doi: 10.1093/intimm/dxq429. [PubMed] [CrossRef] [Google Scholar]
29. Nemeth J., Winkler H.-M., Zwick R.H., Müller C., Rumetshofer R., Boeck L, Burghuber O.C., Winkler S. Specific cytokine patterns of pulmonary tuberculosis in Central Africa. *Clin. Immunol.* 2012;138:50–59. doi: 10.1016/j.clim.2010.09.005. [PubMed] [CrossRef] [Google Scholar]
30. Tang S., Xiao H., Fan Y., Wu F., Zhang Z., Li H., Yang Y. Changes of pro-inflammatory cytokines and their receptors in serum from patients with pulmonary tuberculosis. *Zhonghua Jie He He Hu Xi Za Zhi.* 2002;25:325–329. [PubMed] [Google Scholar]
31. Kart L., Buyukoglan H., Tekin I.O., Altin R., Senturk Z., Gulmez I., Demir R., Ozesmi M. Correlation of serum tumornecrosisfactor-alpha, interleukin-4 and soluble interleukin-2 receptor levels with radiologic and clinical manifestations in active pulmonary tuberculosis. *Mediators Inflamm.* 2003;12:9–14. doi: 10.1080/0962935031000096926. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
32. Joshi L., Ponnana M., Sivangala R., Chelluri L.K., Nallari P., Penmetta S., Valluri V., Gaddam S. Evaluation of TNF- α , IL-10 and IL-6 Cytokine Production and Their Correlation with Genotype Variants amongst Tuberculosis Patients and Their Household Contacts. *PLoS ONE.* 2015;10:e0137727. doi: 10.1371/journal.pone.0137727. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
33. Ameglio F., Casarini M., Capoluongo E., Mattia P., Puglisi G., Giosue S. Post-treatment changes of six cytokines in active pulmonary tuberculosis: Differences between patients with stable or increased fibrosis. *Int. J. Tuberc. Lung Dis.* 2005;9:98–104. [PubMed] [Google Scholar]
34. Gallegos A.M., van Heijst J.W., Samstein M., Su X., Pamer E.G., Glickman M.S. A gamma interferon independent mechanism of CD4 T cell mediated control of *M. tuberculosis* infection in vivo. *PLoS Pathog.* 2011;7:e1002052. doi: 10.1371/journal.ppat.1002052. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
35. Töttemeyer S., Sheppard M., Lloyd A., Roper D., Dowson C., Underhill D., Murray P., Maskell D., Bryant C. IFN- γ enhances production of nitric oxide from macrophages via a mechanism that depends on Nucleotide Oligomerization Domain-2. *J. Immunol.* 2006;176:4804–4810. doi: 10.4049/jimmunol.176.8.4804. [PubMed] [CrossRef] [Google Scholar]
36. Hoeksema M.A., Scicluna B.P., Boshuizen M.C., vander Velden S., Neele A.E., Vanden Bossche J., Matlung H.L., van den Berg T.K., Goossens P., de Winther M.P. IFN- γ priming of macrophages represses a part of the inflammatory program and attenuates neutrophil recruitment. *J. Immunol.* 2015;194:3909–3916. doi: 10.4049/jimmunol.1402077. [PubMed] [CrossRef] [Google Scholar]
37. Sasindran S.J., Torrelles J.B. *Mycobacterium Tuberculosis* Infection and Inflammation: What is Beneficial for the host and for the bacterium? *Front. Microbiol.* 2011;2:2. doi: 10.3389/fmicb.2011.00002. [PMC free article] [PubMed] [CrossRef] [Google Scholar]
38. Achmad H. Horax S, Ramadhany S, et.al. 2019. Resistivity of Ant Nest (*Myrmecodia pendans*) On Ethanol Fraction Burkitt's Lymphoma Cancer Cells (Invitro) Through Interleukin 8 Angiogenesis Obstacles (II-8). *Journal of International Dental and Medical Research* ., ISSN 1309-100X. Vol 12 No. (2) pp.516-523.
39. Ramadhany S, Achmad H, Singgih MF, Ramadhany YF, Inayah NH, Mutmainnah N. A Review: Knowledge and Attitude of Society toward Tuberculosis Disease in Soppeng District. *Systematic Reviews in Pharmacy.* 2020; 11(5): 57-62. doi: 10.31838/srp.2020.5.10
40. Achmad, H. Khairunnisa, P. Mardiana, Karni Aulia, A. Potentially Of Extracted Papua's Anthill (*Myrmecodia pendans*) As Antitumor To Emphasis The Expression Of Vascular Endothelial Growth Factor Cell Burkitt's Lymphoma Cancer, *Asian Journal. of Microbiol. Biotech. Env Sc.* © Global Science Publications pp. 108-112, Vol. 20, No. (1): 2018 . ISSN-0972-3005.
41. Russeng SS, Salmah AU, Saleh LM, Achmad H, Rosanita ANR. The Influence of Workload, Body Mass Index (BMI), Duration of Work toward Fatigue of Nurses in Dr. M. Haulussy General Hospital Ambon. *Systematic Reviews in Pharmacy.* 2020; 11(4): 288-292. doi: 10.31838/srp.2020.4.41.
42. Singgih MF, Huldani, Achmad H, Sukmana BI, Carmelita AB, Putra AP, Ramadhany S, Putri AP. A Review of Nonsteroidal Anti-Inflammatory Drugs (NSAIDs) Medications in Dentistry: Uses and Side Effects. *Systematic Reviews in Pharmacy.* 2020; 11(5): 293-298. doi: 10.31838/srp.2020.5.43

Plasma Levels of Cytokines (IL-10) in Multidrug Resistant Tuberculosis– A literature review

ORIGINALITY REPORT

24%

SIMILARITY INDEX

12%

INTERNET SOURCES

16%

PUBLICATIONS

10%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Higher Education Commission Pakistan Student Paper	3%
2	jnk-signaling.com Internet Source	2%
3	link.springer.com Internet Source	1%
4	Yue Chen, Wen Hui Peng, Shi Feng Lai, Fang Luo, Dan Luo, Bao Guo Wang. "Association of gene polymorphisms and environmental factors in tuberculosis patients and their household contacts", Transactions of The Royal Society of Tropical Medicine and Hygiene, 2021 Publication	1%
5	Submitted to Foundation for Professional Development Student Paper	1%
6	Submitted to Johns Hopkins University Student Paper	1%

7	Erika Cavalheiro Skupien, Tássia Kirchmann Lazzari, Sandra Eugênia Coutinho, Denise Rossato Silva. "The relation between leptin and inflammatory markers with respiratory and peripheral muscle strength in tuberculosis: A case-control study", The Clinical Respiratory Journal, 2018 Publication	1 %
8	Submitted to London School of Hygiene and Tropical Medicine Student Paper	1 %
9	www.duo.uio.no Internet Source	1 %
10	www.hindawi.com Internet Source	1 %
11	Shilpy Sharma. "Genetic polymorphisms in TNF genes and tuberculosis in North Indians", BMC Infectious Diseases, 2010 Publication	1 %
12	Khalid Hussain Bhat, Asma Ahmed, Santosh Kumar, Pawan Sharma, Sangita Mukhopadhyay. "Role of PPE18 Protein in Intracellular Survival and Pathogenicity of Mycobacterium tuberculosis in Mice", PLoS ONE, 2012 Publication	1 %

13 V. P. Mohan, C. A. Scanga, K. Yu, H. M. Scott, K. E. Tanaka, E. Tsang, M. C. Tsai, J. L. Flynn, J. Chan. "Effects of Tumor Necrosis Factor Alpha on Host Immune Response in Chronic Persistent Tuberculosis: Possible Role for Limiting Pathology", *Infection and Immunity*, 2001
Publication

14 J. -S. LEE. "Profiles of IFN-gamma and its regulatory cytokines (IL-12, IL-18 and IL-10) in peripheral blood mononuclear cells from patients with multidrug-resistant tuberculosis", *Clinical & Experimental Immunology*, 6/2002
Publication

15 iai.asm.org
Internet Source

16 www.jove.com
Internet Source

17 Submitted to Isra University
Student Paper

18 abdominalkey.com
Internet Source

19 www.thecochranelibrary.com
Internet Source

20 Submitted to Massey University

<1 %

21

N. Range, P. Magnussen, A. Mugomela, W. Malenganisho et al. "HIV and parasitic co-infections in tuberculosis patients: a cross-sectional study in Mwanza, Tanzania", *Annals of Tropical Medicine & Parasitology*, 2013

Publication

<1 %

22

docplayer.net

Internet Source

<1 %

23

A. Hernández Cruz, S. Garcia-Jimenez, R. Zucatelli Mendonça, V. L. Petricevich. " Pro- and Anti-Inflammatory Cytokines Release in Mice Injected with Venom ", *Mediators of Inflammation*, 2008

Publication

<1 %

24

Submitted to University of Derby

Student Paper

<1 %

25

article.ijimmunology.org

Internet Source

<1 %

26

Shakila Harshavardhan, Kevin Kumar Vijayakumar, Vignesh Sounderrajan, Pushparaju Ramasamy, Sam Ebenezer Rajadas. "Basics of tuberculosis disease and principles of treatment and their effects", Elsevier BV, 2021

Publication

<1 %

27	serval.unil.ch Internet Source	<1 %
28	www.jci.org Internet Source	<1 %
29	www.ncbi.nlm.nih.gov Internet Source	<1 %
30	Lee, Hyo-Ji, Hyun-Jeong Ko, and Yu-Jin Jung. "Insufficient Generation of Mycobactericidal Mediators and Inadequate Level of Phagosomal Maturation Are Related with Susceptibility to Virulent Mycobacterium tuberculosis Infection in Mouse Macrophages", <i>Frontiers in Microbiology</i> , 2016. Publication	<1 %
31	"Posters", <i>Clinical Microbiology and Infection</i> , 05/2009 Publication	<1 %
32	Alice M. Biggane, Eleanor Briegal, Angela Obasi. "Interventions for adolescent alcohol consumption in Africa: protocol for a scoping review including an overview of reviews", <i>Systematic Reviews</i> , 2021 Publication	<1 %
33	Isabel SADA-OVALLE. " <i>In vitro</i> cytotoxicity of CD8+ T cells in multi-drug-resistant	<1 %

tuberculosis. A preliminary report",
Respirology, 05/2009

Publication

34

Park, S.K.. "Pulmonary resection combined with isoniazid- and rifampin-based drug therapy for patients with multidrug-resistant and extensively drug-resistant tuberculosis", International Journal of Infectious Diseases, 200903

Publication

35

Roberto Suárez-Méndez, Idrian García-García, Norma Fernández-Olivera, Magalys Valdés-Quintana et al. "Adjuvant interferon gamma in patients with drug – resistant pulmonary tuberculosis: a pilot study", BMC Infectious Diseases, 2004

Publication

36

Johannes Nemeth, Heide-Maria Winkler, Ralph H. Zwick, Catharina Müller, Rudolf Rumetshofer, Lucas Boeck, Otto C. Burghuber, Stefan Winkler. "Peripheral T Cell Cytokine Responses for Diagnosis of Active Tuberculosis", PLoS ONE, 2012

Publication

37

Sushma K. Cribbs, David Rimland. "Chapter 15 Alcohol and HIV: Experimental and Clinical Evidence of Combined Impact on the Lung",

<1 %

<1 %

<1 %

<1 %

Springer Science and Business Media LLC, 2014

Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On