

[Skip to Main Content](#)

# Infectivity

*by* Sitti Nuraenii

---

FILE	SITTI_NURAENI,_INFECTIVITY.PDF (407.71K)		
TIME SUBMITTED	09-JUL-2019 09:20AM (UTC+0700)	WORD COUNT	2487
SUBMISSION ID	1150342703	CHARACTER COUNT	12920



Original Research Article

doi: <http://dx.doi.org/10.20546/ijrbp.2016.305.001>

## Infectivity of Two Isolates of Microsporidia on Silkworm (*Bombyx mori* L.) in South Sulawesi

Sitti Nuraeni<sup>1\*</sup>, Djamal Sanusi<sup>1</sup>, Tutik Kuswinanti<sup>2</sup> and Andi Nasaruddin<sup>2</sup>

<sup>1</sup>Technology Non Timber Forest Products, Faculty of Forestry, Hasanuddin University, Makassar, South Sulawesi 90425, Indonesia

<sup>2</sup>Plant Protection, Faculty of Agriculture, Hasanuddin University, Makassar, South Sulawesi, 90425 Indonesia

\*Corresponding author.

### Abstract

Pebrine disease caused by *Nosema bombycis* is a detrimental disease of *Bombyx mori* and a serious limiting factor in sericulture industry that can even thwart the cocoon harvest. This study aims to determine the infectivity of two isolates of *N. bombycis* from two different seed sources (locally produced, PebLok and imported seeds, PebCin) in South Sulawesi. Spore suspension of *N. bombycis* ( $1 \times 10^8$  spores/ml aquadest) was evenly smeared on mulberry leaf surfaces. Early third and fourth larval instars of *B. mori* were fed with the smeared leaves. Insect mortality and symptom development were observed every 24 hr. Research results show that PebCin and PebLok were more infective at instar 3 and instar 4, respectively, although virulence of both isolates were relatively equal with cumulative mortality of more than 60%. Infection of both isolates caused a longer period of larval development for more than 10 days compared to uninfected larvae which was related to increased duration for each instar.

### Article Info

Accepted: 04 April 2016

Available Online: 06 May 2016

### Keywords

*Bombyx mori*  
Infectivity  
*Nosema bombycis*  
PebCin  
PebLok

### Introduction

South Sulawesi is the center of silk production in Indonesia, but for the last five years, production was declining steadily. One of the main problems is the presence of pebrine disease. The disease caused by *Nosema bombycis* Naegeli (Microsporidia: Nosematidae). Pebrine disease had also destroyed the natural silk industry in Europe, particularly France and Italy in the mid-19<sup>th</sup> century (Franzen, 2008; Plian, 1991; Rahmathulla et al., 2012). In 1845, France cocoon production was 26,000 tons and fell to 4,000 tonnes in 1853, later in 1865 only then the cause of the France and Italy silk industry collapse was revealed which related to the pebrine disease (Bhat et al., 2009). More than 40% of infection on female parents of silkworm were reported in Southeast Asia including Indonesia and Vietnam (Japan

Overseas Cooperation Volunteers, 1975). Furthermore, the level of attacks in the last five years has reached more than 60% and even it frequently causes harvest failures in Indonesia (BPA, 2015).

Pathogenicity is absolute whether pathogenic organisms causing disease in its host or not, while virulence is a quantitative measure of the ability to cause disease. Infectivity is the ability of an organism to cause infection (Shapiro-Ilan et al., 2012). Silkworm farmers in South Sulawesi use locally produced and imported eggs. In both of seed sources, infection of the microsporidia is unavoidable. Therefore, a study was conducted to determine the infectivity of microsporidian obtained from the two sources of silkworm seeds.

## Materials and methods

### Isolation and purification of microsporidian spores

Two isolates of microsporidia were isolated from silkworm larvae (*N. bombycis*) grown from seeds obtained from two different sources. Microsporidian PebLok isolate was from local seed source and PebCin isolate was from import seed source. Heavily infected larvae were crushed in distilled water and large silkworm and mulberry debris were removed by filtration on 2 cheesecloth layers. The supernatant was discarded and centrifuged 3000 rpm for 15 min, and the spore pellet was re-suspended to 10 ml using sterile distilled water, and then the suspension was re-centrifuged. This procedure was repeated 3 times, spores were washed in distilled water until the pellet was white. Finally, 10 ml spore solution was pipetted and poured into a hemocytometer for spore counting. The spores were counted under a light microscope at x40 magnifications (Armani et al., 2013). Spore suspensions of  $1 \times 10^8$  ml<sup>-1</sup> were prepared by diluting with distilled water, and then stored at 4°C until further use.

### Preparation of silkworms infected with Microsporidia

The silkworm larvae of BS09 were reared under room conditions with an average temperature of 25.9°C and humidity of 87.4%, 12 hrs of light and 12 hrs of dark. The silkworm larvae of the 2<sup>nd</sup> molt (third instar) and 3<sup>rd</sup> molt (fourth instar) were orally inoculated by feeding with mulberry leaves that were previously smeared with *N. Bombycis* spores.

## Results and discussion

### Disease occurrence

A day after fed with *N. bombycis* spore, no obvious symptoms were shown by the larvae. On the third day after infection then the larvae showed symptoms, i.e., slower larval movements, sluggish, reduced appetite and stunted development. These symptoms continued until the next moulting period. In the third instar inoculation, mortality occurred at the third day after infection (DAI) and was earlier in PebCin compared to PebLok (Fig. 1).




Fig. 1: Silkworm mortality after infection of two isolates of *N. bombycis* with different strains at third instar inoculation.

Death of larvae were observed almost every day, but the highest mortality occurred in PebCin at 14<sup>th</sup> DAI or in the late instar 4 period and at 33<sup>rd</sup> or in the late spinning, while the highest mortality in PebLok was at 15<sup>th</sup> DAI. In the fourth instar inoculation (Fig. 2), the opposite happened on mortality where highest mortality in PebLok occurred at third DAI which was earlier than in PebCin. Although the occurrence of larvae death was observed every day, the highest mortality occurred at the time of the last molt. This continued and the highest mortality in PebLok occurred on day 11, 12 and 13 DAI or when entering fifth instar while the highest mortality

in PebCin was at the 14<sup>th</sup> DAI (5<sup>th</sup> instar). The new external symptoms were observed after day 10 DAI and mortality of larvae increased in both inoculation instar 3 or 4. Histopathological study by Jyothi and Patil (2011) shows the midgut has formed sporoblast phase on the third day after infection. Hossain et al. (2012), spore plasma gradually increased in size and developed into meronts 24-48 hrs after inoculation. Infection by the most pathogenic strain of *N. bombycis* in India showed mortality occurred on day 10 after infection and can occur at all stages of larval development which is described by Rao et al. (2004).

The incubation period is the period starting from the entry of the pathogen into the host's body until the occurrence of symptoms or signs of the disease (Nataraju et al., 2005). The incubation period, or the latent period is a period from an infection or pathogen inoculation until the commencement of larval mortality (Begum et al., 2004). Bhat et al. (2009), reported that

the larvae become infected during early instar stages 1 and 2 will be dead in instar 5, but if the infection occurs during the 4<sup>th</sup> or 5<sup>th</sup> instar, larvae can still survive and form cocoons. Further observation by Bhat and Nataraju (2005) showed that inoculated larvae at 4<sup>th</sup> instar with a concentration  $1 \times 10^5$  spores ml<sup>-1</sup> can survive until pupa formation phase.




Fig. 2: Silkworm mortality after infection of two isolates of *N. bombycis* with different strains at fourth instar inoculation.




Fig. 3: Cumulative mortality of silkworms after infection of two isolates of *N. bombycis* with different strains at third instar inoculation.

Cumulative mortality of silkworms after inoculation of two isolates at instar 3 shows higher figure in PebCin (64%) than in PebLok which only showed cumulative mortality of 28% (Fig. 3). Contrary, in the fourth instar inoculation, PebCin cumulative mortality was 65% while PebLok reached 77% (Fig. 4). Infectivity is a manifestation of the severity of the disease that can be measured on the infected host. Estimates of

mortality or the mortality rate of the host is one measure of the infectivity of the pathogen (Thomas and Elkinton, 2004). Sharma et al. (2014) suggests that infection of  $1 \times 10^6$  spores ml<sup>-1</sup> in early instar 3 showed cumulative mortality of larvae to become pupa of 50.66%. Infection of *N. bombycis* on the first day of instar 2 resulted in cumulative mortality of larvae entering the cocoon forming phase of 41.0% (Bhat

et al., 2009). In addition, study of Tu et al. (2011) shows mortality up to 61.11% at infectivity of *N. bombycis* was at concentration of  $10^7$  ml<sup>-1</sup>. This result similar to Rao et al. (2004) that suggests that at a dose of  $10^7$  spores ml<sup>-1</sup> mortality can reach more than 50%

after infection. While studies of Jyothi and Patil (2011) showed that fourth instar mortality was 86% at a concentration of  $1 \times 10^6$  spores larva<sup>-1</sup>, infectivity can reach 90% at a concentration  $1.52 \times 10^6$  spores ml<sup>-1</sup> (Chakrabarty et al., 2012).




Fig. 4: Cumulative mortality of silkworm after infection of two isolates of *N. bombycis* with different strains on fourth instar inoculation.

### Disease progression

As response to the infection of microspores, a change occurred on the duration of each instar development stage. Table 1 shows that the duration of larval development of infected larvae were longer 12 an in the not infected larvae, i.e., longer 2.5 days in the third instar; 3 days in the

instar; 2.5 days in the fifth instar and 2 days in the cocoon formation phase (spinning) resulted in overall increase of 10.5 days compared to the uninfected larvae. According to Solter et al. (2012), effects that are typical on silkworm larvae that sublethal for infection of *N. bombycis* is extended period for larval development due to suspended ecdysis process.

Table 1. Larval and spinning stages of silkworms infected with *N. bombycis* strain PebLok and PebCin on 3<sup>rd</sup> or 4<sup>th</sup> instar.

Instar	Treatment			Control (no infection)		
	Eating stage (day)	Moulting (day)	Larval stage (day)	Eating stage (day)	Moulting (day)	Larval stage (day)
3	6.0	1.5	7.5	4.0	1.0	5.0
4	7.0	2.5	9.5	4.5	1.5	6.0
5	9.5	-	9.5	7.0	-	7.0
Spinning	5.0	-	5.0	3.0	-	3.0
Total			31.5	Total		21.0

### Conclusion

Strain PebCin is more ineffective at the third instar, with cumulative mortality of more than 60%. Strain PebLok infective at the fourth instar but the virulence of two isolates of *N. bombycis* were relatively equal in the cumulative larval mortality of more than 60%. Both isolates resulted in a longer duration of larval development of about 10 days longer compared to uninfected due to increased duration for each instar stage.

### Conflict of interest statement

Authors declare that they have no conflict of interest.

### References

- BPA, 2015. Statistik Pengembangan Persuteraan Alam Tahun 2015. Kementerian Kehutanan. Direktorat Jenderal Bina Pengelolaan DAS dan Perhutanan Sosial.
- Begum, H. A., E. Hassan, and J. Dingle. 2004. Resistance of three different populations of mulberry silkworm (*Bombyx mori* L.) to *Bacillus thuringiensis*. J. Plant Dis. Prot. (3), 231–237.

- Bhat, S.A., Nataraju, B., 2005. Mode of transmission of a newly discovered microsporidian and its effect on fecundity and hatching in silkworm, *Bombyx mori* L. *Int. J. Indust. Entomol.* 11(2), 81-86.
- Bhat, S.A., Bashir, I., Kamili, A.S., 2009. Microsporidiosis of silkworm, *Bombyx mori* L. (Lepidoptera-Bombycidae): A review. *Spl. Rev. Afr. J. Agric. Res.* 4(13), 1519-1523.
- Chakrabarty, S., Manna, B., Saha, A.K., Bindroo, B.B., 2012. Comparative study on effect different types of *Nosema* sp. (Microsporidia: Neosematidae) on mulberry and vanya silkworm. *J. Biol. Sci.* 12(1), 1-14.
- Franzen, C., 2008. Microsporidia: A review of 150 years of research. *Open Parasitol. J.* 2, 1-34.
- Hossain, Z., Gupta, S.K., Chakrabarty, S., Saha, A.K., Bindroo, B.B., 2012. Studies on the life cycle of five microsporidian isolates and histopathology of the mid-gut of the silkworm *Bombyx mori* (Lepidoptera: Bombycidae). *Int. J. Trop. Insect Sci.* 32(4), 203-209.
- Japan Overseas Cooperation Volunteer, 1975. Textbook of Tropical Sericulture, Tokyo. 612p.
- Jyothi, N. B., Patil, C. S., 2011. Light and electron microscopic study on the development of *Nosema bombycis* Naegeli in the midgut of silkworm *Bombyx mori* L. *Int. J. Indust. Entomol.* 23(1), 107-113.
- Kermani, N., Abu-hassan, Z.A., Dieng, H., Ismail, N. F., Attia, M., Ghani, I.A., 2013. Pathogenicity of *Nosema* sp. (Microsporidia) in the Diamondback Moth, *Plutella xylostella* (Lepidoptera: Plutellidae). *Plos One.* 8(5), 1-8.
- Nataraju, B., Sathayaprasad, K., Manjunath, D., Kumar, C.A., 2005. Silkworm Crop Protection. Central Silk Board, Bangalore, India. 412p.
- Rahmathulla, V. K., Kumar, C. M. K., Angadi, B. S., Divaprasad, V., 2012. Influence of weather factors on incidence and intensity of Microsporidiosis in silkworm (*Bombyx mori* L.). *J. Entomol.* 9(5), 266-273.
- Rao, S. N., Muthulakshmi, M., Kanginakudru, S., Nagaraju, J., 2004. Phylogenetic relationships of three new microsporidian isolates from the silkworm, *Bombyx mori*. *J. Invertebr. Pathol.* 86, 87-95.
- Shapiro-Ilan, D.I., Fuxa, J.R., Lacey, L.A., Onstand, D.W., Kaya, H.K., 2012. Definitions of pathogenicity and virulence in invertebrate pathology. *J. Invertebr. Pathol.* 88, 1-7.
- Sharma, S.D., Balavenkatasubbaiah, M., Babu, A.M., Kumar, S.N., Bindroo, B.B., 2014. Impact of a new microsporidian infection on larval and cocoon parameters of the silkworm, *Bombyx mori* L. *Int. J. Plant Anim. Environ. Sci.* 4(1), 82-87.
- Solter, L.F., Becnel, J.J., Oi, D.H., 2012. Microsporidian entomopathogens. In: *Insect Pathology* (Ed.: Vega, F.E., Kaya, H.K.). 2<sup>nd</sup> Edn. Academic Press, San Diego. pp.221-263.
- Thomas, S.R., Elkinton, J.S., 2004. Pathogenicity and virulence. *J. Invertebr. Pathol.* 85, 146-151.
- Yup-lian, L., 1991. Silkworm Disease. Rome: FAO Agricultural Services Bulletin 73/4. 75 p.
- Tu, Z., Liu, Q., Wan, Y., 2011. Reduced infectivity and environmental resistance of microsporidian *Nosema bombycis* after spore wall proteins-removal treatment. *Afr. J. Microbiol. Res.* 5(30), 5319-5324.

**How to cite this article:**

Nuraeni, S., Sanusi, D., Kuswinanti, T., Nasaruddin, A., 2016. Infectivity of two isolates of Microsporidia on silkworm (*Bombyx mori* L.) in South Sulawesi. *Int. J. Curr. Res. Biosci. Plant Biol.* 3(5), 1-5. doi: <http://dx.doi.org/10.20546/ijcrbp.2016.305.001>

# Infectivity

---

## ORIGINALITY REPORT

---

% **19**  
SIMILARITY INDEX

% **13**  
INTERNET SOURCES

% **11**  
PUBLICATIONS

% **8**  
STUDENT PAPERS

---

## PRIMARY SOURCES

---

**1** [ijcrbp.com](http://ijcrbp.com) Internet Source %**4**

---

**2** Nadia Kermani, Zainal-Abidin Abu-hassan, Hamady Dieng, Noor Farehan Ismail, Mansour Attia, Idris Abd Ghani. "Pathogenicity of Nosema sp. (Microsporidia) in the Diamondback Moth, *Plutella xylostella* (Lepidoptera: Plutellidae)", PLoS ONE, 2013  
Publication %**3**

---

**3** [www.ijcrbp.com](http://www.ijcrbp.com) Internet Source %**3**

---

**4** Submitted to Mansoura University Student Paper %**2**

---

**5** [tel.archives-ouvertes.fr](http://tel.archives-ouvertes.fr) Internet Source %**2**

---

**6** Muhammad Aswad, Junya Chiba, Takenori Tomohiro, Yasumaru Hatanaka. "Evaluation of dipole moment and electrophilicity on the nature of click-type coupling reaction between %**1**

thioamide and sulfonyl azide", Tetrahedron Letters, 2016

Publication

- 
- 7 Darui Chen, Zhongyuan Shen, Feng Zhu, Rui Guan, Jiange Hou, Jiao Zhang, Xiaofang Xu, Xudong Tang, Li Xu. "Phylogenetic characterization of a microsporidium (Nosema sp. MPr) isolated from the *Pieris rapae*", Parasitology Research, 2012  
Publication % 1
- 
- 8 Shapiro-Ilan, David I., Denny J. Bruck, and Lawrence A. Lacey. "Principles of Epizootiology and Microbial Control", Insect Pathology, 2012.  
Publication % 1
- 
- 9 S Nuraeni, A Arif, A Sadapotto. "Testing of two microsporidia isolates towards breeds of silkworm resistance", IOP Conference Series: Earth and Environmental Science, 2019  
Publication % 1
- 
- 10 [media.neliti.com](http://media.neliti.com)  
Internet Source <% 1
- 
- 11 "Industrial Entomology", Springer Nature, 2017  
Publication <% 1
- 
- 12 Judith A McBride. "Developmental changes in the caudal and lateral processes of larvae of *Aspidomorpha deusta* (Fabricius) (Coleoptera: Chrysomelidae: Cassidinae)", Australian <% 1

# Journal of Entomology, 7/4/2000

Publication

---

**13** [library.wur.nl](http://library.wur.nl) <% 1  
Internet Source

---

**14** [journal.unhas.ac.id](http://journal.unhas.ac.id) <% 1  
Internet Source

---

**15** Submitted to Miami University of Ohio <% 1  
Student Paper

---

EXCLUDE QUOTES ON

EXCLUDE  
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5  
WORDS