

ia_Singgani_and_Tipo_Arum-
Andi_Ummu-Atjo-Arifin-Apik-
Suriah.pdf

by

Submission date: 24-Dec-2021 01:52PM (UTC+0700)

Submission ID: 1735443595

File name: ia_Singgani_and_Tipo_Arum-Andi_Ummu-Atjo-Arifin-Apik-Suriah.pdf (302.03K)

Word count: 4212

Character count: 19994

Risk factors of anemia among pregnant women in community health center (Puskesmas) Singgani and Puskesmas Tipo Palu¹



Arum Diah Pusporini^{a,*}, Andi Ummu Salmah^b, Atjo Wahyu^c, Arifin Seweng^b, Apik Indarty^b, Suriah^d, Rosmala Nur^e, Aminuddin Syam^f, Mahfudz^g

^a Student, Faculty of Public Health, Hasanuddin University, Indonesia

^b Lecturer in Reproductive Health, Faculty of Public Health, Hasanuddin University, Indonesia

^c Lecturer in Occupational Health and Safety, Faculty of Public Health, Hasanuddin University, Indonesia

^d Lecturer in Health Promotion and Behavioral Sciences, Faculty of Public Health, Hasanuddin University, Indonesia

^e Lecturer in Public Health, Faculty of Public Health, Tadulako University, Indonesia

^f Lecturer in Nutrition Department, Faculty of Public Health, Hasanuddin University, Indonesia

^g Agroecology Department, Faculty of Agriculture, Tadulako University, Indonesia

ARTICLE INFO

30 Article history
Received 28 June 2021
Accepted 30 July 2021

Keywords:
Anemia
Nutritional status
Pregnant women

ABSTRACT

Objective: The aim of this study is to describe the risk factors of anemia among pregnant women.
Method: We used an observational analytic study with a matched case-control study design. The sampling method used in this study is a simple random sampling technique. The sample size in this study is 138 samples that consist of 46 cases and 92 controls. The data obtained from patient medical records and analyzed statistically using the chi-square test.
Results: Nutritional status is a risk factor of anemia among pregnant women in Community Health Center (Puskesmas) Singgani and Puskesmas Tipo. The risk of pregnant women with chronic energy deficiency (CED) developing anemia is higher in Puskesmas Singgani compared to in Puskesmas Tipo.
Conclusion: Prevention can be done by counseling the bride and groom about pregnancy preparation and counseling the pregnant women to pay attention to the nutritional intake, particularly the consumption of folic acid supplements and iron.

© 2021 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

36 Introduction

Anemia in pregnancy is a condition that hemoglobin (Hb) level in blood is less than 11.0gr% as a result of the inability of red blood cell production to maintain normal Hb concentration levels.¹ Anemia in pregnancy is defined as a hemoglobin level of the mother less than 11.0gr% during the first trimester or hemoglobin level lower than 10.5gr% during the second and third trimester.^{2,3}

World Health Organization (WHO) reported the worldwide prevalence of anemia in pregnant women is 40% in 2016.⁴ In Indonesia, based on RISKESDAS 2018, the prevalence of anemia among pregnant women remains as high as 48.9% or nearly half of the pregnant women population in Indonesia have anemia. The number is higher compared to the amount in 2013 of 37.1%. The highest proportion of anemia among pregnant women is in the age of 15–24 years old, which is 84.6%, 33.7% with the age of 25–34 years old, 37.1% with the age of 35–44 years old, and 24% in pregnant women with the age of 45–54 years old.⁵ The proportion of anemia pregnant women in Central Sulawesi, especially Palu in

2016, showed that anemia in pregnant women reached 19.52%, an increase of 7.52% when compared to 2015 of 12%.^{6,7}

Pregnant women who are anemic can be at risk of increased mortality and morbidity in both the mother and the baby.⁸ Anemic pregnant women can be at risk of abortion, preterm labor, premature rupture of membrane, postpartum hemorrhage, and maternal death. Infants with an anemic mother may be at risk for low birth weight, birth defects, perinatal death, and low infant intelligence. The several causes of anemia in pregnancy are iron deficiency, infection, folic acid deficiency, and hemoglobin abnormalities.^{2,5,9,10}

Material and method

We used an observational analytic study with a matched case-control study design. The dependent variable of this study is the anemia status, while the independent variable is the factors that are thought to be affecting the incidence of anemia in pregnant women, which are oxytocin massage and breast care. The sampling method used in this study is a simple random sampling technique. The sample size in this study is 138 samples consisting of 46 cases and 92 control. The data obtained from patient medical records and analyzed statistically using the chi-square test using SPSS software application version 22.

Peer-review under responsibility of the scientific committee of the 3rd International Nursing, Health Science Students & Health Care Professionals Conference. Full text and the content of it is under responsibility of authors of the article.

* Corresponding author.

E-mail addresses: arumdiahts@has.ac.id, pmw@agccornhas.ac.id (A.D. Pusporini).

<https://doi.org/10.1016/j.gaceta.2021.10.010>

0213-9111/© 2021 SESPAS. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Sample characteristic pregnant women in the Singgani health center.

Variable	Anemia		Nonanemia	
	n	%	n	%
Maternal age				
<20 years	3	13.0	0	0
20-35 years	15	65.2	41	89.1
>35 years	5	21.8	5	10.9
Total	23	100	46	100
Occupational				
Didn't work	12	52.2	29	63
Work	11	47.8	17	37
Total	23	100	46	100
Pregnancy spacing				
<2 years	4	17.4	6	13.0
≥2 years	19	82.6	40	87.0
Total	23	100	46	100
History of miscarriage				
Has miscarriage	5	21.7	9	19.6
Hasn't miscarriage	18	78.3	37	80.4
Total	23	100	46	100
Frequency of ANC				
High risk (<1 x tmt1; <1 x tmt2; <2 x tmt3)	14	60.9	19	41.3
Low risk (≥1 x tmt1; ≥1 x tmt2; ≥2 x tmt3)	9	39.1	27	58.7
Total	23	100	46	100
Nutritional status				
CED	8	34.8	1	2.2
Non-CED	15	65.2	45	97.8
Total	23	100	46	100

Results

Sample characteristics

Distribution of the sample characteristics of pregnant women in the Singgani health center includes maternal age, occupational status, pregnancy spacing, history of miscarriage, frequency of antenatal care, and nutritional status can be seen in Table 1.

Based on Table 1, it is known that the proportion of mothers with age less than 20 years old only exists in 13.0% of the case group while in the control group, there is none. Then, the maternal age of more than 35 years old is more in the case group, which is 21.8% compared to the control group is 10.9%. While mothers with the age of 20-35 years old in the case group are 60.9%, and the control group is 89.1%. The proportion of mothers who did not work in the case group was 52.2%, and the control group was 63%.

Most of the pregnancy spacing ≥ 2 years were 19 samples (82.6%) in the case group and 40 samples (87%) in the control group. The proportion of miscarriage history in the case group and the control group is similar in which there is a low more expectant mother with no history of a miscarriage of 80.4% in the control group and 78.3% in the case group. The proportion of the frequency of high-risk pregnancy examinations is the majority in the case group at 60.9% compared to the control group at 41.3%. The proportion of mothers with CED nutritional status in the case group was 34.8%. This number is higher when compared to the control group, which is only 2.2%.

Based on Table 2, it can be seen that the proportion of mothers with the age of fewer than 20 years old in the case group is 21.8%, while in the control group, 6.5%. The mother with the age of more than 35 years old is more in the case group, which is 13.0% compared to the control group, which is 2.2%. While mothers with the age of 20-35 years in the case group are 65.2% and the control group 91.3%.

The proportion of mothers who did not work in the case group was 69.6%, and the control group was 50%. The most common pregnancy spacing in the case group and the control group is the low-risk pregnancy spacing (87%). The proportion of frequency of antenatal care in the case group was more low risk, which was

Table 2
Sample characteristic pregnant women in Tipo Health Center.

Variable	Anemia		Nonanemia	
	n	%	n	%
Maternal age				
<20 years	5	21.8	3	6.5
20-35 years	15	65.2	42	91.3
>35 years	3	13.0	1	2.2
Total	23	100	46	100
Occupational				
Didn't work	16	69.6	23	50
Work	7	30.4	23	50
Total	23	100	46	100
Pregnancy spacing				
<2 years	3	13	6	13
≥2 years	20	87	40	87
Total	23	100	46	100
History of miscarriage				
Has miscarriage	3	13	4	8.7
Hasn't miscarriage	20	87	42	91.3
Total	23	100	46	100
Frequency of ANC				
High risk (<1 x tmt1; <1 x tmt2; <2 x tmt3)	11	47.8	16	34.8
Low risk (≥1 x tmt1; ≥1 x tmt2; ≥2 x tmt3)	12	52.2	30	65.2
Total	23	100	46	100
Nutritional status				
CED	14	60.9	5	10.9
Non-CED	9	39.1	41	89.1
Total	23	100	46	100

52.2%. Similarly, the control group was more low risk at 65.2%. The proportion of case groups with CED nutritional status was 52.2%. This number is greater when compared to the control group, which is only 10.9%.

Bivariate analysis of risk factors for anemia among pregnant women in Community Health Center (Puskesmas) Singgani and Puskesmas Tipo

Risks of each variable in this study can be seen in the following table.

Based on Table 3, the statistical test results of pregnancy spacing variables at the Puskesmas Singgani obtained an OR value of 1.404, but this was not significant because the 95% CI range included one (95% CI = 0.354-5.568). Statistical test results of pregnancy spacing at Puskesmas Tipo obtained OR values of 1.00. It is also not significant because the 95% CI range includes one (95% CI = 0.226-4.420).

Statistical test results of miscarriage history at the Puskesmas Singgani obtained an OR value of 1.142, but this is not significant because the 95% CI range includes one (95% CI = 0.334-3.906). Statistical test results of miscarriage history at Puskesmas Tipo obtained OR values of 1.575. This is also not significant because the 95% CI range includes one (95% CI = 0.322-7.714).

Statistical test results of the frequency of antenatal care variables at Puskesmas Singgani obtained an OR value of 2.211, but this was not significant because the 95% CI range included one (95% CI = 0.795-6.146). Statistical test results on the frequency of antenatal care at the Puskesmas Tipo obtained an OR value of 1.719. It is also not significant because the 95% CI range includes one (95% CI = 0.621-4.760).

Statistical test results of nutritional status variables at Puskesmas Singgani obtained an OR value of 24, which means that the nutritional status variable is a risk factor because the OR value is more than one. This is significant because the 95% CI value range does not include the number one (95% CI = 2.769-207.987). Pregnant women with chronic energy deficiency (CED) are at risk of anemia by 24 times. The statistical test results of nutritional status variables at Puskesmas Tipo obtained an OR value of 12.75, which means that the nutritional status variable is a risk factor because

Table 3
Bivariate analysis of risk factors for anemia among pregnant women in Puskesmas Singgani and Puskesmas Tipo.

Variable	Cases n	Controls n	p	OR	95% CI
Pregnancy spacing					
<i>Singgani Health Center</i>					
High risk (<2 years)	4 (17.4)	6 (13.0)	0.721	1.404	0.354–5.568
Low risk (≥2 years)	19 (82.6)	40 (87.0)			
<i>Tipo Health Center</i>					
High risk (<2 years)	3 (13)	6 (13.0)	1.001	1.00	0.226–4.420
Low risk (≥2 years)	20 (87)	40 (87.0)			
Miscarriage history					
<i>Singgani Health Center</i>					
High risk (has miscarriage)	5 (21.7)	9 (19.6)	1.01	1.142	0.334–3.906
Low risk (hasn't miscarriage)	18 (78.3)	37 (80.4)			
<i>Tipo Health Center</i>					
High risk (has miscarriage)	3 (13)	4 (8.7)	0.679	1.575	0.322–7.714
Low risk (hasn't miscarriage)	20 (87)	42 (91.3)			
Frequency of ANC					
<i>Singgani Health Center</i>					
High risk (<1 × tmt1; <1 × tmt2; <2 × tmt3)	14 (60.9)	19 (41.3)	0.201	2.211	0.795–6.146
Low risk (≥1 × tmt1; ≥1 × tmt2; ≥2 × tmt3)	9 (39.1)	27 (58.7)			
<i>Tipo Health Center</i>					
High risk (<1 × tmt1; <1 × tmt2; <2 × tmt3)	11 (47.8)	16 (34.8)	0.433	1.719	0.621–4.760
Low risk (≥1 × tmt1; ≥1 × tmt2; ≥2 × tmt3)	12 (52.2)	30 (65.2)			
Nutritional status					
<i>Singgani Health Center</i>					
CEd (arm circumference <23.5 cm)	8 (34.8)	1 (2.2)	0.001	24	2.769–207.987
Non-CEd (arm circumference ≥23.5 cm)	15 (65.2)	45 (97.8)			
<i>Tipo Health Center</i>					
CEd (arm circumference <23.5 cm)	14 (50.9)	5 (10.9)	0.001	12.756	3.653–44.534
Non-CEd (arm circumference ≥23.5 cm)	9 (39.1)	41 (89.1)			

the OR value is more than one. This is also significant because the 95% CI value range does not include the number one (95% CI = 3.653–44.534), so it can be concluded that expectant mothers with chronic energy deficiency are at risk of 12.75 times developing anemia in their pregnancy later on.

Discussion

Analysis of risk factors of anemia among pregnant women in Puskesmas Singgani and Puskesmas Tipo

Pregnant women with pregnancy spacing that is too close, which is less than two years, are at risk of anemia because the mother's body has not reserved adequate nutrients after going through the previous pregnancy so that the iron in her body is divided for recovery of her body and to fulfill the needs during the next pregnancy.¹² It takes a minimum of 2 years to restore iron reserves to normal on the condition of consuming foods that contain protein and iron. This period also allows the body to restore its physiological and anatomical functions. But in this study, there was no significant effect between pregnancy spacing and the incidence of anemia in pregnant women. This is because both the case and control groups mostly have a pregnancy interval of more than 2 years.^{13,14}

Based on the bivariate test results, it is known that the pregnancy spacing is not a risk factor for anemia among pregnant women in Puskesmas Singgani and Puskesmas Tipo. The results of this study are in line with the research conducted by Tanzaha et al. (2017) in Indonesia. From the results of the previous study, the author obtained a value of 95% CI, includes the number 1, which means there is no relationship between the pregnancy spacing with the incidence of anemia in pregnant women.¹⁵ Another study conducted by Anggraini et al. (2018) at Tanjung Pinang Health Center showed that pregnancy spacing increased the risk of anemia by 15.483 times.¹⁶ Likewise, the results of the study of Getahun et al. in South Ethiopia with the results showing a pregnancy

spacing ≥2 years are a protective factor of anemia among pregnant women.^{13,17}

A history of miscarriage can increase the risk of anemia in subsequent pregnancies due to an increase in previous blood loss, thereby decreasing iron reserves in the body. Statistical test results showed that a history of miscarriage is not a risk factor for anemia among pregnant women at the Puskesmas Singgani and Tipo. A history of miscarriage was previously investigated by Belete et al. (2019) in Northern Ethiopia; from the results of the study found that pregnant women with a history of miscarriage have a 7.9 times risk of anemia in subsequent pregnancies compared with those who have never miscarried.^{18,19}

Pregnancy examination or antenatal care is an examination given to expectant women by health workers during their pregnancy, with a standard number of visits during pregnancy at least four times, including history taking, general physical and obstetric examination, laboratory examinations for certain indications as well as basic and special indications. An examination of the pregnancy carried out early will allow the discovery of abnormalities or health problems faced by the mother during the process of pregnancy, so that any measures can be taken to save the fetus and the mother.

The ANC service standard includes the number of visits and the type of inspection carried out. ANC is done at least four times during pregnancy, namely once in the first trimester, once in the second trimester, and at least twice in the third trimester. The type of examination consists of TT, namely weight checking, blood pressure measurement, checking the height of the fundus uteri, giving TT immunization, administering supplemental iron tablets, testing for sexually transmitted disease, and colloquium in the context of referral preparation. Routine pregnancy examination can reduce the risk of anemia in pregnant women because it can be detected as early as possible and also given iron tablets at each visit.

Statistical test results showed that the frequency of antenatal care is not a risk factor for anemia in pregnancy at the Puskesmas Singgani and Tipo. The results of this study are in line with a study conducted by Tanzaha et al. (2017), where there is no relationship

between antenatal care or ANC with the incidence of anemia in pregnant women.¹⁵ The results of this study differ from studies conducted by Singal et al. (2018) in India with the results obtained p -value = 0.001, which means there is a relationship between lack of antenatal care (ANC) examination with the incidence of anemia in pregnant women.^{20,21}

During physiological pregnancy, changes occur, one of which is an increase in the volume of fluid and red blood cells and a decrease in the concentration of protein binding nutrients in the blood circulation, as well as a decrease in micronutrients. Pregnancy is a period of growth and development of the fetus towards birth, so that nutritional disorders that occur during pregnancy will have a major impact on the health of the mother and fetus. Therefore chronic energy deficiency status (CED) in pregnant women have an impact on the incidence of anemia in pregnant women as well as the incidence of low birth weight (LBW) and stunting.⁹

Based on the bivariate test results, it is known that nutritional status is a risk factor for anemia among pregnant women at the Puskesmas Singgani and Puskesmas Tipo. Pregnant women with chronic energy deficiency conditions at the Puskesmas Singgani are 24 times at risk of anemia compared to pregnant women who do not experience CED. Whereas pregnant women who developed CED in Puskesmas Tipo have a risk of 12.75 times are developing anemia in their pregnancy. This is reinforced by the results of previous studies conducted by Lestari et al. (2018) in Medan, North Sumatra, where there is an association between chronic energy deficiency with the incidence of anemia in pregnancy.²² The same results were obtained from the study of Tekeste et al. (2015) with a large risk of developing anemia by 2.52 times.²³

Chronic energy deficiency (CED) is a condition of the body that lacks macronutrients (carbohydrates, protein, and fat) for a long period of time, characterized by the size of the upper arm circumference of less than 23.5 cm. Macronutrient deficiencies are associated with micronutrient deficiencies, especially vitamin A, vitamin D, folic acid, iron, zinc, calcium, and iodine. The condition of a pregnant CED can cause anemia in her pregnancy.²⁴ Nutritional status is a risk factor of anemia among pregnant women in Puskesmas Singgani and Puskesmas Tipo. It happens because nutritional status is a factor that plays a direct role in the nutritional needs of pregnant women and their fetuses so that the lack of nutritional intake in pregnant women with CED can cause anemia in pregnancy. The risk of pregnant women with CED to develop anemia in the Puskesmas Singgani is higher than the Puskesmas Tipo. This is influenced by the diet of pregnant women, especially food sources of iron, and adherence to the consumption of supplemental iron tablets during pregnancy.

Conclusion

The results of this study are nutritional status is a risk factor for anemia among pregnant women in Puskesmas Singgani and Tipo. We expected that midwives would work closely with active kader posyandu in screening pregnant women with anemia, especially with chronic energy deficiency conditions. Prevention can be done by counseling the bride and groom about pregnancy

preparation and counseling the pregnant women to pay attention to the nutritional intake, particularly the consumption of folic acid supplements and iron.

Conflict of interest

The authors would like to thank the Dean and the all vice-dean, the Faculty of Public Health at Hasanuddin University, for the research funding.

The authors declare no conflict of interest.

References

1. Organization WH. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. World Health Organization; 2011.
2. Edmonds K. Dewhurst's textbook of obstetrics and gynaecology. London: John Wiley & Sons; 2018.
3. Rosmala-Nur SM, Demak BPK. Mother's behavior in pregnancy-puerperal treatments and reproductive health disorders. *Med Leg Update*. 2020;20:1341-6.
4. The World Bank. Prevalence of anemia among pregnant women (%). Washington, DC: The World Bank Group; 2019.
5. Kemenkes RI. Hasil utama riskesdas 2018. Jakarta: Kemenkes RI; 2018.
6. Dinas Kesehatan Kota Palu. Profil Kesehatan Kota Palu Tahun 2016. 2017.
7. Nur R, et al. The effect of moringa leaf extract on increasing hemoglobin and bodyweight in post-disaster pregnant women. *Enferm Clin*. 2020;30:79-82.
8. Nur R, et al. Detection mapping of women with high-risk pregnancy in antenatal care in Kamoni Public Health Center, Palu City, Indonesia. *Syst Rev Pharm*. 2020;11:642-7.
9. Danforth DN. Danforth's obstetrics and gynecology. Philadelphia: Lippincott Williams & Wilkins; 2008.
10. Daru J, et al. Risk of maternal mortality in women with severe anaemia during pregnancy and post partum: a multilevel analysis. *Lancet Glob Health*. 2018;6:e548-54.
11. Stang. Cara praktis penentuan uji statistik dalam penelitian kesehatan dan kedokteran. Jakarta: Mitra Wacana Media; 2018.
12. Nur R, et al. The smart book effect of pregnancy postpartum care towards the husband's knowledge and reduction of reproductive health problems. *Fam Med Prim Care Rev*. 2020;22:133-9.
13. Getahun W, Bealchew T, Wolde AD. Burden and associated factors of anemia among pregnant women attending antenatal care in southern Ethiopia: cross sectional study. *BMC Res Notes*. 2017;10:1-7.
14. Nur R, Fitriyah S, Mallongi A. Women's reactions and health disorders caused by abuse during the pregnancy-postpartum period. *Med Leg Update*. 2020;20:1329-34.
15. Tanzha I, et al. Faktor risiko anemia ibu hamil di Indonesia. *J Gizi dan Pangan*. 2016;11:143-52.
16. Anggraini PD. Factors related to the work area of the Tanjung Finang Health Center in 2018. *J Midwifery*. 2018;7:33-8.
17. Amri I, et al. Number pregnancy and low birth weight (LBW) babies in Aunatapura Hospital in Palu City in 2016. *Enferm Clin*. 2020;30:219-22.
18. Berhe B, et al. Prevalence of anemia and associated factors among pregnant women in Adigrat General Hospital, Tigrai, northern Ethiopia, 2018. *BMC Res Notes*. 2019;12:1-6.
19. Murtiarsani D, et al. Determinants of the incidence of stunting in the working area of Kinovaro Sigi Health Center. *Enferm Clin*. 2020;30:246-52.
20. Singal N, et al. Factors associated with maternal anaemia among pregnant women in rural India. *Bangladesh J Med Sci*. 2018;17:583-92.
21. Nur R, et al. Early age marriage and the impact of health reproduction women. *J Eng Appl Sci*. 2019;4.
22. Lestari S, et al. The prevalence of anemia in pregnant women and its associated risk factors in North Sumatera, Indonesia. In: IOP conference series: earth and environmental science. IOP Publishing; 2018. p. 12195.
23. Okube OI, et al. Prevalence and factors associated with anaemia among pregnant women attending antenatal clinic in the second and third trimesters at punyani maternity hospital, Kenya. *J Obs Gynecol*. 2016;6.
24. Ministry of Health of the Republic of Indonesia. Guidelines for dealing with chronic energy deficiency in pregnant women. Jakarta: Ministry of Health of the Republic of Indonesia; 2015.

ia_Singgani_and_Tipo_Arum-Andi_Ummu-Atjo-Arifin-Apik-Suriah.pdf

ORIGINALITY REPORT

16%

SIMILARITY INDEX

9%

INTERNET SOURCES

12%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

- 1 S Lestari, I I Fujiati, D Keumalasari, M Daulay, S J Martina, S Syarifah. "The prevalence of anemia in pregnant women and its associated risk factors in North Sumatera, Indonesia", IOP Conference Series: Earth and Environmental Science, 2018
Publication 1%
- 2 journal.ipb.ac.id
Internet Source 1%
- 3 www.ncbi.nlm.nih.gov
Internet Source 1%
- 4 www.dovepress.com
Internet Source 1%
- 5 Fusun Yaris, Cunay Ulku, Murat Kesim, Mine Kadioglu, Mesut Unsal, Mustafa Fevzi Dikici, Nuri Ihsan Kalyoncu, Ersin Yaris. "Psychotropic drugs in pregnancy: a case-control study", Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2005 1%

Publication

6	Submitted to Universitas Hasanuddin Student Paper	1 %
7	ejournal2.litbang.kemkes.go.id Internet Source	1 %
8	Submitted to Universitas Diponegoro Student Paper	1 %
9	A Wahyudi, S Ritohardoyo, Sunarto, A J Pitoyo. "10 years of flood events, study of the adaptation capacity of the people living in the Riparian Area of the Madiun River", IOP Conference Series: Earth and Environmental Science, 2020 Publication	1 %
10	worldnutrijournal.org Internet Source	1 %
11	Taofeeq D. Moshood, Gusman Nawanir, Fatimah Mahmud, Fazeeda Mohamad, Mohd Hanafiah Ahmad, Airin AbdulGhani. "Why do consumers purchase biodegradable plastic? The impact of hedonics and environmental motivations on switching intention from synthetic to biodegradable plastic among the young consumers", Journal of Retailing and Consumer Services, 2022 Publication	<1 %

12 Ebrahim Mohammed. "Magnitude of Anemia and Associated Factors Among Pregnant Women Visiting Public Health Institutions for Antenatal Care Services in Adama Town, Ethiopia", Central African Journal of Public Health, 2018
Publication

13 repository.tnmgrmu.ac.in
Internet Source

14 Ina Kusrini, Donny Kristanto Mulyantoro, Dwi Hapsari Tjandrarini, Hadi Ashar. "Profile of Double of Undernutrition Problem, Coexistence with Anemia among Pregnant Women Indonesia 2018: A Cross Sectional Survey", Open Access Macedonian Journal of Medical Sciences, 2021
Publication

15 Fabio Barili, Nicholas Freemantle, Francesco Musumeci, Barbara Martin et al. "Five-year outcomes in trials comparing transcatheter aortic valve implantation versus surgical aortic valve replacement: a pooled meta-analysis of reconstructed time-to-event data", European Journal of Cardio-Thoracic Surgery, 2021
Publication

16 Mihaela Tomită. "Comparative Analysis of Juvenile Delinquency and Non Delinquency",

Procedia - Social and Behavioral Sciences,
2013

Publication

-
- | | | |
|-----------|--|----------------|
| 17 | Hong Ji, Ronghao Chen, Yong Huang, Wenqin Li, Chunhui Shi, Juan Zhou. "Effect of simulation education and case management on glycemic control in type 2 diabetes", <i>Diabetes/Metabolism Research and Reviews</i> , 2018
Publication | <1 % |
| <hr/> | | |
| 18 | K Sayuti, R Yenrina, Y Febri. "Characteristic of Analogue Jerky Made from Moringa Leaves (<i>Moringa oleivera</i> L) with the addition of Tapioca Flour", <i>IOP Conference Series: Earth and Environmental Science</i> , 2020
Publication | <1 % |
| <hr/> | | |
| 19 | Ulfa Malika Putri Raharja, Waryana Waryana, Almira Sitasari. "The economic status of parents and family food security as a risk factor for stunting in children under five years old in Bejiharjo Village", <i>Ilmu Gizi Indonesia</i> , 2019
Publication | <1 % |
| <hr/> | | |
| 20 | ehjournal.biomedcentral.com
Internet Source | <1 % |
| <hr/> | | |
| 21 | ijop.net
Internet Source | <1 % |
-

- 22 www.ejmjih.com Internet Source <1 %
-
- 23 Darmawati Darmawati, Tongku Nizwan Siregar, Hajjul Kamil, Teuku Tahlil. "Exploring Indonesian mothers' perspective on anemia during pregnancy: A qualitative approach", Research Square, 2020 Publication <1 %
-
- 24 Hong Ji, Ronghao Chen, Yong Huang, Wenqin Li, Chunhui Shi, Juan Zhou. "Effect of simulation education and case management on glycemic control in type 2 diabetes patients", Diabetes/Metabolism Research and Reviews, 2018 Publication <1 %
-
- 25 Mithun Mog, Koustav Ghosh. "Prevalence of anaemia among women of reproductive age (15–49): A spatial-temporal comprehensive study of Maharashtra districts", Clinical Epidemiology and Global Health, 2021 Publication <1 %
-
- 26 Mohammed Hasen Badeso, Falaho Sani Kalili, Mohammed Seid Abdulle. "Determinants of Anemia Among Pregnant Women Attending Antenatal Care in Bale-Robe Town Health Facilities, Bale zone, Southeast Ethiopia: A Case-Control Study", Research Square Platform LLC, 2021 <1 %

Publication

- | | | |
|----|---|------|
| 27 | Wellings, Kaye, Mitchell, Kirstin, Collumbien, Martine. "EBOOK: Sexual Health: A Public Health Perspective", EBOOK: Sexual Health: A Public Health Perspective, 2012
Publication | <1 % |
| 28 | contraceptionmedicine.biomedcentral.com
Internet Source | <1 % |
| 29 | journal2.unusa.ac.id
Internet Source | <1 % |
| 30 | link.springer.com
Internet Source | <1 % |
| 31 | persagi.org
Internet Source | <1 % |
| 32 | research.monash.edu
Internet Source | <1 % |
| 33 | safpj.co.za
Internet Source | <1 % |
| 34 | www.glowm.com
Internet Source | <1 % |
| 35 | Tria Mailan Karemoi, Wiwi Mardiah, Fanny Adistie. "Factors Affecting Nutritional Status of Pregnant Women", Asian Community Health Nursing Research, 2020
Publication | <1 % |

36 Alemneh Mekuriaw Liyew, Getayeneh Antehunegn Tesema, Tesfa Sewunet Alamneh, Misganaw Gebrie Worku et al. "Prevalence and determinants of anemia among pregnant women in East Africa; A multi-level analysis of recent Demographic and Health Surveys", PLOS ONE, 2021
Publication

<1%

37 E.L. Achadi, M.J. Hansell, N.L. Sloan, M.A. Anderson. "Women's nutritional status, iron consumption and weight gain during pregnancy in relation to neonatal weight and length in West Java, Indonesia", International Journal of Gynecology & Obstetrics, 1995
Publication

<1%

38 ijph.tums.ac.ir
Internet Source

<1%

Exclude quotes
Exclude bibliography

Exclude matches < 5 words