

# Effect of Trichoderma and tofu waste based organic fertilizer on the fruit development of chili (*Capsicum annuum* L.)

*by* Katriani Mantja

---

**Submission date:** 24-Oct-2021 08:18AM (UTC+0700)

**Submission ID:** 1682105426

**File name:** izer\_on\_the\_fruit\_development\_of\_chili\_\_Capsicum\_annuum\_L\_.pdf (301.79K)

**Word count:** 3465

**Character count:** 17728

PAPER · OPEN ACCESS

## Effect of Trichoderma and tofu waste based organic fertilizer on the fruit development of chili (*Capsicum annuum* L.)

To cite this article: K Mantja *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **486** 012116



View the [article online](#) for updates and enhancements.

## Effect of Trichoderma and tofu waste based organic fertilizer on the fruit development of chili (*Capsicum annuum* L.)

K Mantja<sup>1</sup>, Fitrianti<sup>1</sup>, M Farid<sup>1</sup>, F Ulfa<sup>1</sup>, A Mollah<sup>1</sup>, A R Amin<sup>1</sup>, I Ridwan<sup>1</sup> and Kasmiaty<sup>2</sup>

<sup>1</sup>Department of Agronomy, Faculty of Agriculture, Universitas Hasanuddin, Jl. Perintis Kemerdekaan KM 10, Makassar, 90245, South Sulawesi, Indonesia.

<sup>2</sup>Universitas Sulawesi barat, Indonesia

E-mail: ifayanti@unhas.ac.id

**Abstract.** The study aimed to determine the effect of the application of Trichoderma and liquid organic fertilizer from tofu wastes on the growth and production of large chilli plants (*Capsicum annuum* L.). The study was conducted at the Faculty of Agriculture Experimental Farm, Universitas Hasanuddin, Makassar, from April to August 2017. This study used a factorial two-factor randomized block design with three replications. The first factor of Trichoderma dose consisted of four levels (control, 10 g plant<sup>-1</sup>, 15 g plant<sup>-1</sup>, and 20 g plant<sup>-1</sup>). The second factor was the concentration of liquid organic fertilizer consisted of four levels (control, 10 ml L<sup>-1</sup>, 15 ml L<sup>-1</sup>, and 20 ml L<sup>-1</sup>). The results show that the interaction between Trichoderma treatment with a dose of 10 g plant<sup>-1</sup> and liquid organic fertilizer 15 ml L<sup>-1</sup> gave a higher number of fruits per plant. Trichoderma treatment with a dose of 20 g plant<sup>-1</sup> gave the heaviest fruit weight and fruit weight per plant. There is no one concentration of liquid organic fertilizer which gives better growth and production of chili.

### 1. Introduction

One type of vegetable that is quite important in Indonesia is red chili (*Capsicum annuum* L.) both as a commodity consumed domestically and as an export commodity. The red chili has a fairly high economic value apart from the nutritional value. Indonesian people are among the biggest fans of chili in the world, so that chili is one of the important products in Indonesian food, it can even affect the rate of inflation. With a population of 237.6 million, Indonesia needs 118,800 tons of chili per year [1]. Despite the demand, increased chili production can have an impact on the environment due to the use of chemical fertilizers. The use of artificial chemical fertilizers without being accompanied by the provision of organic material can cause the soil to become barren and decrease productivity and pest dispersion [2,3].

Organic fertilizers play an essential role in increasing the chemical, physical, and biological content in the soil as well as being a good source of plant nutrition [4]. The use of organic fertilizer which is more effective and efficient is in the form of liquid fertilizer. Liquid organic fertilizer is more easily absorbed by plants because the elements in it have been decomposed. Plants not only absorb nutrients through the roots but can also through the leaves of the plants. The use of liquid fertilizer is easier to work and its use, in one application liquid organic fertilizer performs three kinds of processes at once, namely fertilizing plants, watering plants and treating plants [5]. In the making of liquid organic fertilizer, there are several organic materials that can be used as alternative nutrients including cow



manure, tofu waste, coconut water and rice washing water. Cow manure is manure derived from cow dung which is good for improving fertility, physical, chemical and biological soil properties, increasing macro and micro nutrients, increasing water holding capacity and increasing cation exchange capacity [6]. Besides cow manure, industrial wastes such as tofu waste from tofu factory can be used. With a protein content of 30.2% (wet) and 34.9% (dry) [7], tofu waste contains enough nutrients needed for plant growth [8]. In addition, rice washing water which is commonly known by the term *leri* in Indonesia, still contains carbohydrates, protein and vitamin B which is mostly contained in vitamin B1 or thiamine [9]. Other compounds found in coconut water are protein, fat, minerals, carbohydrates, even complete with vitamin C and B complex. Proteins and carbohydrates needed by plants as food reserves, fats needed by plants as energy reserves, minerals as constituents of the body's plants, and vitamins C and B complex play a role in the metabolic process. Thus, coconut water can be used to spur growth in both shoots and roots in various types of plants [10].

Increased production and productivity strived by the application of innovative technology. One effort that can be done is the use of biofertilizers (utilization of soil microbes and natural fertilizers). One of the functional microorganisms known as biofungicides is the trichoderma fungus. Trichoderma as a decomposing organism, can also function as a biological agent of plant growth stimulators [11]. Trichoderma is able to increase plant growth and development, especially on the growth of roots that are more numerous and stronger because besides living on the root surface, the colonies can enter the epidermal layer of the root even deeper which then produces or releases various substances that can stimulate the formation of the body's defense system in plants [29] it is clear that this fungus is not pathogenic or parasitic to its host plant. Trichoderma isolates are found under the mahogany stand in Takalar District [12]. The results showed that plants containing Trichoderma colonies on the root surface only needed less than 40% nitrogen fertilizer compared to roots without colonies [13]. Giving Trichoderma as much as 15 g/plant gives the best production of growth in tomato plants [14]. *Trichoderma harzianum* functions as a biological agent, improves plant roots and stimulates lateral root formation [15]. While *T. asperellum* produces trichodermaerin antibiotics and several enzymes capable of lysis of cell walls such as lipase, NAGase, B-1.3-glucanase, B-glucosidase, and protease [16,17]. In addition, *T. asperellum* can stimulate plant resilience, growth and development resulting in increased crop production [18].

## 2. Methodology

This research was carried out in the Experimental Farm of the Faculty of Agriculture, Universitas Hasanuddin, Makassar, from April to August 2017 located about 16 meters above sea level (m asl). The research was conducted in the form of a two-factor factorial experiment based on the Randomized Blocked Design pattern. The first factor was the dose of Trichoderma which consisted of four levels, namely: t0 = control, t1 = 10 g plant<sup>-1</sup>, t2 = 15 g plant<sup>-1</sup>, and t3 = 20 g plant<sup>-1</sup>. While the second factor was the concentration of liquid organic fertilizer (P) consisting of four levels, namely: p0 = control, p1 = 10 ml L<sup>-1</sup>, p2 = 15 ml L<sup>-1</sup>, and p3 = 20 ml L<sup>-1</sup>. The chili seeds used in this research was variety of Pillar F1 produced by PT East West Seed Indonesia, while the Trichoderma used was *Trichoderma asperellum* inoculants.

Preparation of 20 liters of the liquid organic fertilizer with tofu waste based was carried out using volume per volume method with a composition of 15% tofu waste (3 liters), 50% cow manure (10 liters), 25% old coconut water (5 liters), and 10% rice washing water (2 liters). Solid material (cow manure and tofu wastes) was mixed then put into a sack with pores. Then the liquid ingredients (coconut water and rice washing water) were put into a container with crushed brown sugar (1 liter) and 12 liters of water added. After that the solid material was put into the container that has been filled with liquid material and fermented for about four weeks. Every two days the solution was stirred and the sack containing solid material was dipped so that the fermentation takes place evenly.

The growing media consists of a mixture of soil and compost in a ratio of 2:1 the filled into 30 x 40 cm polybag. Application of the Trichoderma was carried at planting by placed it at the planting hole in

the seedlings rhizosphere. The liquid organic fertilizer treatment was conducted started at 14 days after planting (DAP) and carried out in the morning by pouring it into the polybag every two weeks

Data collected and processed using analysis of variance (ANOVA) to test the effect of the treatment given. If there is a significant influence then proceed with further tests using the Tukey's Honestly Significant Difference (HSD) test at  $\alpha$  0.05 level.

### 3. Results

10

#### 3.1. Number of fruit per plant

Number of fruit per plant of chilli plant was significantly ( $p \leq 0.05$ ) affected by the interaction of Trichoderma and tofu waste based liquid organic fertilizer treatments. Average number of fruit per plant as affect by both treatment are shown in table 1. The Tukey's ( $\alpha = 0.05$ ) test results in Table 1 show that chilli plants applied with Trichoderma 10 g plant<sup>-1</sup> with a concentration of liquid organic fertilizer 15 ml L<sup>-1</sup> produces the highest value of 10.67 fruit plant<sup>-1</sup> which is no different significantly with the treatment application of Trichoderma 20 g plant<sup>-1</sup> and the concentration of liquid organic fertilizer 10 ml L<sup>-1</sup> the application of Trichoderma 0 g plant<sup>-1</sup> (control) with the concentration of liquid organic fertilizer 20 ml L<sup>-1</sup>, the application of Trichoderma 10 g plant<sup>-1</sup> with concentration of liquid organic fertilizer 10 ml L<sup>-1</sup>, and application of Trichoderma 10 g plant<sup>-1</sup> with concentration of liquid organic fertilizer 0 ml L<sup>-1</sup> (control). The treatment application of Trichoderma dose of 10 g plant<sup>-1</sup> and the concentration of liquid organic fertilizer 20 ml L<sup>-1</sup> gave the lowest yield of 6.00 fruits that were not significantly different from the treatment application of Trichoderma 20 g plant<sup>-1</sup> and the concentration of liquid organic fertilizer 20 ml L<sup>-1</sup>.

32


**Table 1.** Average number of fruit per plant (fruits plant<sup>-1</sup>) of chilli (*Capsicum annum* L.) on different dose of Trichoderma and concentration of liquid organic fertilizer.

Dose of Trichoderma	Concentration of liquid organic fertilizer				Tukey's HSD <sub>0.05</sub>
	0 ml L <sup>-1</sup>	10 ml L <sup>-1</sup>	15 ml L <sup>-1</sup>	20 ml L <sup>-1</sup>	
0 g plant <sup>-1</sup>	7.33 b	8.00 b	7.33 b	9.67 a	1.81
10 g plant <sup>-1</sup>	6.67 c	6.33 c	10.67 a	6.00 c	
15 g plant <sup>-1</sup>	10.00 a	8.67 b	6.33 c	7.67 b	
20 g plant <sup>-1</sup>	7.33 b	8.00 b	7.33 b	9.67 a	

Numbers followed by different letters (a, b, c) are significantly different at Tukey's test ( $\alpha=0.05$ ).

#### 3.2. Fruit length

Analysis of variance on the fruit length observation data show that there was no significant effect of all treatment on the fruit length of chilli plants. Figure 1 shows that the average fruit length in the Trichoderma application treatment dose of 20 g plant<sup>-1</sup> and the concentration of liquid organic fertilizer 20 ml L<sup>-1</sup> tended to give the highest yield of 13.23 cm and the treatment application of Trichoderma dose of 15 g plant<sup>-1</sup> and the concentration 0 ml L<sup>-1</sup> liquid organic fertilizer gives the lowest yield of 9.80 cm.



**Figure 1.** Average fruit length (cm) of chilli (*Capsicum annuum* L.) on different dose of Trichoderma and concentration of liquid organic fertilizer.

3.3. Weight per fruit

Analysis of variance on the observed weight per fruit parameter shows that the Trichoderma dose treatment significantly affected the weight of the fruit. Table 2 shows that weight per fruit of the chilli plant varied with the dose of Trichoderma applied. The average weight per fruit in the Trichoderma control application treatment (0 g plant<sup>-1</sup>) resulted in the heaviest fruit (25.77 g) followed by the application of 20 g per plant Trichoderma. The application of Trichoderma application dose of 10 g plant<sup>-1</sup> showed the lowest weight per fruit (22.01 g).

**Table 2.** Average weight per fruit (g) of chilli (*Capsicum annuum* L.) on different dose of Trichoderma and concentration of liquid organic fertilizer.

Dose of Trichoderma	Concentration of liquid organic fertilizer				Mean	Tukey's HSD <sub>0.05</sub>
	0 ml L <sup>-1</sup>	10 ml L <sup>-1</sup>	15 ml L <sup>-1</sup>	20 ml L <sup>-1</sup>		
0 g plant <sup>-1</sup>	23.07	30.87	24.63	24.50	25.77 a	3.07
10 g plant <sup>-1</sup>	21.07	21.80	22.17	23.00	22.01 b	
15 g plant <sup>-1</sup>	21.82	23.93	22.86	27.08	23.92 ab	
20 g plant <sup>-1</sup>	25.97	24.47	25.90	25.05	25.35 a	

Numbers followed by different letters (a, b, c) are significantly different at Tukey's test ( $\alpha=0.05$ ).

3.4. Fruit weight per plant

Data on fruit weight per large chili plant and its variance indicate that the treatment dose of Trichoderma application had a significant effect while the application of liquid organic fertilizer and its interaction with other treatment had no significant effect. The average fruit weight per plant are shown in table 3. The average fruit weight per plant applied with 20 g plant<sup>-1</sup> Trichoderma resulted in the highest yield of 223.55 g and was not significantly different from the Trichoderma dose of 15 g plant<sup>-1</sup>. Trichoderma application treatment dose of 10 g plant<sup>-1</sup> gave the lowest results ie 154.11 g.

**Table 3.** Average fruit weight per plant (g) of chilli (*Capsicum annuum* L.) on different dose of Trichoderma and concentration of liquid organic fertilizer.

Dose of Trichoderma	Concentration of liquid organic fertilizer				Mean	Tukey's HSD <sub>0.05</sub>
	0 ml L <sup>-1</sup>	10 ml L <sup>-1</sup>	15 ml L <sup>-1</sup>	20 ml L <sup>-1</sup>		
0 g plant <sup>-1</sup>	169.47	254.20	181.77	226.67	208.03 a	57.02
10 g plant <sup>-1</sup>	133.33	133.80	218.63	130.67	154.11 b	
15 g plant <sup>-1</sup>	211.08	203.48	144.47	205.75	191.20 a	
20 g plant <sup>-1</sup>	235.83	247.37	244.10	166.91	223.55 a	

Numbers followed by different letters (a, b, c) are significantly different at Tukey's test ( $\alpha=0.05$ ).

#### 4. Discussion

In the recent study, it is shown that the interaction between *Trichoderma* and liquid organic fertilizer significantly affected the parameters of the number of fruit per plant. The results indicate that the application of *Trichoderma* and liquid organic fertilizer package has a significant effect. The package of *Trichoderma* and liquid organic fertilizer provides the best effect on plant growth factors due to the availability of N nutrients used by the roots of chili plants. This is in line with the opinion of Harman and Uphoff [19] that soil microbes will gather near the roots of plants that produce root exudates and root cap pieces as a source of food for soil microbes. If the microbial population around the plant roots is dominated by beneficial microbes, the plants will get the most benefit from the presence of these microbes. Under normal crop conditions, nitrogen administration will result in a normal growth pattern. Other research shows results present that bacteria inoculated from roots and roots area of chili pepper are potential as Plant Growth Promoting Bacteria [20].

The availability of nutrients for plants is sufficiently available and the natural balance or environmental conditions are sufficient for *Trichoderma* sp to multiply. Microorganisms contained in organic fertilizers especially *Trichoderma* sp have the ability to compete with soil-borne pathogens mainly to get nitrogen and carbon [21]. *Trichoderma* sp response at the beginning of plant growth requires time to multiply in organic fertilizer, as well as acting as a decomposer of organic matter in providing nutrients for plants. On the other side, the microbes that interact with plants are divided into phyllosphere and epiphyte [22]. The more microorganisms present in organic fertilizer can help metabolism in the soil so that the soil is more able to provide nutrients needed by plants [23].

#### 20 Conclusions

Based on the results of research that has been done, it can be concluded as follows:

- a) Interaction of *Trichoderma* dose of 10 g plant<sup>-1</sup> and liquid organic fertilizer 15 ml L<sup>-1</sup> gives a higher number of fruits per plant.
- b) *Trichoderma* application with a dose of 20 g plant<sup>-1</sup> gives the heaviest weight per fruit and fruit weight per plant.
- c) There is no one concentration of liquid organic fertilizer which provides better growth and production of chili.

#### References

- [1] Wahyudi T 2011 *Panen Cabe Sepanjang Tahun* (Jakarta: Agromedia Pustaka)
- [2] Rukmana R 2008 *Cabai Hibrida Sistem Mulsa Plastik* (Yogyakarta: Kanisius)
- [3] Halimah N, Baktir A and Purkan P 2019 Exploration of Cellulolytic Microorganism as A Biocatalyst Candidate for Liquid Fertilizer Production *IOP Conf. Ser. Earth Environ. Sci.* **217**
- [4] Purbaningtias T E, Qayyumah N B and Kurniawati P 2018 Comparative analysis method of C-organic in fertilizers by gravimetry and Spectrophotometry *2nd Int. Conf. Chem. Chem. Process Eng.* **020055**
- [5] Pratama Y S 2008 *Pembuatan Pupuk Organik Dan Anorganik Cair Dari Limbah-Sayuran*
- [6] Hadisumitro, L M 2002 *Membuat Kompos*. (Jakarta: Penebar Swadaya)
- [7] Oey K N *Pengantar Mikrobiologi Umum*. (Bandung: Penerbit Angkasa)
- [8] Desiana C, Banuwa I S, Evizal R and Yusnaini S 2013 Pengaruh pupuk organik cair urin sapi dan limbah tahu terhadap pertumbuhan bibit kakao (*Theobroma cacao* L.) *J. Agrotek Trop.* **1**
- [9] Rachmat A and Agustina F 2009 Pembuatan Nata De Coco dengan Fortifikasi Limbah Cucian Beras Menggunakan *Acetobacter xylinum* *Univ. Diponegoro Semarang*
- [10] Ningsih E M N and Nugroho Y A 2010 Pertumbuhan stek nilam (*Pogostemon cablin*, Benth) pada Berbagai Komposisi Media Tumbuh dan Dosis Penyiraman Limbah Air Kelapa *AGRIKA* **4**
- [11] Charisma A M, Rahayu Y S and Isnawati Pengaruh Kombinasi Kompos *Trichoderma* dan Mikoriza Vesikular Arbuskular (MVA) terhadap Pertumbuhan Tanaman Kedelai (*Glycine max* (L.) Merrill) pada Media Tanam Tanah Kapur *LenteraBio* **1** 111–6

- [12] Larekeng S H, Gusmiaty, Restu M, Tunggal A and Susilowati A 2019 Isolation and identification of rhizospheric fungus under Mahoni (*Swietenia mahagoni*) stands and its ability to produce IAA (Indole Acetic Acid) hormones *IOP Conf. Ser. Earth Environ. Sci.* **343** 012051
- [13] Novandini A 2007 Eksudat akar sebagai Nutrisi *Trichoderma harzianum* DT38 serta aplikasinya terhadap pertumbuhan tanaman tomat *Progr. Stud. Biokimia, Fak. MIPA. IPB. Bogor*
- [14] Esrita B I 2011 Pertumbuhan dan hasil tomat pada berbagai bahan organik dan dosis *Trichoderma J. Penelit. Univ. Jambi Seri Sains* **13**
- [15] Hermosa R, Viterbo A, Chet I and Monte E 2012 Plant-beneficial effects of *Trichoderma* and of its genes *Microbiology* **158** 17–25
- [16] Chantapromma S, Jeerapong C, Phupong W, Quah C K and Fun H-K 2014 *Trichoderma* lactone: a diterpene lactone from *Trichoderma asperellum* *Acta Crystallogr. Sect. E Struct. Reports Online* **70** 0408–9
- [17] Guigón-López C, Guerrero-Prieto V, Lanzuise S and Lorito M 2014 Enzyme activity of extracellular protein induced in *Trichoderma asperellum* and *T. longibrachiatum* by substrates based on *Agaricus bisporus* and *Phymatotrichopsis omnivora* *Fungal Biol.* **118** 211–21
- [18] Naher L, Yusuf U K, Ismail A and Hossain K 2014 *Trichoderma* spp.: a biocontrol agent for sustainable management of plant diseases *Pak. J. Bot* **46** 1489–93
- [19] Harman G E and Uphoff N 2019 Symbiotic root-endophytic soil microbes improve crop productivity and provide environmental benefits *Scientifica (Cairo)*. **2019**
- [20] Masniawaty, Mustari K, Astuti, Gusmiaty, Larekeng H, Yani A and Rahim I 2019 Exploration of bacteria associated with chili peppers' rhizosphere and their capacity to absorb and produce gibberellin hormone *IOP Conf. Ser. Earth Environ. Sci.* **343** 012059
- [21] Djatmiko H A and Rohadi S S 1997 Efektivitas *Trichoderma harzianum* hasil perbanyakan dalam sekam padi dan bekatul terhadap patogenesitas *Plasmodiophora brassicae* pada tanah latosol dan andosol *Maj. Ilm. UNSOED* **2** 10–22
- [22] Akib M A, Mustari K, Kuswinanti T and Syaiful S A 2018 Abundance of arbuscular mycorrhizal fungi in rehabilitation area of nickel post-mining land of Sorowako, South Sulawesi *IOP Conf. Ser. Earth Environ. Sci.* **157**
- [23] Handayanto E 1998 Pengelolaan kesuburan tanah secara biologi untuk menuju sistem pertanian sustainabel *Habitat* **10** 1–7

# Effect of Trichoderma and tofu waste based organic fertilizer on the fruit development of chili (*Capsicum annuum* L.)

## ORIGINALITY REPORT

17%

SIMILARITY INDEX

12%

INTERNET SOURCES

10%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Lisa Septiani, RR Yudhy Harini Bertham, Hesti Pujiwati, Bilman Wilman Simanihuruk. "Growth and Yield of Soybean with Application of Liquid Organic Fertilizer and Arbuscular Mycorrhizal Fungi in Ultisols", *TERRA : Journal of Land Restoration*, 2021  
Publication 2%
- 2 [digilib.unhas.ac.id](http://digilib.unhas.ac.id)  
Internet Source 1%
- 3 [oasis.postech.ac.kr](http://oasis.postech.ac.kr)  
Internet Source 1%
- 4 [pr.r.hec.gov.pk](http://pr.r.hec.gov.pk)  
Internet Source 1%
- 5 [www.lib.unair.ac.id](http://www.lib.unair.ac.id)  
Internet Source 1%
- 6 Tri Esti Purbaningtias, Nursi Biwi Qayyumah, Puji Kurniawati, Bayu Wiyantoko, Alfa Akustia Widati. "Comparative analysis method of C-

# organic in fertilizers by gravimetry and spectrophotometry", AIP Publishing, 2018

Publication

---

7	Submitted to Universitas Hasanuddin Student Paper	1 %
8	Submitted to Universitas Sultan Ageng Tirtayasa Student Paper	1 %
9	pure.hud.ac.uk Internet Source	1 %
10	Submitted to Higher Education Commission Pakistan Student Paper	1 %
11	Submitted to Universitas Brawijaya Student Paper	1 %
12	aip.scitation.org Internet Source	1 %
13	edufersa.ufersa.edu.br Internet Source	1 %
14	protan.studentjournal.ub.ac.id Internet Source	1 %
15	ortus.rtu.lv Internet Source	1 %
16	digilib.uinsgd.ac.id Internet Source	<1 %

---

17 [ejurnal.unikarta.ac.id](http://ejurnal.unikarta.ac.id) <1 %  
Internet Source

---

18 G. R. Guimarães, Faber S. Pereira, Fabio S. Matos, Sueli C. M. Mello, Daniel D. C. Carvalho. "Supression of seed borne Cladosporium herbarum on common bean seed by Trichoderma harzianum and promotion of seedling development", Tropical Plant Pathology, 2014 <1 %  
Publication

---

19 [bioone.org](http://bioone.org) <1 %  
Internet Source

---

20 [e-journal.uajy.ac.id](http://e-journal.uajy.ac.id) <1 %  
Internet Source

---

21 Peri Hardiansyah, Uswatun Nurjanah, Widodo Widodo. "Growth Response and Yield of Pakcoy (Brassica rapa L.) on Various Concentrations Liquid Organic Fertilizer of Jiringa Hulls [Phithecellobium jiringa (Jack) Prain Ex King]", Akta Agrosia, 2019 <1 %  
Publication

---

22 Sigit Soeparjono. "The Effect of Media Composition and Organic Fertilizer Concentration on the Growth and Yield of Red Ginger Rhizome (Zingiber officinale Rosc.)", Agriculture and Agricultural Science Procedia, 2016 <1 %

---

23	<a href="http://journal.ugm.ac.id">journal.ugm.ac.id</a> Internet Source	<1 %
24	<a href="http://repositori.usu.ac.id">repositori.usu.ac.id</a> Internet Source	<1 %
25	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet Source	<1 %
26	Sri Lestari Galagi, Aiyen Aiyen, Muhammad Anshar Pasigai. "GROWTH AND YIELD OF ONION ( <i>Allium ascalonicum</i> L.) AGAINST VARIOUS CONCENTRATIONS OF LIQUID ORGANIC FERTILIZER", <i>AGROLAND: The Agricultural Sciences Journal</i> , 2018 Publication	<1 %
27	<a href="http://anzdoc.com">anzdoc.com</a> Internet Source	<1 %
28	<a href="http://garuda.ristekdikti.go.id">garuda.ristekdikti.go.id</a> Internet Source	<1 %
29	<a href="http://repository.usu.ac.id">repository.usu.ac.id</a> Internet Source	<1 %
30	<a href="http://www.bioflux.com.ro">www.bioflux.com.ro</a> Internet Source	<1 %
31	<a href="http://www.neliti.com">www.neliti.com</a> Internet Source	<1 %

---

32

Moshe Lapidot, Michael Friedmann, Oded Lachman, Avidan Yehezkel, Saadia Nahon, Shlomo Cohen, Meir Pilowsky. "Comparison of Resistance Level to Tomato Yellow Leaf Curl Virus Among Commercial Cultivars and Breeding Lines", Plant Disease, 1997

Publication

<1 %

33

"Microbial-mediated Induced Systemic Resistance in Plants", Springer Science and Business Media LLC, 2016

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On