

Immune Response (Cortisol, TNFA, HMGB1) in Trained and Untrained Adolescent after 12 Minutes Run Exercise

by Prof. Muh. Nasrum Massi, Ph.d

Submission date: 27-Nov-2021 05:15PM (UTC+0700)

Submission ID: 1713560744

File name: 777-1534-1-SM.pdf (706.69K)

Word count: 8130

Character count: 36929

Open Access Article

Immune Response (Cortisol, TNFA, HMGB1) in Trained and Untrained Adolescent after 12 Minutes Run Exercise

Ilhamjaya Pattelongi^{1*}, Huldani^{2*}, Muhammad Nasrum Massi³, Irfan Idris¹

¹ Department of Physiology, Faculty of Medicine, Hasanuddin University, Makassar, South Sulawesi, Indonesia

² Department of Physiology, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, South Kalimantan, Indonesia

³ Department of Medical Microbiology, Faculty of Medicine, University Hasanuddin, Makassar, South Sulawesi, Indonesia

Abstract: This study aims to explain the immune response (cortisol, TNF Alpha, HMGB1) after 12 minutes of moderate-intensity aerobic exercise. This exercise is a modification of Cooper's 12-minute running test that can be done by running 20 meters for 12 minutes according to an individual's ability with moderate intensity 60-80% maximum pulse rate. The research was conducted on 15 trained and 15 untrained students at SMAN 1 Banjarbaru. The sampling technique was the purposive sampling method. The value of VO₂ max was measured on the first day with Multistage Fitness Test. Cortisol, TNF Alfa, HMGB1 was measured by taking blood test. Data analysis used the Mann-Whitney test. The results showed that the average VO₂ max, cortisol, and HMGB1 scores of students with basketball achievement were higher than those of untrained students. TNF alpha level that did not increase was higher. The data analysis showed the difference in measurements between trained and untrained students ($p < 0.05$) on VO₂ max, Cortisol, and TNF Alpha. There was no significant difference in HMGB1 ($p > 0.05$). There is a positive correlation between Cortisol and TNF Alpha, Cortisol and HMGB1, HMGB1 and TNF Alpha in adolescents. It can be concluded that trained adolescents' Cortisol levels increased higher. Trained adolescents' TNF alpha levels were lower. It was not proven that trained students' HMGB1 levels differed from those of untrained students. In adolescents, there is an increase in cortisol levels followed by an increase in TNF Alpha levels, an increase in cortisol levels followed by an increase in HMGB1 levels, and an increase in HMGB1 levels followed by an increase in TNF Alpha levels. Moderate-intensity aerobic exercise (12 minutes of running) can be recommended as simple exercises to maintain fitness and boost the immune system or exercise health during a pandemic.

Keywords: immune response, 12 minutes of running, trained adolescents, untrained adolescents.

受过训练和未经训练的青少年跑步12分钟后的免疫反应 (皮质醇、肿瘤坏死因子 α 、HMGB1)

摘要:

本研究旨在解释中等强度有氧运动12分钟后的免疫反应 (皮质醇、肿瘤坏死因子 α 、HMGB1)。这个练习是对库珀12分钟跑步测试的修改, 可以根据个人的能力以中等强度60-80%的最大脉搏率跑20米12分钟来完成。该研究是在斯曼1班加巴鲁对15名受过培训和15名未受过培训的学生进行的。抽样技术是有目的的抽样方法。在第一天使用多阶段体能测试测量最大摄氧量的值。皮质醇、肿瘤坏死因子阿尔法、HMGB1是通过血液测试来测量的。数据分析使用曼惠特尼检验。结果表明, 有篮球成绩的学生的平均VO₂最大限度、皮质醇和HMGB1分数高于未

Received: June 12, 2021 / Revised: August 15, 2021 / Accepted: September 11, 2021 / Published: October 30, 2021

About the authors: Ilhamjaya Pattelongi, Department of Physiology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia; Huldani, Department of Physiology, Faculty of Medicine, Lambung Mangkurat University, Banjarmasin, Indonesia; Muhammad Nasrum Massi, Department of Medical Microbiology, Faculty of Medicine, University Hasanuddin, Makassar, Indonesia; Irfan Idris, Department of Physiology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

Corresponding authors Ilhamjaya Pattelongi, ilham_pt@yahoo.com; Huldani, huldani@gmail.com

受过训练的学生。没有增加的肿瘤坏死因子 α 水平更高。数据分析显示了受过训练和未受过训练的学生之间在最大摄氧量、皮质醇和肿瘤坏死因子 α 方面的测量差异(磷 <0.05)。HMGB1没有显著差异(磷 >0.05)。青少年皮质醇与肿瘤坏死因子 α 、皮质醇与HMGB1、HMGB1与肿瘤坏死因子 α 呈正相关。可以得出结论,受过训练的青少年的皮质醇水平升高得更高。受过训练的青少年的肿瘤坏死因子 α 水平较低。没有证明受过训练的学生的HMGB1水平与未受过训练的学生不同。在青少年中,皮质醇水平升高后肿瘤坏死因子 α 水平升高,皮质醇水平升高后HMGB1水平升高, HMGB1水平升高后肿瘤坏死因子 α 水平升高。中等强度的有氧运动(跑步12分钟)可以被推荐为简单的运动,以在大流行期间保持健康和增强免疫系统或锻炼健康。

关键词: 免疫反应, 12分钟跑步, 受过训练的青少年, 未受过训练的青少年。

1. Introduction

WHO states that in 2019, most people around the world are less physically active, thus endangering their health in the future, so WHO recommends carrying out physical exercise, especially moderate intensity, for 150 minutes a week for someone aged 18-64 years [1]. Regular exercise decreases the risk of developing various diseases and increases life expectancy [2, 3]. One form of physical exercise is a game of basketball. Although not as popular as football, basketball is popular among Indonesians [4]. Basketball uses alternating aerobic-anaerobic metabolism, although anaerobic metabolism is predominant.

Aerobic exercise is an activity that utilizes the continuous and normal contractions of large muscle groups for some time and utilizes aerobic metabolism as a source of energy. One of the components of aerobic exercise is intensity. Moderate-intensity aerobic exercise, an activity that consumes energy comparable to brisk walking, is the aerobic exercise most frequently studied [5]. Physical exercise is related to a person's ability to perform activities through its effect on VO₂ max. 12-minute running was used as a way to measure a person's VO₂ max value. In addition, exercise is a stressor that stimulates the neuroendocrine system and immune response. The correlation between the immune system and exercise was first studied in David Nieman's study, which showed a lack of reports of ARI complaints in individuals who routinely do moderate-intensity physical exercise. Physical exercise drastically affects the number of leukocytes circulating in the peripheral blood. Leukocytosis that accompanies acute exercise is a transient phenomenon because the number and composition of the leukocytes usually return to normal values within 6-24 hours after an exercise session. Leukocytosis that occurs in physical exercise is mostly caused by the activation of neutrophils and lymphocytes

and a small proportion of monocytes. Physical exercise activates the HPA response, which causes the hypothalamus to release corticotropin-releasing hormone (CRH) to cells in the anterior pituitary, which stimulates the release of adrenocorticotropic hormone (ACTH) and stimulates the adrenal cortex to release glucocorticoids (cortisol) into the bloodstream.

2. Methods

Cross-sectional research was conducted on the population of all adolescents at Senior High School 1 (SMAN 1) Banjarbaru. The sample in this study was divided into two; namely, the sample was taken from 15 basketball-trained students and 15 untrained basketball students. The sample was cooperative and had filled out a consent form to be the subject of the study. The purposive sampling method was used to take samples according to the inclusion criteria, namely (a) male, (b) aged 15-18 years, (c) physically healthy, meaning that at the time of the study, the proband was not sick or infected, had no history of heart or lung disease, and allergies, (d) cooperative, research subjects could be invited to work together to carry out research procedures, (e) not smoking, (f) not consuming drugs that affect the number of leukocytes, monocytes, and neutrophils at least two days before blood collection, (g) having a normal body mass index (BMI) (20-25). The sample of basketball-trained adolescents is students of SMAN 1 Banjarbaru, members of a basketball sports club, who routinely practice basketball at least three times a week for one hour per training session for a year. The study will be stopped on subjects who experience signs of fatigue while doing the exercises and cannot complete the exercise.

The research was conducted for three days. VO₂ max measurements were carried out on the first day using the

MFT (Multi stage fitness test). The subject was rested on the second day. Blood samples were taken for analysis of levels of cortisol, Tnf alpha, and HMGB1 were checked on the third day. After running twelve minutes of moderate-intensity aerobic exercise, a 5 cc blood sample was taken from the brachial vein. Before training, the research subjects will have their maximum pulse rate (MHR) calculated using the Tanaka formula. After knowing MHR, research subjects will use pulse oxymetry and warm-up to run in groups of 3 people per group until they reach the MHR target of 70-79%. After reaching the target of 70-79% MHR, the subject will continue to run for twelve minutes at the same rhythm. Blood sampling and subject analysis were carried out by trained personnel from the Prodia Banjarmasin laboratory. Trained health workers always accompany training sessions to prevent life-threatening things from occurring. The Mann-Whitney test was used for VO₂ max, cortisol, TNF Alpha, whereas the unpaired Student T-test for HMGB1.

3. Results

Sample characteristics included age, oxygen saturation, body mass index, pulse, height, weight, systolic and diastolic blood pressure. The average sample was 17 years. The mean value of pulse rate, systolic and diastolic blood pressure, body weight, body mass index of students trained in basketball was lower than students who were not trained in basketball. Pulse rate, systolic and diastolic blood pressure, body mass index, oxygen saturation were within normal limits. Table 1 shows that students trained in basketball were fitter than students who were not trained in basketball.

Table 1 Characteristics of research subjects

Characteristic (Mean ± SD)	Basket group (Trained Basketball Student n=15)	Non Basket Group (Untrained Basketball Student n=15)
Age (Years)	16.93±0.258	17.07±0.495
SO ₂ (%)	97.27±2.576	96.93± 3.731
Pulse per minute	84.20±11.384	92.93±12.981
BMI (kg / m ²)	21.65±2.104	21.68±5.911
Height (cm)	167.37±8.067	172.00±5.305
Weight (kg)	61.13±10.034	64.33±18.289
Systolic blood pressure (mmHg)	127.53±14.282	132.20±13.078
Diastolic blood pressure (mmHg)	78.73±7.001	82.87±9.797

Table 2 Research results

Marker	Trained Basketball Students (n = 15)		Untrained Basketball Students (n = 15)	
	Mean	SD	Mean	SD

VO ₂ max	37,49	26,1	28,85	18,8
Cortisol	1440,1	1863,1	26,3	16,1
TNF Alpha	532,0	667,2	963,7	1144,8
HMGB1	2180,6	166,4	2168,9	295,8

The mean VO₂ max, cortisol, and HMGB1 concentrations of basketball-trained students were higher than untrained. In contrast, the untrained TNF alpha score was higher than the untrained.

3.1. VO₂ Max

The mean VO₂ max of basketball-trained students was 37.49 ± 26.1 ml/kg/minute higher than the untrained students, namely 28.85 ± 18.8 ml/kg/minute. With the Mann-Whitney test, the VO₂ max value obtained was a statistically significant difference in the VO₂ max value after 12 minutes of moderate-intensity running training for basketball-trained students and untrained p = 0.000 (p < 0.05). Spearman's rho correlation test showed a significant correlation p = 0.000, with a very strong positive correlation (0.870 **) between VO₂ max and whether adolescents were trained in basketball. The more trained a teenager is, the greater the VO₂ max value.

3.2. Cortisol Concentration

The average cortisol concentration of basketball-trained students was greater than that of untrained students. In basketball-trained students, the mean blood cortisol concentration was 1,440.1 ng/ml higher than in the untrained, 26.3 ng/ml. In this study, statistically significant results were obtained using the Mann-Whitney test. After twelve minutes of running with moderate-intensity training in basketball-trained and untrained students, there was a significant difference in cortisol levels (p = 0.000 (p < 0.05). Spearman's rho correlation test showed a significant correlation p = 0.000, with a very strong positive correlation (0.720 **) between cortisol and whether adolescents were trained in basketball. The more trained a teenager is, the greater cortisol levels will be.

3.3. TNF Alpha Concentration

The mean TNF Alpha concentration of basketball-trained students was lower than that of untrained students. The mean TNF Alpha concentration for basketball-trained students was 532 pg/ml and untrained 963.7 pg/ml. Mann Whitney test showed a significant difference between the two after 12 minutes of moderate-intensity training for basketball-trained and untrained students p = 0.000 (p < 0.05). There was no significant relation between Spearman's rho test, p = 0.077, with a weak negative correlation (-0.327) between TNF Alpha and whether adolescents were trained or not in

basketball. The more trained a teenager will have lower TNF Alpha levels.

3.4. Concentration of HMGB1

The mean HMGB1 concentration of basketball-trained students 2180.6 ± 166.4 pg/ml ⁴⁷ higher than the untrained 2168.9 ± 295.7 pg/ml. There was no significant difference in HMGB1 levels after 12 minutes of moderate-intensity exercise in basketball-trained and untrained students $p = 0.895$ ($p > 0.05$). There is no significant correlation between Pearson's correlation test $p = 0.895$, with a very weak positive correlation (0.025) between HMGB1 and whether or not teenagers are in basketball. The more trained a respondent is, the greater the HMGB1 level will be.




Fig. 1 Boxplot graph of VO₂ max Value, Cortisol, TNF Alpha, and HMGB1 concentration

4. Discussion

The statistical tests in this study showed significant differences in VO₂ max, cortisol, and TNF alpha levels between basketball-trained and untrained ⁴⁹ students. While another aspect, namely HMGB1, there was no statistically significant difference between students trained in basketball and those who were not trained.

Physical activity causes changes in the body's physiological functions, both temporary changes (response) and permanent changes [6]. This also occurs in people who do sports; there is a physiological response to maintain the body's homeostasis. An increase in workload due to exercise will cause an increase in the workload of the organ systems, for example, in contracting skeletal muscles, an increase in glucose

uptake from the blood, an increase in body metabolism, one of which is the release of heat as compensation characterized by sweating, an increase in the amount of energy needed, and the cardiovascular there is an increase in heart rate due to pumping blood to supply more to all organs. The occurrence of physiological response in the musculoskeletal system as an adaptation to physical exercise occurs by involving many molecular pathways that play a role in regulating muscle contraction and biosynthetic pathways for ATP formation, such as AMP-activated protein kinase (AMPK), Sirtuins (SIRT6), mitogen-activated protein kinases (MAPKs), and oxygen sensor prolyl hydroxylases (PHDs) [7-8]. All these changes require a good balance so that the work of the organs and body condition is not disturbed; that is called homeostasis. Thus, if the homeostatic response fails, it will reduce their physical performance and performance in athletes.

The VO₂ max score of basketball-trained adolescents was higher than the untrained. This result is in line with the research of Buchan et al. In this study, it was found that physical exercise three times a week for seven weeks at moderate intensity increased VO₂ Max values compared to those who did not exercise in adolescents (p = 0.000). The study explains that this is likely due to moderate intensity physical exercise on cardiac output, where physical exercise causes an increase in cardiac output due to an increase in stroke volume.

A research conducted in 2011 described two factors that causes an increase in VO₂ max, namely central factors and peripheral factors. Centrally, the increase in VO₂ max was due to increased stroke volume and a slight increase in maximal heart rate. Peripherally, the increase in VO₂ max occurs due to increased arterial-venous oxygen differences that are influenced by oxygen transport to active muscle fibers, local enzyme adaptation, and mitochondrial density. Peripheral VO₂ max increase occurs in sprint interval training [9].

According to [10], in healthy subjects aged 60-71 years, 53 men and 57 women and a control group did not follow any exercise. The VO₂ max value was measured before and after aerobic exercise in the form of treadmill walking and running for 9-12 months. The results showed an improvement in the VO₂ value after a series of exercises in the male and female groups by 21 and 19%, respectively. It was provided that both elderly women and men can obtain VO₂ max improvement during routine exercise stimuli. Statistically, this study showed that the improvement in VO₂ max level does not depend on age, gender, VO₂ max value before undergoing exercise, or the intensity of the exercise undertaken.

Research was conducted on 34 physically healthy children aged 10-18 years who participated in strength training [11]. The results showed an increase in VO₂ max values after training. This study revealed no significant difference in VO₂ max levels based on age. In compliance with [12], no significant difference in VO₂ max values between athletes and non-athletes after exercising. Meanwhile, as stated in [13], greater VO₂ max improvement can be obtained with exercise for more than 20 weeks and intensity of around 60% and 70% of VO₂ max.

4.1. Cortisol

This study indicated a significant difference in cortisol levels after 12 minutes of physical exercise running between trained basketball and not (P = 0.000). The basketball-trained cortisol value (1440.1 ng/μL) was higher than the untrained (26.3 ng/μL); this may be a homeostatic process not due to a stressor. Moreover, the average value of HMGB1 is also bigger. Therefore, cortisol here functions as an anti-inflammatory and suppresses pro-inflammatory cell cytokines such as IL6, TNF Alpha, Leukocytes, Neutrophils, and Monocytes. In contrast, untrained cortisol values were lower; therefore, the anti-inflammatory function was not strong enough to suppress the inflammatory reaction, as indicated by a greater increase in the TNF alpha concentration. Sports and physical activity can be a source of stress for the body and impact other body systems and have the potential to cause homeostatic disorders.

Previous research has consistently shown that exercise with an intensity of more than 60% VO₂ max stimulates the release of higher cortisol concentrations in adults [14]. Studies show that every teenager has the same body response to an increase in the HPA axis reaction and the cortisol response after exercise. In adolescents aged 15 to 16 years who exercise for 12 minutes with an intensity of 70-85% of the maximum pulse, there will be a striking increase in cortisol levels compared to the group that only does moderate-intensity exercise (maximum 50-65% pulse rate).

Exercise with VO₂ max intensity of 60% (pre-intervention 12.3 ± 4.1 and post-intervention 20.1 ± 6.0) and 80% (pre-intervention 12.9 ± 6.3 and post-intervention) resulted in an increase in cortisol levels, significantly greater than the session with 40% exercise intensity (pre-intervention 12.2 ± 4.3 and post-intervention 10.8 ± 5.4). That is, cortisol levels in plasma increase in moderate and high-intensity exercise. Conversely, exercise with light intensity did not show a significant increase in cortisol levels but decreased.

Cortisol concentration will increase with increasing intensity and duration of exercise [15]. Different

exercises have different effects on the hormonal system; a greater response is shown in strength training. Cortisol levels will increase according to the level of stimulation given. High-intensity exercise will cause an increase in the activity of stress hormones such as cortisol, ACTH, and catecholamines, which inhibit protein synthesis and trigger protein degradation resulting in the breakdown of skeletal muscle protein.

The level of cortisol secretion is influenced by the circadian cycle. Serum cortisol secretion begins to increase at midnight, peaking in the morning. In addition, this increase may be determined by other factors that can increase cortisol secretion, namely psychological stress, which also triggers the release of cortisol [16].

After acute physical exercise, the increase in cortisol levels has been proven in several studies [17, 18]. The increase in cortisol levels is strongly influenced by the intensity of physical exercise and the subject's physical exercise habits. Hackney et al.'s research showed that cortisol levels increased with increasing training load given to subjects aged 16-21 years. In addition, the research [17] showed differences in the increase in cortisol levels in athletes and non-athletes according to the intensity of the physical exercise given. Increased cortisol levels provide an anti-inflammatory effect by emphasizing the expression of proinflammatory cytokines, thereby inhibiting the increase in the number of leukocytes [19].

The release of cortisol is one form of body adaptation to stress due to exercise; this can be seen from the cortisol levels in blood plasma, as found in the results of this study. Cortisol causes an increase in fat content in the blood due to an increase in amino acids and fat mobilization, which, if excessive, can disrupt tissue structure and function and even cell death.

Beserra et al. conducted a review of five articles and seven analyzes to look at post-exercise cortisol levels in 463 respondents with major depressive disorder. Respondents did the exercises with different frequencies, ways of measuring cortisol, and control groups. The results found that physical exercise caused a decrease in cortisol levels. However, this decrease in cortisol depends on the type, frequency, and measurement of cortisol taken. It is known that the decrease in cortisol levels can occur effectively after doing aerobic exercise with a frequency of 5 times per week [20].

On the other hand, the study by Wang et al. [21] showed the opposite result with the research above. The study aimed to assess the comparability of the effectiveness of aerobic exercise and resistance training in improving cognitive function in 42 young adults who were asked to do Stroop exercises after undergoing aerobic exercise, resistance training, and sedentary conditions a few days later. The saliva samples taken

were at baseline conditions before exercise, immediately after exercise, and 30 minutes afterward. The results showed that cortisol levels increased after exercise, compared with cortisol levels during the sedentary phase and baseline levels. Cortisol levels immediately after exercise were also higher 30 minutes afterward. From this, it is concluded that an increase in cortisol levels can occur after exercise but is influenced by the type and intensity of exercise.

4.2. TNF Alpha

The results of this study indicated a significant difference in the levels of TNF Alfa after 12 minutes of running physical exercise between trained basketball and non-trained (P = 0.000).

Downregulation of monocytic TNF production during acute exercise was mediated by high epinephrine levels [22].

According to the study [23], covering 50 men (mean age 58.9 ± 9.9 years) and women (mean age 50.9 ± 11.2 years), all observed cytokine concentrations after exercise (IL6, IL1 β , IL10, IL1 beta, and TNF alpha) increased from baseline values (P <0.001). Then the concentration decreased from day one to day two (P <0.01). Exercise induces an increase in cytokines, but these levels will decrease in the following days even though the exercise is still carried out with the same training intensity and load [23].

A study was conducted on 11 healthy males of middle school age, ranging from 14 to 18.5 years, after 1.5 hours of single wrestling exercise [24]. There was an increase in proinflammatory cytokines after exercise, namely IL6, TNF alpha and IL1 beta.

A study on six healthy male subjects who performed knee extensor exercises for 180 minutes [25] revealed that IL-6 and TNF-mRNA were both detected in the muscle sample at rest. IL-6 levels increased 100-fold during exercise, and there was no significant increase in arterial plasma TNF-alpha mRNA.

Direct muscle contraction stimulates the release of IL-6, an anti-inflammatory cytokine, decreasing the production of TNF alpha and IL1 beta in the acute phase and during cell proliferation. Moderate intensity exercise (MIT) effectively reduces body fat, which can prevent fat cell damage and cell hypoxia, so that proinflammatory cytokines (IL6 and TNF) are reduced through increased adiponectin secretion and increased anti-inflammatory cytokines [26].

Comparison of inflammatory response rates and muscle damage indexes between four popular elite-level team sports is reported in [27]. Thus, 72 elite male players from four sports: soccer, basketball, volleyball, and handball (18 in each), were asked to finish the match, while 18 non-athletes were tested as a control. Blood

samples were drawn before, immediately, 13 and 37 hours after the match. The results found changes in TNF α levels in all types of sports, including basketball, 4 blood samples taken immediately after, 13 hours after, and 37 hours after a competitive match, compared to baseline values (measured in the morning about 8 hours before the match). Soccer players showed a 3–4-fold increase immediately after the largest inflammatory cytokine match, namely TNF alpha tumor. Volleyball players showed the smallest increase in inflammation and markers of muscle damage compared to the other three sports.

4.3 HMGB1

The results of this study indicated no significant difference in HMGB1 levels after doing physical exercise in the form of running 12 minutes between trained basketball and not.

The human body has endogenous danger signals to prevent inflammatory responses secondary to releasing intracellular inflammatory factors into the extracellular area, namely the molecular pattern of associated damage (DAMP). High Molecular Weight 1 (HMGB1) is a protein from DAMP which is a sign of muscle cell damage and causes the mobilization of immune cells to the site of trauma [7]. HMGB1 levels will return to their original concentration after 30 minutes of rest after exercising [28].

Overall, DAMP triggers the release of massive cytokines, including TNF- α , IL-6, and IFN. These mediators enhance the activation, proliferation, maturation, and recruitment of immune cells at the site of trauma, causing indirect activation of adaptive and innate immune cells such as DC or T cells [29].

HMGB1 exposure to human monocyte cultures will stimulate proinflammatory cytokines such as IL6, IL-8, TNF, and inflammatory protein macrophages-1 (MIP-1). The kinetic responses to the LPS mediated release of TNF and HMGB1 were very different. HMGB1 stimulate biphasic TNF release with the second wave is delayed, whereas LPS mediated TNF release occurred in the initial monophasic mode [30].

A study was conducted for three weeks with a high-intensity training program in 7 healthy young men [31]. Aerobic and endurance training is done for three consecutive days each week at the same time. Saliva and blood samples were collected Pre and Post, 30 minutes after exercise each week and 24 hours after exercise at the end of the session at week 3. The results showed plasma HMGB1 increased from Pre- to Post-exercise ($P < 0.05$).

A similar study was performed on 40 male Wistar rats (weight 220 ± 10.2 g) which were randomly divided into

four groups: diabetic neuropathy, diabetic neuropathy + exercise, healthy + exercise, and healthy controls, diabetes was induced by injection of STZ (50 mg/kg) [32]. After confirming the development of diabetic neuropathy with a behavioral test, the exercise group received six weeks of continuous aerobic exercise at an average intensity of 15 m / min for 30 minutes on a treadmill. The serum HMGB1 level was measured by ELISA, and the concentration of malondialdehyde (MDA) and the action of superoxide dismutase (SOD) and catalase (CAT) enzymes in the spinal cord were determined by biochemical methods. Two-way ANOVA and Tukey's post hoc test were used for statistical analysis. The results showed that aerobic exercise significantly reduced HMGB1 protein levels and MDA concentrations and increased SOD and CAT enzyme activity compared to the diabetic neuropathy group ($p < 0.05$). Also, HMGB1 and MDA levels were increased, and SOD and CAT enzyme activity decreased in the diabetic neuropathy group ($p < 0.05$). Aerobic exercise is known to alter the protein HMGB1 and oxidative stress levels and increase the sensitivity of nociceptors to painful agents. It is recommended to use aerobic exercise as a non-prescriptive therapeutic intervention for diabetes patients to reduce neuropathic pain [32].

In another sports study, 34 and 36 nonprofessional female and male runners and half marathon finished each race; immediately, blood samples were collected 1-2 days before and 2-7 days after the race [33]. After the race HMGB1 serum concentration increased significantly by 1.5 times (half marathon; 3.13 ± 1.63 ng/mL to 4.78 ± 2.1 ng/mL) and 2.3 times (marathon; 2.58 ± 2.58 ng/mL to 6.02 ± 2.18 ng/mL), corresponding to an increase in circulation sRAGE for a half marathon but not a full marathon. During the recovery period, the concentration of HMGB1 and sRAGE increases back to the previous training level.

Not all HMGB1 respond positively to every exercise. In recent research, HMGB1 plasma remains below the detection limit (78 pg/mL) of testing after 1,200 km of plyometric training or cycling [34].

There was a decrease in serum HMGB1 at two weeks of Nordic walking exercise combined with vitamin D in elderly women; this suggests regular exercise can weaken the alarmin response in healthy adults [35].

5. Conclusion

This study showed that basketball-trained students had better VO₂ max scores than untrained basketball students. Twelve minutes of moderate aerobic exercise resulted in higher cortisol levels in basketball-trained students than in untrained students, while TNF alpha

levels were opposite. A very strong positive correlation between VO₂ max and whether or not young people are trained in basketball. The more trained a teenager is, the greater the VO₂ max value. There is a very strong positive correlation between cortisol and whether adolescents are trained in basketball. The more trained a teenager is, the greater cortisol levels will be. Twelve minutes of aerobic exercise can be used to boost the immune system and show that the homeostasis process in adolescents is working well.

The Covid-19 pandemic has forced everyone to move and stay at home. It is difficult to do some exercises outside, and not all types of sports can be done because they are vulnerable to transmits of Covid-19. One of the sports that can be done during the pandemic is a 12-minute run in the morning. The scientific novelty in this research is 12 minutes-run modified from Cooper's 12-minute running test; this can be done on a 20-meter running track for 12 minutes according to each ability, people can run or walk quickly according to their abilities with a target maximum pulse rate according to age (220- Age) with moderate intensity 60-80% of maximum pulse rate.

This research was conducted in July 2020 on adolescents at SMA 1 Banjarbaru, who did a 12-minute running test (modified of Cooper's 12-minute running test) with moderate intensity 60-80% of the maximum pulse rate; the results showed that in people who were trained in sports it did not cause system disorders. Immunity due to increased cortisol occurs not because of a response to inflammatory reactions but due to the influence of sympathetic nerves as a homeostatic response. Normal cortisol levels in untrained people indicate that loading with moderate-intensity aerobic exercise for 12 minutes of running does not cause an inflammatory reaction that causes cell damage. Moderate-intensity aerobic exercise for 12 minutes of running can be used as a weight-bearing exercise that improves fitness and immunity in both trained and untrained people.

The limitations of this study are:

1. This study did not measure Cortisol, TNF- α , and HMGB1 levels before and after exercise, so it could not assess the effect of exercise on Cortisol, TNF- α , and HMGB1 levels.
2. The number of samples in this study is limited.
3. No serial 0, 30, and 60-minute examinations were performed to see the recovery of the immune response.
4. Examination concerned only systemic inflammation markers (using serum and blood), so the local inflammatory response in muscle tissue (macrophages) cannot be evaluated.

References

- [1] BENNIE JA, DE COCKER K, TEYCHENNE MJ, et al. The epidemiology of aerobic physical activity and muscle-strengthening activity guideline adherence among 383,928 U.S. adults. *International Journal of Behavioral Nutrition and Physical Activity*, 2019, 16(1): 1-11.
- [2] SINGH R, PATTISAPU A, and EMERY MS. US Physical Activity Guidelines: Current state, impact and future directions. *Trends in Cardiovascular Medicine*, 2019, 30(7): 407-412.
- [3] MALLO FERRER F. Physical Activity and Exercise. In HUHTANIEMI I, and MARTINI L. (Eds) *Encyclopedia of Endocrine Diseases*, 2nd ed. Academic Press; 2018, 436-441.
- [4] PIEPOLI MF, HOES AW, AGEWALL S, et al. European Guidelines on cardiovascular disease prevention in clinical practice. *European Heart Journal*, 2016, 37(29): 2315-2381.
- [5] PATEL H, ALKHAWAM H, MADANIEH R, et al. Aerobic vs anaerobic exercise training effects on the cardiovascular system. *World Journal of Cardiology*, 2017; 9(2): 134.
- [6] MULYONO M. Pengaruh Manipulasi Sport Massage Terhadap Penurunan Denyut Nadi Setelah Latihan Olahraga. *Jurnal Sportif*, 2016, 2(2): 15-21.
- [7] FAN W and EVANS RM. Exercise mimetics: impact on health and performance. *Cell Metabolism*, 2017, 25(2): 242-247.
- [8] KJØBSTED R, HINGST JR, FENTZ J, et al. AMPK in skeletal muscle function and metabolism. *The FASEB Journal*, 2018, 32(4): 1741-1777.
- [9] LEE DC, PATE RR, LAVIE CJ, et al. Leisure-time running reduces all-cause and cardiovascular mortality risk. *Journal of the American College of Cardiology*, 2014, 64(5): 472-481.
- [10] KOHRT WN, MALLEY MT, COGGAN AR, et al. Effects of gender, age, and fitness level on response of $\dot{V}O_{2\max}$ to training in 60-71 yr olds. *Journal of Applied Physiology*, 1991, 70(5): 2004-2011.
- [11] CARAZO-VARGAS P. and MONCADA-JIMÉNEZ J. A meta-analysis on the effects of exercise training on the VO₂max in children and adolescents. *Retos*, 2015, 27: 184-187.
- [12] BAQUET, G., BERTHOIN, S., DUPONT, G., et al. Effects of high intensity intermittent training on peak VO₂ in prepubertal children. *International Journal of Sports Medicine*, 2002, 23(6): 439-444.
- [13] HUANG G, GIBSON CA, TRAN ZV, OSNESS WH. Controlled Endurance Exercise Training and VO₂max Changes in Older Adults: A Meta-Analysis. *Preventive Cardiology*, 2005, 8(4): 217-225.
- [14] BUDDE H, MACHADO S, RIBEIRO P, and WEGNER M. The cortisol response to exercise in young adults. *Frontiers in Behavioral Neuroscience*, 2015, 9, article ID 13. <https://doi.org/10.3389/fnbeh.2015.00013>
- [15] DUCLOS M, and TABARIN A. Exercise and the Hypothalamo-Pituitary-Adrenal Axis. *Frontiers of Hormone Research*, 2016, 47: 12-26.
- [16] HASLINDA DS, PATELLONGI I, and SINRANG AW. Pengaruh Olahraga Futsal Terhadap Kadar Kortisol Serum

Pada Individu Dewasa Muda. In Seminar Nasional LP2M UNM. 2017, 2(1).

- [17] SATO K, IEMITSU M, KATAYAMA K, et al. Responses of sex steroid hormones to different intensities of exercise in endurance athletes. *Experimental Physiology*, 2016, 101(1): 168-175.
- [18] MUSCELLA A, VETRUGNO C, SPEDICATO M, et al. The effects of training on hormonal concentrations in young soccer players. *Journal of Cellular Physiology*, 2019, 234(11): 20685-20693.
- [19] INCE LM, WEBER J, and SCHEIERMANN C. Control of leukocyte trafficking by stress-associated hormones. *Frontiers in Immunology*, 2019, 10(JAN): 1-9.
- [20] BESERRA AHN, KAMEDA P, DESLANDES AC, et al. Can physical exercise modulate cortisol level in subjects with depression? A systematic review and meta-analysis. *Trends in Psychiatry and Psychotherapy*, 2018, 40(4): 360-368.
- [21] WANG CC, ALDERMAN B, WU CH, et al. Effects of Acute Aerobic and Resistance Exercise on Cognitive Function and Salivary Cortisol Responses. *Journal of Sport & Exercise Psychology*, 2019; 41(2):73-81. <https://doi.org/10.1123/jsep.2018-0244>.
- [22] DIMITROV S, HULTENG E., & HONG S. Inflammation and exercise: Inhibition of monocytic intracellular TNF production by acute exercise via β_2 -adrenergic activation. *Brain, Behavior, and Immunity*, 2017, 61: 60-68. <https://doi.org/10.1016/j.bbi.2016.12.017>
- [23] TERINK R, BONGERS CCWG, WITKAMP RF, et al. Changes in cytokine levels after prolonged and repeated moderate intensity exercise in middle-aged men and women. *Translational Sports Medicine*, 2018, 1(3): 110-119. <https://doi.org/10.1002/tsm2.23>
- [24] NEMET D, OH Y, KIM HS, HILL M, and COOPER DM. Effect of intense exercise on inflammatory cytokines and growth mediators in adolescent boys. *Pediatrics*, 2002, 110(4): 681-689. <https://doi.org/10.1542/peds.110.4.681>.
- [25] STEENSBERG A, KELLER C, STARKIE R L, et al. IL-6 and TNF-alpha expression in, and release from, contracting human skeletal muscle. *American Journal of Physiology, Endocrinology and Metabolism*, 2002, 283(6): E1272-E1278. <https://doi.org/10.1152/ajpendo.00255.2002>
- [26] HULDANI, ACHMAD H, ARSYAD A, et al. Differences in VO2 max based on age, gender, hemoglobin levels, and leukocyte counts in Hajj prospective pilgrims in Hulu Sungai Tengah Regency, South Kalimantan. *Systematic Reviews in Pharmacy*, 2020, 11(4): 9-14.
- [27] SOUGLIS A, BOGDANIS GC, GIANNOPOULOU I, et al. Comparison of Inflammatory Responses and Muscle Damage Indices Following a Soccer, Basketball, Volleyball and Handball Game at an Elite Competitive Level. *Research in Sports Medicine*, 2015, 23(1): 59-72. <https://doi.org/10.1080/15438627.2014.975814>
- [28] HULDANI. Pengaruh Latihan Aerobik Ringan dan Sedang terhadap Kadar Interleukin 8 dan Jumlah Netrofil pada Remaja. *Berkala Kedokteran*. 2016, 12(1):61-7. <http://dx.doi.org/10.20527/jbk.v12i1.357>
- [29] VOURC'H M, ROQUILLY A, and ASEHNOUNE K. Trauma-Induced Damage-Associated Molecular Patterns-Mediated Remote Organ Injury and Immunosuppression in the Acutely Ill Patient. *Frontiers in Immunology*, 2018; 9: article ID 1330, <https://doi.org/10.3389/fimmu.2018.01330>
- [30] YANG H, WANG H, CHAVAN SS, and ANDERSSON U. High Mobility Group Box Protein 1 (HMGB1): The Prototypical Endogenous Danger Molecule. *Molecular Medicine*, 21 (SUPPLEMENT 1), 2015, S6-S12.
- [31] GOH J, HOFMANN P, AW NH, et al. Concurrent high-intensity aerobic and resistance exercise modulates systemic release of alarmins (HMGB1, S100A8/A9, HSP70) and inflammatory biomarkers in healthy young men: a pilot study. *Translational Medicine Communications*, 2020, 5(1), <https://doi.org/10.1186/s41231-020-00056-z>
- [32] KAKI A, NIKBAKHT M, HABIBI A, and FATHI MOGHADAM H. The effect of aerobic exercise on HMGB1 protein levels and some oxidative stress indices in rats with diabetic neuropathic pain. *Daneshvar Medicine: Basic and Clinical Research Journal*, 2020; 27(3): 27-38, <https://doi.org/10.22070/dmed.27.142.27>
- [33] BEKOS CM, ZIMMERMANN L, UNGER S, et al., Non-professional marathon running: Rage axis and st2 family changes in relation to open-window effect, inflammation and renal function, *Scientific Reports*, 2016, 6: 32315.
- [34] BEHRINGER M, KILIAN Y, MONTAG J, et al. Plasma concentration of high-mobility group box 1 (HMGB1) after 100 drop to vertical jumps and after a 1200-km bicycle race. *Research in Sports Medicine*, 2016, 24(2): 119-129, <https://doi.org/10.1080/15438627.2015.1126275>
- [58] GMIAT A, MIESZKOWSKI J, PRUSIK K, et al., **Changes in pro-inflammatory markers and leucine concentrations in response to nordic walking training combined with vitamin d supplementation in elderly women.** *Biogerontology*, 2017, 18: 535-548.

参考文献:

- [1] BENNIE JA, DE COCKER K, TEYCHENNE MJ 等。383,928名美国成年人有氧运动和肌肉强化活动指南遵守情况的流行病学。国际行为营养与身体活动杂志·2019, 16(1): 1-11。
- [2] SINGH R, PATTISAPU A 和 EMERY MS。美国体育活动指南：现状、影响和未来方向。心血管医学趋势, 2019, 30(7): 407-412。
- [3] MALLO FERRER F。体育活动和锻炼。在 HUHTANIEMI I 和 MARTINI L。(编辑)内分泌疾病百科全书·第2版。学术出版社, 2018, 436-441。
- [4] PIEPOLI MF, HOES AW, AGEWALL S 等。欧洲临床实践中心血管疾病预防指南。欧洲心脏杂志·2016, 37(29): 2315-2381。

- [5] PATEL H、ALKHAWAM H、马达尼埃 R 等人。有氧与无氧运动训练对心血管系统的影响。世界心脏病学杂志·2017,9 (2) : 134。
- [6] Mulyono M. 运动按摩手法对运动后脉搏率降低的影响。运动杂志, 2016, 2(2) : 15-21。
- [7] Fan W 和 Evans RM. 运动模拟物：对健康和表现的影响。细胞代谢, 2017, 25(2): 242-247。
- [8] KJØBSTED R、HINGST JR、FENTZ J 等。AMPK 在骨骼肌功能和代谢中的作用。FASEB 期刊, 2018, 32(4): 1741-1777。
- [9] Lee DC、Pate RR、Lavie CJ 等。闲暇时间跑步可降低全因和心血管死亡风险。美国心脏病学杂志·2014, 64 (5) : 472-481。
- [10] Kohrt WN、Malley MT、Coggan AR 等。性别、年龄和健康水平对60-71岁人群伊欧、最大限度对训练反应的影响。应用生理学杂志, 1991, 70(5): 2004-2011。
- [11] Carazo-Vargas P. 和 Moncada-Jiménez J. 一项关于运动训练对儿童和青少年最大摄氧量影响的荟萃分析。雷托斯·2015, 27 : 184-187。
- [12] Baquet, G., Berthoin, S., Dupont, G., 等。高强度间歇训练对青春期前儿童VO₂峰值的影响。国际运动医学杂志·2002, 23 (6) : 439-444。
- [13] Huang G, Gibson CA, Tran ZV, Osness WH. 老年人的受控耐力运动训练和VO₂最大限度变化：荟萃分析。预防心脏病学·2005, 8(4) : 217-225。
- [14] Budde H, Machado S, Ribeiro P 和 Wegner M. 年轻人对运动的皮质醇反应。行为神经科学前沿·2015, 9, 文章 ID 13. <https://doi.org/10.3389/fnbeh.2015.00013>
- [15] Duclos M 和 Tabarin A. 运动和下丘脑-垂体-肾上腺轴。激素研究前沿·2016, 47 : 12-26。
- [16] Haslinda DS、Patelongi I 和 Sinrang AW. 五人制足球运动对青年人血清皮质醇水平的影响。在LP2 M UNM 国家研讨会上。2017, 2(1)。
- [17] SATO K、Iemitsu M、Katayama K 等。性类固醇激素对耐力运动员不同强度运动的反应。实验生理学, 2016, 101(1): 168-175。
- [18] Muscella A、Vetrugno C、Sedicato M 等。训练对年轻足球运动员荷尔蒙浓度的影响。细胞生理学杂志, 2019, 234(11): 20685-20693。
- [19] Ince LM、Weber J 和 Scheiermann C. 压力相关激素对白细胞运输的控制。免疫学前沿·2019, 10 (1月) : 1-9。
- [20] Beserra Ahn、Kameda P、Deslandes AC 等。体育锻炼可以调节抑郁症患者的皮质醇水平吗？系统评价和荟萃分析。精神病学和心理治疗趋势·2018, 40(4) : 360-368。
- [21] Wang CC, Alderman B, Wu CH, 等。急性有氧运动和阻力运动对认知功能和唾液皮质醇反应的影响。运动与运动心理学杂志·2019 ; 41 (2) : 73-81. <https://doi.org/10.1123/jsep.2018-0244>。
- [22] Dimitrov S, Hulten E., 和 Hong S. 炎症和运动：急性运动通过β₂-肾上腺素能激活抑制单核细胞细胞内肿瘤坏死因子的产生。大脑、行为和免疫·2017, 61 : 60-68. <https://doi.org/10.1016/j.bbi.2016.12.017>
- [23] Terink R, Bongers CCWG, Witkamp RF 等。中年男性和女性长期和重复中等强度运动后细胞因子水平的变化。转化运动医学·2018, 1(3) : 110-119, <https://doi.org/10.1002/tsm2.23>
- [24] Nemet D, Oh Y, Kim HS, 等。剧烈运动对青春期男孩炎症细胞因子和生长介质的影响。儿科学·2002, 110(4) : 681-689. <https://doi.org/10.1542/peds.110.4.681>。
- [25] Steensberg A, Keller C, Starkie R L 等。伊利诺伊州-6和肿瘤坏死因子-α在收缩的人类骨骼肌中的表达和释放。美国生理学、内分泌学和代谢杂志·2002, 283(6) : E1272-E1278. <https://doi.org/10.1152/ajpendo.00255.2002>
- [26] Huldani, Achmad H, Arsyad A 等。南加里曼丹 葫芦双溪 叠加摄政朝觐预期朝觐者的VO₂最大限度差异基于年龄、性别、血红蛋白水平和白细胞计数。药学系统评价·2020, 11(4) : 9-14。
- [27] Souglis A, Bogdanis GC, Giannopoulos I 等。在精英竞争级别的足球、篮球、排球和手球比赛后炎症反应和肌肉损伤指数的比较。运动医学研究, 2015, 23(1): 59-72. <https://doi.org/10.1080/15438627.2014.975814>
- [28] Huldani. 轻度和中度有氧运动对青少年白细胞介素8水平和中性粒细胞数量的影响。医学期刊. 2016, 12(1): 61-67. <http://dx.doi.org/10.20527/jbk.v12i1.357>
- [29] Vourc'h M, Roquilly A 和 Asehounne K. 急性病患者的创伤诱导损伤相关分子模式介导的远程器官损伤和免疫抑制。免疫学前沿·2018 ; 9 : 文章ID 1330, <https://doi.org/10.3389/fimmu.2018.01330>
- [30] Yang H, Wang H, Chavan SS 和 Andersson U.

高迁移率群蛋白1(HMGB1)：原型内源性危险分子。分子医学·21(补充1)，2015，S6-S12。

[31] GOH J、HOFMANN P、AW NH 等。同时进行高强度有氧运动和抗阻运动可调节健康年轻男性警报素(HMGB1、秒100一种8/一种9、热电偶70)和炎症生物标志物的全身释放：一项初步研究。转化医学通讯·2020，5(1)，<https://doi.org/10.1186/s41231-020-00056-z>

[32] KAKI A、NIKBAKHT M、HABIBI A 和 FATHI MOGHADAM H。有氧运动对糖尿病神经性疼痛大鼠HMGB1蛋白水平和一些氧化应激指数的影响。达内什瓦尔药物：基础和临床研究杂志·2020；27(3)：27-38，<https://doi.org/10.22070/dmed.27.142.27>

[33] BEKOS CM、ZIMMERMANN L、UNGER S 等。非专业马拉松赛跑：与开窗效应、炎症和肾功能相关的愤怒轴和英石2家族变化。科学报告·2016，6：32315。

[34] BEHRINGER M、KILIAN Y、MONTAG J 等。100次垂直跳跃和1200公里自行车比赛后高机动性组箱1(HMGB1)的血浆浓度。运动医学研究，2016，24(2)：119-129，<https://doi.org/10.1080/15438627.2015.1126275>

[35] GMIAT A、MIESZKOWSKI J、PRUSIK K 等。老年妇女对北欧步行训练结合维生素d补充剂的促炎症标志物和亮氨酸浓度的变化。生物老年学·2017，18：535-548。

Immune Response (Cortisol, TNFA, HMGB1) in Trained and Untrained Adolescent after 12 Minutes Run Exercise

ORIGINALITY REPORT

22%

SIMILARITY INDEX

19%

INTERNET SOURCES

14%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	www.sysrevpharm.org Internet Source	7%
2	Jorming Goh, Michael Behringer. "Exercise alarms the immune system: A HMGB1 perspective", Cytokine, 2018 Publication	1%
3	johuns.net Internet Source	1%
4	www.tandfonline.com Internet Source	1%
5	pdf.aminer.org Internet Source	1%
6	sciencepublishinggroup.com Internet Source	1%
7	www.science.gov Internet Source	1%
8	ir.nknu.edu.tw Internet Source	1%

9

e-journal.unair.ac.id

Internet Source

<1 %

10

transmedcomms.biomedcentral.com

Internet Source

<1 %

11

E Supriyono, D Lesmana, K Nirmala, K Adiyana, N A R Setyawidati. "Evaluation of physiological responses and production performance of spiny lobster with different density treatments in floating net cage", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

12

Robert Stawski, Konrad Walczak, Ewelina Perdas, Anna Prymont-Przymińska et al. "Increased Circulating H3 Histone in Response to Repeated Bouts of Exercise Does Not Associate with Parallel Alterations of Cell-Free DNA", Biology, 2021

Publication

<1 %

13

dict.cnki.net

Internet Source

<1 %

14

Jiby Jolly Benjamin, Maheshkumar Kuppusamy, Teena Koshy, Maruthy Kalburgi Narayana, Padmavathi Ramaswamy. "Cortisol and polycystic ovarian syndrome – a systematic search and meta-analysis of case-

<1 %

control studies", Gynecological Endocrinology,
2021

Publication

15

www.ncbi.nlm.nih.gov

Internet Source

<1 %

16

"Basketball Sports Medicine and Science",
Springer Science and Business Media LLC,
2020

Publication

<1 %

17

www.mdpi.com

Internet Source

<1 %

18

"Endocrinology of Physical Activity and Sport",
Springer Science and Business Media LLC,
2020

Publication

<1 %

19

Andac Uzdogan, Meryem Kuru Pekcan, Aylin
Pelin Cil, Ucler Kisa, Filiz Akbiyik. "Progranulin
and tumor necrosis factor-alpha in lean
polycystic ovary syndrome patients",
Gynecological Endocrinology, 2021

Publication

<1 %

20

Stoyan Dimitrov, Elaine Hulteng, Suzi Hong.
"Inflammation and exercise: Inhibition of
monocytic intracellular TNF production by
acute exercise via β 2 -adrenergic activation",
Brain, Behavior, and Immunity, 2017

Publication

<1 %

21	Submitted to University of Central Lancashire Student Paper	<1 %
22	hk.sports.yahoo.com Internet Source	<1 %
23	jurnal.umk.ac.id Internet Source	<1 %
24	www.jove.com Internet Source	<1 %
25	dwherold.de Internet Source	<1 %
26	smallque.com Internet Source	<1 %
27	www.jonuns.com Internet Source	<1 %
28	Ines Ben Ayed, Naomie Castor-Guyonvarch, Souad Amimour, Salma Naija et al. "Acute Exercise and Cognitive Function in Alzheimer's Disease", Journal of Alzheimer's Disease, 2021 Publication	<1 %
29	Jun-Zhao Ye, Bi-Hui Zhong. "Landmarks in research of nonalcoholic fatty liver disease", World Chinese Journal of Digestology, 2017 Publication	<1 %
30	Parsa Gholipour, Alireza Komaki, Mahdi Ramezani. "Effects of the Combination of	<1 %

High-intensity Interval Training and Ecdysterone on Learning and Memory Abilities, Antioxidant Enzymes Activities, and Neuronal Population in an Amyloid-beta-induced Rat Model of Alzheimer's Disease", Research Square Platform LLC, 2021

Publication

31

caod.oriprobe.com

Internet Source

<1 %

32

www.f0cpb3.in

Internet Source

<1 %

33

www.x-mol.com

Internet Source

<1 %

34

Submitted to University of Leeds

Student Paper

<1 %

35

omade.com.tw

Internet Source

<1 %

36

repository.lib.cuhk.edu.hk

Internet Source

<1 %

37

"Wednesday, 7 September 2005", European Heart Journal, 09/02/2005

Publication

<1 %

38

Ali. "A Comparison of Small-Side Games and Interval Training on Same Selected Physical Fitness Factors in Amateur Soccer Players", Journal of Social Sciences, 2011

<1 %

39

G. D. Marsh, D. H. Paterson, D. Govindasamy, D. A. Cunningham. "Anaerobic Power of the Arms and Legs of Young and Older Men", *Experimental Physiology*, 1999

Publication

<1 %

40

Marcin Studencki, Angelina Ignatjeva, Magdalena Nitychoruk, Artur Gołaś, Wojciech Smółka, Adam Maszczyk. "Effect of bench press at a specified movement tempo on post-exercise testosterone and cortisol levels", *Physical Activity Review*, 2021

Publication

<1 %

41

Molecular Medicine, 2015.

Publication

<1 %

42

Shi-Jian Yi, Yang Wu, Lan-Lan Li, Qian-Kun Liang, Yue Xiao. "Compound amino acid combined with high-dose vitamin B6 attenuate traumatic coagulopathy via inhibiting inflammation by HMGB1/TLR4/NF- κ B pathway", *Journal of Inflammation*, 2020

Publication

<1 %

43

Submitted to University of Edinburgh

Student Paper

<1 %

44

big5.39kf.com

Internet Source

<1 %

45	Internet Source	<1 %
46	hdl.handle.net Internet Source	<1 %
47	ijdo.ssu.ac.ir Internet Source	<1 %
48	journals.sagepub.com Internet Source	<1 %
49	medsci.org Internet Source	<1 %
50	onlinelibrary.wiley.com Internet Source	<1 %
51	vb.bioscientifica.com Internet Source	<1 %
52	www.en8848.com.cn Internet Source	<1 %
53	www.nature.com Internet Source	<1 %
54	www.pingxiti.com Internet Source	<1 %
55	www.researchsquare.com Internet Source	<1 %
56	E Suhartono, I Thalib, I Aflanie, Z Noor, R Idroes. "Study of Interaction between	<1 %

Cadmium and Bovine Serum Albumin with UV-Vis Spectroscopy Approach", IOP Conference Series: Materials Science and Engineering, 2018

Publication

57

Zong-Shuai Li, Lei-Wang, Hai-Rong Zhang. "Hepatic oval cells and liver diseases", World Chinese Journal of Digestology, 2015

Publication

<1 %

58

link.springer.com

Internet Source

<1 %

59

"Abstracts 2014", Brain Injury, 2014

Publication

<1 %

60

"Endocrinology of Physical Activity and Sport", Springer Nature, 2013

Publication

<1 %

61

Kesava Kovanur Sampath, Ramakrishnan Mani, Rajesh Katare, Joshua Neale, James Cotter, Steve Tumilty. "Thoracic Spinal Manipulation Effect on Neuroendocrine Response in People With Achilles Tendinopathy: A Randomized Crossover Trial", Journal of Manipulative and Physiological Therapeutics, 2021

Publication

<1 %

62

R.M. Edinburgh, H.E Bradley, N-F. Abdullah, S.L. Robinson et al. "Lipid metabolism links

<1 %

nutrient-exercise timing to insulin sensitivity
in men classified as overweight or obese",
Cold Spring Harbor Laboratory, 2019

Publication

63

Sarah K. Wise, Sandra Y. Lin, Elina Toskala,
Richard R. Orlandi et al. "国际过敏与鼻科学共
识声明：变应性鼻炎", International Forum of
Allergy & Rhinology, 2019

Publication

<1 %

64

cofacts.gov.tw

Internet Source

<1 %

65

opendata.epa.gov.tw

Internet Source

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On