

The_Effect_of_the_Extracts_of_Betel_Piper_betel_Leaves_on. pdf

by Andi Makbul Aman

FILE	THE_EFFECT_OF_THE_EXTRACTS_OF_BETEL_PIPER_BETEL_LEAVE S_ON.PDF (282.36K)		
TIME SUBMITTED	14-JAN-2020 11:21AM (UTC+0700)	WORD COUNT	4816
SUBMISSION ID	1241722768	CHARACTER COUNT	25898



28

The Effect of the Extracts of Betel (*Piper betel*) Leaves on the Content of IL-1 β as the Inflammatory Cytokine in Experimental Mice Using an Acute Wound Healing Modeling

Maryunis Maryunis^{a*}, Syakib Bakri^b, Ilhamjaya Patellongi^c, Makbul Aman^d, Takdir Tahir^e, Azniah Syam^f

^aDr. Wahidin Sudirohusodo General Hospital, Makassar, Departement of Nursing, Universitas Muslim Indonesia

^bDepartement of Internal Disease, Medical Faculty, Hasanuddin University, Indonesia

^cDepartement of Physiology, Medical Faculty, Hasanuddin University, Indonesia

^dDepartement of Nursing, Medical Faculty, Hasanuddin University, Indonesia

^eDepartement of Nursing, Nani Hasanuddin Health Science Academy, Indonesia

Abstract

32

This study aims to determine the effect of ethyl acetate derived from the leaves extraction of betel (*Piper betel*) on the content of Interleukin-1 β (IL-1 β) in experimental mice using an acute wound healing modeling. The experimental design used randomized post test only group with 72 experimental mice which divided into 4 groups and each group consisted of 6 mice with three replications. The negative control group (I) was given vaseline after incision. The positive control group (II) was given bioplasenton. The third group (III) was topically given the extracts of betel leaves at 2% concentration and the fourth group (IV) was given the extracts of betel leaves at 3% concentration after incision respectively. The determination of IL-1 β applied Immunoassay Quantikine Elisa (IQE).

* Corresponding author.

The quantitative analysis (statistical test) used complete randomized design using one-way anova with Spearman's test and expressed as means \pm standard deviation, whereas, the qualitative analysis of the data was presented with the figure of time period of the concentration of IL-1 β for the negative control and control groups as well as both the extracts of betel leaves at 3rd day, 7th day and 14th day respectively. Results revealed that the content of IL-1 β was not statistically correlated with both the negative and positive control groups for both the treatment of vaseline ($\rho < 0,217$, $r = 0,218$) and bioplasenton ($\rho < 0,249$, $r = 0,190$) respectively. In addition, the content of IL-1 β was also not statistically correlated with the treatment of the extracts of betel leaves at 2% concentration ($\rho < 0,060$, $r = 0,420$). In contrast, the content of IL-1 β was statistically correlated with the treatment group of the extracts of betel leaves at 3% concentration ($\rho < 0,002$, $r = 0,701$). Results of the study concluded that the treatment of extracts of betel leaves at 3% concentration affected wound healing process in experimental mice and increased in line with the ascending time in wound healing process, whereas, both the negative and positive control groups as well as the treatment group of the extracts of betel leaves at 2% concentration did not affect the wound healing process among experimental mice.

Keywords: Piper betel; interleukin-1; wound healing.

1. Introduction

The study of the effect of betel (*Piper betel*) in wound healing process is feasible to know among nurse practitioners. It is considered important for the reason that the responsibility boundaries of nurse practitioners are changing, not only because of increased demands, but also they should demonstrate and develop their competence in varied extended and expanded practice roles [1].

Betel plant (*Piper betel*) is popularly regarded as a medicinal plant in Southeast Asia. Its leaves benefit for wound healing process and their extracts contain bioactive molecules like polyphenols, alkaloids, steroids, saponins and tannins [2]. The chief constituents of the leaves are volatile oils that contain phenol, betel-phenol, chavibetol and chavicol, cadinene, and hydroxychavicol which possess anti-oxidant and anti-carcinogenic activities [3,4,5,6,7]. The extracts of betel leaves also exhibits biological capabilities of detoxication, antioxidation, and antimutation against various ailments including liver fibrosis and carcinoma [8]. In addition, it is used for antidiabetic, antiulcer, antiplatelet aggregation, anti fertility, cardiotoxic, antitumour, antimutagenic, respiratory depressant and antihelminthic [9-15], antimicrobial [16], gastroprotective [17], hepatoprotective [18] and antimotility effects on washed human spermatozoa [19].

The health benefits of betel leaves in wound healing process can be identified by investigating the mechanism of wound healing process in the skin of the living species. Wound healing is a complex process involving several overlapping stages that include inflammation, formation of granulation tissue, reepithelialization, matrix formation and remodeling. Upon injury to the skin, the epidermal barrier is disrupted and keratinocytes release prestored interleukin-1 (IL-1). IL-1 is the first signal that alerts surrounding cells to barrier damage [20-30].

IL-1 β is a pro-inflammatory cytokine. It is expressed by many cells including macrophage, NK cells, monocytes, and neutrophils. It belongs to the IL-1 family that includes the IL-1a and IL1-RN genes. The

caspase 1 (CASPI/ICE) gene proteolitically activates IL-1 β . This gene is involved in the proliferation, differentiation and apoptosis of cells that associates with wound healing [31].

Early reports of the effect of the extracts of betel leaves as the inflammatory cytokine in experimental mice prompted the authors to evaluate its effect as the inflammatory cytokine in experimental mice. Therefore, the aim of this study is to analyze the effect of the extracts of betel leaves on IL-1 β as the cytokine inflammation in experimental mice.

2. Materials and Methods

2.1 Plant materials

Piper betel leaves which popularly known as "sirih" in Indonesia was taken from a 300-1000 m elevation above sea level in Soppeng Regency of South Sulawesi Province in the view of the fact that such an elevation was favorable environmental condition for the growth of *Piper betel*.

2.2 Experimental animals

The healthy adult Wistar mice weighing approximately 250-300 grams with the 4-to-6 week interval of age were used. Wistar strain is one the mice strains originated from the American continent. The experimental mice were obtained from the Veterinary Laboratory A of Gadjah Mada University, and then they were transported to the veterinary laboratory of Medicine Faculty of Hasanuddin University.

2.3 Experimental design

The experimental Wistar mice were divided into 4 groups and they were kept in their cages (30 cm x 50 cm x 15 cm) for at least five days prior to commencement of the experiment to allow for acclimatization to the laboratory conditions [32] and they were handled daily during this period.

During the experiment period, the experimental mice were put in the cages at 22 \pm 2 $^{\circ}$ C with 50%-60% air moisture and fed on a standard pellet diet (300 gr/mouse/day) and water *ad libitum* made at the veterinary laboratory of Medicine Faculty, Hasanuddin University. The cages were labeled and cleaned routinely for identification purposes and the light cycle was maintained in a 12 hour light/12 hour dark cycle in their respective groups for 15 days. There were 6 mice per group of treatment and the total experimental animals were 72 mice and they were grouped into 4 random groups with three replications which one mouse was used as demanding supply [33]. The four groups of experimental mice were divided as follows: 1) the negative control group was treated with vaseline (K-), 2) the positive control group was treated with bioplasenton (K+), 3) the treatment group given by the extracts of betel leaves at the 2% concentration (P1), and 4) the treatment group given by the extracts of *Piper betel* leaves at the 3% concentration (P2). Results of the extracts of *Piper betel* leaves in the form of acetyl acetate were given at 3rd day, 7th day and 14th day after excision by examining the concentration of IL-1 β respectively.

2.4 Wound creation

The hair on the back of each of the rats was removed with electrical clippers and bare areas were disinfected using a solution of 0.5% chlorhexidine in 70% alcohol, which was allowed to dry after application. The rats were anesthetized with isoflurane (0.01 µg/kg-0.05 µg/kg) and the fixation was conducted at pronation position. Evenly spaced wounds were made in the skin within the bare area of the back of mice using a punch to obtain biopsies with a diameter of 6 mm [34]. The hair on the back of each of the mice was removed with clippers and bare areas were cleaned using hair removal cream (Veet), which then disinfected using 0.5% chlorhexidine in 70% alcohol, which allowed to dry after application. Excision was made at the back area of experimental mice (0,8 x 0,8 cm) using a punch to obtain biopsies [35].

2.5 Extraction of Piper betel leaves

Betel leaves were chosen and chopped into small sizes, which then left to dry at room temperature in order to preserve the compounds in betel leaves for 7 days. Subsequently, 180 grams of dried betel leaves were put into 3 Erlenmeyer flasks; each flask contained 60 grams of dried betel leaves. After then, dried betel leaves were soaked into 96% ethanol where the volume of each ethanol solution was 600 ml, and the extracts were macerated for 48 hours. Later, the extracts in Erlenmeyer flask were filtered using muslin cloth and then evaporated until dry.

The preparation procedure of topical cream was conducted by suspended its cream base. The suspension of cream base was performed for all kinds of cream bases, viz. absorption base, hydrocarbon and percolating water in hot mortars at ± 60°C while stirring them until achieved homogeneous extracts and left to dry. The ethanol extracts of *Piper betel* leaves were made through the re-suspension of the prepared cream bases and then adding the extracts a little bit until achieved homogeneous extracts and formed cream.

2.6 Preparation of tissues for determination of the content of IL-1β

The extracts of betel were given orally for 15 days; the mice were fasted overnight and sacrificed by cervical decapitation. Afterward, the hairs on the back of each of the rats were removed with electrical clippers and bare areas were disinfected using a solution of 0.5% chlorhexidine in 70% alcohol, which were allowed to dry after application. The excision of biopsy was made approximately 1.5 x 2.5 cm². Samples for the examination of tissues of wound were analyzed using ELISA. The determination of the content of IL-1β used Rat IL-1β Immunoassay Quantikine Elisa.

2.7 Statistical analysis

All values were expressed as the mean±standard error. The quantitative analysis was performed using SPSS statistical package for WINDOWS (SPSS version 17). Results related to the content of IL-1β were statistically analyzed using Spearman's test. Data were presented with figures and tables and expressed as means±standard deviation (SD). Statistical test was repeated one-way ANOVA, *p*-values ≤0.05 were considered significant. The qualitative analysis was the time period of the content of IL-1β for four groups at 3rd day, 7th day and 14th day

respectively.

3. Results

Based on the experiment with three replications conducted among 72 experimental mice, mean values for 4 groups of experimental mice treated by vaseline (the negative control group), bioplasenton (the positive control group), the extracts of *Piper betel* leaves at both 2% and 3% concentrations were illustrated in Figure 1.




Figure 1: Comparison of mean values of the concentration of IL-1β among four groups of experimental mice (n=6) consisted of the negative control group (vaseline), the positive control (bioplasenton), the extracts of *Piper betel* leaves (2% and 3% concentrations) consecutively. Each concentration of IL-1β for each group of experiment was determined for 3rd day, 7th day and 14th day respectively. Data were presented as mean±standard deviation ($p \leq 0.05$).

The data in Figure 1 were then transformed into Table 1, where the lowest concentration of IL-1β at 3rd day was the extracts of betel leaves at 3% concentration (0.97 ± 0.46), whereas, the highest concentration of of IL-1β at the 3rd day was the positive control group (1.83 ± 0.81).

At 7th day after the treatment, the lowest concentration of IL-1β at the 3rd day was the positive control or bioplasenton (0.92 ± 0.55), whereas, the highest concentration of IL-1β was the extracts of betel leaves at 3% concentration (2.67 ± 1.71).

As illustrated in Table 2, the mean differences of the concentration of IL-1β for the negative control were -0.35 (3rd day and 7th day), -0.112 (7th day and 14th day), and -0.770 (3rd day and 14th day) respectively, whereas, the mean values of the concentration of IL-1β for the positive control was 0.90 (7th day and 14th day), -1.19 (3rd day and 14th day), and -0.290 (3rd day and 14th day) respectively. On the other hand, the mean differences of the concentration of IL-1β for the extracts of *Piper betel* leaves at 2% concentration were -0.49 (3rd day and 14th day), -0.73 (7th day and 14th day), and -1.23 (3rd day and 14th day) respectively, whereas, the mean differences of the concentration of IL-1β for the extracts of *Piper betel* leaves at 3% concentration were -0.72 (3rd day and 7th day), 0.90 (7th day and 14th day), and -1.26 (3rd day and 14th day) respectively.

Table 1: The values of mean ± SD of the concentration of IL-1β for experimental mice using an acute wound healing modeling based on the division of groups and time period.

Groups	n	Concentration of IL-1β		
		(Mean ± SD)		
		3rd day	7th day	14th day
Negative control (vaseline)	6	1,49±0.93	1,15±0.86	2.67±1.71
Positive control (bioplasenton)	6	1.83±0.81	0.92±0.55	2.11±0.70
Extracts of <i>Piper betel</i> leaves (2%)	6	1.07±0.65	1,57±0,41	2,30±1.66
Extracts of <i>Piper betel</i> leaves (3%)	6	0.97±0.46	1.68±0.67	2.58±1.05

Table 2: Mean differences of the concentration of IL-1β for experimental mice using an acute wound healing modeling based on the division of groups and time period.

Groups	Mean Differences		
	3rd-7th	7th-14th	3rd-14th
Control			
Negative	-0,35	-1,12	-0,77
Positive	-0,90	-1,19	-0,29
Exp.			
<i>Piper betel</i> (2%)	-0,49	-0,73	-1,23
<i>Piper betel</i> (3%)	-0,72	-0,90	-1,62

As shown in Table 3, *p*-value for the negative control group (vaseline) was 0.217. Hence, the correlation between the overall time period (3rd day, 7th day and 14th day) was not statistically significant with the concentration of IL-1β for the negative control group ($r = 0.218$). Therefore, there was probability the regular

increase of the concentration of IL-1 β in wound healing process. On the other hand, p -value for the positive control group (bioplasenton) was 0.249. Hence, the correlation between the overall time period (3rd day, 7th day and 14th day) was not statistically significant with the concentration of IL-1 β for the positive control group ($r = 0.190$). Accordingly, there was probability the regular increase of the concentration of IL-1 β in wound healing process.

Table 3: The correlation between the overall time period and the concentration of IL-1 β for experimental mice based on an acute wound healing modeling.

Groups	Correlations	
	p^*	r^*
Negative	0,217	0,218
Positive	0,249	0,190
<i>Piper betel</i> (2%)	0,060	0,420
<i>Piper betel</i> (3%)	0,002	0,701

p^* One-Way Anova Statistical Test.

r^* Spearman's correlation

p -value for the treatment group for the extracts of *Piper betel* leaves (2% concentration) was 0.060. Hence, the correlation between the overall time period (3rd day, 7th day and 14th day) was not statistically significant with the concentration of IL-1 β for this treatment group ($r = 0.420$). Therefore, there was probability the regular increase of the concentration of IL-1 β in wound healing process. On the contrary, p -value for the treatment group for the extracts of *Piper betel* leaves (3% concentration) was 0.002. Hence, the correlation between the overall time period (3rd day, 7th day and 14th day) was statistically significant with the concentration of IL-1 β for the treatment group for the extracts of *Piper betel* leaves at 3% concentration ($r = 0.701$). Therefore, there was probability the regular increase of the concentration of IL-1 β in wound healing process.

4. Discussion

Both p -value and r -value for the negative control group based on statistical test using Spearman's correlation were 0.217 and 0.218 respectively. Hence, there was not any correlation between the overall time period (3rd, 7th and 14th) and the content of IL-1 β for the negative control group. Therefore, there was probability the decrease of IL-1 β content at the overall time period (3rd, 7th and 14th) in wound healing process. On the other hand, both p -value and r -value for the positive control group based on statistical test using Spearman's correlation were 0.249 and 0.190 respectively. Hence, there was not any correlation between the overall time period (3rd, 7th and 14th) and the content of IL-1 β for the positive control group. Therefore, there was probability the decrease of IL-1 β content at the overall time period (3rd, 7th and 14th) in wound healing process.

Both p -value and r -value for the treatment group of the extracts of *Piper betel* leaves at 2% concentration based on statistical test using Spearman's correlation were 0.060 and 0.420 respectively. Hence, there was not any correlation between the overall time period (3rd, 7th and 14th) and the content of IL-1 β for this group. Therefore, there was probability the decrease of IL-1 β content at the overall time period (3rd, 7th and 14th) in wound healing process. On the other hand, both p -value and r -value for the treatment group of the extracts of *Piper betel* leaves at 3% concentration based on statistical test using Spearman's correlation were 0.002 and 0.701 respectively. Hence, there was correlation between the overall time period (3rd, 7th and 14th) and the content of IL-1 β for this control group. Therefore, there was probability the increase of IL-1 β content at the overall time period (3rd, 7th and 14th) in wound healing process.

14 IL-1 β is a member of the interleukin 1 family of cytokines. This cytokine is produced by activated macrophages, leukocytes, keratinocytes, and fibroblasts. Its target cells are endothelial cells, macrophages, leukocytes, keratinocytes and fibroblasts. Its signaling/receptors are ICAM-1, VCAM-1, IL-1 in which such receptors have pivotal roles in inflammation, angiogenesis, re-epithelization, and remodeling of tissues [36].

8 *Piper betel* is used in the traditional medicine as diuretic as the antioxidant activity of ethanol, chloroform and ethyl acetate extract of leaves and stem of *Piper betel* where the ethanol extracts of leaves and stems of betel have antioxidant activity [37].

Collagen is the key component in the phases of wound healing. As soon as after wounding, the exposure of fibroblastic collagens in blood streams leads to the aggregation and activation of thrombocyte and releasing chemotactic factors beginning the process of wound healing. Fragments of collagens release leukocyte collagenase to pull fibroblasts into the injury areas. After then collagens are used to form new extracellular matrix [38].

13 A previous study [39] that investigates the inflammatory cytokines such as IL-1 α and IL-1 β , IL-6 and TNF (tumor necrosis factor- α) during wound healing process in experimental rats using enzyme-linked immunosorbent assay and immunostaining for an observation interval of 0-240 hours (three days) provides facts that the concentrations of TNF- α and IL-1 β begin to rise after the onset of wounding and achieve the peak period for 12 hours from the onset of wounding. Infiltration of various leukocytes shows acute inflammation at three hours and six hours, and the main source of this cytokine is immunohistochemical substance identified as neutrophil. Such a study provides fact that both TNF and IL-1 β have pivotal roles in triggering the onset of inflammation. Rebound of the concentration of cytokines is 72 hours from the onset of wounding. 40 Histological examination of the wound at 72 hours shows the migration of fibroblasts and the formation of new granulation as the evidence of the proliferative phase in wound healing process. This is positively correlated with the result of our study where the treatment of *Piper betel* leaves at 3% concentration increased the wound healing process at third day (72 hours). This is also correlates with the study conducted by another study [40] that investigates the effect of the extraction of *Piper betel* leaves on the erythrocytes of experimental mice showing that there is significant reduction in TBARS and significant increase in ascorbic acid, vitamin E, super oxide dismutase, catalase and membrane-bound ATPases among experimental mice fed with 0.2 gm of *Piper betel* leaves per day, the lower concentration than the concentration of *Piper betel* leaves used in our study (3% concentration).

5. Conclusion

The evidence from this study revealed that the extracts of *Piper betel* leaves at 3% concentration could increase the wound healing process among experimental mice than the treatment of vaseline and bioplasenton as well as the treatment of the extracts of *Piper betel* at 2% concentration where the wound healing process increased based on the ascending time at 3rd day, 7th day and 14th day. Results of this study showed the potency of *Piper betel* leaves in improving wound healing process. Further studies should be conducted in investigating the role of *Piper betel* leaves on the physiological and immunological mechanisms.

References

- [1] MacLellan, L., Gardner, G., & Gardner, A. (2002). Designing the future in wound care: the role of the nurse practitioner. *Primary Intention, Vol.10 No.3* , 97-112.
- [2] Alam, B., Akter, F., Parvin, N., Pia, R. S., Akter, S., Chowdhury, J., et al. (2013). Antioxidant, analgesic and anti-inflammatory activities of the methanolic extract of *Piper betel* leaves. *Avicenna J Phytomed, Vol. 3, No. 2, Spring 2013* , 112–125.
- [3] Bhide SV, Zariwala MBA, Amonlar AJ, Azuine MA. Chemo-preventive efficacy of betel leaf extract against benzo(a)pyrene induced fore-stomach tumors in mice. *J Ethnopharmacol.* 1991;34:207–213. [PubMed].
- [4] Garg SC, Jain R. Biological activity of the essential oil of *Piper betel* L. *J Essen Oil Res.* 1992;4:601–606.
- [5] Singh M, Shakya S, Soni VK, Dangi A, Kumar N, Bhattacharya SM. The n-hexane and chloroform fractions of *Piper betel* L. trigger different arms of immune responses in BALB/c mice and exhibit antifilarial activity against human lymphatic filarid *Brugia malayi*. *Int Immunopharmacol.* 2009;9:716–728. [PubMed]
- [6] Choudhary D, Kale RK. Antioxidant and non toxic properties of *Piper betel* leaf extract: in vitro and in vivo studies. *Phytother Res.* 2002;16:461–466. [PubMed]
- [7] Santhakumari P, Prakasam A, Pugalendi KV. Modulation of oxidative stress parameters by treatment with *Piper betel* leaf in streptozotocin induced diabetic rats. *Indian J Pharmacol.* 2003;35:373–378.
- [8] Shun CY, Chau JW, Jing JL, Pei LP, Jui LH, Fen PC. Protection effect of *Piper betel* leaf extract against carbon tetrachloride induced liver fibrosis in rats. *Archives of Toxicology.* 2007; 81:45-55.
- [9] Lei D. Antioxidant and antiplatelet effect of aqueous inflorescence *Piper betel* extract. *Journal of Agricultural Food and Chemistry.* 2003; 51: 2083-8.
- [10] Majumdar B, Chaudhuri SR, Roy A. Potent antiulcerogenic activity of ethanol extract of leaf of *Piper betel* Linn. by antioxidative mechanism. *Indian Journal of Clinical Biological Chemistry.* 2002; 17: 49-57.
- [11] Adhikari P, Chowdhury D, Banerji J, Chatterjee A. Antifertility effect of crude alcoholic extract of *Piper betel* stalk. *Indian Journal of Physiology and Allied Sciences.* 1998; 52: 22-7.
- [12] Adhikary P, Banerji J, Choudhuri D, Das AK, Deb CC, Mukherjee P, Chatterjee SR. Effect of oral administration of stalk of leaves of *Piper betel* Linn on oestrous cycle and its antifertility activity in rats. *Indian Journal of Physiology and Allied Sciences.* 1990; 44:116-23.

- [13] Chen SJ, Wo BN, Yeh JL, Lo YC., Chen IS, Chen IJ. C-fiber evoked autonomic cardiovascular effect of after injection of *Piper betel* inflorescence extracts. *Journal of Ethnopharmacology*. 1995;43: 183-8.
- [14] Amonkar AJ, Padma PR, Bhide S. Protective effect of hydroxychavicol, a phenolic component of *Piper betel* leaf against the tobacco-specific carcinogens. *Mutation Research. Fundamental and Molecular Mechanisms of Mutagenesis*. 1989;210: 249-53.
- [15] Varier PS. *Piper betel* Linn. In: Varier PS. (ed) *Indian medicinal plants*. Kottakkal : Oreint Longman. 1997; 8: 279-283.
- [16] Agarwal T, Singh R, Shukla AD, Waris I, Gujrati A. Comparative analysis of antibacterial activity of four *Piper betel* varieties. *Adv Appl Sc Res*. 2012;3:698–705.
- [17] Majumdar B, Chaudhuri SGR, Ray A, Bandyopadhyay SK. Effect of ethanol extract of *Piper betel* Linn leaf on healing of NSAID-induced experimental ulcer a novel role of free radical scavenging action. *Indian J Exp Biol*. 2003;41:311–315. [PubMed].
- [18] Saravanan R, Prakasam A, Ramesh B, Pugalendi KV. Influence of *Piper betel* on hepatic marker enzymes and tissue antioxidant status in ethanol-treated wistar rats. *J Med Food*. 2002;5:197–204. [PubMed].
- [19] Ratnasooriya WD, Jayawardena KGI, Premakumara GAS. Antimotility effects of *Piper betel* (L) leaf extract on washed human spermatozoa. *J Nat Sci Council*. 1990;18:53–60.
- [20] Freedberg IM, Tomic-Canic M, Komine M, Blumenberg M. Keratins and the keratinocyte activation cycle. *J Invest Dermatol* 2001; 116: 633–40.
- [21] Kupper TS, Deitch EA, Baker CC, Wong WC. The human burn wound as a primary source of interleukin-1 activity. *Surgery* 1986; 100: 409–15.
- [22] Murphy GM, Dowd PM, Hudspith BN, Brostoff J, Greaves MW. Local increase in interleukin-1-like activity following UVB irradiation of human skin in vivo. *Photodermatol* 1989; 6: 268–74.
- [23] Bochner BS, Charlesworth EN, Lichtenstein LM, Derse CP, Gillis S, Dinarello CA, Schleimer RP. Interleukin-1 is released at sites of human cutaneous allergic reactions. *J Allergy Clin Immunol* 1990; 86 (6 Pt 1): 830–9.
- [24] Mizutani H, Schechter N, Lazarus G, Black RA, Kupper TS. Rapid and specific conversion of precursor interleukin 1 beta (IL-1 beta) to an active IL-1 species by human mast cell chymase. *J Exp Med* 1991; 174: 821–5.
- [25] Chan LS, Hammerberg C, Kang K, Sabb P, Tavakkol A, Cooper KD. Human dermal fibroblast interleukin-1 receptor antagonist (IL-1ra) and interleukin-1 beta (IL-1 beta) mRNA and protein are co-stimulated by phorbol ester: implication for a homeostatic mechanism. *J Invest Dermatol* 1992; 99: 315–22.
- [26] Wood LC, Elias PM, Calhoun C, Tsai JC, Grunfeld C, Feingold KR. Barrier disruption stimulates interleukin-1 alpha expression and release from a pre-formed pool in murine epidermis. *J Invest Dermatol* 1996; 106: 397–403.
- [27] Lundqvist EN, Egelrud T. Biologically active, alternatively processed interleukin-1 beta in psoriatic scales. *Eur J Immunol* 1997; 27: 2165–71.
- [28] Zepter K, Haffner A, Soohoo LF, De Luca D, Tang HP, Fisher P, Chavinson J, Elmetts CA. Induction of biologically active IL-1 beta-converting enzyme and mature IL-1 beta in human keratinocytes by

- inflammatory and immunologic stimuli. *J Immunol* 1997; 159: 6203–8.
- [29] Murphy JE, Robert C, Kupper TS. Interleukin-1 and cutaneous inflammation: a crucial link between innate and acquired immunity. *J Invest Dermatol* 2000; 114: 602–8.
- [30] Corsini E, Primavera A, Marinovich M, Galli CL. Selective induction of cell-associated interleukin-1 α in murine keratinocytes by chemical allergens. *Toxicology* 1998; 129: 193–200.
- [31] Giraldo S, Jesus Shancez, Quentin Felty, Deodutta Roy. 2008. Atlas of Genytogenetics in Oncology and Haematology.
- [32] Spielmann, H.E., Genschow, M., Leibsch, M. & Halle, W., 1999, 'Determination of the starting dose for acute oral toxicity. LD50 testing the up and down procedures from cytotoxicity data', *Alternatives to Laboratory Animals* 27, 957–966.
- [33] WHO. (2000). General Guidelines for Methodologies on Research and Evaluation of Traditional Medicine. Geneva: World Health Organization.
- [34] Simonsen, L., Petersen, M.B. & Groth, L., 1992, 'In vivo skin penetration of salicylic compounds in hairless rats', *European Journal of Pharmaceutical Sciences* 17, 95–104.
- [35] Kun, Li. et al. (2011). Tannin Extract From Immature Fruits of Terminalia Chebula Fructuz Retz. Promote Cutaneous Wound Healing In Rats. *BMC Complementary and Alternative Medicine* , 11: 86-95.
- [36] Behm. 2011.
- [37] Patil, D. D., & Wadhawa, G. C. (2014). Antioksidant Effect of The Stem And Leaves of Piper Betle. *International Journal of Pharmacy*, 4(2) , 73-75.
- [38] Mercandetti, M. (2015). *Wound Healing and Repair*. Accessed on 27-7-2016 from <http://emedicine.medscape.com>: <http://emedicine.medscape.com/article/1298129-overview>.
- [39] Kondo & Oshima, 1996. The Dynamics of Inflammatory Cytokines in the Healing Process of Mouse Skin Wound: A Preliminary Study for Possible Wound Age Determination. Springer.
- [40] Chitra and Vidya.2006. Dose Dependent Effect of Piper Betel Linn Leaf Extract on Erythrocytes of Experimental Mice. Sri Ramachandra Journal of Medicine Vol. 1. 1 September 2006.

The_Effect_of_the_Extracts_of_Betel__Piper_betel__Leaves_on.

ORIGINALITY REPORT

%24
SIMILARITY INDEX

%16
INTERNET SOURCES

%16
PUBLICATIONS

%10
STUDENT PAPERS

PRIMARY SOURCES

1	repository.up.ac.za Internet Source	%4
2	Submitted to University Tunku Abdul Rahman Student Paper	%1
3	onlinelibrary.wiley.com Internet Source	%1
4	International Journal of Sociology and Social Policy, Volume 19, Issue 7-8 (2012-04-14) Publication	%1
5	www.sriramachandra.edu.in Internet Source	%1
6	krishikosh.egranth.ac.in Internet Source	%1
7	www.ijppsjournal.com Internet Source	%1
8	www.ijpjournal.org Internet Source	%1
9	A H A Aziz, M A C Yunus, N H Arsad, N Y Lee,	

Z Idham, A Q A Razak. "Optimization of supercritical carbon dioxide extraction of Piper Betel Linn leaves oil and total phenolic content", IOP Conference Series: Materials Science and Engineering, 2016

Publication

% 1

10

Takeuchi, Hisanao, Kozo Hukaya, Katsuji Narita, and Keiichiro Muramatsu. "Changes in Creatine and Urea Formation in Rats Fasted and Fed Low Protein Diets", Agricultural and Biological Chemistry, 1970.

Publication

% 1

11

www.science.gov

Internet Source

% 1

12

Submitted to CSU, San Marcos

Student Paper

% 1

13

Ohshima, T.. "Forensic wound examination", Forensic Science International, 20000911

Publication

% 1

14

en.wikipedia.org

Internet Source

% 1

15

pure.uva.nl

Internet Source

% 1

16

www.spandidos-publications.com

Internet Source

<% 1

17

Submitted to Universitas Hasanuddin

Student Paper

<% 1

18

E.T. Pavlidis, K.D. Ballas, K. Psarras, N.G. Symeonidis et al. "Intraperitoneal Administration of Bevacizumab Intraoperatively Does Not Affect Abdominal Wound Healing in Rats", European Surgical Research, 2011

Publication

<% 1

19

Meghna Singh, Shilpy Shakya, Vishal Kumar Soni, Anil Dangi, Nikhil Kumar, Shailja-Misra Bhattacharya. "The n-hexane and chloroform fractions of Piper betle L. trigger different arms of immune responses in BALB/c mice and exhibit antifilarial activity against human lymphatic filarid Brugia malayi", International Immunopharmacology, 2009

Publication

<% 1

20

www.scielo.br

Internet Source

<% 1

21

A. A. Shahin, A. M. Saleh, H. A. Moustafa, A. A. Aboul-Ella, H. M. Gabr, M. M. Ismail. "Protective effect of folic acid on low-dose methotrexate genotoxicity", Zeitschrift für Rheumatologie, 2001

Publication

<% 1

22

D. A. RIBEIRO. "Biocompatibility of glass?"

<% 1

ionomer cements using mouse lymphoma cells
in vitro", Journal of Oral Rehabilitation, 12/2006

Publication

23

Submitted to Institute of Graduate Studies, UiTM

Student Paper

<% 1

24

tessera.spandidos-publications.com

Internet Source

<% 1

25

D. G. Kanjwani. "Evaluation of
Immunomodulatory Activity of Methanolic
Extract of Piper betel", Scandinavian Journal of
Immunology, 6/2008

Publication

<% 1

26

iovs.arvojournals.org

Internet Source

<% 1

27

lib.bioinfo.pl

Internet Source

<% 1

28

repository.um.edu.my

Internet Source

<% 1

29

Submitted to University of Hertfordshire

Student Paper

<% 1

30

pubs.acs.org

Internet Source

<% 1

31

www.i-scholar.in

Internet Source

<% 1

32

Submitted to De La Salle University - Manila

Student Paper

<% 1

33

Submitted to Sekolah Pelita Harapan

Student Paper

<% 1

34

vseryukzaki.ru

Internet Source

<% 1

35

Submitted to Higher Education Commission
Pakistan

Student Paper

<% 1

36

Le Thi Lien, Nguyen Thi Tho, Do Minh Ha,
Pham Luong Hang, Phan Tuan Nghia, Nguyen
Dinh Thang. "Influence of phytochemicals in
piper betle linn leaf extract on wound healing",
Burns & Trauma, 2015

Publication

<% 1

37

Submitted to University of Sheffield

Student Paper

<% 1

38

Submitted to Lovely Professional University

Student Paper

<% 1

39

Siti Imroatul Maslikah, Nuning Wulandari, Sri
Rahayu Lestari, Siti Nur Arifah. " The Effects of
Red Betel () Leaf Extract to Decreasing Profile
Population of CD4 and CD8 Cells in
Rheumatoid Arthritis-Induced Mice ", IOP
Conference Series: Earth and Environmental

<% 1

40 Toshikazu Kondo. "The dynamics of inflammatory cytokines in the healing process of mouse skin wound: A preliminary study for possible wound age determination", International Journal of Legal Medicine, 09/1996

Publication

41 "Drug Discovery and Evaluation: Pharmacological Assays", Springer Nature, 2016

Publication

42 Submitted to The University of Manchester

Student Paper

43 Masoko, Peter, Jacobus N. Eloff, and Jackie Picard. "The use of a rat model to evaluate the in vivo toxicity and wound healing activity of selected Combretum and Terminalia (Combretaceae) species extracts", Onderstepoort Journal of Veterinary Research, 2010.

Publication

44 Submitted to International Islamic University Malaysia

Student Paper

45 Submitted to University of Bedfordshire

Student Paper

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5
WORDS