

THE RELATIONSHIP BETWEEN TRIPLE TUMOR MARKER (CEA, CA 19-9, AND CA 125) AND COLORECTAL CANCER METASTASES AT MAKASSAR, INDONESIA

by Ibrahim Labeda

Submission date: 15-Jun-2021 03:31PM (UTC+0700)

Submission ID: 1606818965

File name: Ibrahim_Labeda_THE_RELATIONSHIP_BETWEEN_TRIPLE_TUMOR.pdf (199.93K)

Word count: 3869

Character count: 18589

1 THE RELATIONSHIP BETWEEN TRIPLE TUMOR MARKER (CEA, CA 19-9, AND CA 125) AND COLORECTAL CANCER METASTASES AT MAKASSAR, INDONESIA

Agung Kurniawan*, Ronald E. Lusikooy*, Mappincara Mappincara*, Murny A. Rauf*, Ibrahim Labeda*, Warsinggih Warsinggih*, Samuel Sampetoding*, M. Ihwan Kusuma*, M. Iwan Dani*, Ilham Jaya Patellongi** and Muhammad Faruk***,1

*Division of Digestive, Department of Surgery, Faculty of Medicine, Hasanuddin University Makassar, Indonesia., ** Department of Biostatistic, Faculty of Public Health, Hasanuddin University, Makassar, Indonesia., *** Department of Surgery, Faculty of Medicine, Hasanuddin University Makassar, Indonesia.

ABSTRACT Introduction: Tumor markers combination (CEA, CA19-9, and CA125) can increase sensitivity due to interactions between several epitopes and tumor cells, especially in colorectal cancer (CRC) diagnosis. Therefore, this study aimed to assess the relationship between triple tumor markers (CEA, CA 19-9, CA 125) and CRC metastases. **Materials and Methods:** The research samples were CRC patients at Dr Wahidin Sudirohusodo Hospital, Makassar from November 2019 to April 2020. The histopathology data were collected through endoscopic or anal biopsy, CEA, CA 19-9, CA 125, Thorax X-Ray, Abdomen CT Scan, Surgery, and data processing. Furthermore, statistical tests were performed with the Mann Whitney difference test (significance = $p < 0.05$). **Results:** 57 samples were obtained, consisting of 19 metastatic CRC and 38 non-metastatic cases with an age range of 18-82 years, and an average of 56.9 years. Furthermore, most of the patients were males with a ratio of 2:1, and the tumors are mostly located in rectum by 68.5%, of which many are in stage II with 52.6%. The CEA value was obtained at ≥ 10 ng/ml (61.4%) and < 10 ng/ml (38.6%), CA 19-9 was ≥ 10 ng/ml (61.4%) and < 10 ng/ml (38.6%), and CA 125 was ≥ 10 ng/ml (61.4%) and < 10 ng/ml (38.6%). Also, most metastases are located in the liver. Triple tumor marker significantly correlated with CRC ($p < 0.05$), and the accuracy of three combinations increased (87.7%), which indicates the markers can establish the presence or absence of metastasis in CRC patients. **Conclusion:** The markers (CEA, CA 19-9 and CA 125) can be used to establish the presence or absence of metastasis in CRC patients.

KEYWORDS Tumor Markers, Metastasis, Colorectal Cancer, CEA

12 Introduction

Colorectal cancer is the third most common malignancy after lung and breast, and it is the second leading cause of cancer

2
Copyright © 2020 by the Bulgarian Association of Young Surgeons
DOI:10.5455/IJMRCR.colorectal-cancer-indonesia

First Received: May 08, 2020

Accepted: May 28, 2020

Associate Editor: Ivan Inkov (BG);

¹ Department of Surgery, Faculty of Medicine, Hasanuddin University Makassar, Indonesia; Email: faroex8283@gmail.com

death after lung [1]. The key to CRC survival is an early and accurate diagnosis. Furthermore, malignant tumors can spread through invasion, or move from its primary site to secondary sites in the body; this process is called metastasis. It is responsible for 90% of cancer patient deaths.

Early and precise diagnosis according to NCCN guidance 2019 consists of history taking, physical examination, tumor markers, colonoscopy, imaging (CT scan, MRI, Pet CT) and biopsy. However, these methods have several drawbacks such as failure of interpretation, the complexity of implementation, high costs, invasive, time-consuming procedures, and repeat requirements (3-5 years), which has led to bad reputation and

poor patient compliance. Therefore, research is still needed on the use of tumor markers as a diagnostic tool for a non-invasive, quick, economical, and simple method of detecting metastasis [2,3].

These markers are detected through blood, urine, and body tissue (tumor tissue) of cancer patients. Furthermore, EGTM 2007, ASCO 2014, and NCCN 2019 on CRC often used and recommended CEA (Carcinoembryonic Antigen) marker. This is an oncofetal antigen, a high molecular weight glycoprotein produced by colon cells and functions as an intercellular adhesion molecule, which promotes CRC cell aggregation. Also, CA 19-9 acts as an intracellular adhesion molecule that affects cellular synthesis in various parts of the digestive system such as gastric, color [16], and especially sensitive to the pancreas. Furthermore, CA (Carbohydrate Antigen) 125 is a sensitive marker in ovarian cancer, and it is also detected in gastrointestinal cancers such as gastric and colorectal. In addition, it is often used to detect peritoneal metastases in ovarian cancer [4].

According to Li et al. (2013), markers combination can increase sensitivity due to the interaction between several epitopes and tumor cells, which allows for clear cell identification [5]. Lalosevic, a pre-operative CEA combination and CA19-9 serum can be used to diagnose lymph node and liver metastases in CRC patients [6]. Furthermore, Li et al. (2017) found 4 markers in combination [17] causing increased sensitivity. According to Gao et al. (2018), CEA, CA19-9, CA72-4, CA125, and Ferritin serum can be used as diagnostic markers for CRC [3, 7]. Based on the facts above, researchers tried to ascertain the relationship between triple tumor markers and CRC metastases using easily obtained markers and often used in Indonesia, which are CEA, CA 19-9 and CA 125.

4 Patients and Samples

This was an observational study with a cross-sectional approach in the Department of Digestive Surgery, Dr Wahidin Sudirohusodo Hospital in Makassar, South Sulawesi. The research lasted for six months from 1 November 2019 to 31 April 2020.



The population was CRC patients that had not undergone surgery, and with a biopsy of Lower Gastrointestinal Endoscopy (LGIE) and had a histopathological examination. The sample was the entire population that met the research criteria by consecutive sampling. Inclusion Criteria: Patients that did not experience total mechanical intestinal obstruction and perforation prior to surgery, and willing to be a research subject. Exclusion criteria: Patients that have undergone surgery, chemotherapy, radiotherapy, and had incomplete clinical and histopathological data.

The principle of taking serum from the patient's blood

The cubital fossa area was disinfected with 70% alcohol swab, then 2 cc blood was taken with a disposable spout, followed by centrifugation at 200 rpm to get the serum.

Examinations of CEA, CA 19-9, and CA 125

This uses the electrochemiluminescent immunoassay (ECLIA) method from the Elicys 2010 with Linearity of the tools: 0.200 - 1000 ng / ml (CEA), 0.200 - 3840 u/ml (CA 19-9), and 0.250 - 19,000 u / ml (CA 125). This technological method is for immunoassay detection, which uses antibodies to detect and measure specific analytes such as peptides, proteins, and hormones.





22 Research Processing and Analysis 9

The data was analyzed using SPSS version 22 (IBM SPSS Statistics for Windows, Version 22.0. (IBM Corp., Armonk, NY). The statistical test used was the Mann Whitney difference test. Meanwhile, a significant result was obtained with P and It value at 0.05.

Results

There were 271 subjects with only 57 meetings the criteria. 19 cases were metastatic, and 38 were non-metastatic with an age range of 20 - 60 years, and an average of 56.9 years. Furthermore, Male and female patients were 66.6% and 33.4% respectively, with a ratio of 2:1. In this study, the tumor was mostly found in the rectum of 68.5%, and the fewest was found in the transverse colon of 5.2%. According to the stage, patients with stage II, III, and IV were 52.6%, 14.1%, and 33.3% respectively and there was no patient with stage I. Based on CEA, 61.4% had a value of \geq





5 10 ng/ml while 38.6% had ≤ 10 ng/ml. Also, for CA 19-9, 61.4% had ≥ 10 ng/ml while 38.6% had ≤ 10 ng/ml. Based on the CA 125, 61.4% had ≥ 10 ng/ml while 38.6% had ≤ 10 ng/ml (Table 1). Based on location, the most liver metastases were 15 patients with CEA of ≥ 10 ng/ml 87%, CA 19-9 ≥ 37 u/ml 66%, and CA 125 ≥ 35 u/ml 40%. Also, CEA levels of <10 ng/ml was 13%, CA 19-9 <37 u/ml was 33%, and CA 125 <35 u/ml was 60%. Based on the location, pulmonary metastases were 2 patients, peritoneal 1, and bone was also 1 (Table 2).

*Mann Whitney test

1. DIFFERENCES IN BIOMARKER LEVELS BETWEEN NON-METASTATIC AND METASTATIC CRC Based on table 3 results, the three markers (CEA, CA 19-9, CA 125) were significantly different ($p < 0.05$) between non-metastatic and metastatic CRCs. Furthermore, the differences in CEA levels (figure 1), CA 19-9 (figure 2), CA 125 (figure 3) seem to be higher in metastatic than in non-metastatic. Also, it was found that the probability value ($p < 0.05$) of CEA, CA 19-9, and CA 125 were <0.001 , 0.019, and 0.002 respectively, which were significantly correlated. This means they have a strong relationship with metastasis. Figure 4 shows the combined ROC curves of biomarkers through multiple logistic regression analysis which indicates higher accuracy.
2. SENSITIVITY COMPARISON AND SPECIFICITY BETWEEN TRIPLE TUMOR MARKER AND SINGLE TUMOR MARKER IN CRC METASTASIS Table 4 shows the diagnostic test analysis results of each biomarker and a combination of the three. CEA, CA 19-9 and CA 125 have accuracy (81.8%, 69.2%, 75.9%), sensitivity (73.7%, 68.4%, 73.7%), specificity (84.2%, 68.4%, 63.2%), PPV (70.0%, 52.0%, 50.0%), and NPV (86.5%, 81.3%, 82.8%) respectively. Based on these values, CEA is the best. When the three biomarkers are combined, the accuracy, sensitivity, and NPV increase to 87.7%, 84.3%; and 90.9% respectively. However, the PPV specificity decrease. Therefore, the results show an increase in sensitivity, a negative predictive value (NPV), a decrease in specificity and a positive predictive value (PPV) of triple tumor marker on CRC metastases.

Discussion

1. Research Characteristics Based on the age group, the subjects were 18-82 years old with a mean of 56.9 years, and those > 60 years old had the highest percentage level of 49.2%. Also, the tendency of CRC is high in the age group of 50-59 years and ≥ 60 years by 29.8% (17/57) and 49.2% (28/57), respectively. Furthermore, the percentage above

and below 50 years was 79% (45/57) and 21% (12/57) respectively. In the past decade, there has been a trend change in Indonesia, where patients above 50 years experienced a decrease, and those below that age experienced an increase. This sharp increase was caused by (1) environmental factors influence which is either direct or indirect through dietary pattern change in young Indonesian people. This is a consequence of both increased prosperity and a shift towards the westerners' way of eating (westernization), which is high fat and low fiber foods. (2) genetic susceptibility factors, such as individuals born with mutations or genetic defects, for example, in hereditary groups such as HNPCC / Lynch syndrome [8].

Based on gender, male and female patients were 66.6% (38/57) and 33.4% (19/57) respectively, with a ratio of 2:1 (Table 7). According to Globocan 2018, CRC is more common in men than women, and 3-4 times common in developed countries. Also, men are more often affected than women with ratio of 7.7 / 100,000: 4.4 / 100,000 [1]. Based on location, the rectum is mostly affected 68.5% (39/57), the left colon of 15.8% (9/57), the right colon of 12.2% (7/57) and the fewest are in the transverse colon of 3.5% (2/57). Also, the patients were dominated by those with rectal cancer, probably due to the time factor.

Based on stage, most of the patients were in stage II 52.6% (30/57), stage IV 33.3% (19/57), and the fewest were in stage III 14.1% (8/57), and none with stage I. Lusikoooy 2013 reported that, out of the 159 CRC patients, the majority were still in stage IIb 27.7% (44/159), stage IIa 22.1% (35/159), stage IIIb 15.1% (24/159), stage IIIa 13.2% (21/159), stage IV 20.7% (33/159) and stage Ia - Ib each 0.6% (1 / 159) [9]. Therefore, the results show that most CRC cases are still in stages IIa and IIb. This might be due to detection from colonoscopy investigations; thus, the cases can be diagnosed early.

According to table 2, the liver is the most metastasis location, which is 78% (15/19). This is following Palaghia (2015) which stated that most CRC metastasis location is in the liver, at the time of initial diagnosis (20-25% of cases) or after primary tumor resection (40% of cases) [10].

2. RELATIONSHIP BETWEEN CEA AND CRC METASTASIS Figure 1 shows CEA levels in metastatic CRCs are higher than in non-metastatic. In the Mann Whitney test analysis, a Probability value of (Sig) <0.05 is <0.001 was found, which means the level has a significant correlation with CRC metastases. This is by Giovanni et al., in a long-term study in the form of 20 years surveillance, which stated that a serial CEA examination was performed on 125 of 239 patients and found that their sensitivity on the liver and non-hepatic metastases was 99% and 94% respectively. Therefore, they concluded that it is useful for grading preoperative colorectal tumors.
3. RELATIONSHIP BETWEEN CA 19-9 AND CRC METASTASIS Figure 2 shows CA 19-9 levels in metastatic CRC are higher than in non-metastatic. In the Mann Whitney test analysis, it was found that the Probability (Sig) value <0.05 was 0.019, which means the level was significantly correlated with CRC metastasis. This is in accordance with Malati et al. in 1996, which stated that CA 19-9 was also found to increase by 20-40% in colorectal cancers [11]. Fur-

Table 1 General Characteristics of Research Subjects.

Variable	Metastatic (n = 19)		Non- Metastatic (n = 38)		Total (n = 57)	
	n	%	n	%	n	%
Age Group (year old)						
< 20	-	-	1	2.6	1	1.7
20-29	-	-	1	2.6	1	1.7
30-39	1	5.2	3	7.8	4	7.1
40-49	1	5.2	5	13.1	6	10.5
50-59	8	42.2	9	23.7	17	29.8
≥ 60	9	47.4	19	50.2	28	49.2
Gender						
Male	14	73.7	24	63.1	38	66.6
Female	5	26.3	14	36.9	19	33.4
Tumor Location						
Right Colon	2	10.5	5	13.1	7	12.2
Transversum Colon	1	5.2	1	2.7	2	3.5
Left Colon	4	21.1	5	13.1	9	15.8
Rectum	12	63.2	27	71.1	39	68.5
Stage						
I	-	-	-	-	-	-
II	-	-	30	78.9	30	52.6
III	-	-	8	21.1	8	14.1
IV	19	100.0	-	-	19	33.3
CEA (ng/ml)						
< 10	2	10.6	20	52.6	22	38.6
≥ 10	17	89.4	18	47.3	35	61.4
CA 19-9 (u/ml)						
< 37	5	26.4	27	71.1	32	56.1
≥ 37	14	73.6	11	28.9	25	43.9
CA 125 (u/ml)						
< 35	11	63.1	30	78.9	41	73.6
≥ 35	8	36.9	8	21.1	16	26.4

Table 2 Characteristics of Metastasis based on marker levels.

Tumor Marker	Metastasis Location				
	Liver (n=15)	Peritoneum (n=1)	Pulmonary (n=2)	Bone (n=1)	Brain (n=0)
CEA (ng/ml)					
< 10	2 (13 %)	-	-	-	-
≥ 10	13 (87 %)	1 (100 %)	2 (100 %)	1 (100 %)	-
CA 19-9 (u/ml)					
< 37	5 (33 %)	-	-	1 (100 %)	-
≥ 37	10 (66 %)	1 (100 %)	2 (100 %)	-	-
CA 125 (u/ml)					
< 35	9 (60 %)	-	2 (100 %)	-	-
≥ 35	6 (40 %)	1 (100 %)	-	1 (100 %)	-

Table 3 Differences in biomarker levels between Non-metastatic and Metastatic CRCs.

Biomarker	Group	Description			p*
		Min-Max	Mean (medium)	95% IK	
CEA	Non-Metastatic	0,5 – 134,0	26,47(8,25)	14,35-38,58	<0,001
	Metastatic	3,5 – 139,0	79,35(89,00)	57,68-101,02	
CA 19-9	Non-Metastatic	4,2-156,0	40,87(26,50)	28,19-53,54	0,019
	Metastatic	9,0-203,0	70,35(65,00)	9,00-203,00	
CA 125	Non-Metastatic	2,8-158,5	25,22(14,94)	15,21-35,23	0,002
	Metastatic	11,5-327,525	66,17(104,36)	27,98-104,36	

*Mann Whitney test

Table 4 Value of accuracy, sensitivity, specificity, and prediction of each biomarker and its combination.

Biomarker	Accuracy	Cut off	Sen(Spes)	NPP(NPN)
CEA	81,8%	54,0	73,7%(84,2%)	70,0%(86,5%)
CA 19-9	69,2%	35,0	68,4%(68,4%)	52,0%(81,3%)
CA 125	75,9%	20,0	73,7%(63,2%)	50,0%(82,8%)
Combination-3	87,7%	0,20	84,3%(78,9%)	66,7%(90,9%)

thermore, Zhang et al. in 2015 concluded that it might be a tumor biomarker in addition to CEA for CRC [12].

- RELATIONSHIP BETWEEN CA 125 AND CRC METASTASIS Figure 3 shows the CA 125 level in metastatic CRC is higher than in non-metastatic. In the Mann Whitney test analysis, it was found that the Probability value (Sig) <0.05 was 0.002, which means the level is significantly correlated with the CRC metastasis. This is in accordance with Bast et. al. in 1998, which stated that besides ovaries, it also increase in colorectal malignancies by 24.1% [13]. Furthermore, Huang et al in 2016 found an increase in CA 125 levels in men and women with CRC stage 4 (peritoneal metastasis) with a 57.1% sensitivity and 92.0% specificity [14].
- RELATIONSHIP BETWEEN TRIPLE TUMOR MARKER (CEA, CA 19-9, and CA 125) AND CRC METASTASIS In Figure 5, the combined ROC curve of the three markers through multiple logistic regression analysis shows a higher accuracy. The markers combination was made possible based on Yang et al. in 2011, which found that they (CA19-9, CEA, and CA125) had an independent prognostic value to predict recurrence-free survival (RFS) in 5 years, in which patients with combined serum CA19-9, CEA and CA125 had the highest recurrence rate (100%), and the shortest RFS (median of 4 months) [15]. Therefore, combining CEA, CA19-9, CA72-4, CA125, and Ferritin can be used as a diagnostic marker for CRC [1].
- COMPARISON OF SPECIFICITY AND SENSITIVITY BETWEEN TRIPLE AND SINGLE TUMOR MARKER IN CRC Table 4 shows the diagnostic test analysis results of each biomarker and a combination of the three. CEA, CA 19-9 and CA 125 have accuracy (81.8%, 69.2%, 75.9%), sensitivity (73.7%, 68.4%, 73.7%), specificity (84.2%, 68.4%, 63.2%), NPP (70.0%, 52.0%, 50.0%), and NPV (86.5%, 81.3%, 82.8%) respectively. Therefore, the diagnostic values show CEA as the best biomarker. Also, when the three are combined, the accuracy, sensitivity, and NPV increase to 87.7%, 84.3%, and 90.9%, respectively. However, the specificity and NPP decrease. This shows triple markers can determine the presence or absence of metastasis in CRC patients. This is following Li et al. (2013) which stated that markers combination could increase sensitivity because the interaction between several epitopes and tumor cells allows more clear identification [5]. Further, Gao et al in 2018 found that several combinations (CEA, CA19-9, CA72-4, CA125, and Ferritin) can increase CRC diagnosis sensitivity and reliability, and assess the patient's pathological parameters [1].

Conclusion

This study shows triple markers (CEA, CA 19-9, and CA 125) can determine the presence or absence of metastasis in CRC patients.

Ethics committee approval

Ethics committee approval was received for this study from the ethics committee of Faculty of Medicine, Universitas Hasanuddin gave approval for this study (Ref.No.:1114/UN4.6.4.5.31/PP36/2019).

Funding

The authors declared that this study has received no financial support.

Conflict of interest

There are no conflicts of interest to declare by any of the authors of this study.

References

- Globocan. Fact sheets by population, estimated age-standardized incidence and mortality rates: both sexes (Indonesia).
- Sharma S. Tumor Markers in Clinical Practice – General Principles and Guidelines. Amrita J Med 2017; 10–17.
- Gao Y, Wang J, Zhou Y. Evaluation of Serum CEA, CA19-9, CA72-4, CA125 and Ferritin as Diagnostic Markers and Factors of Clinical Parameters for Colorectal Cancer. Sci Rep 2018; 8: 2732.
- Friedlander M. A Review of The Clinical Utility Of Ca 125 And Other Tumour Markers In Ovarian Cancer. Pathol Updat 2009; 15–20.
- Li X, Lu J, Ren H, et al. Combining multiple serum biomarkers in tumor diagnosis: A clinical assessment. Mol Clin Oncol 2013; 1: 153–160.
- Stojkovic Lalošević M, Stanković S, Stojković M, et al. Can preoperative CEA and CA19-9 serum concentrations suggest metastatic disease in colorectal cancer patients? Hell J Nucl Med 2017; 20: 41–45.
- Li L, Ma X. Study on specificity of colon carcinoma-associated serum markers and establishment of SVM prediction model. Saudi J Biol Sci 2017; 24: 644–648.
- Casciato D. Manual of Clinical Oncology. 5th ed. USA: Lippincott Williams & Wilkins, 2004.
- Lusikooy RE. Faktor Resiko Terjadinya Kanker Kolorektal di Indonesia. Universitas Hasanuddin, 2013.
- Palaghia M, Prelipcean C, Cotea E, et al. Metastatic Colorectal Cancer: Review of Diagnosis and Treatment Options. J Surg 2015; 10: 249–256.
- Malati T, Rajani Kumari G, Yadagiri B, et al. Utilitas diagnostik penanda tumor dalam identifikasi primer yang tidak diketahui. Ind J Clin Biochem 1996; 11: 77–80.
- Zhang S, Lin M, Zhang H. Diagnostic value of carcinoembryonic antigen and carcinoma antigen 19-9 for colorectal carcinoma. Int J Clin Exp Pathol 2015; 8: 9404–9409.
- Bast RCJ, Xu FJ, Yu YH, et al. CA 125: the past and the future. Int J Biol Markers 1998; 13: 179–187.
- Huang C-J, Jiang J-K, Chang S-C, et al. Serum CA125 concentration as a predictor of peritoneal dissemination of colorectal cancer in men and women. Medicine (Baltimore) 2016; 95: e5177.

15. Yang X, Chen C, Wang F, et al. Preoperative Serum Carcinoembryonic Antigen, Carbohydrate Antigen19-9 and Carbohydrate Antigen 125 as Prognostic Factors for Recurrence-free Survival in Colorectal Cancer. *Asian Pacific J Cancer Prev* 2011; 12: 1251–1256.
16. Yang X, Chen C, Wang F, et al. Preoperative Serum Carcinoembryonic Antigen, Carbohydrate Antigen19-9 and Carbohydrate Antigen 125 as Prognostic Factors for Recurrencefree Survival in Colorectal Cancer. *Asian Pacific J Cancer Prev* 2011; 12: 1251–1256.
17. Yang W, Luo Y, Hu S, Li Y, Liu Q. Value of combined detection of serum carcino-embryonic antigen, carbohydrate antigen 19-9 and cyclooxygenase-2 in the diagnosis of colorectal cancer. *Oncology Letters*. 2018. 16: 1551-1556

THE RELATIONSHIP BETWEEN TRIPLE TUMOR MARKER (CEA, CA 19-9, AND CA 125) AND COLORECTAL CANCER METASTASES AT MAKASSAR, INDONESIA

ORIGINALITY REPORT

14%

SIMILARITY INDEX

11%

INTERNET SOURCES

8%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	www.bibliomed.org Internet Source	4%
2	Submitted to Universitas Hasanuddin Student Paper	1%
3	Warsingih, Amir Fajar, Ibrahim Labeda, Julianus Aboyaman Uwuratuw, Prihantono, Muhammad Faruk. "Bilioenteric bypass stricture type II with hepatolithiasis: A case report", <i>Annals of Medicine and Surgery</i> , 2020 Publication	1%
4	repo.unand.ac.id Internet Source	1%
5	Kent T. Perry, Catherine T. Anthony, Tom Case, Mitchell S. Steiner. "Transforming growth factor beta as a clinical biomarker for prostate cancer", <i>Urology</i> , 1997 Publication	1%
6	dergipark.org.tr Internet Source	1%

7	ejm.org Internet Source	1 %
8	www.apocp.org Internet Source	1 %
9	www.anaisdedermatologia.org.br Internet Source	<1 %
10	www.annaliitalianidichirurgia.it Internet Source	<1 %
11	C. D. WEIMANN. "Spectrophotometric assessment of peritoneal fluid haemoglobin in colic horses: an aid to selecting medical vs. surgical treatment", Equine Veterinary Journal, 01/05/2010 Publication	<1 %
12	Nehal Samir, Marwa Matboli, Hanaa El - Tayeb, Ahmed El - Tawdi et al. "Competing endogenous RNA network crosstalk reveals novel molecular markers in colorectal cancer", Journal of Cellular Biochemistry, 2018 Publication	<1 %
13	Heijs, Bram, Stephanie Holst, Inge H. Briaire-de Bruijn et al. "Multimodal Mass Spectrometry Imaging of N-glycans and Proteins from the Same Tissue Section.", Analytical Chemistry Publication	<1 %

14

www.spandidos-publications.com

Internet Source

<1 %

15

Thangapandi Kalyani, Arumugam Sangili, Amalesh Nanda, Sengodu Prakash, Ajeet Kaushik, Saikat Kumar Jana. "Bio-nanocomposite based highly sensitive and label-free electrochemical immunosensor for endometriosis diagnostics application", *Bioelectrochemistry*, 2021

Publication

<1 %

16

synapse.koreamed.org

Internet Source

<1 %

17

www.nature.com

Internet Source

<1 %

18

Adnan Aydiner, Erkan Topuz, Rian Discli, Vildan Yasasever, Maktav Dincer, Koray Dincol, Nijad Bilge. "Serum Tumor Markers for Detection of Bone Metastasis in Breast Cancer Patients", *Acta Oncologica*, 2009

Publication

<1 %

19

Rongrong Ren, Guoneng Cai, Zhenzhong Yu, Yongyi Zeng, Dianping Tang. "Metal-Polydopamine Framework: An Innovative Signal-Generation Tag for Colorimetric Immunoassay", *Analytical Chemistry*, 2018

Publication

<1 %

real-d.mtak.hu

20

Internet Source

<1 %

21

Rizha Anshori Nasution, Andi Assadul Islam, Mochammad Hatta, Prihantono, Cahyono Kaelan, Jeni Poniman, Harakati Wangi. "Modification of the Marmarou model in developing countries", *Annals of Medicine and Surgery*, 2020
Publication

<1 %

22

eurjther.com
Internet Source

<1 %

23

www.researchsquare.com
Internet Source

<1 %

24

Cengiz, B.. "Effect of hormone replacement therapy on serum levels of tumor markers in healthy postmenopausal women", *Maturitas*, 20031210
Publication

<1 %

25

Kwon, O.E.. "Dimeric sesquiterpenoids isolated from *Chloranthus japonicus* inhibited the expression of cell adhesion molecules", *Journal of Ethnopharmacology*, 20060308
Publication

<1 %

26

Yufei Yang, Xuebing Yan, Xinxiang Li, Yanlei Ma, Ajay Goel. "Long non-coding RNAs in colorectal cancer: Novel oncogenic

<1 %

mechanisms and promising clinical applications", Cancer Letters, 2021

Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On