

Chronic hepatitis B in pregnant women: is hepatitis B surface antigen quantification useful for viral load prediction?

by Siti Maisuri Tadjuddin Chalid

Submission date: 25-Jun-2021 11:58AM (UTC+0700)

Submission ID: 1611887271

File name: 2015_Chronic_Hepatitis_B.pdf (793.81K)

Word count: 5660

Character count: 28495



Chronic hepatitis B in pregnant women: is hepatitis B surface antigen quantification useful for viral load prediction?



Masita Fujiko^{a,1}, Maisuri T. Chalid^{a,1}, Turyadi^b, Susan I. Ie^b, Maghfira^a, Syafri^a,
Ridha Wahyuni^a, Martono Roni^b, Ilhamjaya Patellongi^a, M. Nasrum Massi^a,
David H. Muljono^{a,b,c,*}

^a Faculty of Medicine, Hasanuddin University, Makassar, South Sulawesi, Indonesia

^b Eijkman Institute for Molecular Biology, Jl. Diponegoro 19 Jakarta Pusat 10430, DKI Jakarta, Indonesia

^c Sydney Medical School, University of Sydney, Sydney, New South Wales, Australia

ARTICLE INFO

Article history:

Received 27 July 2015

Received in revised form 20 October 2015

Accepted 4 November 2015

Corresponding Editor: Eskild Petersen, Aarhus, Denmark.

34 words:

HBsAg level

HBV DNA level

Pregnant women

HBV vertical transmission

Mother-to-child-transmission

SUMMARY

Background: New cases of hepatitis B virus (HBV) infection continue to occur worldwide. Most of these are due to mother-to-child transmission (MTCT), with maternal viraemia as the most important contributing factor. The hepatitis B surface antigen (HBsAg) level, which correlates positively with viral load, has been used for treatment monitoring in chronic hepatitis B. This study evaluated the usefulness of quantitative HBsAg for viral load prediction in HBsAg-positive pregnant women.

Methods: A total of 943 pregnant women in Makassar, Indonesia were screened for HBsAg. Sixty-four women were HBsAg-positive and investigated. HBsAg level and hepatitis B e antigen (HBeAg)/hepatitis B e antibody (anti-HBe) status were determined serologically. Viral load was measured by real-time PCR. HBV DNA was sequenced and analysed for identification of genotype and basal core promoter (BCP)/precore (PC) mutations.

Results: Of 64 subjects, 54 (84.4%) were HBeAg-positive and 10 (15.6%) were HBeAg-negative. HBsAg and HBV DNA levels were significantly higher in the HBeAg-positive group ($p < 0.001$). HBsAg and HBV DNA levels were positively correlated in the HBeAg-positive group ($r = 0.359$; $p = 0.02$), but not in the HBeAg-negative group ($r = 0.194$; $p = 0.168$). Low HBsAg levels ($< 3.0 \log_{10}$ IU/ml) corresponded with HBV DNA levels $< 6.0 \log_{10}$ IU/ml ($r = 0.404$; $p = 0.001$), a recognized threshold for MTCT. Genotype C was more prevalent than genotype B, but not associated with HBsAg level, viral load, or HBeAg status. Two-thirds of HBeAg-negative subjects with high HBV DNA levels harboured BCP (A1762T/G1764A) and/or PC (G1896A) variants.

Conclusions: HBsAg levels provide a good viral load predictor in HBeAg-positive but not HBeAg-negative pregnant women. The HBeAg-negative group had a frequent occurrence of BCP/PC variants, which may have contributed to the lack of correlation observed. Samples with a low HBsAg level, which is associated with a low risk of MTCT, do not require HBV DNA measurement.

© 2015 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

More than 240 million people worldwide are chronically infected with the hepatitis B virus (HBV), and about 780 000 die

from hepatitis B annually. In highly endemic areas of Asia, the Pacific, and Sub-Saharan Africa, most HBV infection occurs perinatally or during early childhood. This is associated with a high rate of persistent infection and increased risk of morbidity and mortality from cirrhosis and hepatocellular carcinoma later in life.¹

Efforts in the prevention of hepatitis B have focused on the immunization of infants implemented in 183 World Health Organization (WHO) member states. As of 2012, 94 member states including Indonesia had introduced the hepatitis B birth

* Corresponding author. Tel.: +62 21 3148695; fax: +62 21 3147982.
E-mail addresses: davidhm@eijkman.go.id, dhmuljono@gmail.com

(D.H. Muljono).

¹ Authors Masita Fujiko and Maisuri T. Chalid contributed equally to this manuscript.

<http://dx.doi.org/10.1016/j.ijid.2015.11.002>

© 2015 The Authors. Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

dose immunization.¹ This program has markedly decreased the disease burden, carrier rate, and HBV-related morbidity and mortality.² However, 50 million new cases of HBV infection continue to be diagnosed annually, with the highest incidence due to mother-to-child transmission (MTCT). Despite administration of hepatitis B immunization (active) or active hepatitis B immune globulin (HBIG) at birth, at least 10% of infants born to HBV-carrying mothers still suffer HBV infection.³ Several factors such as maternal serum HBV DNA level, hepatitis B e antigen (HBeAg) status, HBV S gene variation, mode of delivery, and neonatal immune deficiency have been related to MTCT. Of these factors, maternal HBV DNA level has been identified as the most relevant.⁴ Practice guidelines from major professional associations address the decision for antiviral treatment in pregnant women based on the HBV DNA threshold.^{4–6}

Assays for HBV DNA quantification with high sensitivity and specificity are currently available. However, the routine application of these methods for screening pregnant women is hampered by cost and limited resources. In recent years, the hepatitis B surface antigen (HBsAg) level, which correlates positively with viral load, has been used as a biomarker to predict disease status and to monitor the treatment response in chronic hepatitis B (CHB).^{7,8} The HBsAg level correlates with covalently closed circular DNA (cccDNA) in hepatocytes and can be considered a surrogate marker of infected cells.⁹ One attraction of the use of the HBsAg level is that the assay is less costly, is suitable for high-throughput screening, and its platforms are commonly used in many laboratories. However, some studies reported that HBsAg quantification correlates poorly with the HBV DNA level. HBsAg and HBV DNA levels may vary during different phases of CHB and the correlation is associated with the HBeAg status of the patient.^{10–12} Certain HBV variants with basal core promoter (BCP) A1762T/G1764A or precore (PC) G1896A mutations could influence the synthesis of HBeAg.¹³ The presence of these variants needs to be taken into account since they often occur in endemic areas and may be associated with certain HBV genotypes that differ among geographical regions.^{14,15}

This study aimed to evaluate the usefulness of quantitative HBsAg as a viral load predictor in pregnant women with CHB in Makassar, Indonesia, and to analyse the association of the HBsAg level and viral load with the HBeAg status, as well as the molecular characteristics of HBV variants defective for HBeAg production.

2. Materials and methods

2.1. Study population

This cross-sectional study was carried out from January to July 2014 in the antenatal care units of Wahidin Sudirohusodo Hospital, Hasanuddin University Hospital, Fatimah Mother and Child Hospital, Pertiwi Mother and Child Hospital, Labuang Baji Hospital, and Ibnu Sina Hospital, as well as several maternity clinics, in Makassar, South Sulawesi. A total of 943 pregnant women were screened for HBV infection; 64 of them were HBsAg-positive and considered eligible for enrolment. For inclusion it was required that the woman had been HBsAg-positive for >6 months without prior antiviral therapy.¹⁶ Subjects co-infected with hepatitis A virus, hepatitis C virus, or human immunodeficiency virus, as well as those with evidence of liver diseases, were excluded. This study was approved by the Ethics Committee of the Faculty of Medicine, Hasanuddin University, Makassar, Indonesia. Written informed consent was obtained from each patient.

2.2. Serological examination

HBsAg status was determined by VIDIA HBsAg immunoassay (bioMérieux SA, Marcy l'Etoile, France). HBeAg and hepatitis B e

antibody (anti-HBe) were tested using Monalisa HBeAg-Ab PLUS immunoassay (Bio-Rad, Marnes-la-Coquette, France). HBsAg quantification was done using Elecsys HBsAg Quant II (Roche Diagnostics, Indianapolis, USA) on a Roche Cobas e411 Immunoanalyzer following the manufacturer's protocol.

2.3. HBV DNA detection and analysis

The HBV DNA level was determined from 500 µl of serum by quantitative real-time PCR (CobasTaqman HBV Test; Roche Diagnostics, Indianapolis, USA) with a range of linearity between 6 and 1.1×10^8 IU/ml. HBV DNA for molecular analysis was obtained by extracting the DNA from 140 µl of serum using the QIAamp DNA Mini Kit (Qiagen, Valencia, CA, USA) and amplification by nested PCR using primers S2-1/S1-2 for the first-round and S88/S2-2 for the second-round (Supplementary Material, Table S1).¹² Amplicons were purified using a PCR purification column (Qiagen, Valencia, CA, USA) and subjected to direct sequencing on a DNA sequence analyzer ABI 3130xl (Applied Biosystems, Carlsbad, CA, USA).

The HBV genotype was determined by phylogenetic analysis based on the 226-nucleotide sequences of the S gene compared with 70 reference sequences of known genotypes (A–H) retrieved from GenBank, using Phylip 3.68 software with the Kimura 2-parameter model, neighbour-joining algorithm, and 1000 bootstrapping.¹²

2.4. Identification of BCP and PC mutations

Amplification of BCP and PC regions was done by nested PCR using primers PC1/PC2 for the first-round and S012/S013 for the second-round (Supplementary Material, Table S1). Amplicons were purified and sequenced as described previously.¹² The sequences were aligned with reference sequence M54923 retrieved from GenBank.

2.5. Statistical analysis

The baseline data were summarized descriptively. Continuous and categorical variables were compared between groups using the Mann-Whitney test and Chi-square/Fisher's exact test, respectively. Pearson's correlation coefficient was used to describe the correlation between two continuous, normally distributed variables. Spearman's correlation was used for categorical variables or continuous variables that were not normally distributed.^{10,12,13} Statistical analyses were performed using IBM-SPSS v. 21 software (IBM Corp., Armonk, NY, USA). All statistical significance values were assessed at $p < 0.05$.

3. Results

3.1. Characteristics of study subjects

Among 943 pregnant women attending several antenatal clinics in Makassar, 44 (6.8%) were HBsAg-positive. Of these women, 12 (18.8%) were HBeAg-positive and 52 (81.3%) were HBeAg-negative. HBV DNA levels were significantly higher in the HBeAg-positive group (median 7.43 log₁₀ IU/ml) than in the HBeAg-negative group (median 1.55 log₁₀ IU/ml) ($p < 0.001$). Similarly, HBsAg levels were significantly higher in the HBeAg-positive group (median 4.21 log₁₀ IU/ml) than in the HBeAg-negative group (median 2.91 log₁₀ IU/ml) ($p < 0.001$). Age, alanine aminotransferase (ALT) levels, and HBeAg genotype distribution were comparable in the two groups (Table 1 and Supplementary Material Table S2).

Table 1
Baseline characteristics of HBsAg-positive pregnant women^a

Parameter	Overall (n = 64)	HBeAg-positive (n = 12)	HBeAg-negative (n = 52)	p-Value ^b
Age (years)	29 (18–42)	28.5 (22–42)	30 (18–41)	0.564
ALT (IU/l)	24.5 (9–129)	27 (16–129)	24.5 (9–88)	0.129
HBV DNA (\log_{10} IU/ml)	1.71 (0.78–8.05)	7.43 (1.54–8.05)	1.54 (0.78–6.48)	<0.001
HBsAg (\log_{10} IU/ml)	3.03 (0.70–4.11)	4.21 (3.25–4.91)	2.91 (0.70–4.11)	<0.001
Ratio HBsAg/HBV DNA (\log_{10} IU/ml)	1.25 (0.40–5.30)	0.61 (0.56–0.7556)	1.43 (0.90–2.78)	<0.001
Genotype (n = 47)				
B	8 (17.0%)	2 (22.2%)	6 (15.8%)	0.889
C	39 (83.0%)	7 (77.8%)	32 (84.2%)	

HBsAg, Hepatitis B e antigen; ALT, alanine aminotransferase; HBV, hepatitis B virus; HBsAg, hepatitis B surface antigen.

^a Results are presented as the median (minimum–maximum), or number (percentage) of detected samples.

^b Comparison between HBeAg-positive and HBeAg-negative groups (Mann–Whitney U-test or Chi-square test).

3.2. Distribution of serum HBV DNA levels among HBsAg-positive pregnant women according to HBeAg status

HBV DNA levels were categorized into $<3.0 \log_{10}$ IU/ml (close to $3.3 \log_{10}$ IU/ml or 2000 IU/ml, which is the threshold to define inactive carrier state of CHB), $3.0\text{--}6.0 \log_{10}$ IU/ml, and $\geq 6.0 \log_{10}$ IU/ml (a level associated with HBV immunoprophylaxis failure).¹⁸ The proportions of pregnant women with HBV DNA levels $<3.0 \log_{10}$ IU/ml, $3.0\text{--}6.0 \log_{10}$ IU/ml, and $\geq 6.0 \log_{10}$ IU/ml in the HBeAg-positive group were 9% (1/12), 25% (3/12), and 67% (8/12), respectively; in the HBeAg-negative group, the proportions were 83% (43/52), 13% (7/52), and 4% (2/52), respectively (Figure 1A).

3.3. Distribution of serum HBsAg levels among the HBsAg-positive pregnant women according to HBeAg status

HBsAg levels were categorized into $<3.0 \log_{10}$ IU/ml (a level associated with a lower risk of CHB outcomes), $3.0\text{--}4.0 \log_{10}$ IU/ml, and $\geq 4.0 \log_{10}$ IU/ml.⁵ The proportions of subjects with HBsAg levels $<3.0 \log_{10}$ IU/ml, $3.0\text{--}4.0 \log_{10}$ IU/ml, and $\geq 4.0 \log_{10}$ IU/ml were 0% (0/12), 42% (5/12), and 58% (7/12), respectively, in the HBeAg-positive group, and 58% (30/52), 35% (18/52), and 8% (4/52), respectively, in the HBeAg-negative group (Figure 1B).

3.4. Correlation between HBsAg and HBV DNA levels

In all 64 HBsAg-positive subjects, serum HBsAg and HBV DNA levels showed a significantly moderate correlation ($r = 0.513$; $p < 0.001$) (Figure 2A). When analysed separately according to HBeAg status, there was a strong correlation between HBV DNA and HBsAg levels in the HBeAg-positive group ($r = 0.659$; $p = 0.02$) (Figure 2B), but no correlation was observed in the HBeAg-negative group ($r = 0.194$; $p = 0.168$) (Figure 2C).¹⁹

There were four subjects (M150, M414, M415, and M818) with high HBsAg levels ($>4.0 \log_{10}$ IU/ml) but low HBV DNA levels ($<3.0 \log_{10}$ IU/ml). In contrast, there were four subjects (M167, M173, M258, and M810) with low HBsAg levels ($<3.0 \log_{10}$ IU/ml) but who had moderate viraemia ($3.0\text{--}6.0 \log_{10}$ IU/ml). However, in most cases, low levels of HBsAg were associated with low levels of HBV DNA; HBsAg levels $<3.0 \log_{10}$ IU/ml were significantly correlated to HBV DNA levels $<3.0 \log_{10}$ IU/ml ($r = 0.363$; $p = 0.003$) and to HBV DNA levels $<6.0 \log_{10}$ IU/ml ($r = 0.404$; $p = 0.001$). No subjects with HBsAg levels $<3.0 \log_{10}$ IU/ml had HBV DNA $\geq 6.0 \log_{10}$ IU/ml (Figure 3).

3.5. HBsAg and HBV DNA levels in genotype B and C

HBV genotype was successfully determined in 47 pregnant women based on the S gene sequences (sequencing was not possible in the other subjects because of insufficient HBV DNA

content). The sequences generated have been deposited in the GenBank database (accession numbers **KP241791–KP241837**). Median HBsAg and HBV DNA levels were not significantly different in each genotype. Genotype C was more prevalent than genotype B (83% vs. 17%), but the proportions were comparable between the HBeAg-positive and HBeAg-negative groups (Figure 4).

3.6. Subgroup analysis for BCP and PC mutations in HBeAg-negative pregnant women

Sequencing of the BCP/PC region of the HBV genome was performed successfully in 12 HBeAg-negative subjects (sequencing was not possible in the other subjects because of low HBV DNA content). The sequences generated have been deposited in GenBank (accession numbers **KP241838–KP241844** and **KP241846–KP241850**). Four subjects had the wild-type HBV DNA sequence at both the BCP and PC sites, four had the BCP mutation alone, and two had the PC mutation alone. Concurrent BCP and PC mutations were detected in two subjects. All subjects with the BCP mutation had HBV genotype C, while the PC mutation was detected in two subjects with genotype B and one subject with genotype C (Table 2).

4. Discussion

This study represents one of few reports of HBV infection in pregnant women from Indonesia. Of 943 pregnant women attending several antenatal clinics in Makassar, 64 (6.8%) were HBsAg-positive. This figure is higher than that reported recently from Jakarta (2.2%),²⁰ and other places in Indonesia reported around 1985 (4.7% in West Java, 1.9% in Bali, 3.4% in Mataram).^{21,22} The wide variation in HBV infection rates may be associated with the general HBsAg prevalence in Indonesia (3.4–19.5%), geographical variation, and differences in cultural practices, as well as the methods used to detect HBV infection.^{23,24} This fact is of concern, because it occurs in pregnant women who tend to be in the immune-tolerant phase of CHB with normal physical/laboratory examinations and high-level viraemia, but unaware of their HBsAg-positive status.

Varying thresholds of maternal HBV DNA have been discussed in association with MTCT and immunoprophylaxis failure. Wiseman reported that immunoprophylaxis failure occurred in infants when the maternal viral load was $\geq 8 \log_{10}$ copies/ml ($>1.7 \times 10^7$ IU/ml).²⁵ Zou et al. showed that the immunoprophylaxis failure increased with higher levels of maternal HBV DNA.¹⁸ When the mothers' HBV DNA levels were stratified to $5\text{--}6.99$, $7\text{--}7.99$, and $\geq 8 \log_{10}$ copies/ml, the corresponding rates of immunoprophylaxis failure were 0%, 3.24%, 6.7%, and 7.6%, respectively, and it was concluded that an antenatal HBV DNA level $>6 \log_{10}$ copies/ml ($>200,000$ IU/ml) was the most important predictor of MTCT.¹⁸




Figure 1. Distribution of (A) HBV DNA and (B) HBsAg levels among pregnant women according to HBeAg status: (i) among the HBeAg-positive group ($n = 12$) and (ii) among the HBeAg-negative group ($n = 52$).

The European Association for the Study of the Liver (EASL) and the Asian Pacific Association for the Study of the Liver (APASL) guidelines recommend treating pregnant women when HBV DNA levels are $>2 \times 10^6$ IU/ml in the third trimester for the prevention of MTCT.^{5,6} In the present study, 15.6% (10/64) of all subjects had HBV DNA levels $>6.0 \log_{10}$ IU/ml, distributed in 67% (8/12) of the HBeAg-positive group and 4% (2/52) of the HBeAg-negative group. The fact that the subjects had a skewed distribution toward the higher levels of HBV DNA should be regarded as important, as this shows a higher possibility of MTCT.

There are few studies on the potential applications of quantitative HBsAg in the management of hepatitis B during pregnancy.^{16,26,46} In the best of the authors' knowledge, this is the

first study to evaluate the relationship between HBsAg and HBV DNA levels with regard to HBeAg status in pregnant women with CHB in Indonesia. This study revealed that among all 64 HBsAg-positive pregnant women, the HBsAg level was correlated with the HBV DNA level regardless of age and viral genotype. When stratified based on HBeAg status, the correlation was strong in HBeAg-positive pregnant women but missing in the HBeAg-negative women. A possible explanation for this finding is that HBsAg synthesis has a pathway distinct from HBV DNA synthesis and under the influence of different immune-control mechanisms.^{27,28} HBsAg is present as a component of HBV virions but also as subviral particles, which exceed the number of virions.⁹ Pregnant women with an HBeAg-positive status could be in the




Figure 2. Correlation of HBsAg and HBV DNA levels in pregnant women according to HBeAg status: (A) overall correlation in all pregnant women ($n = 64$); (B) correlation of HBsAg and HBV DNA levels in HBeAg-positive women ($n = 12$); (C) correlation of HBsAg and HBV DNA levels in HBeAg-negative women ($n = 52$). Both the HBV DNA and HBsAg levels are calculated in \log_{10} IU/ml. The correlation is regarded very weak for $r = 0-0.19$, weak for $r = 0.20-0.39$, moderate for $r = 0.40-0.59$, strong for $r = 0.60-0.79$, and very strong for $r = 0.80-1$.




Figure 3. Distribution of HBV DNA levels among pregnant women according to HBsAg levels. In most cases, low levels of HBsAg were associated with low levels of HBV DNA; HBsAg levels $<3.0 \log_{10}$ IU/ml were significantly correlated to HBV DNA levels $<3.0 \log_{10}$ IU/ml ($r = 0.363$; $p = 0.003$) and to HBV DNA levels $<6.0 \log_{10}$ IU/ml ($r = 0.404$; $p = 0.001$). No subjects with HBsAg levels $<3.0 \log_{10}$ IU/ml had HBV DNA $\geq 6.0 \log_{10}$ IU/ml.

immune-tolerant phase where HBV virions and their antigens are minimally subjected to the host immune reaction, while those with an HBeAg-negative status could be in the low-replicative phase of CHB where the number of virions has decreased as a result of successful immune control. Therefore, the reduction in HBsAg levels was not proportional to that of HBV DNA levels. The ratio of HBsAg/HBV DNA reflects the association between HBsAg production and HBV replication; this was significantly higher in HBeAg-negative subjects than in the HBeAg-positive group (median 1.43 vs. 0.61 \log_{10} IU/ml, respectively).

Four pregnant women had high HBsAg levels with low HBV DNA. As explained, this could be due to the larger excess of HBsAg

Table 2 BCP/PC mutations, HBV DNA and HBsAg levels, and HBV genotype in 12 HBeAg-negative pregnant women

Subject code	BCP mutation A1762T/G1764A	PC mutation G1896A	HBV DNA (\log_{10} IU/ml)	HBsAg (\log_{10} IU/ml)	Genotype
M177	+	+	6.48	3.46	C
M384	+	-	6.08	3.04	C
M167	-	+	4.55	2.69	B
M810	-	-	3.50	2.70	C
M218	+	-	3.29	3.56	C
M336	+	+	2.70	3.74	C
M150	-	-	2.62	4.00	C
M898	-	+	2.08	3.04	B
M253	-	-	1.36	3.05	C
M818	+	-	1.34	4.11	C
M19A	+	-	0.78	2.51	C
M212	+	-	0.78	3.55	C

BCP, basal core promoter; PC, precore; HBV, hepatitis B virus; HBsAg, hepatitis B surface antigen; HBeAg, hepatitis B e antigen.

over the number of virions. This implies that a high level of HBsAg cannot be used to predict the HBV DNA level. The serum HBsAg level should thus be used together with, but not as a substitute for, HBV DNA. Four other subjects had low HBsAg levels but moderate viraemia ($>4 \log_{10}$ IU/ml). Possible explanations for the decreased detection of HBsAg include (1) differences in analytical sensitivity and specificity in HBsAg detection of viruses of different genotypes; (2) mutations in the pre-S/S gene that cause HBsAg detection failure; (3) treatment-associated mutations that cause derangement of the P gene with subsequent alteration of the overlapping S gene; or (4) the concomitant presence of hepatitis B surface antibodies (anti-HBs) leading to the formation of immune complexes poorly displaced by HBsAg-capture antibodies. In the majority of subjects, however, low levels of HBsAg ($<3.0 \log_{10}$ IU/ml) correlated significantly with low levels of HBV DNA (<6.0




Figure 4. Distribution of serum HBV DNA (A) and HBsAg (B) levels in subjects with HBV genotype B ($n = 8$) and HBV genotype C ($n = 39$). HBV DNA and HBsAg levels were comparable in the two genotypes. Median values (max, min; \log_{10} IU/ml) with 95% confidence intervals are shown.

\log_{10} IU/ml) and an attendant lower risk of MTCT.^{6,18} This result suggests that HBV DNA quantification may not be necessary for pregnant mothers with low HBsAg levels.

Another important finding from this study was the presence of HBeAg-negative pregnant women with high viraemia. These women, with no evidence of liver disease, were apparently in the inactive carriers of CHB. However, some inactive carriers may have high HBV DNA levels accompanied by persistently normal ALT levels.⁵ Studies have documented that certain HBV variants with nucleotide substitutions in the PC and/or BCP regions could abolish or down-regulate HBeAg production. These variants may replicate rapidly in HBeAg-negative CHB, where HBsAg and DNA levels are preserved.¹³ Analysis for the presence of BCP (A1762T/G1764A) and PC (G1896) mutations was performed on HBeAg-negative subjects who had HBV DNA levels >2000 IU/ml. A substantial portion of isolates analysed, particularly from subjects with HBV DNA levels >6.0 \log_{10} IU/ml, had BCP and PC mutations either alone or in combination. This finding is important because the emergence of these variants frequently occurs in regions with HBV endemicity.^{14,15} Notably, all BCP mutants identified had genotype C, which was prevalent among the subjects studied. It has been recognized that HBV genotype C and BCP mutations are independent risk factors for progression to severe liver disease.^{5,32}

Important limitations of the present study include the relatively small number of pregnant women with chronic HBV infection. Wider clinical and community-based studies in different areas of the Indonesian archipelago will be necessary to estimate the true national burden of HBV infection in pregnant women. Also, the cross-sectional design of the study could not represent the fluctuating profile of CHB. Serial measurements of ALT levels are necessary to distinguish the true inactive carriers from active HBeAg-negative individuals. However, most studies have reported that ALT levels are lower during pregnancy and viral load is more likely to increase due to the natural immune suppression processes linked to pregnancy.³³

In conclusion, this study confirms that serum HBsAg level may be used as a predictor of the serum HBV DNA level in HBeAg-positive pregnant women, but not in HBeAg-negative pregnant women. The measurement of HBV DNA is not necessary if the level of HBsAg is low, since the probability of detecting a high viral load is low. An important role of BCP/PC variants influencing HBeAg status independent of viral load was also identified, providing a cautionary note to the interpretation of negative results of HBeAg testing when classifying HBV-infected individuals. These results offer the promise of practical guidance in using quantitative HBsAg as a tool to better manage HBV-infected pregnant women. Follow-up studies are needed to assess the impacts of the maternal and biological factors discussed in this study on HBV carriage and immunoprophylaxis failure in the infants born to these HBsAg-positive mothers.

Acknowledgements

The authors would like to convey special thanks to the pregnant women who participated in this study. The authors would also like to express their gratitude to the Eijkman Institute for Molecular Biology in Jakarta for the molecular work support, the Hasanuddin University Medical Research Center (HUM-RC) for laboratory facilities, several maternity clinics in Makassar, and the Wahidin Sudirohusodo Teaching Hospital, as well as the Academic Health Center of Hasanuddin University with their staff for the recruitment of study subjects. Special thanks are given to Dr Kevin Baird from the Eijkman-Oxford Clinical Research Unit in Jakarta for proofreading this manuscript.

Funding: This study was supported by an Operational Support Grant for State University (BOPTN) from the Directorate General of Higher Education, Indonesian Ministry of Education (grant number 2058/UN4.20/PL.09/2013). PT Roche Indonesia donated the reagents used for HBsAg quantification.

Conflict of interest: No conflict of interest to declare.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ijid.2015.11.002>.

References

- World Health Organization. Hepatitis B Fact Sheet No 204. Updated July 2015. Geneva: WHO; 2015. Available at: <http://www.who.int/mediacentre/factsheets/fs204/en/> (accessed August 8, 2015).
- Romano L, Paladini S, Van Damme P, Zanetti AR. The worldwide impact of vaccination on the control and protection of viral hepatitis B. *Dig Liver Dis* 2011;43(Suppl 1):S2–7.
- Patton H, Tran TT. Management of hepatitis B during pregnancy. *Nat Rev Gastroenterol Hepatol* 2014;11:402–9.
- Song YM, Sung J, Yang S, Choe YH, Chang YS, Park WS. Factors associated with immunoprophylaxis failure against vertical transmission of hepatitis B virus. *Eur J Pediatr* 2007;166:813–8.
- European Association for the Study of the Liver. EASL clinical practice guidelines: management of chronic hepatitis B virus infection. *J Hepatol* 2012;57:167–85.
- Liaw YF, Kao JH, Piratvisuth T, Chan HL, Chien RN, Liu CJ, et al. Asian-Pacific consensus statement on the management of chronic hepatitis B: a 2012 update. *Hepatol Int* 2012;6:531–61.
- Honer Zu Siederdisen C, Cornberg M. The role of HBsAg levels in the current management of chronic HBV infection. *Ann Gastroenterol* 2014;27:105–12.
- Su TH, Hsu CS, Chen CL, Liu CH, Huang YW, Tseng TC, et al. Serum hepatitis B surface antigen concentration correlates with HBV DNA level in patients with chronic hepatitis B. *Antivir Ther* 2010;15:1133–9.
- Brunetto MR. A new role for an old marker, HBsAg. *J Hepatol* 2010;52:475–7.
- Jaroszewicz J, Calle Serrano B, Wursthorn K, Detering K, Schlie J, Raupach R, et al. Hepatitis B surface antigen (HBsAg) levels in the natural history of hepatitis B virus (HBV)-infection: a European perspective. *J Hepatol* 2010;52:514–22.
- Nguyen T, Thompson AJ, Bowden S, Croagh C, Bell S, Desmond PV, et al. Hepatitis B surface antigen levels during the natural history of chronic hepatitis B: a perspective on Asia. *J Hepatol* 2010;52:508–13.
- Turyadi, Thedja MD, Ie SI, Harahap AR, El-Khobar KE, Roni M, et al. HBsAg, HBeAg and HBV DNA level changes and precore/basal core promoter mutations in the natural history of chronic hepatitis B in Indonesian patients. *Hepatol Int* 2013;7:969–98.
- Thompson AJ, Nguyen T, Iser D, Ayres A, Jackson K, Littlejohn M, et al. Serum hepatitis B surface antigen and hepatitis B e antigen titers: disease phase influences correlation with viral load and intrahepatic hepatitis B virus markers. *Hepatology* 2010;51:1933–44.
- Utama A, Purwantomo S, Siburian MD, Dhenni R, Gani RA, Hasan I, et al. Hepatitis B virus subgenotypes and basal core promoter mutations in Indonesia. *World J Gastroenterol* 2009;15:4028–36.
- Yuen MF, Sablon E, Tanaka Y, Kato T, Mizokami M, Doutreligne J, et al. Epidemiological study of hepatitis B virus genotypes, core promoter and precore mutations of chronic hepatitis B infection in Hong Kong. *J Hepatol* 2004;41:119–25.
- Sun KX, Li J, Zhu FC, Liu JX, Li RC, Zhai XJ, et al. A predictive value of quantitative HBsAg for serum HBV DNA level among HBeAg-positive pregnant women. *Vaccine* 2012;30:5335–40.
- Chen CJ, Yang HI, Su J, Jen CL, You SL, Lu SN, et al. Risk of hepatocellular carcinoma across a biological gradient of serum hepatitis B virus DNA level. *JAMA* 2006;295:65–73.
- Zou H, Chen Y, Duan Z, Zhang H, Pan C. Virologic factors associated with failure to passive-active immunoprophylaxis in infants born to HBsAg-positive mothers. *J Viral Hepat* 2012;19:e18–25.
- Swinscow TD, Campbell MJ. Statistics at square one, 10th ed., London: BMJ Books; 2002.
- Gunardi H, Zaimi LF, Soedjatmiko, Turyadi, Harahap AR, Muljono DH. Current prevalence of hepatitis B infection among parturient women in Jakarta, Indonesia. *Acta Med Indones* 2014;46:3–9.
- Reniers J, Vranckx R, Ngantung W, Sugita E, Meheus A. Prevalence and determinants of hepatitis B virus markers in pregnant women in West Java, Indonesia. *J Trop Med Hyg* 1987;90:249–53.
- Surya IG, Kornia K, Suwardewa TG, Mulyanto, Tsuda F, Mishiro S. Serological markers of hepatitis B, C, and E viruses and human immunodeficiency virus type-1 infections in pregnant women in Bali, Indonesia. *J Med Virol* 2005;75:499–503.

23. Khan M, Dong JJ, Acharya SK, Dhagwahdorj Y, Abbas Z, Jafri SMW, et al. Hepatology issues in Asia: perspectives from regional leaders. *J Gastroenterol Hepatol* 2004;**19**:S419–30.
24. Liaw YF, Brunetto MR, Hadziyannis S. The natural history of chronic HBV infection and geographical differences. *Antivir Ther* 2010;**15**(Suppl 3):25–33.
25. Wiseman E, Fraser MA, Holden S, Glass A, Kidson BL, Heron LG, et al. Perinatal transmission of hepatitis B virus: an Australian experience. *Med J Aust* 2009;**190**:489–92.
26. Belopolskaya M, Avfukin V, Firsov S, Yakovlev A. Relationship between viral load and HBsAg level in pregnant women with chronic hepatitis B virus infection. *Proceeding of EASL Special Conference "Optimal Management of Hepatitis B Virus Infection"*. 2014.
27. Kim YJ, Cho HC, Choi MS, Lee JH, Koh KC, Yoo BC, et al. The change of the quantitative HBsAg level during the natural course of chronic hepatitis B. *Liver Int* 2011;**31**:817–23.
28. Liaw YF. Clinical utility of hepatitis B surface antigen quantitation in patients with chronic hepatitis B: a review. *Hepatology* 2011;**53**:2121–9.
29. Chan HL, Thompson A, Martinot-Peignoux M, Piratvisuth T, Cornberg M, Brunetto MR, et al. Hepatitis B surface antigen quantification: why and how to use it in 2011—a core group report. *J Hepatol* 2011;**55**:1121–31.
30. Candotti D, Allain JP. Transfusion-transmitted hepatitis B virus infection. *J Hepatol* 2009;**51**:798–809.
31. Pollicino T, Amaddeo G, Restuccia A, Raffa G, Alibrandi A, Cutroneo G, et al. Impact of hepatitis B virus (HBV) preS/S genomic variability on HBV surface antigen and HBV DNA serum levels. *Hepatology* 2012;**56**:434–43.
32. Tong MJ, Blatt LM, Kao JH, Cheng JT, Corey WG. Precore/basal core promoter mutants and hepatitis B viral DNA levels as predictors for liver deaths and hepatocellular carcinoma. *World J Gastroenterol* 2006;**12**:6620–6.
33. Jonas MM. Hepatitis B and pregnancy: an underestimated issue. *Liver Int* 2009;**29**(Suppl 1):133–9.

Chronic hepatitis B in pregnant women: is hepatitis B surface antigen quantification useful for viral load prediction?

ORIGINALITY REPORT

24%

SIMILARITY INDEX

14%

INTERNET SOURCES

17%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1

www.birpublications.org

Internet Source

1%

2

Zhirong Liu, Feng Liu, Lei Wang, Youde Liu, Meng Zhang, Tao Li. "Clinical characteristics and outcomes of patients with recurrent chronic hepatitis B after nucleos(t)ide analog withdrawal with stringent cessation criteria: A prospective cohort study", *Hepatology Research*, 2017

Publication

1%

3

En-Qiang Chen, Meng-Lan Wang, Ya-Chao Tao, Dong-Bo Wu, Juan Liao, Min He, Hong Tang. "Serum HBcrAg is better than HBV RNA and HBsAg in reflecting intrahepatic covalently closed circular DNA", *Journal of Viral Hepatitis*, 2019

Publication

<1%

4

Lei Zhang, Xien Gui, Jingyi Fan, Bo Wang et al. "Breast feeding and immunoprophylaxis efficacy of mother-to-child transmission of

<1%

hepatitis B virus", The Journal of Maternal-Fetal & Neonatal Medicine, 2013

Publication

5	Submitted to October University for Modern Sciences and Arts (MSA) Student Paper	<1 %
6	academic.hep.com.cn Internet Source	<1 %
7	jbiomedsci.biomedcentral.com Internet Source	<1 %
8	virologyj.biomedcentral.com Internet Source	<1 %
9	www.intmedpress.com Internet Source	<1 %
10	www.medultrason.ro Internet Source	<1 %
11	Myron John Tong. "A comparison of hepatitis B viral markers of patients in different clinical stages of chronic infection", Hepatology International, 04/07/2010 Publication	<1 %
12	jco.ascopubs.org Internet Source	<1 %
13	pure.uva.nl Internet Source	<1 %

14

Mulyanto, Sulaiman Ngongu Depamede, Kiely Surayah, Andreas Andhry Heru Tjahyono et al. "Identification and characterization of novel hepatitis B virus subgenotype C10 in Nusa Tenggara, Indonesia", Archives of Virology, 2010

Publication

<1 %

15

Qinxu Xie, Xiangyang Hu, Yafei Zhang, Xiaoping Jiang, Xu Li, Jiabin Li. "Decreasing hepatitis B viral load is associated with a risk of significant liver fibrosis in hepatitis B e antigen positive chronic hepatitis B", Journal of Medical Virology, 2014

Publication

<1 %

16

Vutien, P., H. N. Trinh, K. Nguyen, R. T. Garcia, H. A. Nguyen, B. S. Levitt, L. Nguyen, N. B. Ha, A. Ahmed, T. Daugherty, G. Garcia, and M. H. Nguyen. "Precore and basal core promoter mutations in Asian American patients with hepatitis B e antigen-positive chronic hepatitis B", Alimentary Pharmacology & Therapeutics, 2013.

Publication

<1 %

17

natap.org

Internet Source

<1 %

18

www.scribd.com

Internet Source

<1 %

19	ro.ecu.edu.au Internet Source	<1 %
20	tropmedhealth.biomedcentral.com Internet Source	<1 %
21	CHIEN-YI CHEN. "HEPATITIS B AND PREGNANCY, THE SCIENTIFIC BASIS FOR PERINATAL PREVENTION", Fetal and Maternal Medicine Review, 05/2010 Publication	<1 %
22	Maurizia Rossana Brunetto, Francesco Moriconi, Ferruccio Bonino, George K. K. Lau et al. "Hepatitis B virus surface antigen levels: A guide to sustained response to peginterferon alfa-2a in HBeAg-negative chronic hepatitis B", Hepatology, 2009 Publication	<1 %
23	hepatitiscresearchandnewsupdates.blogspot.co.uk Internet Source	<1 %
24	med.unhas.ac.id Internet Source	<1 %
25	pocketdentistry.com Internet Source	<1 %
26	www.researchsquare.com Internet Source	<1 %
27	www.terapeutica.ro Internet Source	<1 %

28 Myoung Hee Lee, Da Mi Lee, Soon Sun Kim, Jae Youn Cheong, Sung Won Cho. "Correlation of serum hepatitis B surface antigen level with response to entecavir in naïve patients with chronic hepatitis B", Journal of Medical Virology, 2011
Publication <1 %

29 www.labome.org
Internet Source <1 %

30 Submitted to Higher Education Commission Pakistan
Student Paper <1 %

31 Kim, Hong, and Bum-Joon Kim. "Association of preS/S Mutations with Occult Hepatitis B Virus (HBV) Infection in South Korea: Transmission Potential of Distinct Occult HBV Variants", International Journal of Molecular Sciences, 2015.
Publication <1 %

32 Viral Infections of Humans, 2014.
Publication <1 %

33 inajog.com
Internet Source <1 %

34 repository.unhas.ac.id
Internet Source <1 %

35 www.natap.org
Internet Source

<1 %

36

Bleich, Lauren M., and Eugene S. Swenson. "Prevention of Neonatal Hepatitis B Virus Transmission :", *Journal of Clinical Gastroenterology*, 2014.

Publication

<1 %

37

Chengcong Chen, Xiaotao Jiang, Xuan Liu, Ling Guo, Weibin Wang, Shuqin Gu, Chunhua Wen, Xuan Yi, Libo Tang, Yongyin Li. "Identification of the association between HBcAg - specific T cell and viral control in chronic HBV infection using a cultured ELISPOT assay", *Journal of Leukocyte Biology*, 2020

Publication

<1 %

38

Holmes, Jacinta A, Tin Nguyen, Dilip Ratnam, Neel M Heerasing, Jane V Tehan, Sara Bonanzinga, Anouk Dev, Sally Bell, Stephen Pianko, Robert Chen, Kumar Visvanathan, Rachel Hammond, David Iser, Ferry Rusli, William Sievert, Paul V Desmond, D Scott Bowden, and Alexander J Thompson. "IL28B genotype is not useful for predicting treatment outcome in Asian chronic hepatitis B patients treated with pegylated interferon- α : does not predict outcome in hepatitis B", *Journal of Gastroenterology and Hepatology*, 2013.

<1 %

39

Jia-Horng Kao. "Role of viral factors in the natural course and therapy of chronic hepatitis B", *Hepatology International*, 11/30/2007

Publication

<1 %

40

Lei Zhang, Xien Gui, Bo Wang, Huiping Ji, Reziyan Yisilafu, Fengliang Li, Yun Zhou, Ling Zhang, Hui Zhang, Xiaohong Liu. "A study of immunoprophylaxis failure and risk factors of hepatitis B virus mother-to-infant transmission", *European Journal of Pediatrics*, 2014

Publication

<1 %

41

www.liverphil.org

Internet Source

<1 %

42

Ching-Sheng Hsu, Wei-Liang Liu, You-Chen Chao, Hans Hsienhong Lin et al. "Adipocytokines and liver fibrosis stages in patients with chronic hepatitis B virus infection", *Hepatology International*, 2015

Publication

<1 %

43

Luo, Z.. "Impact of the implementation of a vaccination strategy on hepatitis B virus infections in China over a 20-year period", *International Journal of Infectious Diseases*, 201202

Publication

<1 %

44	malariajournal.biomedcentral.com Internet Source	<1 %
45	www.mdpi.com Internet Source	<1 %
46	www.spandidos-publications.com Internet Source	<1 %
47	Merrill, R.M.. "Seroprevalence of markers for hepatitis B viral infection", International Journal of Infectious Diseases, 201102 Publication	<1 %
48	Stephen Locarnini, Scott Bowden. "Hepatitis B surface antigen quantification: Not what it seems on the surface", Hepatology, 2012 Publication	<1 %
49	theasianhealthfoundation.org Internet Source	<1 %
50	www.wjgnet.com Internet Source	<1 %
51	A Erhardt. "Response to interferon alfa is hepatitis B virus genotype dependent: genotype A is more sensitive to interferon than genotype D", Gut, 2005 Publication	<1 %
52	Ma, Lin, Nageswara R. Alla, Xiaomao Li, Ospan A. Mynbaev, and Zhongjie Shi. "Mother-to-child transmission of HBV: review of current	<1 %

clinical management and prevention
strategies : Mother-to-child transmission of
HBV", Reviews in Medical Virology, 2014.

Publication

53

hepatmon.com

Internet Source

<1 %

54

medicinamoderna.ro

Internet Source

<1 %

55

www.jsmcentral.org

Internet Source

<1 %

56

www.tandfonline.com

Internet Source

<1 %

57

A.K. Agarwal, Sourav Sen, Debabrata Banerjee, Rakesh Srivastava, A.K. Praharaj. "Distribution of hepatitis B virus genotype and cancer predicting precore and basal core promoter mutations", Medical Journal Armed Forces India, 2015

Publication

<1 %

58

Chen, Huey-Ling, Chien-Nan Lee, Chin-Hao Chang, Yen-Hsuan Ni, Ming-Kwang Shyu, Shih-Ming Chen, Jen-Jan Hu, Hans Hsienhong Lin, Lu-Lu Zhao, Shu-Chi Mu, Ming-Wei Lai, Chyi-Long Lee, Hsien-Ming Lin, Ming-Song Tsai, Jenn-Jeih Hsu, Ding-Shinn Chen, K. Arnold Chan, and Mei-Hwei Chang. "Efficacy of maternal tenofovir disoproxil fumarate in

<1 %

interrupting mother-to-infant transmission of hepatitis B virus", Hepatology, 2015.

Publication

59

Lin, Xiaofang, Yan Guo, Aifen Zhou, Yu Zhang, Jiangxia Cao, Mei Yang, Fang Xiao, Bin Zhang, and Yukai Du. "Immunoprophylaxis Failure against Vertical Transmission of Hepatitis B Virus in the Chinese Population : A Hospital-Based Study and a Meta-Analysis", The Pediatric Infectious Disease Journal, 2014.

Publication

<1 %

60

academic.oup.com

Internet Source

<1 %

61

docplayer.net

Internet Source

<1 %

62

hepatitisb.org.au

Internet Source

<1 %

63

wrap.warwick.ac.uk

Internet Source

<1 %

64

www.menzies.edu.au

Internet Source

<1 %

65

www.yumpu.com

Internet Source

<1 %

66

C. Q. Pan, W. Yi, M. Liu, G. Wan, Y.-H. Hu, M.-F. Zhou. "Lamivudine therapy during the second vs the third trimester for preventing

<1 %

transmission of chronic hepatitis B", Journal of
Viral Hepatitis, 2017

Publication

67

Herbers, U., S. Amini-Bavil-Olyaei, A. Mueller,
T. Luedde, C. Trautwein, and F. Tacke.

"Hepatitis B e antigen-suppressing mutations
enhance the replication efficiency of adefovir-
resistant hepatitis B virus strains :", Journal of
Viral Hepatitis, 2013.

Publication

68

Huey-Ling Chen, Chien-Nan Lee, Chin-Hao
Chang, Yen-Hsuan Ni et al. "Efficacy of
maternal tenofovir disoproxil fumarate in
interrupting mother-to-infant transmission of
hepatitis B virus", Hepatology, 2015

Publication

69

Michael W. Fried. "HBeAg and hepatitis B
virus DNA as outcome predictors during
therapy with peginterferon alfa-2a for HBeAg-
positive chronic hepatitis B", Hepatology,
02/2008

Publication

70

Pivert, A., A. Servant-Delmas, F. Lunel-Fabiani,
H. Le Guillou-Guillemette, S. Laperche, and A.
Ducancelle. "Correlation between the
promoter basal core and precore mutations
and HBsAg quantification in French blood
donors infected with hepatitis B virus : HBV in

<1 %

<1 %

<1 %

<1 %

French Blood Donors", Journal of Medical Virology, 2014.

Publication

71 journals.lww.com <1 %
Internet Source

72 media.neliti.com <1 %
Internet Source

73 onlinelibrary.wiley.com <1 %
Internet Source

74 ssw.unc.edu <1 %
Internet Source

75 www.ashm.org.au <1 %
Internet Source

76 Boccalini, Sara, Elettra Pellegrino, Emila Tiscione, Giovanna Pesavento, Angela Bechini, Miriam Levi, Stefano Rapi, Stefano Mercurio, Francesco Mannelli, Marta Peruzzi, Cesare Berardi, and Paolo Bonanni. "Sero-epidemiology of hepatitis B markers in the population of Tuscany, Central Italy, 20 years after the implementation of universal vaccination", Human Vaccines & Immunotherapeutics, 2013.
Publication

77 Hanah Kim, Mina Hur, Eunsin Bae, Kyung-A Lee, Woo-In Lee. "Performance evaluation of <1 %

cobas HBV real-time PCR assay on Roche cobas 4800 System in comparison with COBAS AmpliPrep/COBAS TaqMan HBV Test", Clinical Chemistry and Laboratory Medicine (CCLM), 2018

Publication

78

Lyu, H., D. Lee, Y.-H. Chung, J. A. Kim, J.-H. Lee, Y.-J. Jin, W. Park, P. Mathews, E. Jaffee, L. Zheng, E. Yu, and Y. J. Lee. "Synergistic Effects of A1896, T1653 and T1762/A1764 Mutations in Genotype C2 Hepatitis B Virus on Development of Hepatocellular Carcinoma", Journal of Viral Hepatitis, 2013.

Publication

79

Man-Fung Yuen. "Hepatitis B virus DNA levels at week 4 of lamivudine treatment predict the 5-year ideal response", Hepatology, 12/2007

Publication

80

Purnima Bhat. "Chapter 6 Maternal-Fetal Transmission of HBV", Springer Science and Business Media LLC, 2012

Publication

81

Yi-Hsing Chen, Sheng-Nan Lu, Jing-Hung Wang, Chao-Hung Hung, Tsung-Hui Hu, Chien-Hung Chen. "Pre-S/Surface and Core Promoter/Precore Mutations in Chronic Hepatitis B Patients with Severe Acute

<1 %

<1 %

<1 %

<1 %

Exacerbation", Digestive Diseases and Sciences, 2019

Publication

82

B. J. Zacher. "Multicenter Evaluation of the Elecsys Hepatitis B Surface Antigen Quantitative Assay", *Clinical and Vaccine Immunology*, 11/01/2011

Publication

<1 %

83

Inês Pita. "Hepatitis B inactive carriers: an overlooked", *Repositório Aberto da Universidade do Porto*, 2014.

Publication

<1 %

84

Julianne Bayliss, Lilly Yuen, Gillian Rosenberg, Darren Wong et al. "Deep sequencing shows that HBV basal core promoter and precore variants reduce the likelihood of HBsAg loss following tenofovir disoproxil fumarate therapy in HBeAg-positive chronic hepatitis B", *Gut*, 2017

Publication

<1 %

85

Martinot-Peignoux, Michelle, Martine Lapalus, Tarik Asselah, and Patrick Marcellin. "The role of HBsAg quantification for monitoring natural history and treatment outcome", *Liver International*, 2013.

Publication

<1 %

86

Tseng, T.-C., C.-J. Liu, C.-L. Chen, H.-C. Yang, T.-H. Su, C.-C. Wang, W.-T. Yang, S. F.-T. Kuo,

<1 %

C.-H. Liu, P.-J. Chen, D.-S. Chen, and J.-H. Kao.
"Risk Stratification of Hepatocellular
Carcinoma in Hepatitis B Virus e Antigen-
Negative Carriers by Combining Viral
Biomarkers", The Journal of Infectious
Diseases, 2013.

Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On