

Correlation of Oxidized-LDL, Resistin and IL-1b in Centrally Obese Men

by Prof. Dr. Muh. Nasrum Massi, Ph.d

Submission date: 01-Dec-2021 01:25AM (UTC+0700)

Submission ID: 1716560834

File name: Correlation_of_Oxidized-LDL.pdf (501.33K)

Word count: 2971

Character count: 16567

RESEARCH ARTICLE

Correlation of Oxidized-LDL, Resistin and Interleukin-1 Beta in Centrally Obese MenTri Nevita Margareth Panjaitan^{1,2,*}, Marita Kaniawati^{1,3}, Ilhamjaya Patellongi⁴¹Postgraduate Program in Clinical Biochemistry, Hasanuddin University, Jl. Perintis Kemerdekaan Km.10, Makassar, Indonesia²Prodia Occupational Health Institute, Jl. Kramat VI No.18, Jakarta, Indonesia³Prodia Clinical Laboratory, Jl. Cisangkuy No.2, Bandung, Indonesia⁴Faculty of Medicine, Hasanuddin University, Jl. Perintis Kemerdekaan Km.10 Makassar, Indonesia

*Corresponding author. E-mail: trinevt@gmail.com

Abstract

BACKGROUND: Obesity is one of serious health problems, which increases the risks of cardiovascular disease, type 2 diabetes mellitus, and cancers. It is strongly associated with changes in the physiological function of adipose tissue, leading to altered secretion of adipokines and activation of inflammatory signaling pathways. This study was aimed to investigate correlation of resistin, interleukin-1 beta (IL-1 β), and oxidized low-density lipoprotein (OxLDL) in centrally obese men.

METHODS: The research was conducted with a cross-sectional design involving 68 centrally obese men aged 31 to 60 years old, with waist circumference (WC) > 90 cm. All subjects fulfilled the inclusion criteria. Anthropometric parameters, creatinine, estimated glomerular filtration rate (eGFR), serum glutamic oxaloacetic transaminase (GOT), serum glutamic-pyruvic transaminase (SGPT), and high-sensitivity C-reactive protein (hs-CRP) were measured. Serum concentrations of resistin, IL-1 β and OxLDL were measured by enzyme-linked immunosorbent assay.

RESULTS: The study results showed there was a significant correlation of WC-OxLDL (r=0.235; p=0.030) and some correlation of WC-resistin (r=0.201; p=0.050). However, we observed no significant correlation of IL-1 β -OxLDL (r=0.042; p=0.369), resistin-OxLDL (r=-0.072; p=0.285) and WC-IL-1 β (r=-0.042; p=0.367).

CONCLUSION: Our data show a relationship between WC and OxLDL, but the mechanism does not appear to be directly related to resistin or IL-1 β .

Abstrak

LATAR BELAKANG: Obesitas merupakan salah satu masalah kesehatan yang cukup serius, yang dapat meningkatkan risiko penyakit kardiovaskular, diabetes melitus tipe 2, kanker, dan lainnya. Hal ini sangat erat kaitannya dengan perubahan fungsi fisiologis jaringan adiposa yang menyebabkan perubahan sekresi adipokin dan aktivasi jalur sinyal inflamasi. Penelitian ini bertujuan untuk menyelidiki kontribusi resistin, interleukin-1 beta (IL-1 β), dan low-density lipoprotein teroksidasi (OxLDL) pada pria obesitas sentral.

METODE: Penelitian ini dilakukan dengan desain potong lintang yang melibatkan 68 subyek pria obesitas berusia 31 sampai 60 tahun, dengan lingkar pinggang (LP) > 90 cm. Semua subyek memenuhi kriteria eksklusi. Dilakukan pengukuran terhadap parameter antropometri, kreatinin, laju filtrasi glomerulus, Serum glutamat oksaloasetat transaminase, serum glutamat piruvat transaminase, dan high-sensitivity C-reactive protein. Konsentrasi serum resistin, IL-1 β dan OxLDL diukur dengan metode enzyme-linked immunosorbent assay.

HASIL: Penelitian ini menunjukkan ada hubungan yang signifikan antar LP-OxLDL (r=0,235; p=0,030), LP-resistin (r=0,201; p=0,050). Namun, tidak terdapat korelasi yang signifikan antara IL-1 β -OxLDL (r=0,042; p=0,369), resistin-OxLDL (r=-0,072; p=0,285) dan LP-IL-1 β (r=-0,042; p=0,367).

KESIMPULAN: Penelitian ini menunjukkan hubungan antara lingkar pinggang dan progresi OxLDL, namun mekanismenya tidak melalui jalur resistin ataupun IL-1 β .



KEYWORDS: resistin, IL-1 β , OxLDL, atherosclerosis

33

Indones Biomed J. 2013; 5(1): 59-64

Introduction

Obesity is a serious health problem that increases the risk not only of cardiovascular disease and type 2 diabetes mellitus (T2DM) but also of various types of cancer. Obesity is strongly associated with changes in the physiological function of adipose tissue, leading to insulin resistance, chronic inflammation, and altered secretion of adipokines.(1) Obesity is a condition in which there is excessive accumulation of body fat.(2) Obesity-associated inflammation occurs as a result of immune cell infiltration of the adipose tissue and increased production of pro-inflammatory cytokines. These changes negatively affect normal adipocyte functions such as triglyceride storage and lipolysis, leading to high circulating levels of free fatty acids and ectopic lipid accumulation. In addition, elevated levels of pro-inflammatory cytokines, including interleukin (IL)-1 β , IL-6, and tumor necrosis factor (TNF)- α , reduce insulin sensitivity, which can lead to the development of T2DM if insulin production by the pancreatic β cells is not sufficiently increased. Although it has been recognized for some time that pro-inflammatory cytokines inhibit insulin signaling, the molecular mechanisms that trigger inflammation and cytokine production in obesity are not well understood.(3)

Obesity-related danger signals, such as palmitate, ceramide, high glucose concentrations, islet amyloid β peptide deposition, defective autophagy, or mitochondrial dysfunction may lead to the generation of reactive oxygen species (ROS) and subsequent activation of the nucleotide-binding oligomerization domain, leucine rich repeat and pyrin domain containing (NLRP3) inflammasome in adipocytes, pancreatic islet cells, or infiltrating macrophages present in these tissues. NLRP3 inflammasome activation promotes the cleavage and activation of Caspase-1, resulting in the secretion of IL-1 β , a pro-inflammatory cytokine that negatively affects adipocyte differentiation, inhibits insulin signaling, and has a cytotoxic effect on insulin-producing pancreatic β cells. Identification of a connection between obesity, the inflammasome, and insulin resistance links inflammasome activation with metabolic tissue dysfunction, which may contribute to the pathogenesis of T2DM, and other diseases

secara langsung.

KATA KUNCI: resistin, IL-1 β , OxLDL, aterosklerosis

associated with obesity-induced inflammation.(3)

IL-1 β , a proinflammatory cytokine with a variety of activities, is strongly induced in monocytes by direct contact with stimulated T lymphocytes. Both these cells involved in immunoinflammatory diseases (such as rheumatoid arthritis, multiple sclerosis, and systemic lupus erythematosus) and atherosclerosis. IL-1 β has been shown to have important effects on the cell types that constitute atherosclerotic lesions.(4)

Resistin belongs to a family of cysteine-rich secretory proteins called resistin-like molecules or is found in inflammatory zones (FIZZ) proteins. In rodents, resistin is derived almost exclusively from fat tissue, and adipose expression and serum levels are elevated in models of obesity and insulin resistance. In contrast to rodents, in humans resistin is expressed primarily in inflammatory cells. Resistin expression in human monocytes was markedly increased by treatment with endotoxin and proinflammatory cytokines. (5) Resistin was found with relevance to the incidence of inflammation and inflammation-related diseases such as atherosclerosis and arthritis.(6)

Experimental studies have shown that the oxidative conversion of low-density lipoproteins (LDL) to oxidized (OxLDL) is considered to be a key event in the biological process that initiates and accelerates the development of the early atherosclerotic lesion, the fatty streak and the unstable and plaque rupture.(7)

Increasing prevalence of obesity has motivated some researchers to do studies to prevent or overcome the negative effects of the disease. It has been known that obesity correlates with atherosclerosis and cardiovascular disease which cause mortality in many countries including Indonesia. This study aimed to investigate the role of resistin and IL-1 β in pathomechanism of atherosclerosis in centrally obese men.

Methods

Subjects

Participants were recruited from the community in Jakarta. The sample population comprised centrally obese men aged 30 to 60 years. WC > 90 cm was determined by

measuring the waist between the lower rib and iliac crest. Interview was done to gather information on medical history, present condition, and drugs consumption. Other characteristics *i.e.* smoking and alcohol consumption habits and medication were investigated. Individual interviews using a structured questionnaire. Subjects with estimated glomerular filtration rate (eGFR) < 60 mL/minute or serum glutamic oxaloacetic transaminase (SGOT) / serum glutamic-pyruvic transaminase (SGPT) more than 2 times the normal value or high-sensitivity C-reactive protein (hs-CRP) \geq 10 mg/L were excluded. This study was approved by the ethics committee of Hasanuddin University Faculty of Medicine (registered number UH1303119), and written informed consent was obtained from each subject.

34. Method collection and biochemical analysis

Blood samples were collected in the morning after overnight fasting. Venous blood was collected, centrifuged, and the

separated serum was frozen immediately at -20°C. Serum levels of SGOT, SGPT, creatinine, and hsCRP were assayed using autoanalyzer Cobas (Roche Diagnostics). Circulating resistin, IL-1 β and OxLDL levels were measured using a commercially available enzyme-linked immunosorbent assay (ELISA) (resistin: Quantikine, catalogue no. DRSN00; IL-1 β : Quantikine, catalogue no. HSLB00C; OxLDL: Mercodia, catalog no. 10-1143-01).

14. Statistical analysis

Statistical analysis was performed using SPSS for Windows version 16.0 (USA, SPSS 101 Inc.). Data were analyzed by Spearman and Pearson bivariate and partial correlation to evaluate the relationships between WC, resistin serum, IL-1 β , and OxLDL. All values were expressed as mean \pm standard deviation (SD), unless otherwise specified. Hypothesis testing was two-tailed at a significance level of 0.05.

Table 1. Subjects Characteristics.

Characteristics	Mean \pm SD	Min	Max	Median
Age (year)	45.960 \pm 8.169	31	60	45.500
Height (cm)	166.200 \pm 5.288	150	176	166
Weight (kg)	79.670 \pm 9.663	62	115	77
WC (cm)	99.220 \pm 6.788	91	130	99
DBP (mmHg)	83.600 \pm 14.295	60	120	80
SBP (mmHg)	124.260 \pm 18.724	90	170	120
BMI (kg/m ²)	28.820 \pm 3.093	23.58	41.23	28.820
SGPT (U/L)	32.290 \pm 17.557	10	122	27
SGOT (U/L)	22.980 \pm 8.010	12	52	21
Creatinine (mg/dL)	0.871 \pm 0.124	0.59	1.20	0.860
eGRF (mL/min/1.73m ²)	102.080 \pm 11.665	67	135	103
hs-CRP (mg/L)	2.950 \pm 2.260	0.30	9.50	2.400
IL-1 β (pg/mL)	0.050 \pm 0.043	0.01	0.28	0.046
Resistin (ng/mL)	8.400 \pm 3.551	2.97	16.74	7.290
OxLDL (U/L)	269.080 \pm 23.805	198.60	308.80	270.400

24. DBP = diastolic blood pressure; SBP = systolic blood pressure

Table 2. Correlation between OxLDL and Other Variables

Variable	OxLDL			
	Bivariate		Partial	
	r	p	r	p
Age	0.260	0.032	0.251	0.022
WC	0.242	0.046	0.235	0.030
IL-1 β	0.058	0.636	0.042	0.369
Resistin	-0.028	0.828	-0.072	0.285

Table 3. Correlation between Resistin, IL-1 β and WC

Variable	WC			
	Bivariate		Partial	
	r	p	r	p
Resistin	0.275	0.010	0.201	0.050
IL-1 β	0.079	0.256	-0.042	0.367

Results

Clinical characteristics of subjects

The anthropometric and clinical characteristics of the participants are shown in Table 1. The subjects comprised centrally obese men aged 46 ± 8.17 (range: 31-60) years and the mean WC was 99 ± 6.78 (range: 91-130) cm.

Correlation between waist circumference and OxLDL

Results of the bivariate and partial correlation analysis of WC, resistin, IL-1 β and OxLDL are presented in Table 2. The circulating OxLDL tended to correlate positively with WC ($r=0.235$) and the correlation was significant ($p=0.030$). IL-1 β also positive correlation with OxLDL, but the correlation was not significant ($r=0.042$ dan $p=0.369$). The correlation of resistin and OxLDL was negative but not significant ($r=-0.072$ dan $p=0.285$).

Correlation between resistin, interleukin-1 beta and WC

Table 3 shows the correlation between resistin, IL-1 β and WC. The partial correlation analysis of WC and resistin shows that both have a significant positive correlation

($r=0.201$; $p=0.050$), while the circulating IL-1 β tended to correlate negatively with WC, but the correlation was not significant ($r=-0.042$; $p=0.367$).

Discussion

This study used bivariate and partial correlation analysis. The partial correlation was done to prevent ambiguous correlation due to some variables, which is called a pure correlation. WC and OxLDL show positive and significant correlation ($r=0.235$; $p=0.030$). It shows that the greater the WC the more the OxLDL concentration increased. This result agrees with the result of the study by Weinbrenner *et al.* which showed >102 cm in men and WC > 88 cm in women with body mass index (BMI)=25.0-29.9 and BMI >30 had strong correlation with elevated OxLDL concentration. Holvoet in 2001 proved that BMI was one of strong predictors in elevating OxLDL concentration. (8, 9) The statistic analysis showed that OxLDL had positive correlation with IL-1 β but not significant ($r=0.042$, $p=0.369$) and a negative non significant correlation with resistin ($r=-0.072$; $p=0.285$).

Several previous studies have been done to investigate correlation between OxLDL and adipokines, but none of them focused on correlation with resistin. This study had the hypothesis that there might be correlation between resistin and OxLDL. The analysis showed that OxLDL and resistin had negative correlation, but not significant. Gazbar *et al.* observed that resistin was significantly more abundant in atherosclerotic lesions ($r=0.822$; $p<0.01$). This correlation was a novel finding though the correlation between resistin and OxLDL was still unclear. It could happen due to proinflammatory characteristic of resistin in human which is a promising biomarker. The different result found in this study could be due to different characteristics of the study subject. Gazbar *et al.* used obese men with diabetic history but this study used obese men with no diabetic history. This might occur due to the fact that diabetic patients have insulin resistance in which resistin has important effect in that condition.(10)

The proatherogenic characteristic of IL-1 is due to cytokine's ability to modulate adhesion cell and monocyte filtration to subendothelium. Kirii *et al.* pointed out that the absence of IL-1 β decreased atherosclerosis progression in mice Apolipoprotein E (ApoE)³ which might exist because of decreased expression of vascular cell adhesion molecule-1 (VCAM-1) and monocyte chemoattractant protein-1 (MCP-1) in aorta.(4) This process might attract monocytes into the artery intima. The adhesion that continues with differentiation into macrophages and forms foam cells, is the key of atherogenesis phase.(11) According to the study by Mojtaba *et al.*, IL-1 β serum concentration in obese men is higher than that in non¹¹ obese men. Nevertheless, not many studies have been done to analyze the role and correlation of IL-1 β in atherosclerosis progression.(12)

Resistin is a proinflammatory cytokine which might have an important role in atherosclerosis and other inflammatory diseases. A recent study has shown that resistin will make worst the atherosclerosis condition by stimulating monocytes, endothelial cells, and vascular smooth muscle cells to induce vascular inflammation.(8)

From our study results we can see that WC has mild correlation with resistin ($r=0.201$; $p=0.050$). It shows that elevated WC in men has some correlation with chronic inflammation process, characterized by positive correlation with inflammation biomarker. Previous studies have proven it already that resistin has positive correlation with body fat. It indicates resistin's elevating expression is a result of adipocyte differentiation.(13)

Results of our statistical tests have shown a negative correlation tendency, but not significant between WC and IL-1 β ($r=-0.042$, $p=0.367$), even it showed reverse correlation.

It means as there is decrease of concentration of IL-1 β , there is increase of the WC. However, not many studies support this result.

Conclusion

According to results of our study, we conclude that there is no significant increase of resistin or significant decrease of IL-1 β in obese men. Regarding this fact, the possible pathways need to be further investigated.

Acknowledgement

This study was supported by the Prodia Education and Research Institute. We thank for the invaluable supports given to this study.

References

1. Kruijsdijk RCM, Wall EVD, Visseren FLJ. Obesity and cancer: the role of dysfunctional adipose tissue. *Cancer Epidem Biomar.* 2009; 18: 2569-78.
2. Soegondo AW, Setiyohadi B, Alwi I, Simadibrata M, Setiati S. *Buku Ajar Ilmu Penyakit Dalam.* 4th ed. Jakarta: Pusat Penerbitan Departemen Ilmu Penyakit Dalam Fakultas Kedokteran Universitas Indonesia; 2007.
3. R&D Systems [http://www.rndsystems.com/index.aspx]. Obesity-Induced Activation of the Nlrp3 Inflammasome Promotes Insulin Resistance; 2013 [updated 2013; cited 2013]. Available from: http://www.rndsystems.com/cb_detail_objectname_cb11i2_obesity_induced_activation_Nlrp3.aspx
4. Kirii H, Niwa T, Yamada Y, Wada H, Saito K, Yoichiro. Lack of Interleukin-1 β decreases the severity of atherosclerosis in ApoE-deficient mice. *Arterioscler Thromb Vasc Biol.* 2003; 23: 656-60.
5. Reilly PM, Lehrke M, Wolfe ML, Rohatgi A, Lazar MA, Rader DJ. Resistin is an inflammatory marker of atherosclerosis in humans. *J Am Heart Assoc.* 2005; 111: 932-9.
6. Pang S and Le Y. Role of resistin in inflammation and inflammation-related diseases. *Cell Mol Immunol.* 2006; 1: 29-34.
7. Mercodia Oxidized LDL ELISA Kit Insert. Enzyme immunoassay. *Mercodia Catalogue.* 2004; No. 10-1143-01.
8. Weinbrenner T, Schroder H, Escurriol V, Fito M, Elosua R, Vila J, *et al.* Circulating OxLDL is associated with increased waist circumference independent of body mass index in men and women. *Am J Clin Nutr.* 2006; 83: 30-5.
9. Holvoet P, Mertens A, Verhamme P, Bogaerts K, Beyens G, Verhaeghe R, *et al.* Circulating oxidized LDL is a useful marker for identifying patients with coronary artery disease. *Arterioscler Thromb Vasc Biol.* 2001; 21: 844-8.
10. Gazbar S, Fethi BS, Abdelmagid T, Chiheb BRM, Omrane B. Relationship between oxidized-LDL and resistin levels in obese diabetic subject. *J Basic Appl Sci Res.* 2013; 4: 313-7.

11. Vicenova B, Vopaslensky V, Burysek L, Pospisek M. Emerging role of interleukin-1 in cardiovascular diseases. *Physiol Res.* 2009; 58: 481-98.
12. Mojtaba E, Mahdi K, Mehdi KJR, Amir S. Serum interleukin-1 beta plays an important role in insulin secretion in type II diabetic. *J Biosci* 2011; 32: 93-9.
13. Kusminski CM, McTeman PG, Kumar S. Role of resistin in obesity, insulin resistance and type II diabetes. *Clin Sci* 2005; 109: 243-56.

Correlation of Oxidized-LDL, Resistin and IL-1b in Centrally Obese Men

ORIGINALITY REPORT

23%

SIMILARITY INDEX

17%

INTERNET SOURCES

15%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

- 1 **Mohammad Nurul Amin, Md. Saddam Hussain, Md. Shahid Sarwar, Md. Mizanur Rahman Moghal et al. "How the association between obesity and inflammation may lead to insulin resistance and cancer", Diabetes & Metabolic Syndrome: Clinical Research & Reviews, 2019** 1%

Publication
- 2 **journals.lww.com** 1%

Internet Source
- 3 **www.researchsquare.com** 1%

Internet Source
- 4 **Terawan Agus Putranto, Irawan Yusuf, Bachtiar Murtala, Andi Wijaya. "Intra Arterial Heparin Flushing Increases Cereberal Blood Flow in Chronic Ischemic Stroke Patients", The Indonesian Biomedical Journal, 2016** 1%

Publication
- 5 **Submitted to Unviersidad de Granada** 1%

Student Paper

6	mafiadoc.com Internet Source	1 %
7	molecularautism.biomedcentral.com Internet Source	1 %
8	Cai, Shuang-Ming, Ren-Qiang Yang, yang Li, Zuo-Wei Ning, Li-Li Zhang, Gao-Su Zhou, Wei Luo, Da-Huan Li, Yan Chen, miao-xia Pan, and Xu Li. "Angiotensin-(1-7) improve liver fibrosis by regulating the NLRP3 inflammasome via redox balance modulation", <i>Antioxidants & Redox Signaling</i> , 2016. Publication	1 %
9	circres.ahajournals.org Internet Source	1 %
10	docplayer.info Internet Source	1 %
11	slideheaven.com Internet Source	1 %
12	www.aasmnet.org Internet Source	1 %
13	Submitted to Mae Fah Luang University Student Paper	1 %
14	spandidos-publications.com Internet Source	1 %
15	media.neliti.com Internet Source	

<1 %

16

repository.ubn.ru.nl

Internet Source

<1 %

17

www.jove.com

Internet Source

<1 %

18

Doddy De Queljoe, Amelia Lorensia, Liana Widharta, Sugiarto Widjaja. "Cost-Effectiveness Analysis of Pharmacotherapy for Hematemesis-Melena Treatment in Hospitalized Patients with Hepatic Cirrhosis", The Indonesian Biomedical Journal, 2013

Publication

<1 %

19

Siti Maryani, Neneng Ratnasari, Siti Nurdjanah. "Correlation between leptin level with lipid profile and free fatty acid in liver cirrhosis patients", Journal of the Medical Sciences (Berkala Ilmu Kedokteran), 2014

Publication

<1 %

20

www.ogscience.org

Internet Source

<1 %

21

Helmut Schröder. "Secular Trends of Obesity and Cardiovascular Risk Factors in a Mediterranean Population*", Obesity, 03/2007

Publication

<1 %

22 Sixto E. Sanchez, Michelle A. Williams, Martin Muy-Rivera, Chunfang Qiu, Surab Vadachkoria, Victor Bazul. "A case-control study of oxidized low density lipoproteins and preeclampsia risk", Gynecological Endocrinology, 2009
Publication

23 bmcpublichealth.biomedcentral.com
Internet Source

24 eprints.mums.ac.ir
Internet Source

25 www.omicsonline.org
Internet Source

26 Dimitrios P. Papadopoulos, Thomas K. Makris, Panagiota G. Krespi, Maria Poulakou et al. "Adiponectin and Resistin Plasma Levels in Healthy Individuals With Prehypertension", The Journal of Clinical Hypertension, 2007
Publication

27 H. MIESZCZANSKA. "Effects of pioglitazone on fasting and postprandial levels of lipid and hemostatic variables in overweight non-diabetic patients with coronary artery disease", Journal of Thrombosis and Haemostasis, 5/2007
Publication

28

Internet Source

<1 %

29

www.unboundmedicine.com

Internet Source

<1 %

30

yoursay.suara.com

Internet Source

<1 %

31

Adriana Todingrante, Mansyur Arif, Uleng Bahrn, Ferry Sandra. "Study of Low-grade Chronic Inflammatory Markers in Men with Central Obesity: Cathepsin S was Correlated with Waist Circumference", The Indonesian Biomedical Journal, 2013

Publication

<1 %

32

H. Asano. "Plasma resistin concentration determined by common variants in the resistin gene and associated with metabolic traits in an aged Japanese population", Diabetologia, 09/01/2009

Publication

<1 %

33

John Wantania, Syakib Bakri, Karel Pandelaki, Maisuri Chalid. "Altered Level of Soluble fms-like Tyrosine Kinase 1 (sFlt1) and Hypoxia Inducible Factor-1alpha (HIF-1alpha) in Normotensive Pregnancy and Preeclampsia", The Indonesian Biomedical Journal, 2013

Publication

<1 %

34 Lapointe, A.. "Circulating oxidized LDL is associated with parameters of the metabolic syndrome in postmenopausal women", *Atherosclerosis*, 200704
Publication

<1 %

35 journal.fkm.ui.ac.id
Internet Source

<1 %

36 journal.gunabangsa.ac.id
Internet Source

<1 %

37 live-look-no.icu
Internet Source

<1 %

38 worldwidescience.org
Internet Source

<1 %

39 Ordonez, F.J.. "Correlation between glutathione peroxidase activity and anthropometrical parameters in adolescents with Down syndrome", *Research in Developmental Disabilities*, 200701/02
Publication

<1 %

40 aimjournal.ir
Internet Source

<1 %

41 nutritionandmetabolism.biomedcentral.com
Internet Source

<1 %

42 patents.google.com
Internet Source

<1 %

- | | | |
|----|---|------|
| 43 | repository.unair.ac.id
Internet Source | <1 % |
| 44 | simdos.unud.ac.id
Internet Source | <1 % |
| 45 | www.science.gov
Internet Source | <1 % |
| 46 | Conceição Calhau. "Oxidative Stress in the Metabolic Syndrome", Oxidative Stress Inflammation and Angiogenesis in the Metabolic Syndrome, 2009
Publication | <1 % |
| 47 | Helmut Schröder, Maria-Isabel Covas. "Reply to P Holvoet", The American Journal of Clinical Nutrition, 2006
Publication | <1 % |
| 48 | Lauri Vehkala, Olavi Ukkola, Y. Antero Kesäniemi, Mika Kähönen et al. "Plasma IgA antibody levels to malondialdehyde acetaldehyde-adducts are associated with inflammatory mediators, obesity and type 2 diabetes", Annals of Medicine, 2013
Publication | <1 % |
| 49 | Mayuko Murakami, Satoko Osuka, Ayako Muraoka, Shotaro Hayashi et al. "Effectiveness of NLRP3 Inhibitor as a Non-Hormonal Treatment for Ovarian | <1 % |

Endometriosis", Research Square Platform LLC, 2021

Publication

50

Anna Meiliana, Nurrani Mustika Dewi, Andi
Wijaya. "Resveratrol: A Sirtuin Activator and
The Fountain of Youth", The Indonesian
Biomedical Journal, 2015

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On