

# Contribution of NRAMP1 gene expression and protein level in pulmonary and latent TB infection in Indonesia

*by Ilhamjaya Patellongi*

---

**Submission date:** 05-Dec-2021 09:41PM (UTC+0700)

**Submission ID:** 1720988536








**File name:** 4.\_Contribution\_of\_NRAMP1\_gene.pdf (467.36K)

**Word count:** 4019

**Character count:** 21686



## Contribution of *NRAMP1* gene expression and protein level in pulmonary and latent TB infection in Indonesia

Irda Handayani<sup>1,2</sup>, Irawaty Djaharuddin<sup>3</sup> , Rosdiana Natzir<sup>4</sup> , Mansyur Arief<sup>1</sup>, Ahyar Ahmad<sup>5</sup> , Mochammad Hatta<sup>6</sup> ,  
Rosana Agus<sup>7</sup>, Ilhamjaya Patellongi<sup>8</sup> , Muhammad Amin<sup>9</sup>, Yuyun Widaningsih<sup>1</sup>, Handayani Halik<sup>2,10</sup>, Najdah Hidayah<sup>2</sup> ,  
Subair Subair<sup>2,10</sup>, Yanti Leman<sup>11</sup>, Wiendra Waworuntu<sup>12</sup>, Muhammad Nasrum Massi<sup>6\*</sup> 

<sup>1</sup>Department of Clinical Pathology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

<sup>2</sup>Postgraduate Program, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

<sup>3</sup>Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

<sup>4</sup>Department of Biochemistry, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

<sup>5</sup>Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Hasanuddin, Makassar, Indonesia.

<sup>6</sup>Department of Clinical Microbiology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

<sup>7</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Hasanuddin, Makassar, Indonesia.

<sup>8</sup>Department of Physiology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

<sup>9</sup>Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Universitas Airlangga, Makassar, Indonesia.

<sup>10</sup>Universitas Mega Rezky, Makassar, Indonesia.

<sup>11</sup>Department of Pharmacology, Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.

<sup>12</sup>Directorate of Traditional Health Services, Directorate General of Healthy Services, Ministry of Health of the Republic of Indonesia.

### ARTICLE INFO

Received on: 19/11/2020  
Accepted on: 22/04/2021  
Available online: 05/08/2021

#### Key words:

*NRAMP1* gene expression,  
protein level, pulmonary TB,  
latent TB.

### ABSTRACT

Natural resistance-associated macrophage protein 1 (*NRAMP1*) gene is regulated to play an important role in human vulnerability against various infections, including mycobacterial infection. This research purpose was to investigate the correlation between *NRAMP1* gene expressions and protein level with susceptibility to tuberculosis (TB) infection and their contacts. *NRAMP1* gene expression and protein level from 30 active TB patients, 27 latent TB infection household contacts, and 33 healthy subjects as control were examined by the real-time polymerase chain reaction and enzyme-linked immunosorbent assay method. *NRAMP1* gene expression value in the healthy group was 4.61 times higher than active TB patients ( $p = 0.025$ ) and the *NRAMP1* expression value in the healthy group was 4.02 times higher than the latent TB group ( $p = 0.117$ ). The mean protein level of *NRAMP1* was 196.41, 199.27, and 224.70 ng/ml in the active TB, latent TB, and healthy groups, respectively ( $p > 0.05$ ). *NRAMP1* expression was downregulated and protein level was lower in active TB patients than latent TB patients and healthy individuals.


### INTRODUCTION

Human genetics that regulates infectious disease vulnerability has been widely researched (Canonne-Hergaux *et al.*, 1999). These research studies are expected to contribute to identifying the high-risk populations being exposed to infectious

diseases, so that prevention and diagnosis can be carried out properly as early as possible. In addition, the study also clarified the pathomechanism of infection and how the host defends against microbes. Several human genes have been studied that correlated to the susceptibility of tuberculosis (TB) infection (Qidwai *et al.*, 2012).

Natural resistance-associated macrophage protein 1 (*NRAMP1*) gene that is also called *SLC11A1* (solute carrier 11a1) has been known to play a role in human susceptibility to various infections, including mycobacterial infection. The human *SLC11A1* gene is arranged on chromosome 2q35 and codes the

\*Corresponding Author

Muhammad Nasrum , Department of Clinical Microbiology,  
Faculty of Medicine, Universitas Hasanuddin, Makassar, Indonesia.  
E-mail: [nasrum2000@yahoo.com](mailto:nasrum2000@yahoo.com)

**9** *NRAMP1*, an integral protein-membrane consisting of 550 amino acids that are widely expressed in the lysosomal compartments of monocytes and macrophages (Dow, 2015). *NRAMP1* is located in the phagosomal membrane of macrophages and is recruited rapidly to the phagosomal membrane after the phagocytosis process. It appears to work as a divalent cation transporter [Mn (II) and Fe (II)] across the phagosomal membrane of macrophages, otherwise known as phagosome waste pumps (Hanikenne *et al.*, 2009; Murray, 2013). The antimicrobial effect of *NRAMP1* is due to the ability of *NRAMP1* to deplete the essential metal nutrients for bacteria in the phagosomes (Zhao and Enns, 2012). In addition, *NRAMP1* also increases the acidity of phagosomes to levels that are lethal to bacteria (Dow, 2015; Flaherty, 2012). For patients with *NRAMP1* deficiency, their phagolysosome pH approaches pH 6.8, which promotes bacterial growth. Mutations in the *NRAMP1* gene are thought to affect alveolar macrophage phagosome function (Flaherty, 2012). *NRAMP1* gene polymorphisms have been reported to be related to tuberculosis (Meilang *et al.*, 2012). However, we did not find any research in patients with TB on the expression of the *NRAMP1* gene.

Understanding the role of human genetic factors in the immune system controlling infectious disease susceptibility/resistance is essential for tuberculosis research because it will enable genetic dissection of antimycobacterial immunity and open up new possibilities for preventive and therapeutic measures. This study aimed to investigate the expression and protein level of *NRAMP1* in patients with pulmonary tuberculosis and household contacts who are latent infected with TB, to contribute to understanding the role of genetic and immunology factors in susceptibility to tuberculosis.

## MATERIALS AND METHODS

### Sample recruitment

In this research, 90 participants were selected and divided into three categories, each with 30 samples of the new smear-positive pulmonary tuberculosis group from the Community Center for Lung Health Makassar or Balai Besar Kesehatan Paru Masyarakat (BBKPM) Makassar as active TB group; 33 subjects of household contact with TB patients and were actively screened using QuantiFERON TB Gold Plus TB test (Qiagen, Germany) to determine their exposure to *Mycobacterium tuberculosis* (Mtb). The group with positive Immunoglobulin Release Assay (IGRA) was categorized as a latent group. The third group included 27 healthy subjects who were also subjected to an IGRA examination first and had negative IGRA results. Positive sputum samples were decontaminated and then continued with the culture process at the Tuberculosis Unit of the Hasanuddin University Medical-Research Center (HUM-RC) Laboratory, Makassar, Indonesia. Blood samples were centrifuged for separation plasma samples at 3,000 rpm for 15 minutes at 4°C. Plasma samples and clotted blood samples were stored at -80°C before ELISA and RNA extraction.

### Ethical approval

This research had ethical approval by the Research Ethics Commission of the Faculty of Medicine, Universitas Hasanuddin, Makassar, South Sulawesi, Indonesia (No. 1125/UN4.6.4.5.31/PP36/2020), and all research subjects had signed the consent.

### *NRAMP1* gene expression measurement

Total RNA was extracted from blood samples using RNA Isolation Kit RNeasy Mini Kit (Qiagen, USA) according to the manufacturer's instructions. In short, the samples were lysed and homogenized first. In order to have optimal binding conditions, ethanol was added to the lysate. The lysate was then placed onto the RNeasy silica membrane. RNA binds, and all contaminants were efficiently washed away. Pure concentrate RNA was eluted in 50 µl RNase free water and used in complementary DNA (cDNA) synthesis.

The iScript cDNA Synthesis Package (Bio-Rad Laboratories, Inc., CA) was used to synthesize the cDNA from 100 ng of total RNA. A mixture of 20 microliters was prepared with 4 µl 5× iScript Reaction blend, 1 µl of iScript reverse, and nuclease-free water. The cycling requirement for the synthesis of cDNA was priming for 5 minutes at a temperature of 25°C, followed by 20 minutes of reverse transcription at 46°C and 1 minute of RT inactivation at 95°C.

For the target genes, the relative quantification was rendered using 12.5 µl Fast Evagreen® qPCR Master Mix (Biotium, Inc., Fremont, CA), 0.5 µl cDNA, and 0.5 µl (0.5 µM concentration) of each *NRAMP* forward and reverse primers for *NRAMP1* (forward 5'-GCATCTCCCAATTCATGGT-3' and reverse 5'-AACTGTCCACTCTATCCTG-3') and glyceraldehyde-3 phosphate dehydrogenase (*GAPDH*) (forward 5'-CCTGCACCACCAACTGCTTA-3' and reverse 5'-GGCATCCACAGTCTTCTGGG-3') and amplified in real-time PCR (Bio-Rad Lab. Inc., Hercules, CA) with cycling conditions at 95°C for 30 seconds (initial denaturation and activation of enzyme) followed by 45 cycles of 95°C for 5 seconds and annealing at 60°C for 30 seconds. The amplicons have been heated at 65°C–95°C with a temperature rise of 0.5°C at 5 seconds per phase, and the melt curve analyses have been completed.

All analyses were performed using Bio-Rad CFX™ Manager Software (version 3.1, Bio-Rad Lab. Inc., Hercules, CA). The level of messenger ribonucleic acid (mRNA) expression was normalized by using *GAPDH* mRNA as a housekeeping gene, and the relative fold induction was determined using the  $2^{-(\Delta\Delta Ct)}$  calculation, where  $\Delta Ct$  is  $Ct(\text{target}) - Ct(\text{GAPDH})$ , and  $Ct$  is the cycle at which an arbitrary detection threshold is crossed.

### Quantification of *NRAMP1* protein level

*NRAMP1* protein level was measured using the Human Natural Resistance-associated Macrophage Protein 1 ELISA Kit (Cat. No. E4411Hu, Bioassay Technology Laboratory, Shanghai, China) according to the manufactory instruction. A linear curve was used to measure the level of *NRAMP1* protein from a standard curve. The standard curve range of the ELISA kit is 15–3,000 ng/l.

### Statistical analysis

All experimental data were analyzed using Bio-Rad CFX™ Manager Software (version 3.1, Bio-Rad Lab. Inc., Hercules, CA) and Statistical Package for the Social Sciences (version 21.0). A statistical analysis using one-way analysis of variance differential test was carried out in order to assess the discrepancy in expression of the *NRAMP1* genes in the three study classes. *p*-values < 0.05 were considered to be statistically significant.

## RESULTS AND DISCUSSION


Of the 90 subjects studied, the number of male respondents was 38 (42.22%) and the number of female respondents was 52 (57.78%). In the active TB group, 17 out of 30 subjects were male (56.67%), and the rest were women (43.33%). In contrast to the latent group and healthy group, female subjects were more than men, as described in Table 1.

The comparison of *NRAMP1* protein level of active TB, latent TB, and healthy groups is shown in Figure 1. Based on the ELISA result, the mean protein level of *NRAMP1* was 196.41, 199.27, and 224.70 ng/ml in the TB, latent TB, and healthy groups, respectively. However, statistical test results (Kruskal-Wallis test) obtained a *p*-value of  $> 0.000$  (0.96) which showed that there was no significant difference in *NRAMP1* protein levels in either active TB, latent TB, or healthy group. It can be stated that the amount of *NRAMP1* protein will decrease if the patient has TB, compared to those who have latent TB only and the amount was even higher in normal people although it was not statistically different.

The expression of *NRAMP1* mRNA genes in the three sample groups can be seen in Figure 2. The mean result of the mRNA expression of the *NRAMP1* genes of the active TB group

**Table 1.** Characteristics of subjects in the active TB, latent TB, and healthy participant groups.

	Active TB n = 30	Latent TB n = 27	Healthy participants n = 33	Total
Gender				
Male	17 (56.7%)	10 (37%)	11 (33.3%)	38 (42.2%)
Female	13 (43.33%)	17 (63%)	22 (66.7%)	52 (57.8%)
Age group				
17–29	7 (23.33%)	7 (25.9%)	5 (15.2%)	19
30–44	10 (33.3%)	13 (48.1%)	12 (36.4%)	35
45–60	11 (36.7%)	6 (22.2%)	14 (42.4%)	31
>60	2 (6.7%)	1 (3.7%)	2 (6.1%)	5




**Figure 1.** Comparison of the mean from *NRAMP1* protein level in the active TB, latent TB, and healthy groups.

was the lowest, followed by the latent TB and healthy group. The differences in mRNA expression of the *NRAMP1* gene of active TB, latent TB, and healthy groups can be seen in Table 2. *NRAMP1* gene expression value in the healthy group was 4.61 times higher than active TB patients ( $p = 0.025$ ) and the *NRAMP1* expression value in the healthy group was 4.02 times higher than those in the latent TB group ( $p = 0.117$ ).

The study involved 90 research subjects. Of the 90 subjects, it was found that active TB patients were 56.7% male and 43.3% female. This is consistent with Uplekar's study in which 64% more men were diagnosed with TB with symptoms and positive acid-fast bacillus (AFB) screening than women which was only 34% (Uplekar *et al.*, 2001). This can happen because more men are working outdoors compared to women; thus, the risk of being exposed to *Mtb* from TB patients TB is considerably large. Active pulmonary TB is more commonly found in working men than women (Htet *et al.*, 2018). However, the biological distinctions between sexes should also be taken into account in the prevalence and diagnosis of TB (Thorson, 2015).

In this study, the active TB and latent TB samples were largely distributed in the age range of 31–45 years, with the active TB category was 33.3% and TB latent 48.1%. This was similar to research conducted by Coppola *et al.* (2020) and Hao *et al.* (2020) in which patients diagnosed with active pulmonary TB 51% and latent TB 48.6% in the age range of 27–40 years. The productive age group is more predominantly exposed to *Mtb* germs because the outdoor activities are higher, and the immunity factor at a productive age is better than the immunity of the child or elderly.

Immunoglobulin Release Assay (IGRA) has become the primary method for clinical diagnosis of a person's exposure to current *M. tuberculosis* infection (Xu *et al.*, 2018). IGRA positiveness rates are higher for individuals who have a history of close contact with active pulmonary TB patients than those in contact with pulmonary disease other than TB (Eom *et al.*, 2018). In our study, it was found that 9 out of 30 healthy people were positive for IGRA and were included in the latent TB group with a total of 27 people with a percentage of 63% more occurring in women. This is in line with the research conducted by Htet *et al.* (2018) where as much as 59.2% of people with a history of active pulmonary TB contact will suffer from TB. This is consistent with



**Figure 2.** Graph of the mean value of *NRAMP1* mRNA expression in the active TB, latent TB, and healthy groups.

**Table 2.** *NRAMP1* expression gene in the active TB, latent TB, and healthy groups.

Target	Biological group	Control	Expression	Expression 95% CI low	Expression 95% CI high	p-value
GAPDH	Latent TB					
GAPDH	Healthy	C				
GAPDH	TB					
<i>NRAMP1</i>	Latent TB		0.24970	0.07341	0.84928	0.117868
<i>NRAMP1</i>	Healthy	C	1.00000	0.29127	3.43322	
<i>NRAMP1</i>	TB		0.21752	0.12723	0.37188	0.025601

the results of our study, where 18 of 30 (60%) household contacts were positive for IGRA, while the rest were negative for IGRA. People who were negative for IGRA can be considered a healthy individual because, despite the history of contact with active pulmonary TB patients, the IGRA results remained negative. It may be due to the fact that the intensity of the exposure was low, or the number of germs that penetrate the body was below the threshold for sensitivity to the applied IGRA method.

A previous study showed that there was a lack or absence of *NRAMP1* expression in macrophages or Langerhans' giant cells found in granulomatous lesions of in Taiwanese indigenous relative to Han subjects, and there was a positive correlation of *NRAMP1* gene polymorphism to tuberculosis susceptibility in Taiwanese indigenous people (Hsu *et al.*, 2006). Gallant *et al.* (2007) also suggested that, in pediatric TB patients with homozygous CC genotype *NRAMP1* 1–274, lower *NRAMP1* activity was found in monocyte-derived macrophages with elevated risk allele than in heterozygous CT genotype patients. Lack of *NRAMP1* protein expression occurred in mouse strains prone to *Mycobacterium bovis* BCG infection (Vidal *et al.*, 1996). However, in bovine TB, *NRAMP1* was high in peripheral blood mononuclear cells; alveolar macrophages (obtained by bronchioalveolar lavage) and Holsten Friesian cattle lymphatic node granulomas with self-proven bovine TB were highly expressed (Estrada-Chávez *et al.*, 2001; Pereira-Suárez *et al.*, 2006). Human and mouse *NRAMP1* proteins share 88% identity and 93% overall sequence similarity, with the predicted N-linked glycosylation signals being precisely conserved in the fourth extracellular loop of both proteins (Vidal *et al.*, 1996).

In our study, we found that the *NRAMP1* expression gene was downregulated in TB compared to latent TB and healthy individuals. Also, the *NRAMP1* protein level in TB is lower than those with latent TB and healthy participants. The decreased *NRAMP1* action may contribute to increased bacterial availability of iron, fostering the replication of mycobacteria in macrophages. *NRAMP1* pumps iron, essential to live from the macrophages via intracellular bacteria like Mtb. Since iron is also essential for the cell to produce reactive oxygen and intermediate nitrogen, loss of intramacrophage iron availability for bacterium and simultaneous weakening of antimicrobial activities of the Fe<sup>2+</sup> ion transport role of the *NRAMP1* protein could thus facilitate infection by Mtb and tuberculosis development (Fernández-Mestre *et al.*, 2015). Many studies have investigated the correlation between *NRAMP1* gene polymorphisms and tuberculosis (Fernández-Mestre *et al.*, 2015; Meilang *et al.*, 2012; Stagas *et al.*, 2011), and the stability and efficiency of mRNA protein translation of these polymorphisms can impact.

## CONCLUSION

*NRAMP1* expression is downregulated and protein level was lower in active TB than latent TB and healthy individuals. Our results demonstrated that *NRAMP1* could contribute to the pathogenesis of TB. This study confirms that *NRAMP1* gene expression correlated with the TB infection.

## ACKNOWLEDGMENTS

This research was supported by an internal grant from Institute for Research and Community Services (Lembaga Penelitian dan Pengabdian Masyarakat) Universitas Hasanuddin (Grant No. UNHAS-LP2M: 12286/UN4.3.2/PL.00.00/2018, Date June 22, 2018; LP2M-Researcher: 3083/UN4.21/PL.00.00/2018, Date June 26, 2018). The authors also thank Community Center for Lung Health Makassar or Balai Besar Kesehatan Paru Masyarakat (BBKPM) Makassar staff for assisting us to obtain the samples in this research. The authors gratefully acknowledge the technician in the Tuberculosis Unit of the HUM-RC Laboratory HUM-RC for helping us in the culture TB examination.

## AUTHOR CONTRIBUTIONS

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; agreed to submit to the current journal; gave final approval of the version to be published; and agree to be accountable for all aspects of the work. All the authors are eligible to be an author as per the international committee of medical journal editors (ICMJE) requirements/guidelines.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

## ETHICAL APPROVALS

This research had ethical approval by the Research Ethics Commission of the Faculty of Medicine, Universitas Hasanuddin, Makassar, South Sulawesi, Indonesia (No. 1125/UN4.6.4.5.31/PP36/2020).

## PUBLISHER'S NOTE

This journal remains neutral with regard to jurisdictional claims in published institutional affiliation.

## REFERENCES

Canonne-Hergaux F, Gruenheid S, Govoni G, Gros P. The *Nramp1* protein and its role in resistance to infection and macrophage function. *Proc Assoc Am Physicians*, 1999; 111(4):283–9.

- Coppola M, Villar-Hernández R, Van Meijgaarden KE, Latorre I, Muriel Moreno B, Garcia-Garcia E, Franken KLMC, Prat C, Stojanovic Z, De Souza Galvão ML. Cell-mediated immune responses to *in vivo*-expressed and stage-specific mycobacterium tuberculosis antigens in latent and active tuberculosis across different age groups. *Front Immunol*, 2020; 11:103.
- Dow CT. Mycobacterium avium subspecies paratuberculosis and human disease: bridging infection and autoimmunity. In: Shoenfeld Y, Agmon-Levin N, Rose N (eds.). *Infection and autoimmunity*. 2nd edition, Academic Press, Amsterdam, The Netherlands, pp 569–81, 2015.
- Eom JS, Kim I, Kim WY, Jo EJ, Mok J, Kim MH, Lee K, Kim KU, Park HK, Lee MK. Household tuberculosis contact investigation in a tuberculosis-prevalent country. *Medicine (Baltimore)*, 2018; 97(3):1–6.
- Estrada-Chávez C, Pereira-Suárez AL, Meraz MA, Arriaga C, García-Carrancá A, Sánchez-Rodríguez C, Mancilla R. High-level expression of *NRAMP1* in peripheral blood cells and tuberculous granulomas from *Mycobacterium bovis*-infected bovines. *Infect Immun*, 2001; 69(11):7165–8.
- Fernández-Mestre M, Villasmil Á, Takiff H, Fuentes Alcalá Z. *NRAMP1* and *VDR* gene polymorphisms in susceptibility to tuberculosis in venezuelan population. *Dis Markers*, 2015; 860628:1–7.
- Flaherty DK. Delayed-type hypersensitivity reactions. In: Flaherty DK (ed.). *Immunology for pharmacy*. Mosby, Saint Louis, MO, pp 152–61, 2012.
- Gallant CJ, Malik S, Jabado N, Cellier M, Simkin L, Finlay BB, Graviss EA, Gros P. Reduced *in vitro* functional activity of human *NRAMP1* (SLC11A1) allele that predisposes to increased risk of pediatric tuberculosis disease. *Genes Immunity*, 2007; 8:691–8.
- Hanikenne M, Merchant SS, Hamel P. Transition metal nutrition: a balance between deficiency and toxicity. In: Harris EH, Stern DB, Witman G (eds.). *The chlamydomonas sourcebook*. Academic Press, London, UK, pp 333–99, 2009.
- Hao X, Bai J, Ding Y, Wang J, Liu Y, Yao L, Pan W. Characterization of antibody response against 16kD and 38kD of M. tuberculosis in the assisted diagnosis of active pulmonary tuberculosis. *Ann Transl Med*, 2020; 8(15):945.
- Hsu YH, Chen CW, Sun HS, Jou R, Lee JJ, Su JJ. Association of *NRAMP 1* gene polymorphism with susceptibility to tuberculosis in Taiwanese aboriginals. *J Formos Med Assoc*, 2006; 105(5):363–9.
- Htet KKK, Liabsuetrakul T, Thein S, McNeil EB, Chongsuvivatwong V. Improving detection of tuberculosis among household contacts of index tuberculosis patients by an integrated approach in Myanmar: a cross-sectional study. *BMC Infect Dis*, 2018; 18(1):660.
- Meilang Q, Zhang Y, Zhang J, Zhao Y, Tian C, Huang J, Fan H. Polymorphisms in the SLC11A1 gene and tuberculosis risk: a meta-analysis update. *Int J Tuberc Lung Dis*, 2012; 16(4):437–46.
- Murray MF. Susceptibility and response to infection. In: Rimoin D, Pyeritz R, Korf B (eds.). *Emery and rimoin's principles and practice of medical genetics*. 6th edition, Academic Press, Oxford, UK, pp 1–24, 2013.
- Pereira-Suárez AL, Estrada-Chávez C, Arriaga-Díaz C, Espinosa-Cueto P, Mancilla R. Coexpression of *NRAMP1*, iNOS, and nitrotyrosine in bovine tuberculosis. *Vet Pathol*, 2006; 43(5):709–17.
- Qidwai T, Jamal F, Khan MY. DNA sequence variation and regulation of genes involved in pathogenesis of pulmonary tuberculosis. *Scand J Immunol*, 2012; 75(6):568–87.
- Stagas MK, Papaetis GS, Orphanidou D, Kostopoulos C, Syriou S, Reczko M, Drakoulis N. Polymorphisms of the *NRAMP1* gene: distribution and susceptibility to the development of pulmonary tuberculosis in the Greek population. *Med Sci Monit*, 2011; 17(1):1–6.
- Thorson A. Gender issues in tuberculosis. In: Klein SL, Roberts, CW (eds.). *Sex and gender differences in infection and treatments for infectious diseases*. Springer, Cham, Switzerland, 2015, pp 231–53.
- Uplekar MW, Rangan S, Weiss MG, Ogden J, Borgdorff MW, Hudelson P. Attention to gender issues in tuberculosis control [Unresolved Issues]. *Int J Tuberc Lung Dis*, 2001; 5(3):220–4.
- Vidal S, Pinner E, Lepage P, Gauthier S, Gros P. Natural resistance to intracellular infections: *Nramp1* encodes a membrane phosphoglycoprotein absent in macrophages from susceptible (*Nramp1* D169) mouse strains. *J Immunol*, 1996; 157:3559–68.
- Xu J, Li Z, Chen X, Shi C, Wu M, Chen H, Zhu X, Song H, Wu M, Xu P. More significance of TB-IGRA except for the diagnose of tuberculosis. *J Clin Lab Anal*, 2018; 32(1):e22183.
- Zhao N, Enns CA. Iron transport machinery of human cells: players and their interactions. *Curr Top Membr*, 2012; 69:67–93.

**How to cite this article:**

Handayani I, Djaharuddin I, Natzir R, Arief M, Ahmad A, Hatta M, Agus R, Patellongi I, Amin M, Widaningsih Y, Halik H, Hidayah N, S Subair N, Leman Y, Waworuntu W, Massi MN. Contribution of *NRAMP1* gene expression and protein level in pulmonary and latent TB infection in Indonesia. *J Appl Pharm Sci*, 2021; 11(08):017–021.

# Contribution of NRAMP1 gene expression and protein level in pulmonary and latent TB infection in Indonesia

## ORIGINALITY REPORT

**21** %  
SIMILARITY INDEX

**12** %  
INTERNET SOURCES

**17** %  
PUBLICATIONS

**4** %  
STUDENT PAPERS

## PRIMARY SOURCES

**1** Submitted to Queen Mary and Westfield College  
Student Paper **2** %

**2** link.springer.com  
Internet Source **1** %

**3** new.esp.org  
Internet Source **1** %

**4** onlinelibrary.wiley.com  
Internet Source **1** %

**5** A.C.C.S. Leandro, M.A. Rocha, C.S.A. Cardoso, M.G. Bonecini-Almeida. "Genetic polymorphisms in vitamin D receptor, vitamin D-binding protein, Toll-like receptor 2, nitric oxide synthase 2, and interferon- $\gamma$  genes and its association with susceptibility to tuberculosis", Brazilian Journal of Medical and Biological Research, 2009  
Publication **1** %

6

Chuan Wang. "Comparative miRNA Expression Profiles in Individuals with Latent and Active Tuberculosis", PLoS ONE, 10/07/2011

Publication

1 %

7

tos.org  
Internet Source

1 %

8

Mercedes Fernández-Mestre, Ángel Villasmil, Howard Takiff, Zhenia Fuentes Alcalá. " and Gene Polymorphisms in Susceptibility to Tuberculosis in Venezuelan Population ", Disease Markers, 2015

Publication

1 %

9

Coad Thomas, Leonardo A.. "Chapter 11 Environmental Triggers of Type 1 Diabetes Mellitus – Mycobacterium Avium Subspecies Paratuberculosis", IntechOpen, 2011

Publication

1 %

10

Valeria Sargentini, Sabrina Mariotti, Stefania Carrara, Maria Cristina Gagliardi et al. "Cytometric detection of antigen-specific IFN- $\gamma$ /IL-2 secreting cells in the diagnosis of tuberculosis", BMC Infectious Diseases, 2009

Publication

1 %

11

0-crossmark-crossref-org.brum.beds.ac.uk  
Internet Source

1 %

[www.spandidos-publications.com](http://www.spandidos-publications.com)

12

Internet Source

1 %

13

M R Milone, B Pucci, F Bruzzese, C Carbone, G Piro, S Costantini, F Capone, A Leone, E Di Gennaro, M Caraglia, A Budillon. "Acquired resistance to zoledronic acid and the parallel acquisition of an aggressive phenotype are mediated by p38-MAP kinase activation in prostate cancer cells", Cell Death & Disease, 2013

Publication

1 %

14

[hdl.handle.net](http://hdl.handle.net)

Internet Source

1 %

15

[journals.lww.com](http://journals.lww.com)

Internet Source

1 %

16

[membranetransport.org](http://membranetransport.org)

Internet Source

1 %

17

Submitted to Indiana University

Student Paper

&lt;1 %

18

Angel H. Alvarez. "Revisiting tuberculosis screening: An insight to complementary diagnosis and prospective molecular approaches for the recognition of the dormant TB infection in human and cattle hosts", Microbiological Research, 2021

Publication

&lt;1 %

- 19 Barokatun Hasanah, Deon Arinaldo, Sepanya Pasaribu, Endon Bharata, Mohammad Ridwan Effendi, Arief Budi Santiko, Achmad Munir. "Verification Design of Up-Down Frequency Converter Based on Double-Balanced Mixer and Its Performance Characterization", 2019 IEEE 13th International Conference on Telecommunication Systems, Services, and Applications (TSSA), 2019  
Publication <1 %
- 
- 20 [www.researchsquare.com](http://www.researchsquare.com)  
Internet Source <1 %
- 
- 21 Nur Hidayati, Tintin Sukartini, Padoli Padoli. "COMPARISON OF TWO TURNING REGIMENS (1 VERSUS EVERY 2 HOURS) IN THE PREVENTION OF CONSTIPATION IN PATIENTS WITH STROKE", Belitung Nursing Journal, 2019  
Publication <1 %
- 
- 22 [pubs.nal.usda.gov](http://pubs.nal.usda.gov)  
Internet Source <1 %
- 
- 23 [www.labome.org](http://www.labome.org)  
Internet Source <1 %
- 
- 24 [www.omicsonline.org](http://www.omicsonline.org)  
Internet Source <1 %
- 
- 25 K M Ibrahim, J A Barrett. "Evolution of mildew resistance in a hybrid bulk population of <1 %

## barley", Heredity, 1991

Publication

26

[aok.pte.hu](http://aok.pte.hu)

Internet Source

<1 %

27

[journals.plos.org](http://journals.plos.org)

Internet Source

<1 %

28

[zagan.unizar.es](http://zagan.unizar.es)

Internet Source

<1 %

29

Sarah Haig, Nadine Kotlarz, Linda Kalikin, Thomas Chen, Seth David Guikema, John LiPuma, Lutgarde Raskin. "Emerging investigator series: Bacterial Opportunistic Pathogen Gene Markers in Municipal Drinking Water are Associated with Distribution System and Household Plumbing Characteristics", Environmental Science: Water Research & Technology, 2020

Publication

<1 %

30

[mro.massey.ac.nz](http://mro.massey.ac.nz)

Internet Source

<1 %

31

"THE GREAT MIMICKER: PULMONARY TUBERCULOSIS WITH AXILLARY MASS",  
Respirology, 2018

Publication

<1 %

32

MICAH J. DRUMMOND. "Human Muscle Gene Expression following Resistance Exercise and

<1 %

Blood Flow Restriction", Medicine & Science in Sports & Exercise, 04/2008

Publication

---

33

P Domenech, A.S. Pym, M Cellier, C.E. Barry, S.T. Cole. " Inactivation of the orthologue ( ) does not affect virulence in a mouse model of tuberculosis ", FEMS Microbiology Letters, 2002

Publication

---

<1 %

34

Upendra Yadav, Pradeep Kumar, Vandana Rai. "FokI polymorphism of the vitamin D receptor (VDR) gene and susceptibility to tuberculosis: Evidence through a meta-analysis", Infection, Genetics and Evolution, 2021

Publication

---

<1 %

35

Yolken, R.H.. "Infectious agents and gene-environmental interactions in the etiopathogenesis of schizophrenia", Clinical Neuroscience Research, 200608

Publication

---

<1 %

36

A.Ejah Umraeni Salam, Muh Tola, Mary Selintung, Farouk Maricar. "A leakage detection system on the Water Pipe Network through Support Vector Machine method", 2014 Makassar International Conference on Electrical Engineering and Informatics (MICEEI), 2014

Publication

---

<1 %

37 Nurhayani Jamil, Werna Nontji, Syafruddin Syarif, Mardiana Ahmad, Suryani As'ad, Azniah Syam, Andi Syarifah Irmadani. "Handling Contraceptive Acceptors' Complaints by Using Android-Based Application", Global Journal of Health Science, 2019  
Publication

---

38 Omics for Personalized Medicine, 2013.  
Publication

---

39 Serene H. John, John Kenneth, Archana S. Gandhe. "Host biomarkers of clinical relevance in tuberculosis: review of gene and protein expression studies", Biomarkers, 2011  
Publication

---

40 [ijdvl.com](http://ijdvl.com)  
Internet Source

---

41 [journals.physiology.org](http://journals.physiology.org)  
Internet Source

---

42 [www.journal.unair.ac.id](http://www.journal.unair.ac.id)  
Internet Source

---

43 V. Berridge. "The importance of the past in public health", Journal of Epidemiology & Community Health, 2004  
Publication

---

---

Exclude quotes      On

Exclude matches      < 5 words

Exclude bibliography      On