

onal_6-
_Asian_Jr_of_Microbiol_Biotech_
_Env_Sc_-Sartini-_20143.pdf
by

FILE	ONAL_6-_ASIAN_JR_OF_MICROBIOL_BIOTECH_ENV_SC_-SARTINI-_20143.PDF (181.93K)	WORD COUNT	2129
TIME SUBMITTED	13-FEB-2021 01:03PM (UTC+0700)	CHARACTER COUNT	11811
SUBMISSION ID	1508589118		

ISOLATION AND SCREENING FUNGAL SYMBIONT IN GREEN ALGA *ULVA RETICULATA* AS CANDIDATE OF ANTIBIOTIC PRODUCER

SARTINI*, SURYADI AND ELMI N. ZAINUDDIN¹

*Faculty of Pharmacy Hasanuddin University, Indonesia

¹Faculty of Fishery and Marine Science Hasanuddin University, Indonesia

(Received 15 February, 2014; accepted 16 March, 2014)

Key words : Fungal symbiont, *Ulva reticulata*, Antibiotic producer

Abstract - The isolation of fungal symbiont from *Ulva reticulata* by direct planting method on Potato Dextrose Agar plate was performed. Preliminary screening of the fungal symbionts as a candidate of antibiotic producer involve²⁴ fermentation process in Potato Dextrose Broth supplemented with 0.5 mg % yeast extract using shaker at 120 rpm for 7 days. A²³ microbial activity of fermentation supernatants was examined using disc agar diffusion method against *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, *Candida albicans* and *Malassezia furfur*. We found that three isolates, identified by macroscopic and microscopic characteristics, were molds genus *Aspergillus*, *Penicillium* and *Cladosporium* and one of them was unidentified fungi. Supernatant of *Aspergillus*, *Penicillium*, *Cladosporium* and the unidentified fungi demonstrated a high antifungal activity against *C. albicans*, verified by the diameter zone of inhibitions of 28.04 mm, 24.98 mm, 25.38 mm, and 27.73 mm, respectively. Only supernatant of the unidentified fungi had antifungal activity against *M. furfur*. A high antibacterial activity against *S. aureus* was shown by the supernatant of *Penicillium* and *Chlad²⁶rium* with diameter zone of inhibitions of 19.15 mm and 16.56 mm, respectively. All isolates had low antibacterial activity against *E. coli* and *S. thyposa*. Fungal symbionts in *Ulva reticulata* from Takalar, South Sulawesi are potential as a candidate of antibiotic producer, especially antibacterial activity against *S. aureus* and *C. albicans*.

INTRODUCTION

Many essential compounds act as antibacterial and antifung⁷ derived from seaweed or marine algae. Marine algae are classified as Rhodophyta (red algae), Phaeo-phyta (brown algae) and Chlorophyta (green algae) (Chanda *et al.*,¹⁶ 0). Some of green algae was more active than other groups of a³ screened for their antibacterial activity and they are potential sources of bioactive compounds²² and should be investigated for natural antibiotics (Kandhasamy and Arunachalam, 2008; Osman *et al.*,⁶ 010). For example, Green algae *Ulva reticulata* and *Ulva lactuca* extracts showed promising antimicrobial activity against bacterial and fungal human pathogens (Kolanjinathan and Stella, 2011). *Ulva fasciata* contains sterol alkaloid, phenolic, flavonoid, terpenoid, glycoside had width

spectrum antibacterial activity (Premalatha *et al.*, 2011; Selvin and Lipton, 2004)

Marine algae live in symbiosis with certain micro¹ganisms such as fungi and bacteria. Epiphytic and endophytic fungi live on the surface and in the inner tissues or even in the c¹⁰ of their hosts, respectively. It is believed that algae and their associated microbial symbionts should represent a good source²¹ of biologically active secondary metabolites (S¹²ulz *et al.* 2008; Suryanarayanan *et al.* 2010). For example, *Ascochyta salicorniae*, an endophytic and obligate marine fungus was isolated¹ from the green algae *Ulva sp.* (Claudia *et al.*, 2000). New marine derived antibiotics, 43 epi- and endophytic fungal strains were also isolated from the surface or the inner tissue of different marine plants and invertebrates. Through preliminary and secondary screening, 10 of them were found to be

*Corresponding author's email : sardj@yahoo.com

able to produce broad-spectrum antimicrobial metabolites (Zhang *et al.*, 2009).

The purpose of this study was to search for antibiotic-producing fungal symbiont in green alga *Ulva reticulata* from Takalar South Sulawesi, Indonesia. In this paper, we report the isolation, identified the fungi by macroscopic and microscopic characteristic and preliminary screening of fungal symbiont from *Ulva reticulata* as antibiotic producer and by diffusion agar assay using several microbial test.

MATERIALS AND METHODS

Collection and Preparation Green Alga

The green alga *Ulva reticulata* were collected from Punaga Coast Takalar Regency, South Sulawesi. Immediately after collection, they were washed in fresh seawater to remove the sand and other extraneous matter. The alga were transported to the laboratory in cool box.

Isolation of Fungal Symbiont

The alga were divided two groups: first group was not subjected to surface sterilization, while second group was subjected to surface sterilization according to Flewelling method with modification (11). The algae was rinsed with sterile sea water, followed by immersion in 1 % sodium hypochlorite for 1 minutes and rinsed again with sterile sea water. After surface sterilization, fragments of approx. 1 cm x 1 cm from sterile algae were inoculated on Petri dishes containing PDA supplemented with chloramphenicol 50 mg/L to suppress bacterial growth. All the plates were incubated at room temperature for up to 12 days. The fungi were transferred to fresh PDA plates, incubated for 1 week and periodically checked for purity. Identification were done by macroscopic and microscopic characteristic.

Production of bioactive compound by submerged fermentation

Production of antimicrobial compound was conducted according to Radu method with modification (2003). The endophytic fungi were fermented with a submerged fermentation with shaking 120 rpm for 7 days using PD-Y medium (2.4 % Potato Dextrose Broth, 0.5 % Yeast extract, 1 % NaCl). After incubation, the media were sonicated for 5 minutes and centrifuged at 3000 rpm for 15

minutes. The supernatant was used as sample test.

Antimicrobial assay by agar diffusion methods

Antibacterial activity was evaluated by agar diffusion method. Supernatant was pipetted 20 µL onto paper disc (diameter 6 mm), each disc was placed on Muller Hinton Agar (pH 7.2) which contained 1 mL bacteria suspension (*Bacillus subtilis*, *Escherichia coli*, *Staphylococcus aureus* and *Salmonella typhi*) and incubated for 24 hours at 37°C. Inhibition results are expressed as diameter of the clear halo surrounding each disc on cultivated agar plates. Tetracycline antibiotic 30 ppm was used as positive control. Antifungal activity was tested using similar method using Potato Dextrose Agar and fungal suspension (*Candida albicans* and *Malessezia furfur*). Incubation time was 24 -48 h at room temperature.

RESULTS AND DISCUSSION

Isolation of Fungal Symbiont

A total of 4 fungal symbiont were obtained from Punaga Coast Takalar Regency, South Sulawesi. The macroscopic and microscopic characteristics are shown in Figure 1.

Strain FSUr-1, FSUr-2, FSUr-3 were identified as *Genus Aspergillus*, *Penicillium*, *Chladosporium*, respectively. FSUr-4 was unidentified. Several studies found that genus *Aspergillus* and *Penicillium* presented in fungal symbiont in marine alga. The genus *Aspergillus* has been known to be a major contributor to the secondary metabolites of marine fungal origin, for example, genus *Aspergillus* has been isolated by Zhang *et al.* (2007) from marine brown alga *Colpomenia sinuosa* similarly, Gamal-Elden *et al.* (2009) has isolated fungal endophyte from green alga *Ulva* sp, which was identified as *Penicillium* sp.. Zhu *et al.* (2009) has also isolated *Penicillium*, from green alga *Blidingia minima*. Gao *et al.* (2011) found that *Penicillium chrysogenum* QEN-24S, an endo-phytic fungus isolated from an unidentified marine red algal species of the genus *Laurencia*, displayed inhibitory activity against the growth of pathogen *Alternaria brassicae* in dual culture test.

Antimicrobial activity of the isolates

Antimicrobial activity obtained from fermentation supernatant of 4 fungal symbionts was examined by disc agar diffusion method against *Staphylococcus*




Fig. 1 Macroscopic and microscopic characteristic of fungal symbiont isolated from *Ulva reticulata*
A. Colony of Culture FSUr-1, FSUr-2, FSUr-3, FSUr-4 grown on PDA for seven days, respectively
B. Microscopic characteristic (conidia and conidiaphores) of FSUr-1, FSUr-2, FSUr-3, respectively

Table 1. Antimicrobial activity of supernatant of Symbiont fungi from green Alga *Ulva reticulata*, in the agar diffusion assay

Sample / (mm) Supernatant of isolates	Diameter zone of inhibition					
	Bacteria			Fungi		
	<i>B.subtilis</i>	<i>S.aureus</i>	<i>E.coli</i>	<i>S.thypi</i>	<i>C.albicans</i>	<i>M.furfur</i>
FSUr-1	11.53±0.6	9.70± 0.33	10.08± 0.86	8.72± 0.42	28.04± 1.26	7.47± 0.52
FSUr-2	22.03± 0.83	19.15± 1.22	15.08± 0.91	13.54± 0.9	24.98± 1.15	10.56± 1.24
FSUr-3	19.95± 0.81	16.56± 1.37	13.74± 0.78	13.34± 0.15	25.38± 1.59	15.16± 1.01
FSUr-4	12.2± 0.07	10.97± 0.53	12.74± 0.75	11.71± 0.05	27.73± 1.72	17.49± 0.94
Chloramphenicol			18.36± 0.77	9.59± 0.09		
Tetracycline	22.01 ± 1.18	11.89 ± 1.47				
Nystatin					8.95 ± 1.30	7.91 ± 0.23

¹ Inhibition zone in mm including disc, expressed as Mean ± SD (n = 3);

aureus, *Escherichia coli*, *Bacillus subtilis*, *Candida albicans* and *Malassezia furfur* The result shows in Table 1 and Fig. 2.

Supernatant of *Aspergillus*, *Penicillium*, *Cladosporium* and the unidentified mold against *C. albicans*, demonstrated a high antifungal activity, verified by the diameter zone of inhibitions of 28.04 mm, 24.98 mm, 25.38 mm, and 27.73 mm, respectively. A high antibacterial activity against *S.aureus* was showed by the supernatant of *Penicillium* and

Cladosporium with diameter zone of inhibition were 19.15 mm and 16.56 mm, respectively. All isolates had low antibacterial activity against *E. coli* and *S. thyposa*. Only supernatant of the unidentified fungi had antifungal activity against *M. furfur*. According to Ibbitassam *et al.* (2009), the antibacterial activity was classified from less active which was (10 mm < diameter of inhibition < 16 mm), to moderately active (16 mm < diameter of inhibition < 20 mm), to highly active (diameter of

inhibition > 27 mm) and they found *Ulva* sp had antibacterial activity against *S. aureus* s higher than *E. coli*.

CONCLUSION

We found four fungal symbiont from green alga *Ulva reticulata* from Punaga Coast Takalar Re-gency, South Sulawesi. Three of them (FSUr-1, FSUr-2, FSUr-3) are potential as candidates for of antibiotic producer.

ACKNOWLEDGEMENT

This article has been orally presented at 5th International Seminar of Indonesian Society for Microbiology Indonesian Society for Microbiology, Manado, 2012.

REFERENCES

- Chanda, S., Dave, R., Kaneria, M. and Nagani, K. 2010. Seaweeds: A novel, untapped source of drugs from sea to combat Infectious diseases. Ed. A. Mendez Vilas, In : *Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology*, pp. 473-480.
- Gamal-Eldeen, A.M., Abdel-Lateff, A. and Okino, T. 2009. Modulation of carcinogen metabolizing enzymes by chromanone A; a new chromone derivative from algicolous marine fungus *Penicillium* sp. *Environ Toxicol Phar*. 28 : 317-322.
- Gao, S.S., Li, X.M., Zhang, Y., Li, C.S. and Wang, B.G. 2011a. Conidiogenones H. and I., two new diterpenes of cyclopiane class from a marine-derived endophyticfungus *Penicillium chrysogenum* QEN-24S. *Chem. Biodivers*. 8 : 1748-1753.
- Gao, S.S., Li, X.M., Du, F.Y., Li, C.S., Proksch, P. and Wang, B.G. 2011b. Secondary metabolites from a marine-derived endophytic fungus *Penicillium chrysogenum* QEN-24S. *Mar Drugs* 9: 59-70.
- Ibtissam, C., Hassane, R., Jose, M.L., Fransisco, D.S.J., Antonio, G.V.J., Hassan, B., dan Mohamed, K. 2009. Screening of antibacterial activity in marine green and brown macroalgae from the coast of morocco. *African Journal of Biotechnology*. 8 (7) : 1258-1262.
- Kandhasamy, M. and Arunachalam, K.D. 2008. Evaluation of in vitro antibacterial property of seaweeds of southeast coast of India. *African Journal of Biotechnology*. 7 (12) : 1958-1961.
- Kolanjinathan, K. and Stella, D. 2011. Comparative studies on antimicrobial activity of *Ulva reticulata* and *Ulva lactuca* against human pathogens. *International Journal of Pharma-ceutical & Biological Archives*. 2 (6) : 1738-1744.
- Osman, M.E.H., Abushady, A.M. and Elshobary, M.E. 2010. *In vitro* screening of antimicrobial activity of extracts of some macroalgae collected from Abu-Qir bay Alexandria, Egypt. *African Journal of Biotechnology*. 9 (12) : 7203-7208.
- Premalatha, M., Dhasarathan, P. and Theriappan, P. 2011. Characterization and antimicrobial efficiency of seaweed samples, *Ulva fasciata* and *Chaetomorpha antennina*. *International Journal Of Pharma And Bio Sciences*. 2 (1) : 288-293.
- Selvin, J. and Lipton, A.P. 2004. Biopotentials of *Ulva fasciata* and *Hypnea musciformis* Col-lected From The peninsular coast of India. *Journal of Marine Science and Technology*. 12 (1) : 1-6.
- Schulz, B., Siegfried, D., dela Cruz, T.E., Rheinheimer, J., Siems, K., Loesgen, S. et al. 2008. Screening strategies for obtaining novel, biologically active, fungal secondary metabolites from marine habitats. *Bot. Mar*. 51 : 219-234.
- Suryanarayanan, T.S., Venkatachalam, A., Thirunavukkarasu, N., Ravishankar, J.P., Doble, M., and Geetha, V. 2010. Internal mycobiota of marine macroalgae from the Tamilnadu coast: distribution, diversity and biotechnological potential. *Bot Mar*. 53 : 457-468.
- Zhang, Y. Mu, J., Feng, Y., Kang, Y., Zhang, J. Gu, P.J., Wang, Y., Ma, L.F. and Zhu, Y.H. 2009. Broad-Spectrum antimicrobial epiphytic and endophytic fungi from marine organisms: isolation, bioassay and taxonomy. *Mar. Drugs*. 7 : 97-112.
- Zhu, T.J., Du, L., Hao, P.F., Lin, Z.J. and Gu, Q.Q. 2009. Citrinal A, a noveltricyclic derivative of citrinin, from an algicolous fungus *Penicillium* sp. i-1-1. *Chinese Chem Lett*. 20 : 917-920.

ORIGINALITY REPORT

% **25**
SIMILARITY INDEX

% **18**
INTERNET SOURCES

% **22**
PUBLICATIONS

% **2**
STUDENT PAPERS

PRIMARY SOURCES

1 www.mdpi.com % **7**
Internet Source

2 www.aensiweb.com % **1**
Internet Source

3 K M, hasamy, D Arunachalam K. "Evaluation of
in vitro antibacterial property of seaweeds of
southeast coast of India", African Journal of
Biotechnology, 2008 % **1**
Publication

4 Peihang Xu, Huaqiang Tan, Weiguang Jin,
Yanfei Li, C. Santhoshkumar, Ping Li, Wenhua
Liu. "Antioxidative and antimicrobial activities of
intertidal seaweeds and possible effects of
abiotic factors on these bioactivities", Journal of
Oceanology and Limnology, 2018 % **1**
Publication

5 studentsrepo.um.edu.my % **1**
Internet Source

6

Internet Source

% 1

7

Submitted to University of Wales Swansea

Student Paper

% 1

8

A. Noorjahan, B. Aiyamperumal, P. Anantharaman. "Isolation and Charecterisation of Seaweed Endophytic Fungi as an Efficient Phosphate Solubiizers", Biosciences, Biotechnology Research Asia, 2019

Publication

% 1

9

mdpi.com

Internet Source

% 1

10

www.koreascience.or.kr

Internet Source

% 1

11

link.springer.com

Internet Source

% 1

12

macau.uni-kiel.de

Internet Source

% 1

13

www.omicsonline.org

Internet Source

% 1

14

Seshagiri Raghukumar. "Fungi in Coastal and Oceanic Marine Ecosystems", Springer Science and Business Media LLC, 2017

Publication

% 1

Marine Vallet, Martina Strittmatter, Pedro Murúa,

15

Sandrine Lacoste et al. "Chemically-Mediated Interactions Between Macroalgae, Their Fungal Endophytes, and Protistan Pathogens", *Frontiers in Microbiology*, 2018

Publication

% 1

16

Samkari , Sakaa Talal Hassan. "Antimicrobial Activity of Marine Macroalgae from the Coast of Jeddah, Saudi Arabia = فعالية المضادات الميكروبية للطحالب البحرية الكبيرة من ساحل مدينة جدة، المملكة العربية السعودية", King Abdulaziz University : Scientific Publishing Centre, 2017

Publication

% 1

17

N B Prihantini. "Morphological identification, isolation, and culturing of cyanobacteria derived from hot spring of Cisolok and Galunggung Mountain based on enrichment method", *Journal of Physics: Conference Series*, 2020

Publication

<% 1

18

www.jmbfs.org

Internet Source

<% 1

19

Kossuga, Miriam H., Stelamar Romminger, Camila Xavier, Marília C. Milanetto, Milene Z. do Valle, Eli F. Pimenta, Raquel P. Morais, Erica de Carvalho, Carolina M. Mizuno, Luís Fernando C. Coradello, Vinícius de M. Barroso, Bruna Vacondio, Darci C. D. Javaroti, Mirna H. R. Selegim, Bruno C. Cavalcanti, Claudia Pessoa,

<% 1

Manoel O. Moraes, Bruna A. Lima, Reginaldo Gonçalves, Rafaella C. Bonugli-Santos, Lara D. Sette, and Roberto G. S. Berlinck. "Evaluating methods for the isolation of marine-derived fungal strains and production of bioactive secondary metabolites", *Revista Brasileira de Farmacognosia*, 2012.

Publication

20

Hanaa H. Ahmed, Muhammad M. Hegazi, Howaida I. Abd-Allac, Emad F. Eskander, Mona S. Ellithey. "Antitumour and Antioxidant Activity of Some Red Sea Seaweeds in Ehrlich Ascites Carcinoma in vivo", *Zeitschrift für Naturforschung C*, 2011

Publication

21

Trichur S. Suryanarayanan, Ambayeram Venkatachalam, Nagamani Thirunavukkarasu, Jagadesan P. Ravishankar et al. "Internal mycobiota of marine macroalgae from the Tamilnadu coast: distribution, diversity and biotechnological potential", *Botanica Marina*, 2010

Publication

22

www.iasj.net

Internet Source

23

"Microbial Biotechnology", Springer Science and Business Media LLC, 2018

Publication

<% 1

<% 1

<% 1

<% 1

24

Aseer Manilal, Sugathan Sujith, Balu Sabarathnam, George Kiran, Joseph Selvin, Chippu Shakir, Aaron Lipton. "Biological activity of the red alga *Laurencia brandenii*", *Acta Botanica Croatica*, 2011

Publication

<% 1

25

Zhang, Yi, Jun Mu, Yan Feng, Yue Kang, Jia Zhang, Peng-Juan Gu, Yu Wang, Li-Fang Ma, and Yan-Hua Zhu. "Broad-Spectrum Antimicrobial Epiphytic and Endophytic Fungi from Marine Organisms: Isolation, Bioassay and Taxonomy", *Marine Drugs*, 2009.

Publication

<% 1

26

moam.info

Internet Source

<% 1

27

Deshmukh, Sunil K., Shilpa A. Verekar, and Sarita V. Bhave. "Endophytic fungi: a reservoir of antibacterials", *Frontiers in Microbiology*, 2015.

Publication

<% 1

28

mafiadoc.com

Internet Source

<% 1

EXCLUDE QUOTES ON

EXCLUDE ON

EXCLUDE MATCHES < 5

WORDS

BIBLIOGRAPHY