

ional_11-
_International_Journal_of_Aquac
ulture-Nasmiah-_2014.pdf
by

FILE	IONAL_11-_INTERNATIONAL_JOURNAL_OF_AQUACULTURE-NASMIAH-_2014.PDF (401.45K)		
TIME SUBMITTED	29-DEC-2019 11:36PM (UTC+0700)	WORD COUNT	3408
SUBMISSION ID	1238655605	CHARACTER COUNT	19418

Characterization and Identification of Bacteria Isolated from Seaweed *Gracilaria verrucosa* (Linn., 1758) Infected by Ice-ice

Nasmia¹, Radjuddin Sya¹, dudin², Alexander Rantetondok^{2*}, Elmi N. Zainuddin²

1. Department of Aquaculture, Faculty of Animal Husbandry and Fisheries, Tadulako University, Palu, Central Sulawesi, Indonesia

2. Faculty of Marine Sciences and Fisheries, Hasanuddin University, Makassar, South Sulawesi, Indonesia

* Corresponding author Email: alexander_tondok@yahoo.com

International Journal of Aquaculture, 2014, Vol.4, No.23 doi: 10.5376/ija.2014.04.0023

Received: 24 Apr., 2014

Accepted: 21 May, 2014

Published: 21 Aug., 2014

Copyright © 2014 Nasmia et al., This is an open access article published under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Preferred citation for this article:

Nasmia et al., 2014, Characterization and Identification of Bacteria Isolated from Seaweed *Gracilaria verrucosa* (Linn., 1758) Infected by Ice-ice, International Journal of Aquaculture, Vol.4, No.23: 1-7 (doi: 10.5376/ija.2014.04.0023)

Abstract Ice-ice disease in cultivated algae occurs due to pathogenic bacterial infections. Generally, ice-ice disease is characterized by whitening of the branches and initiated with the color changes of the thalli becomes transparent. This study was aimed to isolate and identify bacteria on seaweed *Gracilaria verrucosa* infected by ice-ice. Isolated bacteria was inoculated in Tryptic Soy Agar (TSA) and Thiou sulphate Citrate Bile Sucrose (TCBS). Morphological and biochemical characterization of the bacterial isolates revealed eight species of bacteria were found in infected thalli including *Acinetobacter* sp., *Pseudomonas* sp., *Bacillus* sp., *Micrococcus* sp., *Corinoverm* sp., *Cytophaga* sp., *V. mimicus* and *V. Campbelii*, but not all identified bacteria are pathogens on *Gracilaria verrucosa*. The pathogenic bacteria were *Acinetobacter* sp., *Pseudomonas* sp., *Bacillus* sp., *Cytophaga* sp. and *Vibrio* sp.

Keywords *Gracilaria verrucosa*; Pathogenic bacteria; Ice-ice disease; Indonesia

Introduction

Seaweed is an aquaculture commodity that has a high economic value. Production of seaweed *Gracilaria* was estimated to increase concurrent with the increasing demand of seaweed for food industry, pharmacy and cosmetics. *Gracilaria* has been cultivated worldwide. *Gracilaria verrucosa* and *G. gigas* are commonly cultured in brackish water ponds in Indonesia such as in South Sulawesi (Jeneponto, Takalar, Sinjai, Bulukumba, Wajo, Palopo, Bone and Maros Regencies), North Coast of Java (Serang, Tangerang, Bekasi, Karawang, Brebes, Pemalang, Tuban and Lamongan Regencies) and West Lombok (DKP, 2005).

The development of *Gracilaria* cultivation in ponds, in addition to increase the productivity and income of breeding farms, it is also expected to improve the quality of pond environment. By cultivating the *Gracilaria* sp in brackish-water ponds the production of oxygen is increasing due to the increasing of the photosynthesis and the decreasing of the nitrite due to the nitrification process. In the other hand the production of the brackish-water ponds (milkfish and shrimp) can be increased. Seaweed is one of

alternatives that can be used to improve the pond water environment, because it has ability to absorb nutrients, so that can be used as a biofilter, bioakumulator as well as biomonitoring pollution that occurs in aquaculture pond waters (Komarawidjaja, 2005). Thus, the causative agent of ice-ice disease can affect the seaweed as a host of bacteria.

One of the constraints experienced in the process of cultivation of seaweed is disease. The attack of ice-ice disease increased in line with pathogenic bacterial infection of the seaweed thalli. The condition is caused by the increasing of pathogenic bacteria activity in secrete virulence factors. The development of the pathogenic bacteria activity on seaweed thalli result in white patches on the thallus and gradually become brittle and eventually thallus fracture, resulting in decreased production of seaweed range of 70-100% (Vairappan *et al.* 2008).

Results of the identification on the seaweed *Kappaphycus alvarezii* infected by ice-ice disease is caused by a pathogenic bacteria namely; *Vibrio alginolyticus*, *Pseudomonas cepacia*, *Flavobacterium meningosepticum*, *Pseudomonas diminuta* and

Plesiomonas shigelloides (Aris, 2011) [6] Types of bacteria that cause ice-ice disease are *Pseudomonas nigricaciens*, *Pseudomonas fluorescens*, *Vibrio granii*, *Bacillus cercus* and *Vibrio agarliquefaciens* Yulianto (2002).

The [19] management of good seaweed cultivation and free of ice-ice disease are the means in improving production technology of *G.verrucosa*. The disease needs to be studied through isolation, identification and control of pathogenic bacteria in *G. Verrucosa*.

1 Materials and Methods

1.1 Place of Experimental Seaweed

Seaweed *Gracilaria verrucosa* was taken from brackishwater pond at Takalar Regency, South Sulawesi Province, Indonesia. Ice-ice infected thalli of *G.verrucosa* were washed with sterile water and placed into sterile plastic bags. Samples were then placed in the cool box and transported to laboratory. Isolation and biochemical characterization of bacteria were carried out in Marine Microbiology Laboratory, Faculty of Marine Sciences and Fisheries, Microbiology and Immunology Laboratory, Faculty of Medicine, Hasanuddin University, and Institute of Brackishwater Aquaculture Development, Maros.

1.2 Bacterial Isolation and Identification

Isolation and identification of bacteria was conducted from ice-ice infected branches of seaweed *Gracilaria verrucosa*.

One gram of ice-ice infected branches of *Gracilaria verrucosa* was homogenized in 9 ml sterile NaCl 0.9% and dilution series up to 10^{-6} were made up. One milliliter of aliquot [12] in each dilution series was spread-plated onto tryptic soy agar (TSA) [4] and thiosulphatecitrate bile sucrose (TCBS) agar plates and then incubated at 30°C for 24 h. For purification and identification, differentially isolated colonies were randomly picked and streaked onto the TSA medium. Identification of bacteria was conducted based on the morphology of colonies and biochemical characterizations.

1.3 Biochemical Test

Identification of bacteria was done through the biochemical test to find out the behavior or characteristic of the bacteria according to Capuccino and Sherman, (1987), that included the Gram staining,

OF, oxidase test, catalase test, SIM test (indole, motile, gas, H₂S), TSIA test (butt, slant, H₂S, gas), MR, VP, King A and King B. The nature or characteristic of bacteria by using medium TCBS (*Thioisulphate Citrate Bile Salt Sucrose Agar*).

Characterization is one of the activities carried out to observe the results of bacterial isolation (isolates). Characterization activities can be carried out based on the nature of cytology (cell form, movement or motility, Gram nature), morphology, and physiological nature. Test morphological natures include the natures of colonies, such as size, shape, color and edges, while the physiological nature such testing starch hydrolysis test, hydrolysis of fat, protein hydrolysis and catalase test. Test of Gram staining was conducted [14] group bacteria into 2 big groups according to its cell wall structure namely Gram-positive and Gram-negative bacteria. The staining was important stage in the bacteria characterization and identification (Lay, 1994). The catalase test was conducted to find out whether a bacteria has catalase enzyme in which the enzyme oxidized H₂O₂ (Cappuccino and Sherman, 1987).

2 Results

2.1 Bacterial Isolation and Identification

The results of the observation on the thalli showed that the gross signs of the thalli infected with ice-ice disease were initiated with translucent appearance of



Figure 1. Gross signs of the thalli infected with ice-ice disease

the proximal, median or distal end of the thalli leading to whitening of the whole thalli (Figure 1). Santoso (2008) reported that ice ice infection in seaweed attacked the base of thallus, stem and the tip of thallus that caused the tissue become white and soft.

There were eight bacterial isolates with different colonial morphologies found in the infected thalli of the seaweed (Table 1).

Bacterial isolate was successfully isolated from the seaweed *G. verrucosa* infected by ice-ice. Morphology of bacteria colony was almost entirely circular, except at Gv.1 isolates. Almost all isolates were *entire* colony margins, except Gv.2, Gv.3 and Gv.4 which were *serrate*, elevation is generally *convex*, except Gv.4 and Gv.5 are *flat*. The color of the colony on Gv.1 and Gv.5 isolates where white, the isolates Gv.2 was yellow, Gv.4, Gv.7 and Gv.8, were cream, and orange on Gv.3 and Gv.6 isolates. These data were appropriate with funding of Cappucino and Sherman (1987), that bacterial colonies shape in general *circular, irregular, filamentous, rhizoid*, the elevation was *raised, convex, flat, umbonate, crateriform*. Margin shapes *entire, undulate, filiform, curled and lobate*.

2.2 Characteristic of Bacteria

The result of gram staining of ice-ice bacteria isolates

Table 1. Colonial morphology of bacteria isolated from *Gracilaria verrucosa* infected with ice-ice diseases.

Isolates	Media	Colonial morphology			
		Shape	Margin	Elevation	Colour
<i>Gracilaria verrucosa</i> 1 (Gv.1)	TSA	Irregular	Entire	Convex	White
<i>Gracilaria verrucosa</i> 2 (Gv.2)	TSA	Circular	Serrate	Convex	Yellow
<i>Gracilaria verrucosa</i> 3 (Gv.3)	TSA	Circular	Serrate	Convex	Cream
<i>Gracilaria verrucosa</i> 4 (Gv.4)	TSA	Circular	Serrate	Flat	Yellow
<i>Gracilaria verrucosa</i> 5 (Gv.5)	TSA	Circular	Entire	Flat	White
<i>Gracilaria verrucosa</i> 6 (Gv.6)	TSA	Circular	Entire	Convex	Orange
<i>Gracilaria verrucosa</i> 7 (Gv.7)	TCBS	Circular	Entire	Convex	Yellow
<i>Gracilaria verrucosa</i> 8 (Gv.8)	TCBS	Circular	Entire	Convex	Yellow

G. verrucosa was obtained that bacteria isolates Gv.1, Gv.2, Gv.6, Gv.7, and Gv.8 belonged to gram-negative. While isolates Gv.3, Gv.4 and Gv.5 belonged to Gram-positive. The result of catalase test showed that isolates Gv.1, Gv.2, Gv.3 and Gv.4 showed positive catalase reaction, while isolates Gv.5 and Gv.6 showed negative reaction (Table 2).

The SIM test (gas, H₂S) and TSIA (gas, H₂S) showed that all isolates were negative. It was caused by the formation of H₂S marked by the black sign in the medium and the formation of gas marked by the rupture of medium at the lower end of the glass tube.

Biochemical characterization of the isolates showed the presence of *Acinetobacter* sp. at Gv.1, *Pseudomonas* sp. at Gv.2, *Bacillus* sp. at Gv.3, *Micrococcus* sp. at Gv.4, *Corinoverm* sp. at Gv.5, *Cytophaga* sp. at Gv.6 (Table 2), *Vibrio mimicus* at Gv.7 and *Vibrio Campbelii* at Gv.8 (Table 3) in the ice-ice disease infected thalli.

3 Discussion

3.1 *Acinetobacter* sp.

The results of staining obtained show that *Acinetobacter* sp. included in the gram-negative marked in red on the bacterial cell. The results of characteristic (Table 2) showed that surface of the colony is shiny, not motile, oxidase negative, does not produce pigment, catalase positive, and white.

Table 2 Biochemical characterization of bacteria isolated from ice-ice infected seaweed *Gracilaria verrucosa* and inoculated in Tryptic Soy Agar (TSA).

Characteristic	Gv1	Gv.2	Gv.3	Gv.4	Gv.5	Gv.6
Gram's stain	-	-	+	+	+	-
OF	-	-	-	-	-	-
Oxidation	-	-	+	-	+	+
Catalase	+	+	+	+	-	-
SIM	Indol	-	+	-	-	-
	Motile	-	+	+	+	-
	Gas	-	-	-	-	-
TSIA	H ₂ S	-	-	-	-	-
	Butt	Red	Red	Red	Red	Red
	Slant	Red	Red	Red	Red	Red
	H ₂ S	-	-	-	-	-
MR	Gas	-	-	-	-	-
	MR	-	+	-	-	-
VP	-	-	-	-	-	-
King A	-	+	+	+	-	+
King B	-	+	+	+	-	+
Genus	<i>Acinetobacter sp.</i>	<i>Pseudomonas sp</i>	<i>Bacillus sp</i>	<i>Micrococcus sp</i>	<i>Corinoverm sp</i>	<i>Cytopaga sp</i>

Generally, *Acinetobacter* can be found in freshwater (Allen *et al.*, 1983 in Austin and Austin, 1993) and marine environment (Austin, 1982 in Austin and Austin, 1993). These bacteria may have ability to cause diseases in aquatic organisms (Roald and Hastein, 1980 in Austin and Austin, 1993).

2 *Pseudomonas sp.*

Pseudomonas sp. is a rod Gram negative bacteria, motile (some have polar flagellum) and catalase positive. Generally, colonial bacteria are yellow, facultative anaerobic or aerobic, can grow at 4-43 °C. Most species have an optimum temperature around 30 °C (Buchanan and Gibbons, 1974).

Pseudomonas generally can be found in the soil, freshwater and seawater (Buchanan and Gibbons, 1974). These bacteria frequently isolated from plant and animal surfaces. Some members of these genera considered as a true pathogen in plants (Todar, 2013). *Pseudomonas* grows in injured tissues and produces toxins leading to the death of surrounding cells (Moorman, 2013). Some species such as *Pseudomonas aeruginosa*, are opportunistic pathogen releasing extracellular protease that can infect host tissues (Buell *et al.* 2003).

Amiluddin (2007) identified five bacteria causing ice-ice disease on *K. alvarezii* including *Bacillus cereus*,

Table 3. Biochemical characterization of bacteria isolated from ice-ice infected seaweed *Gverrucosa* and inoculated in Thiosulphate Citrate Bile Sucrose Agar (TCBS).

Characteristic	Gv.7	Gv.8
Swarming	-	-
Luminescence	-	-
VP Test	-	-
Arginin Dihydrolase	-	-
Gas From Glucose	-	-
Growth at 400C	-	-
Lysine Decarboxylase	+	+
Pigmentation	-	-
Amylase	+	+
Sucrose	-	-
Indole	-	-
Omithin Decarboxylase	+	+
Putrescine	-	-
Ethanol	-	-
Serine	-	-
Heptaniate	-	-
Xantine	-	-
Aminobutirate	-	-
Arabinose	-	-
Cellubiose	-	-
Glucuronate	-	-
Ketoglutarate	-	-
L-Alanine	-	+
Leucine	-	-
Propionate	-	-
Genus	Vibrio sp.	Vibrio sp.
Species	Vibrio mimicus	Vibrio campbelli

Vibrio granii, *V. liquefaciens*, *P. nigrificiens*, and *P. fluorescens*. Darmayanti *et al.* (2003) also isolated *Pseudomonas* sp., *Vibrio* sp. and *Aeromonas* sp. in healthy and infected thalli of cultured *K. alvarezii* in Pari Island. In addition, *Pseudomonas* sp. was found on cultivated seaweed in coastal water in Takalar Regency, South Sulawesi, Indonesia

3.3 *Bacillus* sp

Bacillus is a Gram negative bacteria, rod, motile (some non-motile), catalase positive and oxidase positive. *Bacillus* may form endospore, aerobic and

facultative anaerobic (Feliatra, 2004). *Bacillus* can be isolated from soil and water including seawater. These bacteria can be found as a pathogen, opportunistic saprophytes. Some species produced extracellular enzymes that have ability to hydrolyze proteins and polysaccharides (Pelczar *et al.* 1975).

Bacillus sp. produced spores that are heat resistant and have ability to degrade xylan and carbohydrates (Cowan and Steel, 1973). *Bacillus* sp. has ability to grow at temperature more than 50 °C and less than 5 °C, at high concentration of salt (> 10%), resistant to pasteurization, produce spores and has higher proteolytic activities than other microbes. *Bacillus* sp. resistant to environmental conditions like resistance of heat, acid and extreme salt content. Kinds of *Bacillus* produce extracellular enzymes which can hydrolyze protein and polysaccharide complex.

3.4 *Micrococcus* sp

Micrococcus is a Gram positive bacteria, sphere (cocci) with a size of 0.5–3.0 µm in diameter. It is aerobic and anaerobic and has an optimum temperature of 25–37°C to grow. These bacteria can be found as a pathogen, opportunistic saprophytes and commensal organism. The role of this bacteria in the ground is as decomposers to decompose organic material into its composing constituent that can be used directly by plants.

3.5 *Cytophaga* sp

Cytophaga is a rod Gram negative bacteria. Some member within this group can cause diseases in fish. These bacteria can utilize cellulose and agar/chitin. This microbe can produce cell membranes containing endogluconases and periplasmic exogluconases that have activity in degrading cellulose. These bacteria can attack cellulose, but they cannot produce cellulose. This enzyme can only bind to the cell envelope, mix with mucus and is released during movement. This bacterium inhibits soil and water. The colony of these bacteria is yellow or orange and can be grown in the medium containing cellulose. Largo *et al.* (1995), who isolated bacteria in *Kappaphycus* obtained 10 bacteria strains *Cytophaga* sp. (*Cytophaga-Flavobactroium*).

3.6 *Vibrio* sp

Vibrio have a characteristic of curved rod Gram negative bacteria with a size of 1 – 3 × 0.4 – 0.6 µm,

non-forming spore and capsule, and motile with a polar flagella. This saprophytic microbe becomes pathogenic when the environment quality depleted. According to Kabata (1988), *Vibrio* is a pathogenic bacteria that commonly cause problems to the aquatic organisms. Largo et al. (1995) isolate *Vibrio* sp from the infected *K. alvarezii*. Recently, a study on ice-ice infected *Kalvarezii* revealed the presence of *V. alginolyticus*, *P. cepacia*, *P. diminuta*, *Plesiomonas shigelloides* and *Flavobacterium meningosepticum* in the infected thalli (Aris, 2011). Furthermore, Bockemuhl et al. (1986), Chowdury et al. (1989) in Eaves (1994) reported that *Vibrio mimicus* was found in brackish water and fresh water.

Largo et al. (1999) explained the mechanism of the *Vibrio* infection in seaweed thallus was when the seaweed under stress. The *Vibrio* would breed in the cell wall by using polysaccharide as the media or carbon source. Furthermore, Lin in Yulianto (2009) explained that after 2-3 days, *Vibrio* entered the tissue until the medullary seam by pumping the caraginase enzyme then caused the thallus became pale/white and its tissue was soft and easily broken.

Acknowledgement

The authors thank the Ministry of Education and Culture of the Republic of Indonesia for provided the funding research, Thank also our friends (Huyyimah, Rusaini, Ph.D. Cand.) for their assistance in the provide.

References

Amiluddin N. M., 2007, Study of Growth and Carragenan Content of Seaweed *Kappaphycus alvarezii* (Linn., 1758). Affected by Ice ice in the Water of Puri island, Seribu Islands, Thesis for M.S, Graduate Scholl of Bogor Agricultural University, Bogor, pp.1-78

Aris M., 2011, Identification, Pathogenicity of Bacteria and the Use of Gene 16S rRNA for Ice-ice Detection on Seaweed Aquaculture . *Kappaphycus alvarezii* (Linn., 1758), Thesis for M.S, Graduate School of Bogor Agricultural University (IPB), Bogor, pp.1-127

Austin B., and Austin D.A., (eds). 1993, Bacterial fish pathogen disease in farmed and wild fish, Second Edition, Ellis Horwood Limited. Department of Biological Sciences, Heriot-Watt University, England, pp.553

Buchanan R.E., and Gibbons N.E., 1974, *Bergey's Manual of Determinative Bacteriology*, 8th ed. USA, The Williams & Wilkins Co, Inc, pp.1272

Buell C.R., Joardar V., and Lindeberg M., 2003, The Complete Genome Sequence of the *Arabidopsis* and *Tomato* Pathogen *Pseudomonas syringae* pv. *Tomato* DC3000, Proc. Natl. Acad. Sci. USA 100:10181-10186
<http://dx.doi.org/10.1073/pnas.1731982100>

Cappuccino J.G., and Sherman N., 1987, *Microbiology a Laboratory Manual* (7th ed.). San Francisco, The Benjamin/Cummings Publishing Company, Inc, pp.458

Cowan S.T., and Steels D., 1973, Manual for identification of medical bacteria, Second Edition, Cambridge University Press, London, pp.352

Darmayanti D. 2001, Fusarium Function,
<http://sciweb.nybg.org/science2/hcol/fusarium3.asp> .(Online).

Directorate General of Aquaculture, 2005, Fisheries Revitalization. Department of Marine and Fisheries, Jakarta

Eaves L.E., and Katterer P.J., 1994, Mortalities in Red Claw Crayfish *Cherax quadricarinatus* Associated with Systematic *Vibrio mimicus* Infection, Diseases of Aquatic Organisms, 19: 233-237
<http://dx.doi.org/10.3354/dao019233>

Feliatra I.E., Efendi I., and Suryadi E., 2004, Isolation and Identification of Probiotic Bacteria Tiger Grouper (*Ephinephelus fuscogutatus*) in the Effort of Fish Feed Efficiency, Indonesian Nature Journal, 6(2): 75-80

Moorman G.W., 2013, *Pseudomonas* Syringae: A Common Pathogen on Woody Plants.
<http://extension.psu.edu/pests/plant-diseases/all-fact-sheets/pseudomonas-syringae-a-common-pathogen-on-woody-plants>.

Kabata Z., 1988, Parasites and diseases of fish culture in the Tropics, Taylor and Francis. London, pp.318

Komarawidjaja W., 2005, Seaweed *Gracilaria* sp. as Fitoremedial Pond Aquaculture Aquatic Organic Materials, Jur.Tekn.Ling. P3TL (Center for Environmental Assessment and Application of the Technology), Jakarta, 6(2): 410-415

Largo D.B., Fukami K., and Nishijima T., 1995, Occasional Pathogenic Bacteria Promoting Ice-ice Disease in the Carrageenan-producing Red Algae *Kappaphycus alvarezii* (Linn., 1758) and *Euchemia denticulatum* (Solieriaceae, Gigartinales, Rhodophyta), Journal of Applied Phycology, 7: 545-554
<http://dx.doi.org/10.1007/BF00003941>

Largo D.B., Fukami K., and Nishijima T., 1999, Time-dependent attachment mechanism of bacterial pathogen during ice-ice infection in *Kappaphycus alvarezii* (Linn., 1758) (*Gigartinales*, Rhodophyta), Journal of Applied Phycology, 11: 129-136
<http://dx.doi.org/10.1023/A:1008081513451>

Lay W., Bibiana., 1994, Microbial Analysis in a Laboratory, PT Raja Grafindo Persada, Jakarta, pp.168

Nurjanna, M., 2008, Bacterial Identification Isolated from Seaweed with Ice ice Disease, Bul. Tech. Lit. Aquaculture, 7(1): 79-81 (in Indonesian)

Pelczar M.J., Chan E.C.S., 2005, Basics of microbiology, 1st and 2nd Volume. Hadioetomo R.S, Imas T, Tjitrosomo S.S, Angka S.L, translator, Jakarta: University of Indonesia Press, Translation from: Element of Microbiology, pp.997

Santoso L., and Nugraha Y.T., 2008, The Control of Ice ice Disease to Improve Seaweed Production in Indonesia. Science and Technology Journal of Fisheries, 3(2): 37-43 (in Indonesian)

Vairappan C. S., Chung C.S., Hurtado A.Q., Soya F.E., Bleicher-Lhonneur G, Critchley A., 2008, Distribution and Symptoms of Epiphyte Infection in Major Carrageenophyte-Producing Farms. J. Appl. Phycol. 20: 477-483
<http://dx.doi.org/10.1007/s10811-007-9299-8>

Yulianto, K and Mira S., 2009, Macro Algae Cultivation of *Kappaphycus alvarezii* (Linn., 1758) Vertically and Symptoms of Ice ice disease in the Water of Pari Island. Pulau Pari, Oceanology and Limnology Journal, 35(3): 323-332 (in Indonesian)

ORIGINALITY REPORT

% **19**
SIMILARITY INDEX

% **15**
INTERNET SOURCES

% **12**
PUBLICATIONS

% **12**
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to The University of the South Pacific **%3**
Student Paper

2 C. N. Ariole, H. A. Nwogu, P. W. Chuku. "Enzymatic Activities of Intestinal Bacteria Isolated from Farmed *Clarias gariepinus*", International Journal of Aquaculture, 2014 **%3**
Publication

3 nopr.niscair.res.in **%2**
Internet Source

4 link.springer.com **%1**
Internet Source

5 Submitted to University of Stirling **%1**
Student Paper

6 id.scribd.com **%1**
Internet Source

7 Ince Khairana Kadriah, Koko Kurniawan, Endang Susianingsih, Muharijadi Atmomarsono. "SENSITIVITAS DETEKSI PENYAKIT" **%1**

VIBRIOSIS PADA UDANG PENAEID DENGAN
PENANDA MOLEKULER SPESIFIK
HAEMOLYSIN (IAVh)", Jurnal Riset Akuakultur,
2017

Publication

8	d-nb.info Internet Source	% 1
9	www.tumtoon.com Internet Source	% 1
10	www.bioflux.com.ro Internet Source	% 1
11	Submitted to Surabaya University Student Paper	% 1
12	doaj.org Internet Source	% 1
13	Submitted to Nevada State College Student Paper	<% 1
14	www.renevanmaarsseveen.nl Internet Source	<% 1
15	krishikosh.egranth.ac.in Internet Source	<% 1
16	Anna E. Kolodenkova, Evgenia R. Muntyan. "Researches of Interaction of Actors with Use Fuzzy Hypergraph and Cognitive Modeling", 2018 XIV International Scientific-Technical	<% 1

Conference on Actual Problems of Electronics Instrument Engineering (APEIE), 2018

Publication

17

Submitted to Universiti Sains Malaysia

Student Paper

<% 1

18

M. R. Mikluscak, B. E. Dawson-Andoh.
"Microbial colonizers of freshly sawn yellow-
poplar (*Liriodendron tulipifera* L.) lumber in two
seasons: Part 2. Bacteria", *Holzforschung*, 2004

Publication

<% 1

19

E Syafitri, S B Prayitno, W F Ma'ruf, O K
Radjasa. " Genetic diversity of the causative
agent of ice-ice disease of the seaweed from
Karimunjawa island, Indonesia ", IOP
Conference Series: Earth and Environmental
Science, 2017

Publication

<% 1

20

iopscience.iop.org

Internet Source

<% 1

21

peerj.com

Internet Source

<% 1

22

Submitted to Mississippi State University

Student Paper

<% 1

23

I. Gancarz. "Modification of polysulfone
membranes. 2. Plasma grafting and plasma
polymerization of acrylic acid", *Acta Polymerica*,

<% 1

09/01/1999

Publication

24

Elmi Nurhaidah Zainuddin, Hilal Anshary, Huyyirnah Huyyirnah, Ridha Hiola, Dolores V. Baxa. "Antibacterial activity of *Caulerpa racemosa* against pathogenic bacteria promoting "ice-ice" disease in the red alga *Gracilaria verrucosa*", *Journal of Applied Phycology*, 2019

<% 1

Publication

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5
WORDS