

# Internasional\_Surabaya\_2010\_V ol\_2\_-Elmi\_-08-11-2019-277- 2871.pdf

*by*

---

FILE	INTERNASIONAL_SURABAYA_2010_VOL_2_-ELMI_-08-11-2019-277-2871.PDF (280.31K)		
TIME SUBMITTED	22-AUG-2020 06:56AM (UTC+0700)	WORD COUNT	5176
SUBMISSION ID	1372442887	CHARACTER COUNT	28859

<sup>44</sup>  
**PRELIMINARY SCREENING OF MARINE ALGAE FROM  
<sup>50</sup>SOUTH SULAWESI COAST FOR CYTOTOXIC ACTIVITY  
USING BRINE SHRIMP *ARTEMIA SALINA* LETHALITY TEST**

**Elmi Nurhaidah Zainuddin**

Marine Science and Fisheries Faculty, Hasanuddin University, e-mail: elmi18id@yahoo.com

<sup>43</sup>  
**Abstract:** Fifteen algae extracts, isolated from seven <sup>43</sup>species of marine algae belonging to the classes **Phaeophyta** (*Padina boergesenii*, *Sargassum prismaticum*, *Rosenvingea orientalis*, *Dictyopteris acrostichoides*), Chlorophyta (*Codium dwarkense*), and Rhodophyta (*Sarconema filiforme*, *Wrangelia tanegana*) were investigated for their cytotoxic activity against brine shrimp <sup>42</sup>*Artemia salina*. The algae extracts were prepared successively first by n-hexan followed by dichloromethane (DCM) and finally with ethyl acetate (EtOAc). Of all the extracts tested, only the dichloromethane (DCM) extract of *Sarconema filiforme* showed no significant activity with LC<sub>50</sub> value >1000 µg/mL. Other 12 extracts such as *Rosenvingea orientalis* (n-hexan, DCM), *Wrangelia tanegana* (n-hexan, DCM), *Padina boergesenii* (n-hexan, DCM, EtOAc), *Codium dwarkense* (n-hexan), *Sargassum prismaticum* (n-hexan, DCM) and *Dictyopteris acrostichoides* (DCM, EtOAc) showed significant activity with LC<sub>50</sub> value below 62.5 µg/mL. The n-hexan extract of *Dictyopteris acrostichoides* and the dichloromethane extract of *Codium dwarkense* exhibited intermediary cytotoxicity with LC<sub>50</sub> value between 62.5 and 250 µg/mL.

**Keywords:** Marine algae, cytotoxicity, brine shrimp *Artemia salina* lethality test.

**INTRO<sup>13</sup>DUCTION**

Over the past several decades, algae and their extracts have generated an enormous amount of interest in the pharmaceutical industry <sup>49</sup>as a fresh source of bioactive compounds with different bioactivity such as cytotoxic (Tang *et al.*, 2002), antibacterial (Vallinayagam *et al.*, 2009), antifungal (Aliya and Shamaeel, 1999), antiviral (Serkedjieva, 2004; Garg *et al.*, 1992), antitumour (K<sup>15</sup>i, 2002), antioxidant (Yuan and Walsh, 2006) and larvasidal (Thangam *et al.*, 1993). Until now more than 2400 marine natural products have been isolated from seaweeds of subtropical and tropical populations (Faulkner, 2001; Munro and Blunt, (1999).

In the world, also in Indonesia, diseases crisis caused by non-controlled cell proliferation as tumour- and cancer cells have a significant position. National Cancer Institute USA reported that almost one third of human death caused by cancer disease have correlation with human diet. As reported by Harvard School of Public Health in USA that pre-menopause women in Japan have a chance that on severe breast cancer three times lower than that severe by American women. This caused by they always include algae on their food menu. Algae have a big potential as food and medicine since long ago since they rich on vitamin, mineral, crude fibrous, protein and polysaccharide (Ito and Hori, 1989; Lahaye, 1991; Darcy-Vrillon, 1993). Algae have a low fat content with unsaturated form which can protect human from cardiovascular pathogen. Their non-toxic phycocolloid role as low calorie nutrition and stabilization agent in food industry (Van den Hoek *et al.*, 1993; Critchley and Ohno, 1998; Lee, 1999). In Japan traditional food, algae were used as sushi wrapper, seasonings, condiments and salad or vegetables which composed 10-25% of food consume from most Japan people (Skibola, 2004; Teas, 1981).

Global economy crisis, high price of medicinal drugs, difficult access of population on medical treatment and pharmacy, and side effect of synthetic chemical drugs, are the reasonable factor to use herbal medicine for disease treatment. Many

experts suggest consuming various foods <sup>4</sup>help to prevent the developing of cancer cells or reduce the tumour cells size. Algae have ingredients <sup>4</sup>like  $\beta$ -carotene, protein, vit B12, fibre <sup>46</sup>chlorophyll, and essential fatty acid which protect women from breast cancer. The omega-3 fatty acids (eicosapentaenoic- and stearidonic acid) <sup>4</sup>and iodine in seaweed is thought to reduce breast cancer risk (Vinayak *et al.*, 2010). Diets with three algae, *Porphyra tenera*, *Laminaria religiosa* and *L. japonica* var. *ochotensis*, showed an inhibitory effect on mammary tumorigenesis with tumor incidences of 35%, 35%, and 50%, respectively, while in control only 69% (Yamamoto *et al.*, 1987).

Most anticancer and antitumour drugs for chemotherapy treatment are synthesized from the synthetic chemical compounds and for the long term chemotherapy, many cancer patients could not be stand from suffering the drugs side-effect and can finally die. It is therefore urgent to find a new anticancer drug which safely for the long term therapy. For this, non-synthetic or natural product especially marine natural product could be a promised resource. Several study showed that marine environment which cover almost 70% of earth is a rich sources on bioactive compounds, many of them have unique structure different with those from terrestrial one. Marine secondary metabolites are organic compounds produced by marine organisms <sup>28</sup>microbes, sponges, and seaweeds. The organism biosynthesizes these compounds to protect themselves and to maintain homeostasis in their environment (Selvin and Lipton, 2004). The problem which always appears in marine drug discovery is raw material supply for industrial scale. Marine algae or seaweeds are the renewable resources that easy to cultivate in Indonesia. With the coast line of  $\pm 81.000$  km, Indonesia has a big challenge or a high potential on marine algae cultivation. Until now there are 555 species of algae that have been found in Indonesia water (Bengen, <sup>22</sup>1) and one of the areas which has a big potential for algae culture is South Sulawesi. The aim of this study was to assess the cytotoxic effects of organic <sup>41</sup>algal extracts of marine algae collected from South Sulawesi coast, towards the nauplii of the brine shrimp *Artemia salina*, as a potential source of marine bioprospecting for anticancer drugs candidate.

Brine shrimp bioassay has been used in this study since it is <sup>1</sup>known as an efficient, safe, fast and reproducible procedure to assess biological and <sup>7</sup>pharmacological potential of new compounds (Meyer *et al.*, 1982; Manilal *et al.*, 2009a) and requires only a small amount of the assay <sup>1</sup>substance (Svensson *et al.*, 2005). This assay has presented satisfactory correlation with cytotoxicity property to some solid human tumours (Badisa <sup>18</sup>*et al.*, 2007). Moreover, it is an inexpensive test and without ethical constraints (Aristides *et al.*, 2008).

## MATERIALS AND METHODS

### Algae Collecting

Algae were collected during the low tide along the coasts of Takalar and Pangkep in South Sulawesi, Indonesia. The materials were washed with cleaned sea water and put into plastic bags before kept in an ice box to prevent photolysis and thermal degradation during transportation.

### Sample Preparation

In the laboratory of Marine Science and Fishery <sup>25</sup>faculty of Hasanuddin University, algae species were washed with filtered seawater to remove the epizooones, epiphytes, animal castings, sand, calcareous and other adhering detritus matters. Small samples of the species were separated for identification and the rest were washed with freshwater to remove the salt. After draining off the water, the cleaned plant materials were wiped with a blotting sheet and <sup>27</sup>sun-dried carefully under shade for 24-48 hour. Dried materials were weighed and cut into small pieces before finely grounded in a mechanical grinder. The powdered algae were kept airtight in plastic bags and put in the room temperature for further experiment.

### Extraction of Algae

Extraction of algal materials was conducted as described previously (Zainuddin, 2006). 50 g of finely powdered algal material were extracted with 500 mL n-hexane in a 1-L capacity round bottom flask (1:10, w/v). The extraction was run on a stirrer plate for 24 h under room temperature. The extracts were filtered through a Whatman no. 1 filter paper then evaporated under reduced pressure in a rotary evaporator until 5-10 mL volume. The concentrated extracts were kept on small vials and let dry under room temperature to yield thick oily crude extract and stored airtight at -20°C for further analysis. The algae residue from n-hexane extraction were dried at room temperature for 24 h and re-extracted successively with higher polarity solvents (dichloromethane and ethyl acetate) using the method as described above.

### Brine Shrimp Lethality Test

Brine shrimp *Artemia salina* eggs were hatched in a flask containing filtered seawater (1 g cyst per litre) under continuous illumination at 27–30°C for 48 h. The air stone was placed in the bottom of the jar to ensure complete hydration of the cysts. After incubation period, newly hatched nauplii were collected for the assay.

Brine shrimp cytotoxicity assay was performed using the freshly hatched free-swimming nauplii of *Artemia salina*. Stock solution was prepared by dissolving ten mg of crude algae extract in 100% methanol in an eppendorf and filled with filtered seawater until 1 mL volume. Test solutions were prepared by diluting the stock solution with filtered seawater to obtain two-fold serial dilution (1000, 500, 250, 125 and 62.5 µg). Ten brine shrimp nauplii were transferred into vials containing 4 mL filtered seawater. One mL of the test solution containing each extract concentration were added into vials to make 5 mL of total solution. Parallel positive (only methanol and seawater) and negative (only seawater) controls were included in experiment set up. The cytotoxicity was determined after 24 h exposure under constant illumination. The number of live brine shrimp nauplii in each vial was determined with a hand lens, and the mortality rate was calculated. The mortality end point of the bioassay was determined if no internal or external movement of larvae was observed during 30 seconds. Each test was run in duplicate and it was repeated until two or three times.








Based on the percent mortality, the results were expressed as LC<sub>50</sub> value. It was defined as the concentration needed to cause half of the tested brine shrimp died within 24 h. If the LC<sub>50</sub> value is <1000 µg/mL the assayed compound was regarded as cytotoxic and if the value is >1000 µg/mL, the assayed compound was regarded as non-toxic (Meyer *et al.*, 1982; Badisa *et al.*, 2007; Parra *et al.*, 2001).

### RESULTS

The algae species are collected along the coasts of Takalar and Pangkep, South Sulawesi Province, Indonesia. They are distributed in the intertidal zone of sandy beaches and belonging to classes Phaeophyta, Chlorophyta and Rhodophyta. Five species consist of four Phaeophyta (*Padina boergesenii*, *Sargassum prismaticum*, *Rosenvingea orientalis*, *Dictyopteris acrostichooides*) and one Chlorophyta (*Codium dwarkense*) were collected from Takalar coast, whereas two Rhodophyta (*Sarconema filiforme* and *Wrangelia tanegana*) were collected from Pangkep coast (Table 1).

Percent of dry biomass ranging from 6.57% to 19.62%. The highest percent were shown by brown alga *Rosenvingea orientalis* and red alga *Sarconema filiforme* (both have dry weight of 19.62%) whereas the lowest was shown by green algae *Codium dwarkense* with 6.57% (Table 2).

**Table 1. Figure and classification of marine algae collected from Takalar and Pangkep coast of Sulawesi Selatan**

No.	Algae Species	Class, Order and Family (Location)	Algae Figure
1.	<i>Padina boergesenii</i>	Phaeophyta, Dictyotales, Dictyotaceae (Takalar)	
2.	<i>Sargassum prismaticum</i>	Phaeophyta, Fucales, Sargassaceae (Takalar)	
3.	<i>Rosenvingea orientalis</i>	Phaeophyta, Scytosiphonales, Chnoosporaceae (Takalar)	
4.	<i>Dictyopteris acrostichoides</i>	Phaeophyta, Dictyotales, Dictyotaceae (Takalar)	
5.	<i>Codium dwarkense</i>	Chlorophyta, Bryopsidales, Codiaceae (Takalar)	
6.	<i>Sarconema filiforme</i>	Rhodophyta, Gigartinales, Solieriaceae (Pangkep)	
7.	<i>Wrangelia tanegana</i>	Rhodophyta, Ceramiales, Wrangeliaceae (Pangkep)	

**Table 2. Wet- and dry-weight of marine algal biomass**

No.	Algae Species	Class	Wet Weight (ww) (g)	Dry Weight (dw) (g)	Percent of dw/ww
1	<i>Padina boergesenii</i>	Phaeophyta	540	58.22	10.78
2	<i>Sargassum prismaticum</i>	Phaeophyta	597.3	74	12.39
3	<i>Rosenvingea orientalis</i>	Phaeophyta	265.04	52	19.62
4	<i>Dictyopteris acrostichoides</i>	Phaeophyta	1008	110,9	11
5	<i>Codium dwarkense</i>	Chlorophyta	1129	74.14	6.57
6	<i>Sarconema filiforme</i>	Rhodophyta	158	31	19.62
7	<i>Wrangelia tanegana</i>	Rhodophyta	514	78.67	15.31

A total of 15 organic extracts of seven marine algae species were obtained by extraction with three different polarities of solvent. The dry weight of crude extracts ranging from 110.5 mg to 662 mg (0.22-1.32%). The highest percent of dry weight was shown by dichloromethane extract of brown alga *Dictyopteris acrostichoides* (1.32%), whereas the lowest one was shown by n-hexane extract of red alga *Wrangelia tanegana* (0.22%) (Table 3).

<sup>36</sup> The fifteen organic extracts were evaluated for their cytotoxicity using brine shrimp *Artemia salina* lethality test. Twelve of 15 organic extracts showed high cytotoxicity with  $LC_{50} < 62.5 \mu\text{g/mL}$ , whereas two extracts only presented intermediary cytotoxicity with  $LC_{50}$  ranging from 62.5 to 250  $\mu\text{g/mL}$ . Of all the organic extracts tested, dichloromethane extract of *Sarconema filiforme* exhibited no cytotoxic activity against *Artemia salina* larvae with  $LC_{50} > 1000 \mu\text{g/mL}$  (Fig. 1).

**Table 3. Percentage of crude extracts obtained from 50 mg of biomass dry weight in 500 mL organic solvent.**

No.	Algae species	Class	Extracts	Crude Extracts (mg)	Crude Extracts (%)
1	<i>Rosenvingea orientalis</i>	Phaeophyta	Hex	218.1	4.36
2	<i>Wrangelia tanegana</i>	Rhodophyta	Hex	110.5	2.21
3	<i>Padina boergesenii</i>	Phaeophyta	Hex	226.9	4.54
4	<i>Codium dwarkense</i>	Chlorophyta	Hex	147.5	2.95
5	<i>Sargassum prismaticum</i>	Phaeophyta	Hex	215.7	4.31
6	<i>Dictyopteris acrostichoides</i>	Phaeophyta	Hex	519.6	10.39
7	<i>Codium dwarkense</i>	Chlorophyta	DCM	222.4	4.45
8	<i>Sarconema filiforme</i>	Rhodophyta	DCM	142.4	2.85
9	<i>Dictyopteris acrostichoides</i>	Phaeophyta	DCM	662	13.24
10	<i>Rosenvingea orientalis</i>	Phaeophyta	DCM	213	4.26
11	<i>Padina boergesenii</i>	Phaeophyta	DCM	144.4	2.89
12	<i>Wrangelia tanegana</i>	Rhodophyta	DCM	361.3	7.23
13	<i>Sargassum prismaticum</i>	Phaeophyta	DCM	238.6	4.77
14	<i>Dictyopteris acrostichoides</i>	Phaeophyta	EtOAc	328.4	6.57
15	<i>Padina boergesenii</i>	Phaeophyta	EtOAc	282.6	5.65

48  
**Table 4. Cytotoxicity of organic extracts of marine algae collected from South Sulawesi coast against *Artemia salina* nauplii.**

No	Algae species	Class	Extracts	LC <sub>50</sub> value	Cytotoxicity Levels
1	<i>Rosenvingea orientalis</i>	Phaeophyta	n-Hexan	<62.5 µg/mL	High Cytotoxicity
2	<i>Wrangelia tanegana</i>	Rhodophyta	n-Hexan	<62.5 µg/mL	High Cytotoxicity
3	<i>Padina boergesenii</i>	Phaeophyta	n-Hexan	<62.5 µg/mL	High Cytotoxicity
4	<i>Codium dwarkense</i>	Chlorophyta	n-Hexan	<62.5 µg/mL	High Cytotoxicity
5	<i>Sargassum prismaticum</i>	Phaeophyta	n-Hexan	<62.5 µg/mL	High Cytotoxicity
6	<i>Dictyopteris acrostichoides</i>	Phaeophyta	n-Hexan	125-250 µg/mL	Intermediary Cytotoxicity
7	<i>Codium dwarkense</i>	Chlorophyta	DCM	62.5-125 µg/mL	Intermediary Cytotoxicity
8	<i>Sarconema filiforme</i>	Rhodophyta	DCM	>1000 µg/mL	No Cytotoxicity
9	<i>Dictyopteris acrostichoides</i>	Phaeophyta	DCM	<62.5 µg/mL	High Cytotoxicity
10	<i>Rosenvingea orientalis</i>	Phaeophyta	DCM	<62.5 µg/mL	High Cytotoxicity
11	<i>Padina boergesenii</i>	Phaeophyta	DCM	<62.5 µg/mL	High Cytotoxicity
12	<i>Wrangelia tanegana</i>	Rhodophyta	DCM	<62.5 µg/mL	High Cytotoxicity
13	<i>Sargassum prismaticum</i>	Phaeophyta	DCM	<62.5 µg/mL	High Cytotoxicity
14	<i>Dictyopteris acrostichoides</i>	Phaeophyta	EtOAc	<62.5 µg/mL	High Cytotoxicity
15	<i>Padina boergesenii</i>	Phaeophyta	EtOAc	<62.5 µg/mL	High Cytotoxicity




Figure 1. Graphic of percent mortality of *Artemia salina* nauplii after 24 h exposed in various concentrations of algae extracts (number on the graphic related to the number in Table 4).

Of six n-hexan extracts, only n-hexan extract of brown alga *Dictyopteris acrostichoides* had intermediary activity towards the nauplii of the brine shrimp *A. salina*

(LC<sub>50</sub> value between 125-250 µg/mL), whereas others showed high cytotoxicity with LC<sub>50</sub> below 62.5 µg/mL. Two dichloromethane extracts obtained from red alga *Sarconema filiforme* and green alga *Codium dwarkense* had no lethal effect (LC<sub>50</sub> >1000 µg/mL) and intermediary cytotoxicity (LC<sub>50</sub> occurred between 62.5 and 125 µg/mL) respectively. Ethyl <sup>5</sup>acetate extracts of two brown algae *Dictyopteris acrostichoides* and *Padina boergesenii* showed a potent cytotoxic effect against *Artemia salina* nauplii (LC<sub>50</sub> value below 62.5 µg/mL) (Table 4).

## DISCU<sup>47</sup>SSION

A number of studies have demonstrated that marine algae secondary metabolites from different geographical region had cytotoxic effects against *Artemia salina*. The algae, *Ulva fasciata* and *Hypnea musciformis* collected from India <sup>10</sup> showed moderately cytotoxicity against brine shrimp (Selvin and Lipton, 2004). El-Baroty *et al.* (2007) demonstrated that the powdered red alga *Asparagopsis taxiformis* showed cytotoxic activities on *Daphnia magna*. In our study, almost all extracts from seven species of marine algae exhibited high and moderate cytotoxic activities against *Artemia salina* nauplii.

In this study, the brown algae were established as a rich source of cytotoxic compounds since 10 of 14 potential extracts were obtained from these species. These results are comparable to the results of other <sup>31</sup> cytotoxic study of algae, which five ethanolic extracts of brown algae species namely *Stoechospermum marginatum*, *Sargassum* <sup>39</sup> *artzii*, *Sargassum binderi*, *Spatoglossum asperum*, *Stokeyia indica* showed significant cytotoxicity on brine shrimp assay (Ara *et al.*, 1999). From the other study with brown algae <sup>35</sup> *Sargassum ringgoldianum* and *Porphyra yezoensis*, a potential antitumor activity to ehrlich carcinoma and meth A fibrosarcoma was detected (Noda *et al.*, 1989). Beside brown algae, many of the secondary metabolites produced by the marine <sup>40</sup> algae are well known for their cytotoxic property against cancer and tumor cells. As noted by Harada and Kamei (1997) that the extract of red alga, *Amphiroa zonata* exhibited strong cytotoxicity to human leukemic cell line.

<sup>8</sup> In this study, from 15 non-polar extracts, almost all were effective against *Artemia salina* nauplii. This may be due to a non-polar structure of cytotoxic compound(s). Of all tested extracts, the hexane extracts performed high and moderate cytotoxic activities against the nauplii of *Artemia salina*. From the study of antitumor activity of 42 Japanese seaweeds species against K562 human leukemia cells, significant cytotoxic activity was found in hexane-extract from *Protomonostroma undulatum*, *Enteromorpha linza*, *Monostroma latissimum*, *Scytosiphon lomentaria*, *Hizikia fusiformis* and *Sargassum thunbergii*, while cytostatic activity was indicated in hexane-extract from *Colpomenia sinuosa*, *E<sup>38</sup>cklonia cava*, *Undaria pinnatifida* and *Sargassum muticum* and *Chondria crassicaulis*. The hexane-extract of *Ecklonia cava* (Phaeophyceae) had the highest antitumor activity which inhibited cell growth by 45, 92.1 and 96.4% at 10, 20 and 50 µg/ml, respectively. Hexane extract of *E. cava* also induced apoptosis of K562 cells (Kaori, 2002). This is also related with our study, where the hexane extracts of brown algae (Phaeophyceae) showed more potential than the other algae. The dichloromethane extracts showed <sup>16</sup> also good potential against *Artemia* nauplii. Five of seven DCM-extracts exhibited high cytotoxic <sup>20</sup> activity against *Artemia salina* nauplii (LC<sub>50</sub> = <62.5 µg/mL). From the other study, the crude dich<sup>20</sup>loromethane:chloroform extract of *Styopodium zonale*, a Brazilian coastal seaweeds, showed good cytotoxic activity against the C32 cell line (Rocha *et al.*, 2006). In this study, all ethyl acetate extracts of brown algae showed high cytotoxicity against the nauplii of *Artemia salina*. <sup>5</sup> This was also performed in the other studies, where the ethyl acetate extracts of <sup>5</sup> *Sarcilaria salicornia* and *Hypnea flagelliformis*, collected from Persian Gulf showed potent cytotoxic effect against *Artem<sup>3</sup>ia salina* nauplii with LC<sub>50</sub> values of 3 and 4 µg/ml, respectively (Saeidnia *et al.*, 2009). The ethyl acetate extracts of *Sargassum thunbergii*

and *Dictyopteris divaricata* showed excellent cytotoxic activity against the HL-60 cell line. Furthermore, the *Sargassum thunbergii* extract also exhibited good cytotoxic activity against the HT-29 and B16F10 cell lines. These results suggest that *S. thunbergii* and *D. divaricata* have great potential value as food additives, medicinal supplements for patients with chronic diseases and preventive agents against cancer (Kim *et al.*, 2009).

Other than the extracts, many specific compounds also have been found in algae. Fucoidan, laminarin and terpenoids stated to possess anticancer, antitumor and antiproliferative properties (Smith, 2004). These cytotoxic compounds could be further explored as novel leads in cancer chemoprevention and complementary chemotherapy and necessitates further investigation (Vinayak *et al.*, 2010). Fucoidan, a sulphated polysaccharides isolated from brown algae *Laminaria cichorioides*, exerts chemopreventive effects. It inhibits neoplastic cell transformation induced by epidermal growth factor or a tumor promoter (12-O-tetradecanoylphorbol-13-acetate) (Anonim, 2009). Beside fucoidan, other highly sulphated polysaccharide isolated from *Gracilaria domingensis* inhibited the transplantation of Ehrlich ascites carcinoma in mice (Fernandez *et al.*, 1989). The polysaccharide DAEB was isolated and purified from *Enteromorpha intestinalis* and had potent antitumor activity which may be associated with its potent immunostimulating effect (Jiao *et al.*, 2009). The *in vitro* and *in vivo* antitumor properties of a sulfated polysaccharide isolated from the alga *C. feldmannii* (Cf-PLS) has some interesting anticancer activity that could be associated with its immunostimulating properties (Lins *et al.*, 2008).

Many species from genus *Sargassum*, such as *Sargassum micracanthum* (Mori *et al.*, 2005), *Sargassum caryophyllum* (Tang *et al.*, 2002) and *Sargassum tomentosum* (Numata *et al.*, 1991) exhibited cytotoxic activity against cancer cell lines. The sulphated polysaccharides of *Sargassum* act as a potent anticancer agent (Dias *et al.*, 2005). In our studies, one of genus *Sargassum* (Sargassaceae), *Sargassum prismaticum* showed high cytotoxic activity on *Artemia* nauplii.

## CONCLUSION

High price of the medicine and difficult access to the pharmacy could be the reason for poor people in the coastal community to provide their drug with marine medicinal plant surround their area. As we know, culture location of algae or seaweeds is in the sub-tidal zone with 2 m depth. Like terrestrial plant, marine algae require sun light for their metabolisms and Indonesia is one of the tropical countries which obtain sun light through the year. This advantage makes the algae is easy to be cultivated in Indonesia regions (marine bioprospecting). With  $\pm 75\%$  of marine area (5,8 million km<sup>2</sup>),  $\pm 81.000$  km coast line and 17.500 big and small island, make the marine and fishery industries have good expectations. Sulawesi Selatan is one of the east Indonesia regions which has a potential area to develop the algae culture. Almost half of total regional productions of algae commercial are come from South Sulawesi area. Therefore, the aim of the study is to explore the bioactive compounds of marine algae from South Sulawesi as marine bioprospecting resources. In conclusion, the results of this study showed that marine algae from South Sulawesi can be used as medicinal marine plant for anticancer drugs candidate.

## ACKNOWLEDGMENTS

We sincerely thank to Indonesia government for the financial support through competitive grant DIPA DP2M-DIKTI Stranas 2010.

## REFERENCES

- Aliya R., Shamael M., 1999, Phytochemical evaluation of four coenocytic green seaweeds from the coast of Karachi. *Pakistan Journal of Marine Biology*, **5**, 65-76.
- Anonim. 2009, Fucoidan in seaweed may prevent cancer by interfering with epidermal growth factor binding, Breast cancer study. AACR International Conference on Frontiers in Basic Cancer Research, October 2009.
- Ara J., Sultana V., Ehteshamul-Haque S., Qasim R., Ahmed V., 1999, Cytotoxic activity of marine macro-algae on *Artemia salina* (brine shrimp). *Phytother. Res.* **13**, 304-307.
- Aristides M. L., Lima E. D. O., De Souza E. L., Diniz M. D. F. F. M., Leite S.P., Xavier A.L., De Medeiros I.A., 2008, Preliminary study of the molluscicidal and larvicidal properties of some essential oils and phytochemicals from medicinal plants. *Revista Brasileira de Farmacognosia, Brazilian Journal of Pharmacognosy*, **19**, 842-846.
- Badisa R. B., Badisa V. L. D., Walker E. H., Latinwo L. M., 2007. Potent cytotoxic activity of *Saururus cernuus* extract on human colon and breast carcinoma cultures under normoxic conditions. *Anticancer Research*, **27**(1A): 189-193.
- Bengen D.G., 2001, Ekosistem dan sumberdaya alam pesisir laut. Pusat Kajian Sumberdaya Pesisir dan Lautan IPB: 25 p.
- Critchley A., Ohno M., 1998, Seaweed resources of the world, Japan International Cooperation Agency.
- Darcy-Vrillon B., 1993, Nutritional aspects of the developing use of marine macro algae for the human food industry. *International Journal of Food Sciences and Nutrition*, **44**, 23-35.
- Dias P. F., Siqueira J. M., Vendruscolo L. F., de Jesus Neiva T., Gagliardi A. R., M. Maraschin, Ribeiro-do-Valle R.M., 2005. Antiangiogenic and antitumoral properties of a polysaccharide isolated from the seaweed *Sargassum stenophyllum*. *Cancer chemoth Pharm.* **56**(Supple 4):436-446.
- El-Baroty G. S., Moussa M. Y., Shallan M. A., Ali M. A., Sabh A.Z., Shalaby E.A., 2007, Contribution to the aroma, biological activities, minerals, protein, pigments and lipid contents of the red alga: *Asparagopsis taxiformis* (Delile) Trevisan. *J. Applied Sci. Res.*, **3**, 1825-1834.
- Faulkner D. J., 2001, Marine natural products, *Natural Product Reports*, **18**, 1-49.
- Fernández L. E., Valiente O. G., Mainardi V., Bello J. L., Vélez H., Rosado A., 2001, Isolation and characterization of an antitumor active agar-type polysaccharide of *Gracilaria dominguensis*. *Carbohydrate Research*, **190**, 77-83.
- Garg H.S., Sharma T., Bhakuni D. S., Pramanik B.N., Bose A.K., 1992, An antiviral sphingosine derivative from green alga *Ulva fusiata*, *Tetrahedron Letters*, **33**, 1641-1644.
- Harada H., Kamei Y., 1997, Selective cytotoxicity of marine algae extracts to several human leukemic cell lines. *Cytotechnology* **25**, 213-219.
- Ito K., Hori K., 1989, Seaweed: Chemical composition and potential foods uses. *Food Rev. Int.*, **5**, 101-104.
- Jiao L., Li X., Li T., Jiang P., Zhang L., Wu M., Zhang L., 2009, Characterization and anti-tumor activity of alkali-extracted polysaccharide from *Enteromorpha intestinalis*. *International Immunopharmacology*, **9**, 324-329.
- Kaori K. (2002). Screening of seaweeds possessing antitumor activity against K562 leukemia cells. *Urakami Foundation Memoirs*, **10**, 44-53.
- Kim K. N., Ham Y. M., Moon J. Y., Kim M.J., Kim D.S., 2009, *In vitro* cytotoxic activity of *Sargassum thunbergii* and *Dictyopteris divaricata* (Jeju Seaweeds) on the HL-60 tumour cell line. *Int. J. Pharmacol.*, **9**, 298-306.

- Lahaye M., 1991, Marine algae as sources of fibres: Determination of soluble and insoluble dietary fibre contents in some 'sea vegetables'. *Journal of the Science of Food and Agriculture*, **54**, 587–594
- Lee R. E., 1999, Phycology. Cambridge, Cambridge University Press.
- Lins K. O. A. L., Bezerra D. P., Alves A. P. N. N., Alencar N. M. N., Lima M. W., Torres V. M., Farias W. R. L., Pessoa C., De Moraes M. O., Costa-Lotufo L. V., 2008, Antitumor properties of a sulfated polysaccharide from the red seaweed *Champia feldmannii* (Diaz-Pifferer). *Journal of Applied Toxicology*, **9**, 20-26.
- Manilal A., Sujith S, Kiran G. S., Selvin J., Shakir C., 2009a, Cytotoxic potentials of red alga, *Laurencia brandenii* collected from the Indian Coast. *Global Journal of Pharmacology*, **3**, 90-94.
- Manilal A., Sujith S, Kiran G. S., Selvin J., Shakir C., Gandhimathi R., Panikkar M. V. N., 2009b, Biopotential of seaweeds collected from Southwest coast of India. *Journal of Marine Science and Technology*, **17**, 67-73.
- Meyer B. N., Ferrigni N. R., Putnam J. E., Jacobsen L. B., Nichols D.E., McLaughlin J. L., 1982, Brine shrimp: a convenient general bioassay for active plant constituents. *Planta Medicine*, **45**, 31-34.
- Mori J., Iwashima M., Wakasugi H., Saito H., Matsunaga T., Ogasawara M., Takahashi S., Suzuki H., Hayashi T., 2005, New plastoquinones isolated from the brown alga, *Sargassum micracanthum*. *Chem Pharm Bull*, **53**, 1159-1163.
- Munro M. H. G., Blunt J. W., 1999, Marinlit, version10.4, Marine Chemical Group, University of Canterbury, Christchurch, New Zealand.
- Noda H., Amano H., Arashima K., Hashimoto S., Nisizawa K., 1989, Antitumour activity of polysaccharides and lipids from marine algae. *Nippon Suisan Gakkaishi*, **55**, 1265-1271.
- Numata A., Kanbara S., Takahashi C., Fujiki R., Yoneda M., Fujita E., Nabeshima Y., 1991, Cytotoxic activity of marine algae and a cytotoxic principle of the brown alga *Sargassum tortile*. *Pharmaceutical Society of Japan*, **39**, 2129-2131.
- Parra A. L., Yhebra R. S., Sardinas I. G., Buela L. I., 2001, Comparative study of the assay of *Artemia salina* L. and the estimate of the medium lethal dose (LD50 value) in mice, to determine oral acute toxicity of plant extracts. *Phytotherapy* **8**, 395–400.
- Rocha F. D., Soares A. R., Houghton P. J., Pereira R. C., Kaplan M. A. C., Teixeira V. L., 2007, Potential cytotoxic activity of some Brazilian seaweeds on human melanoma cells. *Phytotherapy Research*, **21**, 170–175.
- Saeidnia S., Gohari A. R., Shahverdi A. R., Perme P., Nasiri M., Mollazadeh K., Farahani F., 2009, Biological activity of two red algae, *Gracilaria salicornia* and *Hypnea flagelliformis* from Persian Gulf. Published on Pharmacognosy Network Worldwide 1(6): 428-430.
- Selvin J., Lipton A. P., 2004, Biopotentials of *Ulva fasciata* and *Hypnea musciformis* collected from the Peninsular Coast of India. *Journal of Marine Science and Technology*, **12**, 1-6.
- Serkedjiva J., 2004, Antiviral activity of the red marine alga *Ceramium rubrum*. *Phytotherapy Research*, **18**, 480–483.
- Skibola C., 2004, The effect of *Fucus vesiculosus*, an edible brown seaweed, upon menstrual cycle length and hormonal status in three pre-menopausal women: a case report. *BMC Complementary and Alternative Medicine*, **4**, 10–17.
- Svensson B. M., Mathiasson L., Martensson L., Bergstrom S., 2005, *Artemia salina* as test organism for assessment of acute toxicity of leachatewater from landfills. *Environmental Monitoring and Assessment*, **102**, 309-321.
- Tang H. F., Yi Y. H., Yao X. S., Xu Q. Z., Zhang S. Y., Lin H. W., 2002, Bioactive steroids from the brown alga *Sargassum caryophyllum*. *J Asian Nat Product Res.*, **4**, 95-105.

- Teas J., 1981, The consumption of seaweed as a protective factor in the etiology of breast cancer. *Medical Hypotheses*, **7**, 601–613.
- Thangam T. S., Kathiresan K., Mabbett T., 1993, Mosquito larvicidal activity of seaweed extract against *Aedes aegypti* and *Culex quinquefasciatus*. *International Pest Control*, **35**, 94-95.
- Vallinayagam K., Arumugam R., Kannan R. R. R., Thirumaran G., Anantharaman P., 2009, Antibacterial Activity of Some Selected Seaweeds from Pudumadam Coastal Regions. *Global Journal of Pharmacology* **3**, 50-52.
- Van den Hoek C., Jahns H. M., Mann D. G., 1993, Algen. Stuttgart-New York, Georg Thieme Verlag
- Vinayak R. C., Sabu A. S., Chatterji A., 2010, Bio-prospecting of a few brown seaweeds for their cytotoxic and antioxidant activities, Oxford University Press.
- Vinayak R. C., Sabu A. S., Chatterji A., 2010, Bio-prospecting of a few brown seaweeds for their cytotoxic and antioxidant activities. *eCAM* 2010: 1-9.
- Yamamoto I., Maruyama H., Moriguchi M., 1987, The effect of dietary seaweeds on 7,12-dimethylbenz[*a*]anthracene-induced mammary tumorigenesis in rats The effect of dietary seaweeds on 7,12-dimethylbenz[*a*]anthracene-induced mammary tumorigenesis. *Cancer Letters*, **35**, 109-118.
- Yuan Y.V., Walsh N. A., 2006, Antioxidant and antiproliferative activities of extracts from a variety of edible seaweeds. *Food and Chemical Toxicology*, **44**, 1144-1150.
- Zainuddin E.N., 2006, Chemical and Biological Investigations of Selected Cyanobacteria (Blue-green Algae). PhD Thesis, Institute of Pharmaceutical Biology, Ernst Moritz Arndt University of Greifswald, German.

ORIGINALITY REPORT

---

**%24**  
SIMILARITY INDEX

**%16**  
INTERNET SOURCES

**%21**  
PUBLICATIONS

**%3**  
STUDENT PAPERS

---

PRIMARY SOURCES

---

**1** Leite, Aristides M., Edeltrudes de O. Lima, Evandro L. de Souza, Margareth de F. F. M. Diniz, S nia Pereira Leite, Aline L. Xavier, and Isac A. de Medeiros. "Preliminary study of the molluscicidal and larvicidal properties of some essential oils and phytochemicals from medicinal plants", *Revista Brasileira de Farmacognosia*, 2009. **%2**  
Publication

---

**2** Elisabeth Catherina Widjajakusuma, Arijanto Jonosewojo, Lucia Hendriati, Sumi Wijaya et al. "Phytochemical screening and preliminary clinical trials of the aqueous extract mixture of *Andrographis paniculata* (Burm. f.) Wall. ex Nees and *Syzygium polyanthum* (Wight.) Walp leaves in metformin treated patients with type 2 diabetes", *Phytomedicine*, 2019 **%2**  
Publication

---

**3** [www.scialert.net](http://www.scialert.net) **%1**  
Internet Source

---

4	Food Factors for Cancer Prevention, 1997. Publication	% 1
5	<a href="http://www.phcogres.com">www.phcogres.com</a> Internet Source	% 1
6	<a href="http://jmst.ntou.edu.tw">jmst.ntou.edu.tw</a> Internet Source	% 1
7	Kézia O. A. L. Lins. "Antitumor properties of a sulfated polysaccharide from the red seaweed <i>Champia feldmannii</i> (Diaz-Pifferer)", <i>Journal of Applied Toxicology</i> , 01/2009 Publication	% 1
8	<a href="http://www.scielo.cl">www.scielo.cl</a> Internet Source	% 1
9	<a href="http://www.pakbs.org">www.pakbs.org</a> Internet Source	% 1
10	<a href="http://baadalsg.inflibnet.ac.in">baadalsg.inflibnet.ac.in</a> Internet Source	% 1
11	<a href="http://ijpsr.com">ijpsr.com</a> Internet Source	% 1
12	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	% 1
13	<a href="http://www.omicsonline.org">www.omicsonline.org</a> Internet Source	% 1
14	Shenshen Zhang, Shaoping Nie, Danfei Huang,	

Jianqin Huang, Yawei Wang, Mingyong Xie.  
"Polysaccharide from Ganoderma atrum Evokes  
Antitumor Activity via Toll-like Receptor 4-  
Mediated NF-κB and Mitogen-Activated Protein  
Kinase Signaling Pathways", Journal of  
Agricultural and Food Chemistry, 2013

Publication

% 1

15

Submitted to Higher Education Commission  
Pakistan

Student Paper

% 1

16

[www.ijpr.sbmua.ac.ir](http://www.ijpr.sbmua.ac.ir)

Internet Source

<% 1

17

Amal D. Premarathna, T.H. Ranahewa, S.K.  
Wijesekera, D.L. Harishchandra et al.  
"Preliminary screening of the aqueous extracts  
of twenty-three different seaweed species in Sri  
Lanka with in-vitro and in-vivo assays", Heliyon,  
2020

Publication

<% 1

18

"Recent Advances in Environmental Science  
from the Euro-Mediterranean and Surrounding  
Regions", Springer Science and Business Media  
LLC, 2018

Publication

<% 1

19

[www.mdpi.com](http://www.mdpi.com)

Internet Source

<% 1

Chennappan Kamal, Mathur Gopalakrishnan

- 20 Sethuraman. "Caulerpin—A bis-Indole Alkaloid As a Green Inhibitor for the Corrosion of Mild Steel in 1 M HCl Solution from the Marine Alga *Caulerpa racemosa*", Industrial & Engineering Chemistry Research, 2012  
Publication <% 1
- 
- 21 Mookkan Bodhaguru, Santhiyagu Prakash, Ramasamy Ramasubburayan, Natarajan Kumari Ahila et al. "Screening, partial purification of antivibriosis metabolite sterol-glycosides from *Rhodococcus* sp. against aquaculture associated pathogens", Microbial Pathogenesis, 2019  
Publication <% 1
- 
- 22 [file.scirp.org](http://file.scirp.org)  
Internet Source <% 1
- 
- 23 [link.springer.com](http://link.springer.com)  
Internet Source <% 1
- 
- 24 [repository.uin-malang.ac.id](http://repository.uin-malang.ac.id)  
Internet Source <% 1
- 
- 25 [www.idosi.org](http://www.idosi.org)  
Internet Source <% 1
- 
- 26 Sevim Polat, >Yesim Ozogul. "Biochemical composition of some red and brown macro algae from the Northeastern Mediterranean Sea", International Journal of Food Sciences <% 1

---

27 Vimala Balasubramaniam, Suraiami Mustar, Norhayati Mustafa Khalid, Aswir Abd Rashed et al. "Inhibitory activities of three Malaysian edible seaweeds on lipase and  $\alpha$ -amylase", Journal of Applied Phycology, 2013

Publication

---

28 Submitted to University of Central Lancashire

Student Paper

---

29 [www.ijpba.info](http://www.ijpba.info)

Internet Source

---

30 [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)

Internet Source

---

31 Jehan Ara. "Cytotoxic activity of marine macroalgae on Artemia salina (Brine shrimp)", Phytotherapy Research, 06/1999

Publication

---

32 69.164.208.4

Internet Source

---

33 [www.biomedcentral.com](http://www.biomedcentral.com)

Internet Source

---

34 "Final Report on the Safety Assessment of Tocopherol, Tocopheryl Acetate, Tocopheryl Linoleate, Tocopheryl Linoleate/Oleate,

Tocopheryl Nicotinate, Tocopheryl Succinate, Dioleoyl Tocopheryl Methylsilanol, Potassium Ascorbyl Tocopheryl Phosphate, and Tocophersolan", International Journal of Toxicology, 2016

Publication

35

Jinju Jiang, Shaojuan Shi. "Seaweeds and Cancer Prevention", Elsevier BV, 2018

Publication

<% 1

36

[www.jove.com](http://www.jove.com)

Internet Source

<% 1

37

M. Kalaiselvi, R. Narmadha, P. Ragavendran, B. Vidya et al. "Chemopreventive Effect and HPTLC Fingerprinting Analysis of Jasminum sambac (L.) Ait. Extract Against DLA-Induced Lymphoma in Experimental Animals", Applied Biochemistry and Biotechnology, 2013

Publication

<% 1

38

[creativecommons.org](http://creativecommons.org)

Internet Source

<% 1

39

[pjek.org.pk](http://pjek.org.pk)

Internet Source

<% 1

40

Masatoshi Moku, Kouichi Kawaguchi. "Chemical composition of three dominant myctophid fish, , , and , in the subarctic and transition waters of the western North Pacific ",

<% 1

# Journal of the Marine Biological Association of the United Kingdom, 2008

Publication

- 
- 41** Jorge Güette-Fernández, Jesús Olivero-Verbel, Indira O'Byrne-Hoyos, Beatriz Jaramillo, Elena Stashenko. " Chemical Composition and Toxicity Against of the Essential Oil of (Sims) DC. Collected in Bogota (Colombia) ", Journal of Essential Oil Research, 2008  
Publication <% 1
- 
- 42** [www.ukm.my](http://www.ukm.my)  
Internet Source <% 1
- 
- 43** [ijppr.humanjournals.com](http://ijppr.humanjournals.com)  
Internet Source <% 1
- 
- 44** [www.jmbfs.org](http://www.jmbfs.org)  
Internet Source <% 1
- 
- 45** El Gamal, Ali A.. "Biological Importance of Marine Algae", Handbook of Marine Macroalgae Biotechnology and Applied Phycology, 2011.  
Publication <% 1
- 
- 46** Seafood Processing By-Products, 2014.  
Publication <% 1
- 
- 47** Polysaccharides, 2015.  
Publication <% 1
- 
- 48** Hb25\_Springer Handbook of Marine <% 1

49

D. Senthil Rajan, M. Rajkumar, R. Srinivasan, R.P. Harikumar, S. Suresh, Senthil Kumar KL. "Antitumour Activity of Sargassum wightii (Greville) Extracts against Dalton's Ascites Lymphoma", Pakistan Journal of Biological Sciences, 2013

Publication

---

<% 1

50

Rashmi C. Vinayak, A. S. Sabu, Anil Chatterji. "Bio-Prospecting of a Few Brown Seaweeds for Their Cytotoxic and Antioxidant Activities", Evidence-Based Complementary and Alternative Medicine, 2011

Publication

---

<% 1

---

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 5 WORDS