

# Isolation and characterization of N-fixing and IAA producing rhizobacteria from two rice field agro-ecosystems in South Sulawesi, Indonesia

*by* Feranita Feranita

---

**Submission date:** 07-Jan-2022 10:08AM (UTC+0700)

**Submission ID:** 1738354229

**File name:** 04\_BIODIVERSITA-VOL.\_MEI-2021-0423\_.pdf (954.25K)

**Word count:** 5207

**Character count:** 28623

6

## Isolation and characterization of N-fixing and IAA producing rhizobacteria from two rice field agro-ecosystems in South Sulawesi, Indonesia

NINING HAERANI<sup>1,2,\*</sup>, ELKAWAKIB SYAM'UN<sup>3,\*\*</sup>, BURHANUDDIN RASYID<sup>4,\*\*\*</sup>, FERANITA HARING<sup>3,\*\*\*\*</sup>

<sup>1</sup>Program of Agriculture, Graduate School, Universitas Hasanuddin, Jl. Perintis Kemerdekaan Km.10, Tamalanrea, Makassar 90245, South Sulawesi, Indonesia. Tel.: +62-411-587064, \*email: nininghaerani06@gmail.com

<sup>2</sup>Program of Agrotechnology, Faculty of Agriculture, Animal Science and Forestry, Universitas Muslim Maros. Jl. Dr. Ratulangi no. 62 Maros, South Sulawesi, Indonesia

<sup>3</sup>Departemen of Agrotechnology, Faculty of Agriculture, Universitas Hasanuddin, Jl. Perintis Kemerdekaan Km. 10, Tamalanrea, Makassar 90245, South Sulawesi, Indonesia. Tel.: +62-411-587064, \*\*email: elkawakib@yahoo.com, \*\*\*\*email: feraharing20@gmail.com

<sup>4</sup>Department of Soil Science, Faculty of Agriculture, Universitas Hasanuddin, Jl. Perintis Kemerdekaan Km. 10, Tamalanrea, Makassar 90245, South Sulawesi, Indonesia. Tel.: +62-411-587064, \*\*\*email: burrasyid@unhas.ac.id

Manuscript received: 8 December 2020. Revision accepted: 9 April 2021.

**Abstract.** Haerani N, Syam'un E, Rasyid B, Feranita. 2021. Isolation and characterization of N-fixing and IAA producing rhizobacteria from two rice field agro-ecosystems in South Sulawesi, Indonesia. *Biodiversitas* 22: 2497-2503. Nitrogen-fixing bacteria that live in the rhizosphere are microorganisms that can increase N availability for rice plants in a non-symbiotic manner. Nitrogen-fixing bacteria also support growth through the production of growth hormones. These abilities are mainly influenced by the environment and their association with plants. This research aims to determine the microbial characteristics from two different rice fields agro-ecosystems (rainfed and irrigated) in South Sulawesi. Bacteria isolated from the rhizosphere soil of rice in six districts were cultured and isolated using Burk's N-free media for screening purposes. A total of 20 isolates were observed for their morphological and biochemical properties, determined their ability to fix nitrogen and produce IAA. The results showed that all isolates could grow on nitrogen-free media, in which 13 isolates were isolated from rainfed rice fields, and 7 isolates were from irrigated rice fields with various physical and biochemical properties, all other qualities tested also varied. NG4 isolate from Gowa Regency (rainfed rice field) produced the highest nitrogen (0.157%), and NB13 isolate from Barru (rainfed rice field) had the highest concentration of growth hormone (IAA) (1,835 mg L<sup>-1</sup>). All isolates obtained from rainfed rice fields showed a better ability to fix nitrogen and produce growth hormone than those from irrigated rice fields. Bacteria isolates from rainfed rice fields have higher biofertilizers and biostimulant capacity to develop environmentally-friendly rice cultivation.

**Keywords:** IAA, irrigated and rainfed rice fields, nitrogen fixation, rhizobacteria

### INTRODUCTION

Continuous use of land, especially for rice cultivation, will have an impact on decreasing land quality and productivity. The subsequent land management through fertilization often leads to a decrease in organic matter and fertility. It also affects soil physicochemical, biochemical, and microbiological properties. Inorganic nitrogen sources, such as urea, often reduce land efficiency when applied and are not organically bound to N; therefore, the plant becomes deficient in nitrogen (Witte 2011). However, biological approaches to improve soil quality and fertility will increase rice production and reduce synthetic fertilizers consumption (Sun et al. 2019). It also reduces the environmental problems associated with the excessive use of fertilizers. Therefore, improving biological nitrogen fixation on rice fields is very important (Ma et al. 2019).

The use of non-symbiotic nitrogen-fixing bacteria as a biofertilizer agent has been widely developed in agriculture (Pereg et al. 2018). The relevance of manure is increasing rapidly because chemical fertilizers damage the environment (Vinten et al. 2002; Tahovská et al. 2013).

However, biological fertilizer leads to soil enrichment, long-term sustainability, and environmentally friendly products (Shridhar et al. 2012).

Nitrogen comprises about 78% of the atmosphere's gases; however, plants do not use it directly, although it is abundant in the air (Rascio and La Rocca 2018). Therefore, chemical or biological processes are needed to convert N gas into forms utilized by plants (Dong et al. 2019). Soil bacteria carry out the natural fixation of nitrogen from the atmosphere to benefit the plants (Li et al. 2021). Rhizobacteria are microorganisms that live in plants' roots (rhizosphere) (Khatoun et al. 2020). They can reduce free nitrogen in the atmosphere to ammonia compounds (Pas Aksamir et al. 2015).

Previous studies showed that rhizobacteria were reported as plant growth promoters by fixing atmospheric nitrogen and producing hormones (Shridhar et al. 2012). However, further studies are still needed to determine their biofertilization and biostimulant activity in various rice fields agro-ecosystem conditions. Israwan et al. (2015) reported that the surrounding environment strongly influences bacterial strains' ability to survive and produce

metabolites. Munif et al. (2012) also reported that environmental factors such as soil properties, organic matter present, cultivation techniques, fertilization, and pesticide application affect bacterial strain. The same microbe species from different places do not necessarily have the same potential.

Rice fields in Indonesia consist of irrigated, rainfed, tidal, and lowland soil, with a total area of 8.1 million ha (Widiastuti 2017). In South Sulawesi, it is dominated by irrigated and rainfed fields. Therefore, this research aims to determine the bacteria characteristics from two wetland ecosystems (rainfed and irrigated) in South Sulawesi, which can fix nitrogen and produce IAA (Indole Acetic Acid). These bacteria can be recommended as good biofertilizers and bio stimulating agents to develop environmentally friendly and sustainable rice cultivation.

## 43 MATERIALS AND METHODS

### Study area

This research was conducted from July to October 2019 at the Laboratory of Soil Microbiology, Department of Soil Science, Faculty of Agriculture, Hasanuddin University, Makassar, Indonesia. Soil samples were collected from six districts in South Sulawesi. Three samples from rainfed rice field in Gowa Regency (119°22'30.202"E; 5°34'6.423"S), Pangkep (119°30'33.005"E; 4°56'33.936"S), and Barru (119°35'28.889"E; 4°33'8.955"S). Three other samples from irrigated rice fields in Takalar (119°23'1.499"E; 5°12'51.978"S), Maros (119°29'36.983"E; 5°12'11.804"S), and Sidrap (119°39'51.033"E; 3°28'59.109"S) respectively (Figure 1). Soil samples were collected from roots of 25-40 days old of healthy rice in the depth of 0-20 cm at five

points with an X pattern for each location. It was transferred into paper envelopes labeled according to the location, and the mixture was composited. Samples were taken to the laboratory using a cool box for immediate analysis (Kesaulya et al. 2015).

### 24 Isolation and characterization of nitrogen fixing bacteria

The isolation of Nitrogen-fixing bacteria was performed using Burk's N-free media (Park et al. 2005). Burk's N-free solid medium contained 20 g of sucrose, 0.64g of  $K_2HPO_4$ , 0.16 g of  $KH_2PO_4$ , 0.20 g of  $MgSO_4 \cdot 7H_2O$ , and 0.20 g of NaCl, 0.05 g of  $CaSO_4 \cdot 2H_2O$ , 5 mL of  $Na_2MoO_4 \cdot 2H_2O$  (0.05%), 5 mL of  $FeSO_4 \cdot 7H_2O$  (0.3%), 21 g of agar and 1000 mL of aqua dest (Hartono and Oslan Jumadi 2014). Serial sample dilution was carried out by grinding 1 gram of soil using a mortar until it was finely ground and then added with 10 mL of distilled water while stirring till it became a homogeneous solution. Furthermore, graded dilutions from  $10^{-1}$  to  $10^{-6}$  were carried out by taking 1mL of the mixture using a pipette. Then, it was placed in the first test tube containing 9 mL of sterile distilled water, shaking it until it was consistent in composition until a vortex. It was referred to as the first dilution ( $10^{-1}$ ). This procedure was repeated until  $10^{-6}$  (the 6<sup>th</sup> test tube). 0.1 mL of the diluted soil samples  $10^{-3}$ ,  $10^{-4}$ ,  $10^{-5}$ , and  $10^{-6}$  were cultured on the Burk's medium using a triangle spreader rod and incubated for 24 hours at room temperature. Single colonies that grew on the growth media were then cultured by streaking on NA medium and purified 3 times. Pure colonies that grew with different appearances were characterized morphologically. It was performed according to the method proposed by Cappuccino and Sherman (2011).




Figure 1. Location for rice rhizosphere sampling in South Sulawesi, Indonesia

**Gram test**

The Gram test was carried out using the procedure by Pambudi et al. (2017). One loop full of bacterial isolates was taken using a loop needle and placed on an object glass that had been dripped with 10 µL of 3% KOH (Potassium Hydroxide) then stirred repeatedly. If the mixture of bacteria suspension and KOH suspension becomes markedly viscid or forms a gel within 5 to 60 s, the isolate is classified as Gram-negative (-). If no gelling is observed, the isolate is classified as Gram-positive (+).

**Catalase test**

Catalase test was carried out according to the method by Suhartanti et al. (2010) with modification. This test was conducted to determine the ability of microbes to degrade hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>). One loop full of pure single colony culture was smeared on a previously dripped slide with two drops of 3% H<sub>2</sub>O<sub>2</sub>. A positive reaction was characterized by the emergence of gas bubbles from free oxygen.

**The nitrogen fixing ability test**

The bacterial ability to fix nitrogen qualitatively tested on Burk's N-free liquid media according to the method described by Park et al. (2005). Bacterial isolates were grown on Burk's N-free media in glass bottles and placed in an orbital shaker for 24 hours at 28 °C.

The total N content of bacterial cultures was measured quantitatively using the Kjeldahl method (Rao 2010; Ramadhan and Tiwow 2016). As much as 5 mL of supernatant was put into the digestion tube and added 1 g of a mixture of selenium and 3 mL of concentrated sulfuric acid (H<sub>2</sub>SO<sub>4</sub>) and digested for 3-4 hours at 35°C until white steam comes out. Furthermore, the tube was removed and cooled down. Then, the extract was diluted with distilled water 50 mL and shaken until homogeneous. The extract was allowed to stand for 24 hours for the particles to settle. The extract was transferred into the Erlenmeyer flask as a boiling flask containing 10 mL of 1% boric acid and 3 drops of Conway indicator (red-colored) and connected to a distillation device. Ten mL of 40% NaOH was added to the boiling flask closed immediately. The distillation process was carried out until the container's volume reached 50 - 75 mL (green). The nitrogen content (%) was then calculated using the formula:

$$\text{Nitrogen Level (\%)} = \frac{\{(vc - vb) \times N \times bst \times N\}}{c} \times 100 \%$$

Where:

Vc : Distillation titration example

Vb : Blank

N : Normality of H<sub>2</sub>SO<sub>4</sub> standard solution

bst N: nitrogen equivalent weight (14)

C : mass of the sample

100 : conversion to%

**The IAA production test of rhizobacteria**

The ability of rhizobacteria to produce IAA hormone was tested using a spectrophotometer. The microorganisms were cultured on Nutrient Agar (NA) media contained 200 ppm L-tryptophan. Then, 10 mL of the bacterial isolate was taken and placed in a 15 mL centrifuge tube and was centrifuged at 8000 rpm for 10 minutes. Two mL of supernatant was taken and placed in a test tube and added with 2 drops of orthophosphoric acid and 4 mL of Salkowski's reagent (Glickman and Dessaux 1995). Furthermore, it was incubated in a dark room for 24 hours. The change in color to pink indicated the microbe's ability to produce auxin. The pink color is caused by the reaction between Salkowski's reagent and the hormone. The optical density was measured at a wavelength of 535 nm using a UV-VIS spectrophotometer (Genesys IOS UV 840208100). The concentration of IAA was calculated using a linear regression equation from the IAA standard curve (Kesaulya et al. 2015; Israwan and Ardyati 2015).

**RESULTS AND DISCUSSION****Isolation and characterization of nitrogen-fixing bacteria from rice rhizosphere**

Twenty bacterial isolates with the potential to fix nitrogen from 6 sampling locations had various morphological and biochemical characteristics (Table 1). The ability of bacteria to fix nitrogen was confirmed by growing them on Burk's N-free media so that only bacteria with the ability to attach N that grow on it (Stella and Suhaimi 2010). Table 1. shows that 13 isolates were collected from rainfed rice fields (Gowa, Pangkep, Barru), and 7 isolates were from irrigated rice fields (Takalar, Maros, Sidrap).


Colonies with moderate-sized dominated the collected bacteria. Colony sizes ranging from 2.0 mm to 4.0 mm (small to large). The shape was dominated by irregular shapes (13 isolates), 6 rhizoids, and 1 with a circular form. The physical properties based on the edges varied widely; there were some with lobate, filamentous, entire, serrated, and undulate sides. The grouping on elevation showed that 10 isolates had umbonate elevation and the other 10 isolates had altitude. Furthermore, the color was dominated by cream and beige; the rest were white and yellowish. However, the morphological and biochemical characters cannot be used to identify the tested isolates.

Gram test results showed that 13 isolates were Gram-negative, and 7 isolates were Gram-positive (+) bacteria. The results of the catalase test showed that all nitrogen-fixing bacteria were catalase positive. It was indicated by the presence of gas bubbles from free oxygen that produced when the microbe secrete catalase to break down hydrogen peroxide into water and oxygen (Pulungan and Tumangger 2018). Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) is poisonous to bacteria's metabolic system when they cannot degrade the chemical into other harmless compounds (Prathama et al. 2018). However, this is also achievable when a catalase enzyme is available without bacteria (Cappuccino and Sherman 2011).

**1** **Table 1.** The morphological and biochemical characteristics of bacteria isolated from rice rhizosphere at six rice fields in South Sulawesi, Indonesia

Isolate code	Size	Colony morphology				Color	Gram (+/-)	Catalase (+/-)
		Form	Edge	Elevation				
NG1	Moderate	Irregular	Lobate	Raised	Cream	-	+	
NG4	Moderate	Irregular	Entire	Umbonate	Cream	-	+	
NG5	Moderate	Irregular	Serrate	Raised	Beige	+	+	
NG7	Moderate	Irregular	Lobate	34 sed	Beige	+	+	
NG11	Large	19 zoid	Filamentous	Umbonate	Cream	-	+	
NT1	Moderate	Irregular	Lobate	Raised	Cream	+	+	
NT2	Moderate	Irregular	Entire	Raised	Beige	-	+	
NT3	Large	Irregular	Serrate	Umbonate	White	-	+	
NS2	Moderate	Irregular	Serrate	Umbonate	Cream	-	+	
NS13	Small	Irregular	Lobate	Raised	Beige	+	+	
NP1	Large	Rhizoid	Filamentous	Umbonate	Cream	+	+	
NP7	Moderate	Rhizoid	Filamentous	Naik	Beige	-	+	
NP11	Moderate	Irregular	Lobate	Umbonate	Cream	-	+	
NP12	Moderate	Circular	Entire	Umbonate	Cream	-	+	
NM4	Moderate	Circular	Undulate	Raised	White	+	+	
NM5	Small	Irregular	Lobate	Umbonate	Yellow	-	+	
NB8	Large	Rhizoid	Filamentous	Umbonate	White	+	+	
NB10	Moderate	Rhizoid	Filamentous	Umbonate	Yellow	-	+	
NB12	Moderate	Irregular	Serrate	Raised	Cream	-	+	
NB13	Moderate	Rhizoid	Filamentous	Raised	White	-	+	

Notes: NG: isolates collected from Gowa; NT: isolates from Takalar; NS: isolates from Sidrap; NP: isolates from Pangkep; NM: isolates from Maros; NB: isolates from Barru.




**Figure 2.** Pure culture of nitrogen-fixing bacteria from rice rhizosphere collected from several locations in South Sulawesi, Indonesia. A: NG1, B: NG4, C: NT1, D: NT2, E: NP1, F: NP7, G: NB8, H: NB10

### The ability of nitrogen fixation


The quantitative test results of nitrogen-fixing ability were directly proportional to the qualitative examination using Burk's N-free media. All isolates were able to produce nitrogen in varying amounts (Figure 3). The NG4 isolate from Gowa had the highest ability to fix N (0.157%), followed by NP11 isolate from Pangkep (0.153%), and NT2 isolate from Takalar (0.1510%). Meanwhile, the NM5 isolate from Maros produced the lowest (0.132%). Varying ability to fix N caused by the different bacteria species and their capacity to perform this function.

### 9 The ability of rhizobacteria to produce IAA

Bacterial isolates were tested qualitatively and quantitatively for their ability to produce IAA. The results showed that all isolates could produce IAA. The qualitative test results (Figure 4) showed the variations in the color intensity of the supernatant. The result of qualitative analysis was directly proportional to the quantitative test. The NB13 isolate from Barru produced a more intense pink color and had the highest IAA concentration (1,835 mg L<sup>-1</sup>), followed by NG11 isolate from Gowa (1,630 mg L<sup>-1</sup>) and NP1 isolate from Pangkep (1,566 mg L<sup>-1</sup>). Meanwhile, NT3 isolates had the lowest (0.316 mg L<sup>-1</sup>) (Figure 5).



**Figure 3.** Nitrogen fixation ability of bacterial isolates from rice rhizosphere



**Figure 4.** Qualitative analysis of the IAA production by bacterial isolates with Salkowski's reagent. Changes to pink color were determined in the supernatant of: A. NM5, B. NG11, C. NS12, D. NB10, E. NB13, F. NP1, G. NB12, H. NG1, I. NG4, J. NP7

This variation was due to the difference in the ability of the microorganisms to change tryptophan into IAA. According to Lestari et al. (2015), IAA-producing bacteria can be cultivated on growth media containing this amino acid, with the enzyme IAA oxidase. It was synthesized as a secondary metabolite under suboptimal growth conditions (Panigrahi et al. 2019). According to Carlos et al. (2016), L- tryptophan in the soil comes from root exudates or damaged cells.


#### Discussion

The diversity of the collected bacterial isolates indicated the variety of nitrogen-fixing bacteria. Table 1 showed that 13 isolates were collected from 148 fed rice fields, and 7 isolates were from irrigated lands. The results were consistent with previous studies by Hartono and Oslan Jumadi (2014); Herwati et al. (2020); Sukmawati et al. (2020); Tarigan et al. (2013).

Morphological characteristics of the 20 collected bacteria differ from one another. Further analysis was still needed to determine the genus and species of these bacterial isolates. Based on the Gram test, 6 isolates were Gram-positive (+), and 14 isolates were Gram-negative (-). Grouping of bacteria by Gram test is the basis for

identifying and classification of bacteria (Radji 2006). This difference was due to the cell wall structure. The cell wall structure of Gram-positive bacteria consists of peptidoglycan, while the cell wall structure of Gram-negative (-) consists of lipids (Sudewi et al. 2020). Rainfed rice fields were dominated by Gram-negative bacteria. The results of the catalase test showed that all isolates produce catalase. The catalase enzyme was essential for aerobic growth because  $H_2O_2$ , a respiratory byproduct, is toxic to microbial cells (Sahu et al. 2019). Non-pathogenic bacteria, both Gram-positive and Gram-negative, have a vital role as biocontrol agents in controlling plant diseases (Amaria et al. 2019). Gram-negative *Azotobacter* and *Azospirillum* are non-pathogenic. One species of *Azotobacter*, i.e., *Azotobacter vinelandii*, produces an anti-fungal that suppresses the development of *Fusarium oxysporum*, a causative agent of wilt disease (Hindersah et al. 2018).

Agro-ecosystems and technological inputs in rice fields affect bacterial diversity. The highest varieties were obtained in rainfed lands by 65% compared to irrigated fields with only 35%. Kurniawati et al. (2013) reported that the highest diversity of microbial populations was obtained in rainfed rice fields by 37.5%, in the swampy area by 31.75%, and then 30.75% irrigated lands.



**Figure 5.** Quantitative analysis of IAA production by bacterial isolates using a UV-VIS spectrophotometer 535 nm

All of the collected microorganisms were a non-symbiotic nitrogen-fixing potential that capable of growing on N-free media. The nitrogen-fixing ability was varied. NG4 isolate had the highest ability to fix nitrogen (0.157%), followed by NP11(0.153%) and NT2 (0.150%). Meanwhile, the NM5 isolate produced the lowest nitrogen (0.132%).

Nitrogen fixing bacteria perform this function because they have a specific enzyme in the cell known as nitrogenase composed of two mutually supporting components: Fe and Mo-Fe protein (Geddes et al. 2015). Sella and Suhaimi (2010) stated that non-symbiotic bacteria in the rhizosphere of Gramineae plants, such as *Azotobacter paspali* and *Beijerinckia* spp. are aerobic groups that colonized the root surface. The isolates from rainfed ricefield usually have better nitrogen-fixing ability than that of irrigated fields. The irrigated field has a high level of intensive chemical N fertilizers, which reduces the diversity and activity of these microorganisms since it makes their work unnecessary (Martínez-Romero 2009).

A qualitative test of bacterial isolates' ability to produce IAA was carried out by observing the change in color of the supernatant to pink. The higher the color intensity, the higher the hormone concentration. Hormone concentrations vary due to the differences in agro-ecosystem conditions and cultivation techniques at the sampling locations. Israwan et al. (2015) stated that the surrounding environment greatly influences bacterial ability to survive and produce IAA. Munif et al. (2012) indicated that soil properties, organic matter, cultivation techniques, fertilization, and pesticide application affect the presence of bacteria.

The results showed that isolate NB13 from Barru produced a more intense pink color and had the highest IAA concentration (1,835 mg L<sup>-1</sup>), followed by NG11 from Gowa (1,630 mg L<sup>-1</sup>) and NP1 from Pangkep (1,566 mg L<sup>-1</sup>). Meanwhile, the NT3 isolate had the lowest IAA concentration of 0.316 mg L<sup>-1</sup>. Isolates with high IAA production were isolated from rainfed rice fields (Barru, Pangkep, and Gowa). It might be concluded that bacteria

isolated from rainfed rice fields have a higher IAA production than those from irrigated rice fields. The IAA production was strongly influenced by biochemical characteristics and environmental factors (Numan et al. 2018). Rainfed fields are not flooded all year round so that allowing aerobic bacteria to grow. Moreover, agro-ecosystems and cultivation techniques also greatly influence an isolate's type and ability (Kurniawati et al. 2013). Intensive agricultural practices that utilize irrigation extensively and the increasing use of chemicals and heavy machinery causes degradation that affects biological life in the soil (Zalidis et al. 2002; Li J et al. 2015; Pereg et al. 2018).

This study concluded that microbes from rainfed rice fields in South Sulawesi were potential as biofertilizers and biostimulant agents than those from irrigated rice fields.

#### ACKNOWLEDGEMENTS

The authors are grateful to the LPDP/Indonesia Endowment Fund for Education for funding this research and all the parties involved in publishing this article.

#### REFERENCES

- Amaria W, Kasim NN, Munif A. 2019. Kelimpahan populasi bakteri filofser, rizosfer dan endofit tanaman kemiri sunan (*Reutealis trisperma* (Blanco) Airy Shaw), serta potensinya sebagai biokontrol. Tabaro 3 (1): 305-317. [Indonesian]
- Cappuccino JG, Sherman N. 2011. Microbiology: A Laboratory Manual, 9th ed. Pearson, London.
- Carlos MHJ, Stefani PVY, Janette AM, Melani, MSS, Gabriela PO. 2016. Assessing the effects of heavy metals in ACC deaminase and IAA production on plant growth-promoting bacteria. Microbiol Res 188 (2016): 53-61. DOI: 10.1016/j.micres.2016.05.001.
- Dong L, Tong X, Li X, Zhou J, Wang S, Liu B. 2019. Some developments and new insights of environmental problems and deep mining strategy for cleaner production in mines. J Cleaner Prod 210: 1562-1578. DOI: 10.1016/j.jclepro.2018.10.291.
- Geddes BA, Ryu MH, Mus F, Garcia Costas A, Peters JW, Voigt CA, Poole P. 2015. Use of plant colonizing bacteria as chassis for transfer

- of N<sub>2</sub>-fixation to cereals. *Curr Opin Biotechnol* 32: 216-222. DOI: 10.1016/j.copbio.2015.01.004.
- Glickman E, Dessaux Y. 1995. A critical examination of the specificity of the Salkowski reagent for indolic compounds produced by phytopathogenic bacteria. *Appl Environ Microbiol* 61: 793-796.
- Hartono, Jumadi O. 2014. Seleksi dan karakterisasi bakteri penambat nitrogen non simbiotik pengkalsium amonium pada tanah pertanian jagung (*Zea mays L.*) dan Padi (*Oryza sativa L.*) Asal Kabupaten Barru, Sulawesi Selatan, Indonesia. *Jurnal Sainsmat* 3 (2): 143-153. [Indonesian]
- Herwati A, Patandjengi B, Jayadi M, Masniawati. 2020. Isolation and characterization of nitrogen-fixing bacteria and producing IAA (Indole acetic acid) from rice rhizosphere from the Sopeng regency. *Int J Pharm Res* 13 (1): 808-815. DOI: 10.31838/ijpr.2021.13.01.138
- Hindersah R, Kalay M, Talahaturuson A, Lakburlawal Y. 2018. Bakteri pemfiksasi nitrogen *Azotobacter* pada tanaman kacang panjang. *Agric* 30 (1): 25-32. DOI: 10.24246/agric.2018.v30.i1.p25-32. [Indonesian]
- Irawan RF, Ardyati T. 2015. Eksplorasi bakteri pemfiksasi nitrogen non simbiotik penghasil IAA dan pelatun fosfat asal rhizosfer tanaman apel Kota Batu, Jawa Timur. *Biotropika* 3 (2): 55-59. [Indonesian]
- Kesaulya H, Baharuddin, Zakaria B, Syaiful SA. 2015. Isolation and physiological characterization of PGPR from potato plant rhizosphere in medium land of Bumi Island. *Procedia Food Sci* 3: 190-199. DOI: 10.1016/j.profoo.2015.01.021.
- Khatoon Z, Huang S, Rafique M, Fakhar A, Kamran MA, Santoyo G. 2020. Unlocking the potential of plant growth-promoting rhizobacteria on soil health and the sustainability of agricultural systems. *J Environ Manag* 273 (August): 111118. DOI: 10.1016/j.jenvman.2020.111118.
- Kumiawati S, Muttaqin HK, Giyanto. 2013. Keragaman bakteri pada pertanian padi di lahan sawah irigasi, tadah hujan dan rawa. *Ketahanan Pangan* 259-266. [Indonesian]
- Lestari P, Suryadi Y, Susilowati DN, Priyatno TP, Samudra IM. 2008. Karakterisasi bakteri penghasil asam indol asetat dan pengaruhnya terhadap vigor benih padi. *Berita Biologi* 14 (1): 19-28. [Indonesian]
- Li J, Deng J, Gu Q, Wang K, Ye F, Xu Z, Jin S. 2015. The accelerated urbanization process: A threat to soil resources in eastern China. *Sustainability (Switzerland)* 7 (6): 7137-7155. DOI: 10.3390/su7067137.
- Li Y, Yuan L, Xue S, Liu B, Jin G. 2021. Artificial root exudates excite bacterial nitrogen fixation in the subsurface of mine soils. *Appl Soil Ecol* 157 (September 2020): 103774. DOI: 10.1016/j.apsoil.2020.103774.
- Ma J, Bei Q, Wang X, Lan P, Liu G, Lin X, Liu Q, Lin Z, Liu B, Zhang Y, Jin H, Hu T, Zhu J, Xie Z. 2019. Impacts of Mo application on biological nitrogen fixation and diazotrophic communities in a flooded rice-soil system. *Sci Total Environ* 649: 686-694. DOI: 10.1016/j.scitotenv.2018.08.318.
- Martínez-Romero E. 2009. Coevolution in rhizobium-legume symbiosis. *DNA Cell Biol* 28 (8): 361-370. DOI: 10.1089/dna.2009.0863.
- Munif A, Wiyono S, Suwarno S. 2012. Isolasi bakteri endofit asal padi gogo dan potensinya sebagai agens biokontrol dan pemacu pertumbuhan. *Jurnal Fitopatologi Indonesia* 8 (3): 57-64. DOI: 10.14692/jfi.8.3.57. [Indonesian]
- Numan M, Bashir S, Khan Y, Mumtaz R, Shinwari ZK, Khan AL, Khan A, AL-Harrasi A. (2018). Plant growth promoting bacteria as an alternative strategy for salt tolerance in plants: A review. *Microbiol Res* 209 (Jan): 21-32. DOI:10.1016/j.micres.2018.02.003
- Pambudi A, Susanti S, Priambodo, TW. 2017. Isolasi dan karakterisasi bakteri tanah sawah di Desa Sukawali dan Desa Belimbing, Kabupaten Tangerang. *Al-Kaunyah: Jurnal Biologi* 10 (2): 105-113. DOI: 10.15408/kaunyah.v10i2.4907. [Indonesian]
- Panigrahi S, Mohanty S, Rath CC. 2019. Characterization of endophytic bacteria *Enterobacter cloacae* MG00145 isolated from *Ocimum sanctum* with Indole Acetic Acid (IAA) production and plant growth promoting capabilities against selected crops. *South Afr J Bot* 000: 1-10. DOI: 10.1016/j.sajb.2019.09.017
- Park M, Kim C, Yang J, Lee H, Shin W, Kim S, Sa T. 2005. Isolation and characterization of diazotrophic growth promoting bacteria from rhizosphere of agricultural crops of Korea. *Microbiol Res* 160 (2): 127-133. DOI: 10.1016/j.micres.2004.10.003
- Pas Aksarah A, Sopandje D, Trikoesoemaningtyas, Santosa DA. 2015. Aplikasi konsorsium mikroba filofosfer dan rizosfer untuk meningkatkan pertumbuhan dan hasil tanaman padi. *Institut Pertanian Bogor (IPB) Kampus*, 15-24. [Indonesian]
- Pereg L, Morugán-Coronado A, McMillan M, García-Oreñe F. 2018. Restoration of nitrogen cycling community in grapevine soil by a decade of organic fertilization. *Soil Tillage Res* 179 (July 2017): 11-19. DOI: 10.1016/j.still.2018.01.007
- Prathama Y, Nelvia N, Amri AI. 2018. Pemberian amelioran dan isolat bakteri fiksasi nitrogen non simbiotik untuk meningkatkan pertumbuhan dan serapan N tanaman padi gogo (*Oryza sativa L.*) pada medium ultisol. *Jurnal Solum* 15 (2): 40. DOI: lum.15.2.40-49.2018. [Indonesian]
- Pulungan AS, Tumangger, DE. 2018. Isolasi dan Karakterisasi bakteri endofit penghasil enzim katalase dari daun Buasbuas (*Premna pubescens* Blume) Isolation and characterization of endophytic bacteria producing catalase enzyme from Buasbuas (*Premna pubescens* Blume) leaves. *Jurnal Biologi Lingkungan, Industri, Kesehatan* 5 (1): 72-80. DOI: 10.31289/biolink.v5i1.1665 [Indonesian]
- Radji M. 2006. Penuntun Praktikum Mikrobiologi Farmasi, Edisi 2, Departemen Farmasi, FMIPAUI, Depok. [Indonesian]
- Ramadhan S dan Tiwow VMA 2016. Analisis kadar unsur nitrogen (N) dan fosfor (P) dalam lamun (*Enhalus acoroides*) di wilayah perairan Kabupaten Donggala. *J Akad Kimia* 5 (1): 37-43. DOI: 10.22487/j24775185.2016.v5.i1.7998
- Rao S. 2010. Mikroorganisme tanah dan pertumbuhan tanaman. *terjemahan Herawati Susilo. Universitas Indonesia (UI-Press), Jakarta.* [Indonesian]
- Rascio N, La Rocca N. 2018. Biological nitrogen fixation. *Encyclopedia Ecol* 2 (Jul): 264-279. DOI: 10.1016/B978-0-444-63768-0.00685-5
- Sahu AK, Kar BRK, Sahu SS, Achary, Pragnya P, Harichandan PS. 2019. Isolation, characterization of endophytic bacteria from *Psidium guajava* plant and screening of their antibiotic activities. *Int J Res Appl Sci Eng Technol* 7 (1): 345-355. DOI: 10.22214/ijraset.2019.1061.
- Shridhar BS, Shrimant Shridhar B. 2012. Review: Nitrogen fixing microorganisms. *Int J Microbiol Res* 3 (1): 46-52. DOI: 10.5829/idosi.ijmr.2012.3.1.61103.
- Stella M, Suhaimi M. 2010. Selection of suitable growth medium for free-living diazotrophs isolated from compost. *J Trop Agric Fd Sc* 38 (2): 211-219.
- Sudewi S, Ala A, Baharuddin, Farid M. 2020. The isolation, characterization endophytic bacteria from roots of local rice plant Kamba in, Central Sulawesi, Indonesia. *Biodiversitas* 21 (4): 1614-1624. DOI: 10.13057/biodiv/d210442.
- Suhartanti M, Sarjono PR, Aminin ALN. 2010. Studi filogeni dan uji potensi enzim ekstraseluler (amilase, β-galaktosidase, protease, katalase) Isolat *Allicyclobacillus* sp. Gedong Songo. *Jurnal Kimia Sains dan Aplikasi* 13 (3) 80-87. DOI: 10.14710/jksa.13.3.80-87. [Indonesian]
- Sukmawati, Ala A, Patandjengi B, Gusli S. 2020. Exploring of promising bacteria from the rhizosphere of maize, cocoa and lamtoro. *Biodiversitas* 21 (12): 5665-5673. DOI: 10.13057/biodiv/d211224.
- Sun X, Zhong T, Zhang L, Zhang K, WuW. 2019. Reducing ammonia volatilization from paddy field with rice straw derived biochar. *Sci Total Environ* 660: 512-518. DOI: 10.1016/j.scitotenv.2018.12.450.
- Tahovská K, Kaňa J, Bárta J, Oulehle F, Richter A, Šantrůčková H. 2013. Microbial N immobilization is of great importance in acidified mountain spruce forest soils. *Soil Biol Biochem* 59: 58-71. DOI: 10.1016/j.soilbio.2012.12.015.
- Tarigan RS, Jamilah IDE. 2013. Seleksi bakteri penambat nitrogen dan penghasil hormon IAA (Indole Acetic Acid) dari rizosfer tanah perkebunan kedelai (*Glycine max L.*). *Saintia Biologi* 1 (2): 42-48. [Indonesian]
- Vinten AJA, Ball BC, O'Sullivan MF, Henshall JK. 2002. The effects of cultivation method, fertilizer input and previous sward type on organic C and N storage and gaseous losses under spring and winter barley following long-term leys. *J Agric Sci* 139 (3): 231-243. DOI: 10.1017/S0021859602002496.
- Widiastuti F. 2017. Lahan sawah sebagai pendukung ketahanan pangan serta strategi pencapaian kemandirian pangan. *Lahan Sawah Sebagai Pendukung Ketahanan Pangan Serta Strategi Pencapaian Kemandirian Pangan* 8 (3): 17-30. DOI: 10.2018/jSDL.v8i3.6479. [Indonesian]
- Witte CP. 2011. Urea metabolism in plants. *Plant Sci* 180 (3): 431-438. DOI: 10.1016/j.plantsci.2010.11.010.
- Zalidis G, Stamatiadis S, Takavakoglou V, Eskridge K, Misopolinos N. 2002. Impacts of agricultural practices on soil and water quality in the Mediterranean region and proposed assessment methodology. *Agric Ecosyst Environ* 88 (2): 137-146. DOI: 10.1016/S0167-8809(01)00249-3.

# Isolation and characterization of N-fixing and IAA producing rhizobacteria from two rice field agro-ecosystems in South Sulawesi, Indonesia

## ORIGINALITY REPORT

**20%**  
SIMILARITY INDEX

**13%**  
INTERNET SOURCES

**15%**  
PUBLICATIONS

**4%**  
STUDENT PAPERS

## PRIMARY SOURCES

- 1** [www.hindawi.com](http://www.hindawi.com) Internet Source **2%**
- 2** Rajidah Arsita, Hilda Karim, Yusminah Hala, Neni Iriany, Oslan Jumadi. " Isolation and identification of nitrogen-fixing bacteria in the corn rhizosphere ( I.) originating from Jeneponto Regency, South Sulawesi ", IOP Conference Series: Earth and Environmental Science, 2020 Publication **1%**
- 3** N. Tejera. "Isolation and characterization of Azotobacter and Azospirillum strains from the sugarcane rhizosphere", Plant and Soil, 03/2005 Publication **1%**
- 4** [journal.uin-alauddin.ac.id](http://journal.uin-alauddin.ac.id) Internet Source **1%**
- 5** [link.springer.com](http://link.springer.com) Internet Source **1%**

6

[www.ikprress.org](http://www.ikprress.org)

Internet Source

1 %

7

Aizuddin, MBAB, NT Athena, and W Nyet Kui. "Antioxidant and antimicrobial properties of marine actinomycetes from sipadan Island, Sabah", *International Journal of Biological and Chemical Sciences*, 2013.

Publication

1 %

8

David A. Mormak, L. E. Casida. " Study of Endospores in Soil by Use of a Modified Endospore Stain ", *Applied and Environmental Microbiology*, 1985

Publication

1 %

9

Gusmiaty, M Restu, B Bachtiar, SH Larekeng. "Gibberellin And IAA Production by Rhizobacteria From Various Private Forest", *IOP Conference Series: Earth and Environmental Science*, 2019

Publication

1 %

10

Justyna Nykiel-Szymańska, Przemysław Bernat, Mirosława Słaba. "Biotransformation and detoxification of chloroacetanilide herbicides by *Trichoderma* spp. with plant growth-promoting activities", *Pesticide Biochemistry and Physiology*, 2020

Publication

1 %

11

[ppsub.ub.ac.id](http://ppsub.ub.ac.id)

Internet Source

1 %

---

12	<a href="http://ojs.uma.ac.id">ojs.uma.ac.id</a> Internet Source	1 %
13	Submitted to Centre for Nutrition Education & Lifestyle Management (CNELM) Student Paper	<1 %
14	<a href="http://e-jmsb.id">e-jmsb.id</a> Internet Source	<1 %
15	Héctor Antonio Cristóbal. "Diversity of protease-producing marine bacteria from sub-antarctic environments", <i>Journal of Basic Microbiology</i> , 12/2011 Publication	<1 %
16	<a href="http://www.news-medical.net">www.news-medical.net</a> Internet Source	<1 %
17	Submitted to Higher Education Commission Pakistan Student Paper	<1 %
18	Microbial Diversity and Biotechnology in Food Security, 2014. Publication	<1 %
19	Submitted to Universiti Teknologi MARA Student Paper	<1 %
20	<a href="http://academicjournals.org">academicjournals.org</a> Internet Source	<1 %

---

21

H Kesaulya, A Talahaturuson, A M Kalay, E Matatula, I J Lawalatta, M L Hehanussa, S J Nendissa. "Characterization of plant growth promoting rhizobacteria of maize", IOP Conference Series: Earth and Environmental Science, 2021

Publication

---

<1 %

22

Gaurav Sood, Rajesh Kaushal, Girish Panwar, Meenakshi Dhiman. " Effect of Indigenous Plant Growth-Promoting Rhizobacteria on Wheat ( L.) Productivity and Soil Nutrients ", Communications in Soil Science and Plant Analysis, 2018

Publication

---

<1 %

23

Hoang Minh Tam. "Isolation and Identification of Rhizospheric Bacteria in Sugarcane (Saccharum spp. L.) Cultivated on Acrisols and Ferrasols of Dong Nai Province, the Southeast of Vietnam", American Journal of Life Sciences, 2015

Publication

---

<1 %

24

Karina Elwanda Saputri, Nora Sari Idiawati, Mega Sari Juane Sofiana. "Isolasi dan Karakterisasi Bakteri Penambat Nitrogen dari Rizosfer Mangrove di Kuala Singkawang", Jurnal Laut Khatulistiwa, 2021

Publication

---

<1 %

25 Saleema S Saleh. "Involvement of *gacS* and *rpoS* in enhancement of the plant growth-promoting capabilities of *Enterobacter cloacae* CAL2 and UW4", Canadian Journal of Microbiology, 08/2001  
Publication

---

26 [brage.bibsys.no](http://brage.bibsys.no)  
Internet Source

---

27 [idosi.org](http://idosi.org)  
Internet Source

---

28 "Carbon and Nitrogen Cycling in Soil", Springer Science and Business Media LLC, 2020  
Publication

---

29 Hadija, T. Kuswinanti, M. Jayadi, S.H Larekeng. "Isolation, characterization and identification of nitrogen fixing bacteria with organic fertilizer applications in paddy soil", IOP Conference Series: Earth and Environmental Science, 2021  
Publication

---

30 Seo-Jung Jang, Yu-Jin Kim, Sul-Hee Lee, Young-Seo Park, Jung-Min Park, Dong-Hoon Bai. "Oceanobacillus gochujangensis sp. nov., isolated from gochujang a traditional Korean fermented food", Journal of Microbiology, 2014  
Publication

---

31 [core.ac.uk](http://core.ac.uk) <1 %  
Internet Source

---

32 B. Cibichakravarthy, R. Preetha, S. P. Sundaram, K. Kumar, D. Balachandar. <1 %  
"Diazotrophic diversity in the rhizosphere of two exotic weed plants, Prosopis juliflora and Parthenium hysterophorus", World Journal of Microbiology and Biotechnology, 2011  
Publication

---

33 D Lestari, L Asrul, T Kuswinanti, Y Musa, Sulfahri. "Selection of fungi that potentially produces IAA (Indole Acetic Acid) hormone origin of Takalar sugar factory waste", IOP Conference Series: Earth and Environmental Science, 2021 <1 %  
Publication

---

34 Submitted to Portage Northern High School <1 %  
Student Paper

---

35 [dokumen.pub](http://dokumen.pub) <1 %  
Internet Source

---

36 [iosrjournals.org](http://iosrjournals.org) <1 %  
Internet Source

---

37 [s3.amazonaws.com](http://s3.amazonaws.com) <1 %  
Internet Source

---

38 [www.ijcmas.com](http://www.ijcmas.com) <1 %  
Internet Source

---

39	<a href="http://www.neliti.com">www.neliti.com</a> Internet Source	<1 %
40	<a href="http://www.tandfonline.com">www.tandfonline.com</a> Internet Source	<1 %
41	"Biocommunication in Soil Microorganisms", Springer Science and Business Media LLC, 2011 Publication	<1 %
42	<a href="http://journal.uinjkt.ac.id">journal.uinjkt.ac.id</a> Internet Source	<1 %
43	<a href="http://jurnal.permi.or.id">jurnal.permi.or.id</a> Internet Source	<1 %
44	<a href="http://sfamjournals.onlinelibrary.wiley.com">sfamjournals.onlinelibrary.wiley.com</a> Internet Source	<1 %
45	<a href="http://www.bluepenjournals.org">www.bluepenjournals.org</a> Internet Source	<1 %
46	<a href="http://www.scribd.com">www.scribd.com</a> Internet Source	<1 %
47	Park, M.. "Isolation and characterization of diazotrophic growth promoting bacteria from rhizosphere of agricultural crops of Korea", Microbiological Research, 20050425 Publication	<1 %
48	Wei Shi, Hui-Yu Zhao, Yin Chen, Jin-Song Wang, Bing Han, Cong-Ping Li, Jun-Yuan Lu, Li-	<1 %

Mei Zhang. "Organic manure rather than phosphorus fertilization primarily determined asymbiotic nitrogen fixation rate and the stability of diazotrophic community in an upland red soil", Agriculture, Ecosystems & Environment, 2021

Publication

---

49

Juan Fu, Xudong Xu. "The functional divergence of two glgP homologues in *Synechocystis* sp. PCC 6803", FEMS Microbiology Letters, 2006

Publication

---

<1 %

50

Sunarti Sunarti, Romelos Untailawan. "UTILIZATION OF TOFU AND SAGO PULP AS ORGANIC LIQUID FERTILIZER AND ITS APPLICATION TO MUSTARD PLANT", EDU SCIENCES JOURNAL, 2020

Publication

---

<1 %

51

Thanakorn Saengsanga. "Isolation and Characterization of Indigenous Plant Growth-Promoting Rhizobacteria and Their Effects on Growth at the Early Stage of Thai Jasmine Rice (*Oryza sativa* L. KDML105)", Arabian Journal for Science and Engineering, 2017

Publication

---

<1 %

Exclude bibliography  On