

The_Effect_of_Giving_Rice_Bran _Milk_on_Blood_Glucose.pdf

by

Submission date: 04-Mar-2021 02:08PM (UTC+0700)

Submission ID: 1523911074

File name: The_Effect_of_Giving_Rice_Bran_Milk_on_Blood_Glucose.pdf (516.92K)

Word count: 4622

Character count: 23738

Research Article

The Effect of Giving Rice Bran Milk on Blood Glucose Levels and Body Weight in Hyperglycemic Primary School Teachers in Makassar City

SAIFUDDIN SIRAJUDDIN^{*1}, MASNI², ABDUL SALAM¹¹ Nutrition Department, Faculty of Public Health, Hasanuddin University, City of Makassar, Indonesia² Faculty of Public Health, Hasanuddin University, City of Makassar, Indonesia

*Corresponding author:

Email:saifuddinsirajuddin.unhas@yahoo.com

Received: 14.10.20, Revised: 26.11.20, Accepted: 17.12.20

ABSTRACT

This study aims to determine the effect of rice bran milk on fasting blood glucose levels and body weight of primary school teachers in Makassar City. This study was quasi-experimental with pretest-posttest with control group design. The population is all primary school teachers in Makassar City who have abdominal circumference > 80 cm for woman and >90 for men. The samples are those who meet the inclusion criteria who had fasting blood glucose levels >100 mg/dL, do not consume drugs. Sample selected then randomly divided into experimental and control group, each group consist of 12 participants. Provision of rice bran milk was given once a day (30 gram) and nutritional education for one month for experimental group and only nutritional education once a day through WhatsApp media for the control group. Blood Glucose level, body weight and 24H dietary recall was reported before and after study both for experiment and control groups. The data were analyzed by paired t-test and experimental independent samples t-test, using SPSS. The results of this study showed that, in experimental group, the fasting blood glucose level reduced significantly ($p < 0.05$) but increased in the control group after intervention there were also a significance differences between fasting blood glucose level in experimental and control group ($p < 0.05$). There were a significance differences between body weight before and after intervention in intervention group ($p < 0.05$) but not in control group and there were also significance differences of body weight between experimental and control groups at posttest. Conclusion, rice bran milk can be used as nutritional supplements for the control of hyperglycemia and body weight of overweight and diabetic individuals.

Keywords: rice bran milk, hyperglycemia, diabetes, blood glucose, body weight**INTRODUCTION**

Non-Communicable Diseases (NCD) has become a major concern in public health issues, this is marked by the shift in epidemiological patterns of disease from infectious diseases to non-communicable diseases that continue to increase both nationally and globally, one of which is Diabetes Mellitus (DM). DM is a degenerative disease whose prevalence expected to continue to increase from year to year. According to the International of Diabetic Federation (IDF), in 2019, approximately 463 million adults (20-79 years) worldwide were living with diabetes this is predicted to rise to 578 million by 2030 and to 700 million by 2045 (International Diabetes Federation, 2019).

NCD, including Diabetes, have now become a serious threat to global health. 90-95% of cases of Diabetes are Type 2 Diabetes which is largely preventable because it is caused by an unhealthy lifestyle. Indonesia also faces a situation with the threat of diabetes similar to the world. The

International Diabetes Federation (IDF) Atlas 2017 reports that, the Diabetes epidemic in Indonesia is still showing an increasing trend. Indonesia is the sixth ranked country in the world after China, India, the United States, Brazil and Mexico with the number of people with diabetes aged 20-79 years, around 10.3 million people (International Diabetes Federation, 2017).

Based on Basic Health Research shows a significant increase in the prevalence of Diabetes, from 6.9% in 2013 to 8.5% in 2018; so, the estimated number of sufferers in Indonesia reached more than 16 million people who were then at risk of diabetic complications, such as: heart attack, stroke, blindness and kidney failure can even death. Diabetes is a global epidemic problem which if not treated immediately seriously will result in an increase in the impact of significant economic losses, especially for developing countries in Asia and Africa. IDF data also shows that the direct costs of managing diabetes reach more than 727 billion USD per

year or around 12% of global health financing. The National Health Insurance Data (JKN) also shows an increase in the number of cases and financing of Diabetes services in Indonesia from 135,322 cases with funding of Rp 700.29 billion in 2014 to 322,820 cases with funding of Rp 1,877 trillion in 2017 (Kementerian Kesehatan Republik Indonesia, 2018).

The increasing prevalence of NCD, has led to changes in people's attitudes, which tend to be more trying to prevent disease and implement healthy living. The emergence of various functional food products can be an alternative in preventing several types of diseases that arise both due to genetic factors, lifestyle and diet. One ingredient that has the potential to be used as functional food is rice bran (Tuarita et al., 2017).

Rice bran is a by-product of rice milling process is abundant in Indonesia. Rice production in 2016 reached 79.141 million tons of Dry Grain Paddy and in 2017 rice production reached 81.38 million tons (Indonesian Agricultural Departemen, 2017). The percentage of bran from milled dry grain is around 10%. So far, most of it is only used as animal feed (Indonesian Agricultural Departemen, 2018). The results research showed that, rice bran is a good source of bioactive phytochemicals, namely oryzaol, vitamin E group (tocopherols and tocotrienols) and flavonoids as well as macro nutrient content such as protein, fiber, fat and carbohydrates. The phytochemical compound have health beneficial properties and antioxidant activities^{4,5,6}. (Moongnarm et al., 2012; Rohman et al., 2014; Watchararuij et al., 2008). With this nutritional content, efforts can be made to diversify food that can provide added value to rice bran as an animal feed. Some research found that the content of gamma oryzaol and fiber in rice bran can reduce blood glucose levels (Devarajan et al., 2016; Qureshi et al., 2002; Sivamaruthi et al., 2018; Tazakori et al., 2007) and other research found that rice bran can increase blood glucose levels (Lai et al., 2011).

Besides can decrease blood glucose level, rice bran also found as a source of fiber which have a functions to decrease blood pressure, control blood glucose level, decrease body weight and increase the immunity (Anderson et al., 2009). Many studies support the inverse relationship of dietary fiber and the risk for CHD. However, more recent studies found interesting data illustrating that for every 10 g of additional fiber added to a diet, the mortality risk of CHD decreased by 17-35% (Pereira et al., 2004; Streppel et al., 2008). Risk factors for CHD include hypercholesterolemia, hypertension,

obesity and type two diabetes. The results of one larger clinical study, supports this theory in their finding that for every 20 g/d increase in consumption of bran, body weight decreased by 0.80 lbs (Khan-Banerjee et al., 2004).

The main aim of this study was to investigate the effect of rice bran milk on glucose levels and body weight in hyperglycemic.

MATERIALS AND METHODS

Materials

Rice bran used in this research was obtained from rice mill PB. Rice bran taken directly from the rice mill was directly roasted using low heat for approximately 10 minutes, then sieved using a 60-mesh sieve. The rice bran was then made a rice bran milk by adding Arabic gum as an emulsifier, corn sugar as a sweetener, sodium bisulfite as a preservative and melon flavoring as flavor and odor.

Methods

This study was Quasi experimental design with pretest-posttest with control group design. A total of 24 primary school teachers with hyperglycemic were screen from all primary school teachers in Makassar City who meet the criteria: have abdominal circumference >80 cm for woman and >90 for men. For those who had fasting blood sugar levels >100 mg/dl. Fasting blood glucose levels was determined by Prodia Laboratory after fasting for 8 hours. Exclusion criteria: liver disease, using anti diabetic medicines, and pregnancy or lactation for women participants. Those subjects who have blood glucose levels >100 mg/dl randomly divided to experimental and control group; each group consist of 12 participants. Provision of rice bran milk was given once a day (30 gram) for one month for experimental group and nutrition education for the control group also once a day through WhatsApp media. Fasting blood glucose levels were measured twice, before and after the study and 24-hour recall of food were checked before and after study. The data were analyzed by paired t-test and independent sample t-test using SPSS as Food processor.

The study was approved by the Institutional Review Board of Faculty of Public Health Hasanuddin University, (Approval number: 6156/UN4.14.8/TP.02.02/2019 - July 25, 2019). Each participant provided written consent before participating in the study.

RESULTS

Characteristics of Subjects

Out of 24 subjects entered the study, both group experiment and control, dominated by women, average age for group were lower (about 48

years old) while control group (about 54 years old) (Table 1).

Table 1. Characteristics of respondent based on sex and age

Variable	Experimental Group (n=12)	Control Group (n=12)
Sex		
- Men	5	1
- Women	7	11
Age (Years)		
- Minimum	29	49
- Maximum	60	57
- Average	48.75	54.58
- Std	8.29	3.31

The effect of rice bran milk on fasting of Blood Glucose

Fasting blood glucose level examination results are shown in Table 2. The average fasting blood glucose levels in experimental group, before intervention were 217,00 ± 88,24 mg/dl and 182,75 ± 61,46 mg/dl after the intervention, it means that there was a decrease value equal to 34.25 ± 26,78 mg/dl (15,78%); while in the control group were 141,17 ± 48,46 before intervention and 143,58 ± 54,59 after the intervention period, it means that there was an increase value of fasting blood glucose levels

equal to 2.41 ± 6.13 mg/dL. The results of statistical test (paired t-test) showed that there was a significant decrease of fasting blood glucose before and after intervention in the experimental Group (p<0.05), but in control groups there was no significance differences. The results also showed that, there was a significance difference of fasting blood glucose levels on pre-test between experiment and control group (p=0.018) but there were no significance differences of fasting blood glucose level on post-test between experimental group and control groups (p>0.05).

Table 2. The Effect of Rice Bran Milk on Fasting Blood Glucose Levels of Hyperglycemic and controls Elementary School Teachers in Makassar City

	Experimental Group		Control Group		p
	Pre	Post	Pre	Post	
Fasting blood glucose (mg/dL)	217,00 ± 88,24	182,75 ± 61,46	141,17 ± 48,46	143,58 ± 54,59	0.018 ^a 0.113 ^b
P	0.020 ^c		0,569 ^d		

^a Two independent sample t-test – the differences between fasting blood glucose levels in pre-test

^b Two independent sample t-test – the differences between fasting blood glucose levels in post-test

^c Paired t-test – differences between fasting blood glucose levels before and after test for experimental group

^d Paired t-test – differences between fasting blood glucose levels before and after test for control group

The effect of rice bran milk on Body Weight

Table 3. The Effect of Rice Bran Milk on Body Weight of Hyperglycemic and controls Elementary School Teachers In Makassar City

Weight (Kg)	Experimental Group		Control Group		P
	Pre	Post	Pre	Post	
Average ± std	72,79 ± 12,41	70,73 ± 11,82	58,12 ± 9,81	57,55 ± 9,80	0.004 ^a 0.007 ^b
P	0.042 ^c		0,130 ^d		

^a Two independent sample t-test – the differences of body weight between intervention and control groups at pre-test

^b Two independent sample t-test – the differences of body weight between experimental and control groups at post-test

^c Paired t-test – differences between body weight before and after test for intervention group

⁵ Paired t-test – differences between body weight before and after test for control group Table 3 showed that, the experimental group had a significance loss in weight ($p < 0,05$). The average reduction in weight was about 2,06 kg after administration rice bran milk for one month, namely $72,79 \pm 12,41$ kg in pretest to $70,73 \pm 11,82$ kg in post-test, it means that, the reduction about 2,83%, while in control group there was also reduction but not significant between pre and post-test ($p = 0,130$). If we compare body weight between groups at posttest, there was also a significant difference ($p < 0,05$).

The results of 24H Recall Diet

The reported data of the respondent's food intake through a 1 x 24-hour food recall were analyzed with a nutrisurvey. Recall was done twice, before the intervention and the day after the intervention ends. The results of the analysis including energy (kcal), carbohydrate (g), protein (g), fat (g), fiber (g) and cholesterol (mg) which are shown in Table 4. The results showed that, dietary intake did not change over the study interval. The mean energy, protein, fat, carbohydrate, cholesterol and fiber

intakes were not significantly different between pre and post-test, and also between groups ($p > 0.05$). The measurement results show that, in experimental group, there were a decrease in the average intake of energy, protein, fat, carbohydrate and cholesterol before and after intervention, while for the fiber where was increased. In control group (nutritional education group), all item was decrease but also not significant.

Table 4. The results of 24-hour Food Recall before and after intervention in Experimental and Control groups

Daily Intake	Experimental Group n=12			Control Group n=12			p-value Experimental Vs Control
	Pre	Post	p-	Pre	Post	p	
Energy (kcal)	1127.52±426	1151.67±531.89	0.838	1184.73±434.14	1047.09±533.33	0.332	0.529
Protein (g)	45.52±21.77	52.78±29.64	0.416	46.94±22.29	41.23±22.23	0.454	0.894
Fat (g)	39.92±29.27	42.73±28.75	0.690	42.14±30.96	35.73±33.85	0.483	0.748
Carbohydrate (g)	147.86±67.27	141.53±63.31	0.624	156.18±65.98	140.41±69.69	0.256	0.229
Fiber (g)	7.03±4.21	7.30±5.98	0.878	8.30±5.28	5.70±3.19	0.135	0.352
Cholesterol	186.96±186.91	230.28±188.88	0.450	211.94±190.29	222.342±234.34	0.884	0.544

DISCUSSION

The aims of this study was to determine the effect of rice bran milk on fasting blood glucose levels and body weight of hyperglycemic primary school teachers. Antioxidant compounds contained in rice bran can be grouped into 6 groups, including phenolic acids, flavonoids, anthocyanins and proanthocyanins, tocopherols and tocotrienols, γ -oryzanol and phytic acids (Goufo & Trindade, 2014). The main antioxidant in rice bran is γ -oryzanol² (Tuarita et al., 2017). Antioxidants are able to neutralize excess free radical compounds in pancreatic β cells by donating electrons or cutting off free radical chain reactions and causing free radicals to become stable. So that this can stop or inhibit oxidative damage to

pancreatic β cells (Dewi et al., 2014). γ -oryzanol can also increase glucokinase activity and inhibit glucose 6-phosphatase and phosphoenolpyruvate carboxinase in the liver. This increased activity can increase the utilization of blood glucose into energy or glycogen storage in the liver, thereby causing decrease blood glucose concentration (Son et al., 2015). Other antioxidants, namely flavonoids contained in rice bran, can prevent complications or progression of diabetes mellitus by clearing excessive free radicals, breaking the chain of free radical reactions, binding of metal ions (chelating), and blocking the polyol pathway by inhibiting the enzyme aldose reductase. Flavonoids also have an inhibitory effect on the alpha glucosidase enzyme through hydroxylation

bonds and substitution in the β ring. The principle of inhibition is similar to acarbose which has been used as a drug for the treatment of diabetes mellitus, namely by producing carbohydrate and disaccharide delays and glucose absorption and inhibiting the metabolism of sucrose into glucose and fructose (Prameswari & Widjanarko, 2014; Salehi & Sardarodiyani, 2016).

As we found in this study, there were a significant decrease of fasting blood glucose levels on experimental group ($p=0.02$) and in control group the blood glucose levels tend to increase but not significant. There were also found that in pre-test, there were a significance differences of fasting blood glucose between experiment and control group ($p=0.016$) and in post-test there were also differences but not significant. Base on this results we can say that, provision of rice bran milk were good because can decrease the blood glucose levels. This is in line with the research results of Cheng et al. (2010) that provision of rice bran 20 g/day for 12 weeks can reduce glucose levels (Cheng et al., 2010). This also in line with Devarajan et al. (2016) that provision of rice bran oil blend with sesame oil can reduce fasting blood glucose levels (Devarajan et al., 2016). In control group, there was an increase in fasting blood glucose level after the intervention period. Increasing fasting blood glucose levels in control group was in line with the research results of Lai, M.H. et al. (2012) that consumption of 18 g rice bran oil modified milk daily for five weeks could increase 11 g/dL fasting blood glucose levels (Lai et al., 2011). The results of this study showed that, there were just a little decrease in blood glucose levels in the experimental groups, these maybe caused by an increase fat consumption of the experimental group. This results in line with the research results of Kim et al. (2010) that, high fat-feed resulted in increase in the blood glucose levels in mice (Kim et al., 2010).

The effect of rice bran administration on weight loss in this research not provide adequate results, because the weight loss was too small, although the results of the reduction were significant ($p<0,05$), this may be due to the short time of administration, which only 30 days. The results of this study are in line with the results of the study of Edrisi et al. (2017) which shows that, by administered rice bran for 12 weeks can significantly reduce weight ($p < 0.01$) (Edrisi et al., 2018). The same results also found by Zavoshy et al. (2012) that body weight were significantly reduced after four weeks consumption of rice bran oil (30g/day) (Zavoshy et al., 2012). The research results by Hongu et al. (2014) showed a decrease body weight for about 4.7 kg after the

administration of a snack bar which was a mixture of rice bran and plant sterol for eight weeks (Hongu et al., 2014).

The strengths of this study were used quasi experimental design with pre test-post test with control group design, for the exclusion of individuals who were consuming drugs or conditions that may have altered metabolism. Our study population was hyperglycemic and obese adults. A limitation of this study is the relatively low number of participants through the study was powered based on expected reduction of blood glucose and weight loss. Another potential limitation of this study is the relatively short dietary intervention period (4 weeks). Though this interval did allow for an average weight loss of 2,06 kg, it did not allow for the assessment of long-term effects, which should be considered in future studies with rice bran.

CONCLUSION

Results of this study showed that, giving rice bran milk 30 gram for one month, once a day can decrease significantly blood glucose levels of hyperglycemic people and also can reduce body weight.

CONFLICT OF INTEREST STATEMENT

The authors declare that there is no conflict of interest

ACKNOWLEDGMENTS

The authors gratefully acknowledge the Hasanuddin University for providing funds for conducting this research

REFERENCES

1. Anderson, J. W., Baird, P., Davis, R. H., Ferreri, S., Knudtson, M., Koraym, A., Waters, V., & Williams, C. L. (2009). Health benefits of dietary fiber. *Nutrition Reviews*, 67(4), 188–205. <https://doi.org/10.1111/j.1753-4887.2009.00189.x>
2. Cheng, H.-H., Huang, H.-Y., Chen, Y.-Y., Huang, C.-L., Chang, C.-J., Chen, H.-L., & Lai, M.-H. (2010). Ameliorative Effects of Stabilized Rice Bran on Type 2 Diabetes Patients. *Annals of Nutrition and Metabolism*, 56(1), 45–51. <https://doi.org/10.1159/000265850>
3. Devarajan, S., Chatterjee, B., Urata, H., Zhang, B., Ali, A., Singh, R., & Ganapathy, S. (2016). A blend of sesame and rice bran oils lowers hyperglycemia and improves the lipids. *The American Journal of Medicine*, 129(7), 731–739.
4. Dewi, Y. F., Anthara, M. S., & Dharmayudha, A. (2014). Efektifitas ekstrak daun sirih merah (*Piper crocatum*) terhadap penurunan kadar glukosa darah tikus putih jantan (*Rattus novergicus*) yang

- di induksi aloksan. *Buletin Veteriner Udayana*, 6(1).
- Edrisi, F., Salehi, M., Ahmadi, A., Fararoei, M., Rosta, F., & Mahmoodianfard, S. (2018). Effects of supplementation with rice husk powder and rice bran on inflammatory factors in overweight and obese adults following an energy-restricted diet: A randomized controlled trial. *European Journal of Nutrition*, 57(2), 833–843. <https://doi.org/10.1007/s00394-017-1555-3>
 - Goufo, P., & Trindade, H. (2014). Rice antioxidants: Phenolic acids, flavonoids, anthocyanins, proanthocyanidins, tocopherols, tocotrienols, γ -oryzanol, and phytic acid. *Food Science & Nutrition*, 2(2), 75–104. <https://doi.org/10.1002/fsn3.86>
 - Hongu, N., Kitts, D. D., Zawistowski, J., Dossett, C. M., Kopeć, A., Pope, B. T., & Buchowski, M. S. (2014). Pigmented Rice Bran and Plant Sterol Combination Reduces Serum Lipids in Overweight and Obese Adults. *Journal of the American College of Nutrition*, 33(3), 231–238. <https://doi.org/10.1080/07315724.2013.869772>
 - Indonesian Agricultural Departemen. (2017). Indonesian Rice Production. <https://www.pertanian.go.id>
 - International Diabetes Federation. (2017). IDF Diabetes Atlas eighth Edition. <https://diabetesatlas.org/en/>
 - International Diabetes Federation. (2019). IDF Diabetes Atlas ninth Edition. <https://diabetesatlas.org/en/>
 - Kementerian Kesehatan Republik Indonesia. (2018). Laporan Hasil Riset Kesehatan Dasar (Riskesdas) Nasional. Badan Litbangkes, Depkes RI.
 - Kim, S. M., Rico, C. W., Lee, S. C., & Kang, M. Y. (2010). Modulatory Effect of Rice Bran and Phytic Acid on Glucose Metabolism in High Fat-Fed C57BL/6N Mice. *Journal of Clinical Biochemistry and Nutrition*, advpub, 1004080015–1004080015. <https://doi.org/10.3164/jcfn.09-124>
 - Koh-Banerjee, P., Franz, M., Sampson, L., Liu, S., Jacobs, D. R., Spiegelman, D., Willett, W., & Rimm, E. (2004). Changes in whole-grain, bran, and cereal fiber consumption in relation to 8-y weight gain among men. *The American Journal of Clinical Nutrition*, 80(5), 1237–1245. <https://doi.org/10.1093/ajcn/80.5.1237>
 - Lai, M.-H., Chen, Y.-T., Chen, Y.-Y., Chang, J.-H., & Cheng, H.-H. (2011). Effects of rice bran oil on the blood lipids profiles and insulin resistance in type 2 diabetes patients. *Journal of Clinical Biochemistry and Nutrition*, 51(1), 15–18. <https://doi.org/10.3164/jcfn.11-87>
 - Moongngarm, A., Daomukda, N., & Khumpika, S. (2012). Chemical Compositions, Phytochemicals, and Antioxidant Capacity of Rice Bran, Rice Bran Layer, and Rice Germ. *APCBEE Procedia*, 2, 73–79. <https://doi.org/10.1016/j.apcbee.2012.06.014>
 - Pereira, M. A., O'Reilly, E., Augustsson, K., Fraser, G. E., Goldbourt, U., Heitmann, B. L., Hallmans, G., Knekt, P., Liu, S., & Pietinen, P. (2004). Dietary fiber and risk of coronary heart disease: A pooled analysis of cohort studies. *Archives of Internal Medicine*, 164(4), 370–376.
 - Prameswari, O. M., & Widjanarko, S. B. (2014). Test Effect of Pandan Wangi Leaf Water Extract on Reduction of Blood Glucose Level and Histopathology of Diabetes Mellitus Rat Uji Efek Ekstrak Air Daun Pandan Wangi Terhadap Penurunan Kadar Glukosa Darah dan Histopatologi Tikus Diabetes Mellitus. *Jurnal Pangan Dan Agroindustri*, 2(2), 16–27.
 - Qureshi, A. A., Sami, S. A., & Khan, F. A. (2002). Effects of stabilized rice bran, its soluble and fiber fractions on blood glucose levels and serum lipid parameters in humans with diabetes mellitus Types I and II. *The Journal of Nutritional Biochemistry*, 13(3), 175–187. [https://doi.org/10.1016/S0955-2863\(01\)00211-X](https://doi.org/10.1016/S0955-2863(01)00211-X)
 - Rohman, A., Helmiyati, S., Hapsari, M., & Setyaningrum, D. L. (2014). Rice in health and nutrition. *International Food Research Journal*, 21(1), 13–24.
 - Salehi, E. A., & Sardarodiyani, M. (2016). Bioactive phytochemicals in rice bran: Processing and functional properties. *BioChemistry: An Indian Journal*, 10(3), 1–10.
 - Sivamaruthi, B. S., Kesika, P., & Chaiyasut, C. (2018). A comprehensive review on anti-diabetic property of rice bran. *Asian Pacific Journal of Tropical Biomedicine*, 8(1), 79. <https://doi.org/10.4103/2221-1691.221142>
 - Son, M. J., Miura, Y., & Yagasaki, K. (2015). Mechanisms for antidiabetic effect of gingerol in cultured cells and obese diabetic model mice. *Cytotechnology*, 67(4), 641–652. <https://doi.org/10.1007/s10616-014-9730-3>
 - Streppel, M. T., Ocké, M. C., Boshuizen, H. C., Kok, F. J., & Kromhout, D. (2008). Dietary fiber intake in relation to coronary heart disease and all-cause mortality over 40 y: The Zutphen Study. *The American Journal of Clinical Nutrition*, 88(4), 1119–1125. <https://doi.org/10.1093/ajcn/88.4.1119>
 - Tazakori, Z., Dehghan, M. H., Iranparvar, M., Zare, M., Foladi, N., & Mohmmadi, R. (2007). Effect of rice bran powder on blood glucose levels and serum lipid parameters in diabetes patient II. *Res. J. Biol. Sci*, 2, 252–255.
 - Tuarita, M. Z., Sadek, N. F., Sukarno, S., Yuliana, N. D., & Budijanto, S. (2017). Rice Bran Development as Functional Foods: The Opportunities, Obstacles, and Challenge. *Jurnal Pangan*, 26(2), 1–11.
 - Watchararujji, K., Goto, M., Sasaki, M., & Shotipruk, A. (2008). Value-added subcritical

Saifuddin Sirajuddin et al/ The ¹Effect of Giving Rice Bran Milk on Blood Glucose Levels and Body Weight ⁱⁿ Hyperglycemic Primary School Teachers in Makassar City

⁸ water hydrolysate from rice bran and soybean meal. *Bioresource Technology*, 99(14), 6207–6213.

<https://doi.org/10.1016/j.biortech.2007.12.021>

27. Zavoshy, R., Noroozi, M., & Jahanihashemi, H. (2012). Effect of low calorie diet with rice bran

oil on cardiovascular risk factors in hyperlipidemic patients. *Journal of Research in Medical Sciences: The Official Journal of Isfahan University of Medical Sciences*, 17(7), 626–631.

ORIGINALITY REPORT

24%

SIMILARITY INDEX

19%

INTERNET SOURCES

21%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|----|
| 1 | www.jstage.jst.go.jp
Internet Source | 3% |
| 2 | E Peniati, E Setiadi, R Susanti, R S Iswari. "Anti-hyperglycemic effect of peel extract on blood sugar level of alloxan-induced Wistar rats ", Journal of Physics: Conference Series, 2018
Publication | 2% |
| 3 | www.ncbi.nlm.nih.gov
Internet Source | 2% |
| 4 | Alireza Jashni Motlagh, Reyhane Asgary, Kouros Kabir. "Evaluation of Clinical Risk Index for Babies to Predict Mortality and Morbidity in Neonates Admitted to Neonatal Intensive Care Unit", Electronic Journal of General Medicine, 2020
Publication | 1% |
| 5 | "Abstracts of the IDF Congress in Paris 2003", Diabetologia, 2003
Publication | 1% |

6	Internet Source	1 %
7	krishikosh.egranth.ac.in Internet Source	1 %
8	Milena Álvarez-Viñas, Paula Rodríguez-Seoane, Noelia Flórez-Fernández, Ma Dolores Torres et al. "Subcritical Water for the Extraction and Hydrolysis of Protein and Other Fractions in Biorefineries from Agro-food Wastes and Algae: a Review", Food and Bioprocess Technology, 2020 Publication	1 %
9	Nurliyani Nurliyani, Eni Harmayani, Sunarti Sunarti. "Synbiotic Goat Milk Kefir Lowered Peroxisome Proliferator Activated Receptor Gamma (PPAR γ) Gene Expression in Rat Adipose and Liver Tissue", Research Square, 2021 Publication	1 %
10	academic.oup.com Internet Source	1 %
11	sharc.sums.ac.ir Internet Source	1 %
12	repositori.usu.ac.id Internet Source	1 %
13	www.jfrailtyaging.com Internet Source	

1 %

14

Kim, Ja-Young, Malshick Shin, and Young-Ran Heo. "Effects of Stabilized Rice Bran on Obesity and Antioxidative Enzyme Activity in High Fat Diet-induced Obese C57BL/6 Mice", Journal of the Korean Society of Food Science and Nutrition, 2014.

Publication

<1 %

15

worldnutrijournal.org

Internet Source

<1 %

16

srv4.eulc.edu.eg

Internet Source

<1 %

17

Nakane, A.. "Effect of testicular biopsy in childhood on spermatogenesis, fertility, and paternity in adulthood-A mouse model study", Urology, 200509

Publication

<1 %

18

www.i-scholar.in

Internet Source

<1 %

19

archive.org

Internet Source

<1 %

20

jdisabilstud.org

Internet Source

<1 %

21

ppjp.ulm.ac.id

Internet Source

<1 %

22

www.scribd.com

Internet Source

<1 %

23

Sumarsono, Yafizham, D W Widjajanto. "The level of organic rice farming technology at farmer group in Ketapang village, Susukan sub-district, Semarang district, Central Java Province, Indonesia", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

24

Myoung Jin Son. "Effect of Oryzanol and Ferulic Acid on the Glucose Metabolism of Mice Fed with a High-Fat Diet", Journal of Food Science, 01/2011

Publication

<1 %

25

garuda.ristekbrin.go.id

Internet Source

<1 %

26

link.springer.com

Internet Source

<1 %

27

www.frontiersin.org

Internet Source

<1 %

28

www.ifrj.upm.edu.my

Internet Source

<1 %

29

Anna McDaniel, Jean Kristeller, Deborah Hudson. "Chart reminders increase referrals for inpatient smoking cessation intervention", Nicotine & Tobacco Research, 1999

Publication

<1 %

30

www.tandfonline.com

Internet Source

<1 %

31

Nurtaç Canpolat, Tacettin Pınarbaşı, Samih Bayrakçeken, Omer Geban. "The conceptual change approach to teaching chemical equilibrium", Research in Science & Technological Education, 2006

Publication

<1 %

32

www.mdpi.com

Internet Source

<1 %

33

dokumen.pub

Internet Source

<1 %

34

www.sportpedagogy.org.ua

Internet Source

<1 %

35

Bhupinder Kaur, Viren Ranawana, Jeyakumar Henry. "The Glycemic Index of Rice and Rice Products: A Review, and Table of GI Values", Critical Reviews in Food Science and Nutrition, 2015

Publication

<1 %

36

www.scialert.net

Internet Source

<1 %

37

Shanggong Yu, Zachary T. Nehus, Thomas M. Badger, Nianbai Fang. "Quantification of Vitamin E and γ -Oryzanol Components in Rice

<1 %

Germ and Bran", Journal of Agricultural and Food Chemistry, 2007

Publication

38

K. Higashiura, N. Ura, T. Takada, Y. Li, T. Torii, N. Togashi, M. Takada, H. Takizawa, K. Shimamoto. "The effects of an angiotensin-converting enzyme inhibitor and an angiotensin II receptor antagonist on insulin resistance in fructose-fed rats", American Journal of Hypertension, 2000

Publication

<1 %

39

hdl.handle.net

Internet Source

<1 %

40

www.hindawi.com

Internet Source

<1 %

41

www.brsfoundation.org

Internet Source

<1 %

42

www.theijmed.com

Internet Source

<1 %

43

www.jchr.org

Internet Source

<1 %

44

www.myfoodresearch.com

Internet Source

<1 %

45

repository.uhamka.ac.id

Internet Source

<1 %

46

Jianping Zhai, Li Li, Lihou Dong, Kelly Dong, Shensi Xiang, Luolan Gui, Jiaying Zhang, Haifeng Song, Zhiqiang Ge. "Simultaneous quantitative determination of liraglutide and insulin degludec in rat plasma by liquid chromatography–tandem mass spectrometry method and its application", *Biomedical Chromatography*, 2020

Publication

<1 %

47

heanoti.com

Internet Source

<1 %

48

"Dietary fibre", Wageningen Academic Publishers, 2010

Publication

<1 %

49

Kangyu Zheng, Zeju Huang, Jiaying Huang, Xiangmei Liu et al. "Effect of A 2-HP- β -Cyclodextrin Formulation on the Biological Transport and Delivery of Chemotherapeutic PLGA Nanoparticles", *Research Square*, 2020

Publication

<1 %

50

Yurike Septianingrum, Nety Mawarda Hatmanti, Andikawati Fitriasisari. "THE EFFECT OF MUROTTAL QUR'AN ON MENSTRUAL PAIN IN NURSING STUDENT OF UNIVERSITAS NAHDLATUL ULAMA SURABAYA", *Nurse and Health: Jurnal Keperawatan*, 2019

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On