

ect_of_Vitamin_A_Supplementa
tion,_Cooking_Oil_Fortification,.
pdf
by

Submission date: 19-Oct-2021 06:30PM (UTC+0700)

Submission ID: 1678001729

File name: ect_of_Vitamin_A_Supplementation,_Cooking_Oil_Fortification,.pdf (552.24K)

Word count: 4818

Character count: 23339

Vitamin A. To overcome this situation, food fortification is done to increase daily intake of Vitamin A [8]. One type of food considered to be quite potential and meet the requirements for fortification is cooking oil. Results of study in Tasikmalaya and Ciamis districts show that fortified cooking oil with 45 SI retinol/g has increased retinol levels of breastfeeding mothers from 20.5 $\mu\text{g}/\text{dl}$ at baseline to 32.5 $\mu\text{g}/\text{dl}$ at end line [9]. This study investigated the effects of giving two forms of Vitamin A supplementation ($\times 200\ 000$ SI dose and $1 \times 200\ 000$ SI dose), giving fortification of Vitamin A in cooking oil, and nutritional education on breast milk retinol levels.

Materials and Methods

Study design and site

The research used a quasi-experiment design. This research was carried out in seven public health center (Puskemas) in Gowa district, South Sulawesi Province, Indonesia from June 2017 to March 2018.

Sample

Subjects in this study were postpartum mothers in working area of Gowa District's Public Health Center selected based on criteria: (1) giving birth normally and at exact month, (2) normal birth weight, and (3) maximum 3rd parity. The minimum number of subjects needed for this study was calculated using mean different hypothesis test formula with 80% test strength and 5% significance, an average standard deviation of breast milk retinol levels is 10 $\mu\text{g}/\text{dl}$ and an average difference of breast milk retinol levels is 7.2 $\mu\text{g}/\text{dl}$. Based on calculation results, a minimum number of subjects for each treatment group is 24 people. There were four treatment groups, which meant the minimum number of subjects in this study was 96 people. At the beginning of the study, as much as 160 subjects were met the criteria, but at the end only 129 of postpartum mothers could be analyzed, while the rest of the subjects were drop out because they moved to another area (two people) or they did not have breast milk in the 2nd and 3rd collected of the sample (29 people).

Intervention

There were four treatment groups in this study, they were Group A who received 2 tablets of Vitamin A 200 000 SI dose, Group B received 1 tablet of Vitamin A 200 000 SI dose, Group C received Vitamin A fortification in cooking oil, and Group D educated about nutrition and health.

Group A was given two tablets of Vitamin A 200 000 SI dose which consumed in 2 steps. The first

tablet was given right after the mother gave a birth, and the second tablet was given minimum 24 h after the first tablet. Breast milk sample was taken before consuming the first tablet of Vitamin A.

Meanwhile, Group B was given 1 tablet of Vitamin A 200 000 SI dose on the 6th month after the mother gave a birth which was administrated by enumerator when they visited the mother's house. Yet, interviewing and collecting the first breast milk sample was done on day-3 after gave a birth.

As for Group C, they were given cooking oil fortified with Vitamin A for 3 months after gave a birth. Distribution was done along with collected baseline data. Subjects were given 1 l of cooking oil every week to consume for 12 weeks (3 months). The oil was a branded manufacturer oil that has been fortified with 62 SI of retinol palmitate.

Finally, it was Group D which was educated for 5 times that divided into two stages. Firstly was divided into two sessions when the subjects were pregnant, secondly was a moment when the subjects gave a birth (especially postpartum) for 3 times. Educating pregnant mothers was done in a group (15 pregnant mothers maximum) using pocket book and power point that has been arranged by researcher team for about 30–60 min. Educating postpartum mothers was done by visiting the subjects' houses and using leaflet containing information about Vitamin A consumption during postpartum moment and how vital is exclusive breast milk for infants. The first visiting was done in day-3 after the subject gave a birth (along with step 1 of collecting breast milk sample), the second was given in week 2–4, and the third was in week 5–6 after gave birth. Each stages took for about 10–15 min, focusing on food sources of Vitamin A, consuming Vitamin A during pregnancy and breastfeeding is essential, impact of Vitamin A deficiency, how to breastfeeding infant, and how to keep healthiness after bring forth.

Data collection

Data of subjects' characteristics were age of mother, parity, education, and employment, while characteristics of babies born were collected in this study using structured questionnaire. Food consumption's data collected were nutrition daily intake (protein, fat, Vitamin A, zinc, and iron) by 2 \times 24 h recall method (represents work day and off day). This method was done before intervention (after gave birth) and after intervention. Anthropometry measurement done in this study was body weight and height of postpartum mothers. Body weight was measured by using digital scale with 0.1 kg accuracy, while body height measured using microtoise with 0.1 cm accuracy.

Breast milk sample of postpartum mothers collected 3 times, which were: After birth (before intervention), a month after birth (during intervention),

and 3 months after birth (after intervention). The breast was cleaned first using a wet cloth before breast milk was taken as a sample. Breast milk was not taken from the breast that still full, but from the one that has not been touched by the infant for at least 30 min. The sample was taken only at the time between 09.00 and 12.00 a.m using manual pump as much as 3 and 5 ml [10]. It then placed in a vial tube and wrapped up with aluminum foil before labeled with respondent's code and date of taken. The tube was stored in refrigerator at 20°C. Levels of retinol in breast milk was analyzed using high performance liquid chromatography method [11]. This was done in Laboratory of Gizi Terpadu Badan Penelitian dan Pengembangan Kesehatan in Bogor.

Data analysis

Data were processed and analyzed using SPSS program for windows. Subject characteristics analyzed using Kruskal–Wallis to know differences in the proportion of subject characteristics between to pups A, B, C, and D. Anthropometry results were used to calculate body mass index (BMI) of breastfeeding mothers using this formula: Body weight (in kg) divided with body height square (in m). This BMI result reflected nutritional status of breastfeeding mothers. Category of BMI was wasting (<18.5 kg/m²), normal (18.5–24.9 kg/m²), overweight (25–26.9 kg/m²), and obese (≥27 kg/m²).

Nutrient intake and levels of adequacy of the subject's nutritional intake from the results of 2 × 24 h recall processed using Table of Indonesian Food Composition of 2017. The results then compared to recommended dietary allowances (RDA) of 2013. Normality test was done to ensure the normality of data distribution. Some numerical data such as levels of nutritional RDA and levels of retinol in breast milk analyzed using Analysis of variance (ANOVA) to identify the average differences of variables A, B, C, and D. ANOVA test results continued with Duncan test. To identify average differences of levels of adequate intake of nutrients before and after intervention, paired sample t test used in this study. To adjust confounder variable to the amount of difference of retinol levels in breast milk after intervention, for instance (1) retinol levels of breast milk before intervention, (2) retinol levels during intervention, and (3) adequate levels of intake of protein, fat, Vitamin A, zinc, and iron, researchers was using ANCOVA test.

Results

The Kruskal–Wallis results on Table 1 show that there were no significant differences of subject's characteristics in each treatment group (p > 0.05).

Subjects aged 26–35 years old was higher in proportion (56.6%) compared to those aged 20–25 years old (43.4%). As for levels of education, subjects graduated from high school as much as 40.3%, and most of the postpartum mothers in this study did not work (housewives) (93.8%). In parity level, 76% subjects were in the multipara category (parity ≥2). Results of BMI measurements show that normal nutritional status of postpartum mothers identified as much as 60.5%, this is dominating compared to wasting, overweight, and obesity.

Table 1. Distribution of subject in group based on characteristics

Variable	Intervention groups				Total	p
	A	B	C	D		
	n = 29	n = 30	n = 25	n = 35	n = 129	
	n (%)	n (%)	n (%)	n (%)	n (%)	
Age						
20–25 yrs	11 (37.9)	15 (50.0)	13 (48.0)	16 (45.7)	55 (42.4)	0.758
26–35 yrs	18 (62.1)	15 (50.0)	12 (44.7)	19 (54.3)	73 (56.6)	
Education						
High school	10 (34.5)	9 (30.0)	8 (27.7)	11 (31.4)	38 (29.4)	0.941
Junior High school	5 (17.2)	7 (23.3)	10 (37.0)	8 (22.9)	29 (22.5)	
Senior High school	11 (37.9)	25 (83.3)	7 (24.1)	14 (40.0)	57 (44.3)	
Diploma/Bachelor	4 (13.6)	0 (0)	4 (13.8)	2 (5.7)	10 (7.8)	
Occupation						
Unemployed	27 (93.1)	28 (93.3)	27 (93.1)	34 (97.1)	116 (90.7%)	0.993
Housewife						
Working	2 (6.9)	3 (10)	2 (6.9)	1 (2.9)	6 (4.6)	
Parity						
Primipara	9 (31.0)	8 (26.7)	10 (34.5)	7 (20.0)	34 (26.4)	0.119
Multipara	20 (69.0)	22 (73.3)	15 (50.0)	28 (80.0)	85 (65.6)	
Nutritional status						
Wasting (<18.5)	4 (13.8)	4 (13.3)	2 (6.9)	2 (5.7)	12 (9.3)	0.929
Normal (18.5–24.9)	15 (51.7)	23 (76.7)	17 (64.0)	22 (62.9)	77 (59.7)	
Overweight (25–26.9)	6 (20.7)	5 (16.7)	5 (17.3)	6 (17.1)	22 (17.1)	
Obesity (≥27.0)	4 (13.6)	3 (10)	4 (13.8)	5 (14.3)	16 (12.4)	

Table 2 shows the average levels of adequate intake of after and before interventions. Before intervention illustrates that almost all of nutritional intake of subjects between treatment groups were different (p < 0.05), except for zinc intake (p > 0.05). The nutrients intake experienced downward trends in all intervention groups unless for Vitamin A in Group C which shows an increase.

Table 2. Adequacy level of nutrients intake of subjects based on groups before and after intervention

Nutrients	Time	Groups of intervention				p
		A	B	C	D	
Protein (g)	Before	72.0 ± 0.11*	83.9 ± 0.12	75.9 ± 0.12	75.7 ± 0.10*	0.000*
	After	49.9 ± 0.13*	62.8 ± 0.13	50.5 ± 0.14*	66.7 ± 0.15	0.001*
	p [†]	0.000*	0.000*	0.000*	0.042*	
Lipid (g)	Before	50.3 ± 0.20*	72.1 ± 0.21	55.7 ± 0.22	50.7 ± 0.21	0.001*
	After	29.7 ± 0.25*	50.7 ± 0.12	32.7 ± 0.29	35.2 ± 0.29	0.001*
	p [†]	0.003*	0.000*	0.000*	0.634	
Vitamin A (µg)	Before	53.5 ± 0.24*	67.7 ± 0.15*	46.7 ± 0.24*	58.5 ± 0.27*	0.004*
	After	50.8 ± 0.34	63.5 ± 0.12	50.7 ± 0.19	55.5 ± 0.17*	0.000*
	p [†]	0.004*	0.000*	0.000*	0.001*	
Zinc (µg)	Before	32.8 ± 0.14*	40.5 ± 0.15	38.5 ± 0.18*	37.1 ± 0.18*	0.100
	After	22.5 ± 0.13*	29.4 ± 0.12*	25.2 ± 0.14*	28.4 ± 0.15*	0.000*
	p [†]	0.000*	0.000*	0.000*	0.000*	
Iron (µg)	Before	55.0 ± 0.20*	55.7 ± 0.24	52.1 ± 0.19	50.7 ± 0.22	0.000*
	After	35.8 ± 0.18*	29.8 ± 0.21	28.9 ± 0.19*	33.1 ± 0.18	0.785
	p [†]	0.811	0.000*	0.317	0.169	

* Significant difference (p < 0.05) between before and after intervention. † Significant difference (p < 0.05) between before (2 days after delivery) and after (3 months after delivery).

The subjects' average retinol level before treatment was 52.15 µg/dl and the highest score was found in D group as much as 61.25 µg/dl (Table 3). ANOVA test results show no significant differences at the time before, during, and after interventions, similar case happened for differences of average retinol levels of breast milk between intervention groups (p > 0.05).

Table 3: Breast milk retinol levels of subjects based on group before, during, and after 3 months intervention

Retinol (µg/dl)	Intervention Group				p
	A	B	C	D	
Before	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	
During	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	
After	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	
Difference					

ANOVA results show no significant differences of retinol levels of breast milk at the time before, during, and after interventions, it is similar to the difference (decreasing) between intervention groups (p > 0.05). It means that intervention such as high dosage of Vitamin A supplementation (2 capsules and 1 capsule), fortified cooking oil, or nutritional education has similar effect on breast milk retinol levels. Other studies by Bezerra et al. [12] and Tomiya et al. [13] show that consumption of 2 capsules and 1 capsule of Vitamin A did not have a differences on retinol levels between groups for 1 month and 4 months of interventions.

A significant reducing on retinol levels of breast milk was also found after intervention in A and D groups (p < 0.05), but this was not found in B and C groups (p > 0.05). Figure 1 shows that subject with normal category (>30 µg/dl) of breast milk retinol levels before intervention was higher rather than low category (found in all intervention groups).




Figure 1: Percentages of subjects with normal category of breast milk retinol levels before, during, and after 3 months of interventions

Discussion

In this study, it was found that there were no differences of breast milk retinol levels between intervention groups at the time before, during, and after intervention. Moreover, it also known that subjects given fortified cooking oil has higher retinol levels rather than other groups after intervention. This is similar to a study conducted by Sandjaja et al. who found that giving cooking oil at baseline became 32.5 µg/dl at end line [9].

ANOVA results show no significant differences of retinol levels of breast milk at the time before (0 month), during (1 month), and after (3 months) interventions, it is similar to the difference (decreasing) between intervention groups (p > 0.05). It means that intervention such as high dosage of Vitamin A supplementation (2 capsules and 1 capsule), fortified cooking oil, or nutritional education has similar effect on breast milk retinol levels. Other studies by Bezerra et al. [12] and Tomiya et al. [13] show that consumption of 2 capsules and 1 capsule of Vitamin A did not have a differences on retinol levels between groups for 1 month and 4 months of interventions.

Theoretically, retinol levels in breast milk will decrease as the infant is growing [14]. This study shows a decreasing average levels of breast milk retinol in all intervention groups, whether 1 month or 3 months after mother's gave birth. Nevertheless, the levels were still categorized as normal (>30 µg/dl). When they were analyzed based on the groups (intragroup), a significant decrease of retinol levels are shown at the time before, during, and after interventions in A and D groups (p < 0.05) (Table 3). It does not identified in B and C Groups (p > 0.05). This might happen in Group B due to the supplementation of Vitamin A was done in 6th week after gave birth, or because of the average intake of nutrition before treatment was better than after that. Different results were obtained by Tomiya et al. [13], where they found a significant decrease of retinol levels of breast milk at the end of their treatment targeting postpartum mothers given 1 capsule of Vitamin A (p < 0.05). Different time of consuming Vitamin A by the subject can be the reason of the results. The study of Tomiya et al. 2015 [13] given the subjects 1 capsule of Vitamin A at 6th week after gave birth.

Furthermore, this study also found that there was no significant decrease of breast milk retinol levels in C group (fortified cooking oil group). This might be due to intervention of cooking oil every week that affected intake of Vitamin A of mothers. Sandjaja et al. found fortified cooking oil with Vitamin A could increase retinol levels of mothers' breast milk who breastfeed babies aged 6–12 months [9].

Decreasing of retinol levels of breast milk experienced by A group which significance 3 months after gave birth. It indicates that supplementation of Vitamin A did not affect breast milk retinol levels. Theoretically, it might happen because when mothers gave birth, it can trigger acute response that can increase C-protein reactive levels. This is believed that it affects metabolism and transportation of Vitamin A in our body, so that consumption of high dose of Vitamin A after gave birth might not affect Vitamin A levels of breast milk [13].

The examination of breast milk retinol levels on the 1st month after delivery (during intervention) shows that there has been a decrease in the proportion of mothers with normal categories of breast milk retinol. The largest decrease occurred in group D (around 40%). This decline was probably due to the intervention in the form of education that did not directly affect levels of retinol in breast milk. It is contrast to other interventions, educational effect given to the subject requires time to be able to change the habit of consuming food sources of Vitamin A which is expected to affect the retinol levels of breast milk.

At the end of the study, it can be seen that the proportion of breast milk retinol levels in the normal category has decreased (this occurred in all treatment groups). The highest percentage of subjects with normal breastfeeding retinol category was found in Group C

while the lowest in Group A. However, the Kruskal–Wallis test results showed no significant difference in the proportion of subjects with normal breastfeeding categories between the treatment groups before, during, and after intervention ($p > 0.05$). Overall, it can be seen in Picture 1 that about 1 in 2 breastfeeding mothers experience vitamin deficiency problems (breast milk retinol $<30 \mu\text{g/dl}$). Based on the WHO, the problem of Vitamin A deficiency in breastfeeding mothers in this study area is included in the severe category of public health problems because the percentage is more than 25% [15].

Conclusions

It can be concluded that the intervention given to the subject had the same effect on retinol levels in breast milk. In this study, it was found that the administration of cooking oil fortified with 62 SI vitamin A and supplementation with 1 capsule of vitamin A tended to reduce the proportion of mothers with low retinol levels compared to supplementation with 2 capsules of vitamin A and nutrition education.

References

- World Health Organization. Guideline: Vitamin A Supplementation in Postpartum Women. Geneva: World Health Organization; 2011.
- Humphrey JH, Agoestina T, Wu L, Usman A, Nurachim M, Subardja D, et al. Impact of neonatal Vitamin A supplementation on infant morbidity and mortality. *J Pediatr*. 1996;128(4):489-96. [https://doi.org/10.1016/s0022-3476\(96\)70359-1](https://doi.org/10.1016/s0022-3476(96)70359-1) PMID:8618182
- Roy S, Islam A, Molla A, Akramuzzaman S, Jahan F, Fuchs G. Impact of a single megadose of Vitamin A at delivery on breastmilk of mothers and morbidity of their infants. *Eur J Clin Nutr*. 1997;51(5):302-7. <https://doi.org/10.1038/sj.ejcn.1600398> PMID:9152680
- Klemm RDW, Labrique AB, Christian P, Rashid M, Shamim AA, Katz J, et al. Newborn Vitamin A supplementation reduced infant mortality in rural Bangladesh. *Pediatrics*. 2008;122(1):e243-50. <https://doi.org/10.1542/peds.2007-3448> PMID:18595969
- Rotendi MA, Khebbi N. Vitamin A supplementation and neonatal mortality in the developing world: A meta-regression of cluster-randomized trials. *Bull World Health Organ*. 2010;88(9):697-702. <https://doi.org/10.2471/blt.09.068080> PMID:20865075
- Bhaskaran P, Balakrishna N. Effect of administration of 200,000 IU of Vitamin A to women within 24 hrs after delivery on response to OPV administered to the newborn. *Indian Pediatr*. 1998;35(3):217-22. PMID:9707874
- Kemenkes. Kementerian Kesehatan. Laporan Hasil Riset Kesehatan Dasar 2013. Jakarta: Kemenkes; 2013.
- Dary O, Mora JO. Food fortification to reduce Vitamin A deficiency: international Vitamin A consultative group recommendations. *J Nutr*. 2002;132 (Suppl. 9):2927S-53S. <https://doi.org/10.1093/jn/132.9.2927s> PMID:12221271
- Sandjaja, Jus'at I, Jahari AB, Iffrad, Hiet MK, Tilden RL, et al. Vitamin A-fortified cooking oil reduces Vitamin A deficiency in infants, young children and women: Results from a programme evaluation in Indonesia. *Public Health Nutr*. 2015;18(14):2511-22. <https://doi.org/10.1017/s136898001400322x> PMID:25391926
- Stoltzfus RJ, Underwood B. Breast-milk Vitamin A as an indicator of the Vitamin A status of women and infants. *Bull World Health Organ*. 1995;73(5):703-11. PMID:8846497
- Espesito M, Sakurai E, Laineurier JA, Teixeira RA, Bonomo E, da Silva CA, et al. Retinol and fat from breast milk of Brazilian mothers at high risk for food unsafe. *Ann Public Health Res*. 2017;4(3):1063-8.
- Bezerra DS, de Araujo KF, Azevêdo GM, Dimenstein R. A randomized trial evaluating the effect of 2 regimens of maternal Vitamin A supplementation on breast milk retinol levels. *J Hum Lact*. 2010;6(2):148-56. <https://doi.org/10.1177/0890334409356445> PMID:20110563
- Tomiya MT, de Arruda K, da Silva Diniz A, Santana RA, da Silveira KC, Andreto LM. The effect of Vitamin A supplementation with 400,000 IU vs 200,000 IU on retinol concentrations in the breast milk: A randomized clinical trial. *Clin Nutr*. 2015;36(1):100-6. <https://doi.org/10.1016/j.clnu.2015.11.018> PMID:26725194
- Newman V. Vitamin A and Breastfeeding: A Comparison of Data from Developed and Developing Countries. San Diego: US: Wellstart International; 1993.
- World Health Organization. Indicator for Assessing Vitamin A Deficiency and their Application in Monitoring and Evaluating Intervention Programmes. Geneva: World Health Organization; 1998.

ect_of_Vitamin_A_Supplementation,_Cooking_Oil_Fortificatio...

ORIGINALITY REPORT

12%
SIMILARITY INDEX

10%
INTERNET SOURCES

8%
PUBLICATIONS

1%
STUDENT PAPERS

PRIMARY SOURCES

1 media.neliti.com
Internet Source **1%**

2 www.repositorio.ufpa.br
Internet Source **1%**

3 zenodo.org
Internet Source **1%**

4 www.e3s-conferences.org
Internet Source **1%**

5 f1000research.com
Internet Source **<1%**

6 sites.kowsarpub.com
Internet Source **<1%**

7 www.science.gov
Internet Source **<1%**

8 Cristiane Santos Sânzio Gurgel, Larisa Alves de Araújo Pereira, Aldiane de Assis Costa, Mayara Adja da Silva Souza et al. "Effect of routine prenatal supplementation on vitamin

concentrations in maternal serum and breast milk", Nutrition, 2017

Publication



www.frontiersin.org

Internet Source

<1 %



"IUNS. 21st International Congress of Nutrition. Buenos Aires, Argentina, October 15-20, 2017: Abstracts", Annals of Nutrition and Metabolism, 2017

Publication

<1 %



www.unu.edu

Internet Source

<1 %



newinera.com

Internet Source

<1 %



ijae.springeropen.com

Internet Source

<1 %



www.frontiersin.org

Internet Source

<1 %



Audrey Messelt, Lauren Thomaier, Patricia I. Jewett, Heewon Lee et al. "Comparisons of emotional health by diagnosis among women with early stage gynecological cancers", Gynecologic Oncology, 2021

Publication

<1 %



annalsofrscb.ro

Internet Source

<1 %

17

edepot.wur.nl

Internet Source

<1 %

18

www.journals.aiac.org.au

Internet Source

<1 %

19

www.researchsquare.com

Internet Source

<1 %

20

enutrition.fnri.dost.gov.ph

Internet Source

<1 %

21

Erny Y. Mulyani, Mardiansyah, Dodik Briawan, Dudi I. Santoso, Idrus Jus'at. "Effect of dehydration during pregnancy on birth weight and length in West Jakarta", *Journal of Nutritional Science*, 2021

Publication

<1 %

22

Ulbricht, Catherine, Ethan Basch, Wendy Chao, Julie Conquer, Dawn Costa, Samantha Culwell, Kelly Flanagan, Jackie Guilford, Paul Hammerness, Sadaf Hashmi, Richard Isaac, Erica Rusie, Jill M. Grimes Serrano, Christine Ulbricht, Mamta Vora, Regina C. Windsor, Matt Woloszyn, and Sara Zhou. "An Evidence-Based Systematic Review of Vitamin A by the Natural Standard Research Collaboration", *Journal of Dietary Supplements*, 2012.

Publication

<1 %

23

journal.unhas.ac.id

Internet Source

<1 %

24

www.yumpu.com

Internet Source

<1%

25

[Carotenoids and Human Health, 2013.](#)

Publication

<1%

26

[NIKMAN UTAMI DEWI, TRIAS MANMUDJONO,](#)

["EFFECTIVENESS OF FOOD FORTIFICATION IN IMPROVING NUTRITIONAL STATUS OF MOTHERS AND CHILDREN IN INDONESIA". International Journal of Environmental Research and Public Health 2021](#)

Publication

<1%

27

[Rajiv Bahl, Nita Bhandari, Mohammed A. Wahed, Geeta T. Kumar, Maharaj K. Bhan.](#)

["Vitamin A Supplementation of Women Postpartum and of Their Infants at Immunization Alters Breast Milk Retinol and Infant Vitamin A Status", The Journal of Nutrition, 2002](#)

Publication

<1%

28

[T M Martins. "Impact of maternal vitamin A supplementation on the mother-infant pair in Brazil", European Journal of Clinical Nutrition, 09/15/2010](#)

Publication

<1%

29

academic.oup.com

Internet Source

<1%

eprint.ncl.ac.uk

30	Internet Source	<1 %
31	internationalbreastfeedingjournal.biomedcentral.com Internet Source	<1 %
32	ir.lib.uwo.ca Internet Source	<1 %
33	scielosp.org Internet Source	<1 %
34	www.karger.com Internet Source	<1 %
35	www.scielo.br Internet Source	<1 %
36	Mojtaba Keikha, Ramin Shayan-Moghadam, Maryam Bahreynian, Roya Kelishadi. "Nutritional supplements and mother's milk composition: a systematic review of interventional studies", International Breastfeeding Journal, 2021 Publication	<1 %
37	Ye Ding, Ping Hu, Yue Yang, Mei Ye et al. "Daily Oral Low-Dose Vitamin A Supplementation Increases the Retinol Levels in Serum and Breast Milk of Lactating Mothers and Has No Effect on Infant Health Status: A Randomized Placebo-Controlled Trial in China", Research Square, 2020	<1 %

Publication

Exclude quotes

Exclude matches

Exclude bibliography