

Lipocalin 2 could predict circulating MMP9 levels in patients with breast cancer

by Liong Boy Kurniawan

Submission date: 11-Jul-2021 06:47PM (UTC+0700)

Submission ID: 1618136813

File name: bd-40-bd219017-2.pdf (43.75K)

Word count: 1664

Character count: 8894

Lipocalin 2 could predict circulating MMP9 levels in patients with breast cancer

Ulung Bahrun^{a,b,c}, Wildana Wildana^d, Hegaria Rahmawati^e, Liong Boy Kurniawan^{a,b,*} and William Hamdani^{b,c,f}

^aDepartment of Clinical Pathology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

^bHasanuddin University Hospital, Makassar, Indonesia

^cDr. Wahidin Sudirohusodo Hospital, Makassar, Indonesia

^dLamadukkelleng Hospital, Sengkang, South Sulawesi, Indonesia

^ePKU Muhammadiyah Hospital, Gombong Kebumen, Central Java, Indonesia

^fDivision of Oncology, Department of Surgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

Abstract.

OBJECTIVES: Breast cancer is the most prevalent carcinoma found in Indonesian women, and its incidence remains high worldwide. Lipocalin 2 has been linked with the progression of breast cancer. Matrix metalloproteinase 9 (MMP9) is an enzyme that has an important role in angiogenesis. We investigated the relationship between lipocalin 2 and MMP9 and the ability of lipocalin 2 for predicting MMP9 levels in female patients with breast cancer.

METHOD: A total of 55 female patients with breast cancer were enrolled in this cross-sectional study. Lipocalin 2 and MMP9 were measured by enzyme-linked immunosorbent assay.

RESULTS: Lipocalin 2 was significantly correlated with MMP9 levels ($r = 0.756$, $p < 0.001$). Lipocalin 2 levels could describe the MMP9 levels ($\beta = 0.76$, $p < 0.001$, $R^2 = 56.9\%$).

CONCLUSION: Higher lipocalin 2 levels in female patients with breast cancer indicate higher MMP9 levels. Lipocalin 2 can be used to predict MMP9 levels.

Keywords: Lipocalin 2, MMP9, breast cancer, female

1. Background

Breast cancer has the highest prevalence among other types of cancer in adult females. In women in the USA, new breast cancer cases are the highest compared to other cancer types, while the estimated death due to breast cancer is ranked second after lung and bronchial cancer [1]. In Indonesia, the incidence of breast cancer was highest (30.9%) compared to that of other types of cancer among the female population in 2018 [2].

Lipocalin 2, also known as neutrophil gelatinase-associated lipocalin (NGAL), is a secreted glycoprotein of the adipokine superfamily produced by various cells and tissues, including neutrophils adipose, bone

marrow, and spleen. Neoplastic tissues from several organs, including the breast, colon, lung, express higher lipocalin 2 levels compared to the normal tissues [3–5]. Lipocalin 2 promotes breast cancer progression [6]. Conversely, inhibiting lipocalin 2 activity via inhibitory monoclonal antibodies reduced the metastasis progression of breast cancer in mice with breast tumors [7]. Therefore, lipocalin 2 is proposed as a potential target in the therapy of breast cancer metastasis [8].

Matrix metalloproteinases (MMPs) are a family of endopeptidases containing zinc and can degrade various components of the extracellular matrix. MMPs have important roles in the development of angiogenesis [9]. MMP9 is a member of MMPs with gelatinase activity and the ability to break down type IV collagen of the basement membrane; thus, it has an important role in tumor cell invasion and metastasis. Patients with breast cancer had higher serum MMP9 levels than those

*Corresponding author: Liong Boy Kurniawan, Department of Clinical Pathology, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia. E-mail: liongboykurniawan@yahoo.com.

15

Table 1

Characteristics of the patients with breast cancer

| Variable | n (%) | Mean ± SD | Median | Min. | Max. |
|---------------------|---------|-----------------|--------|--------|---------|
| Stage | | | | | |
| Early | 11 (20) | | | | |
| Late | 44 (80) | | | | |
| Age (years) | | 43.64 ± 9.05 | 42 | 31 | 68 |
| Lipocalin 2 (ng/mL) | | 104.86 ± 73.51 | 79.58 | 33.43 | 347.95 |
| MMP9 (ng/mL) | | 766.34 ± 408.43 | 671.98 | 159.25 | 2400.81 |

with benign breast disease and normal controls [10]. Higher plasma MMP9 levels have been associated with lymph node metastasis and higher tumor stage [10,11]. Lipocalin 2 may be present with MMP9 in the form of a complex via the disulfide bridge and increases the gelatinolytic activity of MMP9 [1].

In the present study, we investigated the relationship between lipocalin 2 and MMP9 and the ability of lipocalin 2 for predicting MMP9 levels among female patients with breast cancer.

2. Methods

2.1. Study population and ethics approval

This cross-sectional study was conducted from June 13 to September 2013. The ethical recommendation was approved by the Health Research Ethical Committee of the Faculty of Medicine, Hasanuddin University, Makassar, Indonesia, following the Declaration of Helsinki.

The participants were women with breast cancer hospitalized at Dr. Wahidin Sudirohusodo Hospital, Makassar, Indonesia. Breast cancer was confirmed based on the histopathological diagnosis. Patients who had previous breast cancer therapy, recent infection and a history of kidney and liver disorders were excluded from the study.

2.2. Laboratory procedures

A venous blood sample (3 ccs) was obtained from each participant in a tube without anticoagulant, followed by serum separation with 5-minute centrifugation at 300 pm. The serum was stored at -20 °C until lipocalin 2, and MMP9 testing was performed. Lipocalin 2 and MMP9 were measured by enzyme-linked immunosorbent assay (ELISA) (Quantikine, Minneapolis).

Table 2
Correlation between Age and Lipocalin 2 with MMP9 levels

| Age | | Lipocalin 2 | |
|--------|-------|-------------|--------|
| r | p* | r | p* |
| -0.163 | 0.234 | 0.756 | <0.001 |

*Spearman correlation test.

2.3. Statistical analysis

Data distribution normality was tested with the Kolmogorov-Smirnov test. The correlation between parameters was analyzed with either the Pearson or Spearman test. The association between lipocalin 2 and MMP9 levels was determined with linear regression testing.

3. Results

A total of 55 women with breast cancer were recruited to this study, consisted of 11 patients (20%) with early-stage disease (stage 1 and 2) and 44 patients (80%) with late-stage disease (stage 3 and 4). The mean patient age was 43.64 ± 9.05 years (Table 1).

There was a significant correlation between lipocalin 2 with MMP9 levels ($r = 0.756$, $p < 0.001$), but age was not correlated with MMP9 levels (Table 2).

Linear regression analysis showed that lipocalin 2 could describe 56.9% of serum MMP9 levels (Table 3).

4. Discussion

Lipocalin 2 has been used to predict the progression of several cancers, including breast cancer. Lipocalin 2 is overexpressed in breast cancer cells and

Table 3
Linear regression analysis of Lipocalin 2 and MMP9 levels

| | Variable | β | <i>p</i> |
|----------------------|-------------|---------|----------|
| Model $R^2 = 56.9\%$ | Lipocalin 2 | 0.76 | <0.001 |

can upregulate several mesenchymal markers, including fibronectin and vimentin, and downregulates the epithelial marker E-cadherin, resulting in increased tumor cell motility and invasiveness [6]. The HER2 upregulates the high expression of lipocalin 2 in breast cancer cells—phosphoinositide 3-kinase (PI3K)—AKT—NF- κ B pathway [7].

Here, we show that serum lipocalin 2 was significantly correlated with circulating MMP9 levels in women with breast cancer. Interestingly, the highlight of our findings is that lipocalin 2 levels alone could predict 56.9% of serum MMP9 levels. There are several possible explanations for this. Lipocalin 2 may form a complex with MMP9 via a disulfide bridge, rendering MMP9 less prone to degradation by tissue inhibitor of metalloproteinases 1 (TIMP1) due to the protective effect of lipocalin 2 on the structural stability of MMP9 in the form of the complex [1]. Lipocalin 2 can induce increased MMP9 enzymatic activity, thus contributing to MMP9 levels. Lipocalin 2 also regulate pro-MMP9 by inducing the activation of pro-MMP9. These actions may explain the strong effect of serum lipocalin 2 on MMP9 levels in patients with breast cancer [4]. A study on gastric cancer reported that hepatic growth factor produced by mesenchymal cells upregulated lipocalin 2 expressions. The upregulation of lipocalin 2 regulated MMP9 expression through the PI3K—AKT—NF- κ B path [12].

This study has some limitations. First, the cross-sectional design cannot describe the causality interaction between the variables observed. Second, the number of patients in this study was limited; therefore, future studies should involve a larger sample size.

5. Conclusion

Higher lipocalin 2 levels in women with breast cancer indicate higher MMP9 levels. Lipocalin 2 can be used for predicting MMP9 levels.

Conflict of interest

None.

Funding source

None.

References

- [1] Siegel RL, Miller KD, Jemal A. Cancer statistics, *CA Cancer J Clin*, 69(1): 7–34, 2019.
- [2] Organization WH. International Agency for Research on Cancer. Published online 2019, 2019.
- [3] Roldo S, Abrams SL, Steelman LS et al., Roles of NGAL and MMP-9 in the tumor microenvironment and sensitivity to targeted therapy, *Biochim Biophys Acta (BBA)-Mol Cell Res*, 1863(3): 438–448, 2016.
- [4] Santiago-Sánchez GS, Pita-Grisanti V, Quiñones-Díaz B, Sampedro K, Cruz-Monserrate Z, Vivas-Mejía PE, Biological functions and therapeutic potential of Lipocalin 2 in cancer, *Int J Mol Sci*, 21(12): 4365, 2020.
- [5] Yang J, Moses MA, Lipocalin 2: a multifaceted modulator of cancer, *Cell Cycle*, 8(15): 2347–2352, 2009.
- [6] Wang J, Bielenberg DR, Rodig SJ et al., Lipocalin 2 promotes breast cancer progression, *Proc Natl Acad Sci USA*, 106(10): 3913–3918, 2009.
- [7] Wang X, Ding T, Lin H et al., Inhibition of lipocalin 2 impairs breast tumorigenesis and metastasis, *Cancer Res*, 69(22): 8579–8584, 2009.
- [8] Chen C, Yang K, Li M, Huang W, Zhang F, Wang H, Lipocalin 2: a potential therapeutic target for breast cancer metastasis, *Oncotargets Ther*, 11: 8099, 2018.
- [9] Srinivasan J, Matrix metalloproteinases and angiogenesis, *J Cell Mol Med*, 9(2): 267–285, 2005.
- [10] Wu Z, Wu Q, Yang J et al., Prognostic significance of MMP-9 and TIMP-1 serum and tissue expression in breast cancer, *Int J Cancer*, 122(9): 2050–2056, 2008.
- [11] Hemati S, Sadeghi M, Motovali-Bashi M, Sahebi A, Malekshahi ZV, Higher plasma MMP-9 level in breast cancer patients, *J Biol Res*, 13: 113, 2010.
- [12] Lee KH, HGF mediated upregulation of lipocalin 2 regulates MMP9 through nuclear factor- κ B activation, *Oncol Rep*, 34(4): 2179–2187, 2015.

Lipocalin 2 could predict circulating MMP9 levels in patients with breast cancer

ORIGINALITY REPORT

18%

SIMILARITY INDEX

10%

INTERNET SOURCES

17%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|--|----|
| 1 | Semon Wu, Lung-An Hsu, Ming-Sheng Teng, Jeng-Feng Lin, Hsien-Hsun Chang, Yu-Chen Sun, Hsuan-Pu Chen, Yu-Lin Ko. "Association of SELE genotypes/haplotypes with sE-selectin levels in Taiwanese individuals: interactive effect of MMP9 level", BMC Medical Genetics, 2012 Publication | 2% |
| 2 | ar.iarjournals.org Internet Source | 2% |
| 3 | www.mdpi.com Internet Source | 2% |
| 4 | "Encyclopedia of Cancer", Springer Science and Business Media LLC, 2017 Publication | 2% |
| 5 | mafiadoc.com Internet Source | 1% |
| 6 | translational-medicine.biomedcentral.com Internet Source | 1% |

7

Rikki A. M. Brown, Kirsty L. Richardson, Tasnuva D. Kabir, Debbie Trinder, Ruth Ganss, Peter J. Leedman. "Altered Iron Metabolism and Impact in Cancer Biology, Metastasis, and Immunology", *Frontiers in Oncology*, 2020

Publication

1 %

8

Sayan Ghosh, Nadezda Stepicheva, Meysam Yazdankhah, Peng Shang et al. "The role of lipocalin-2 in age-related macular degeneration (AMD)", *Cellular and Molecular Life Sciences*, 2020

Publication

1 %

9

Jun Wu, Jing Wang, Qiang Su, Wei Ding, Teng Li, Junxian Yu, Bangwei Cao. "Traditional Chinese medicine *Astragalus* polysaccharide enhanced antitumor effects of the angiogenesis inhibitor apatinib in pancreatic cancer cells on proliferation, invasiveness, and apoptosis", *OncoTargets and Therapy*, 2018

Publication

1 %

10

diposit.ub.edu

Internet Source

1 %

11

Sina Rahimi, Amaneh Mohammadi Roushandeh, Ebrahim Ahmadzadeh, Ali Jahanian-Najafabadi, Mehryar Habibi Roudkenar. "Implication and role of neutrophil gelatinase-associated lipocalin in cancer: lipocalin-2 as a potential novel

1 %

emerging comprehensive therapeutic target for a variety of cancer types", Molecular Biology Reports, 2020

Publication

12

Zheng-sheng Wu. "miR-340 inhibition of breast cancer cell migration and invasion through targeting of oncoprotein c-Met", Cancer, 07/01/2011

Publication

1 %

13

www.frontiersin.org

Internet Source

1 %

14

Mustafa Erman, Esmen Baltali, Aziz Karaoglu, Huseyin Abali et al. "A Phase II Study on the Safety and Efficacy of 5-Fluorouracil, Epirubicin, Cyclophosphamide (FEC) Followed by Paclitaxel in the Adjuvant Treatment of Breast Cancer", Cancer Investigation, 2009

Publication

1 %

15

A. Oshima. "Survival of Male Breast Cancer Patients: A Population-Based Study in Osaka, Japan", Japanese Journal of Clinical Oncology, 09/25/2006

Publication

1 %

16

koreascience.or.kr

Internet Source

1 %

17

tessera.spandidos-publications.com

Internet Source

1 %

18 www.futuremedicine.com 1 %
Internet Source

19 www.tandfonline.com 1 %
Internet Source

20 Gerasimos Socrates Christodoulatos, Nikolaos Spyrou, Jona Kadillari, Sotiria Psallida, Maria Dalamaga. "The Role of Adipokines in Breast Cancer: Current Evidence and Perspectives", Current Obesity Reports, 2019 <1 %
Publication

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On