

Component, Fatty Acid And Mineral Composition Of Rice Bran Oil Extracted By Multistage With Hexane And Ethanol

by Zainal 04

FILE	COMPONENT , FATTY ACID AND MINERAL COMPOSITION OF RICE BRAN OIL EXTRACTED BY MULTISTAGE WITH HEXANE AND ETHANOL .PDF (511.45K)	WORD COUNT	6169
TIME SUBMITTED	13-APR-2019 03:45PM (UTC+0700)	CHARACTER COUNT	32954
SUBMISSION ID	1111640379		

Component, Fatty Acid And Mineral Composition Of Rice Bran Oil Extracted By Multistage With Hexane And Ethanol

Fajriyati Mas'ud, Meta Mahendradatta, Amran Laga, Zainal Zainal

Abstract: Rice bran oil (RBO) has been extracted from Celebes rice bran by multistage extraction with hexane solvent followed by ethanol to see the component, profile of fatty acids and mineral contained in both of them. As a comparison, RBO directly extracted with ethanol was also presented. Extraction process was performed using reflux method at 55°C, for 5 hours with bran and solvent ratio of 1:7. Analysis of components and fatty acids of RBO was conducted with GC-MS QP 2010 Shimadzu. Oleic, linoleic and palmitic were found dominant in first stage extraction by hexane with concentration of 3716.56, 1630.78 and 1021.89 mg/L, respectively. Palmitic (6.34 mg/L), lauric (4.78 mg/L), and linoleic (3.52 mg/L) were dominant in the second stage extraction by ethanol. Linoleic (28.85 mg/L), stearic (2.88 mg/L) and myristic (2.02 mg/L) were found in extracted directly by ethanol. RBO extracted with hexane had 18.6% of saturated fatty acid and 81.4% of unsaturated fatty acids, with ratio of saturated fatty acids : monounsaturated fatty acids: polyunsaturated fatty acids of approximately 1 : 2.3 : 1.3. It contained about 56.7% of monounsaturated, 24.7% of polyunsaturated, and 18.6% of saturated fatty acids. In the present paper, we provide also an analysis of mineral composition of RBO by X-ray Spectrometer and melting point of RBO by Differential Scanning Calorimeter (DSC) instrument.

Keywords: Edible oil, fatty acid, mineral, multistage extraction, solvent extraction

1 INTRODUCTION

Rice bran, a by-product which is obtained from the rice milling processes, it can be used as a major source of RBO [1], [2]. The utilization of RBO has increased considerable due to its well-known for wealth of phytochemical [3] and its fatty acid profile. Fatty acids played an important role in metabolism, which was the main fuel for the metabolism, storage and transportation of energy, as an essential component of all membranes, and as a general regulator [4]. The nutritional recommendations indicated that 25% of energy should be covered by lipid [5] and total fat daily intake of 20%-35% of energy was recommended by Food and Agriculture Organization (FAO) of the United Nations and the World Health Organization (WHO) [6].

Oil consisted of fatty acids which were classified based on the presence or absence of double bonds as saturated fatty acids (SFA), without double bonds; monounsaturated fatty acids (MUFA), with one double bond; and polyunsaturated fatty acids (PUFA), with two up to six double bonds [7]. The human body cannot synthesize PUFA with the first double bond on C3 and C6 of the methyl-end due to the absence of appropriate enzymes. Thus, these fatty acids must be obtained from diet [8], [9], [10]. Analysis of fatty acids in the food was very important because of the implications on nutrition and its effect on health [11]. Oil fractionation to determine the fatty acid composition was quite urgent to produce a product with physical or nutritional properties of interest to the food industry [12]. The content of fatty acids and the ratio of saturated and unsaturated fatty acids were important parameters determining the nutritional value of the edible oil, because the latest trends in the food processing industry to inform the composition of fatty acids in the oil. Related with it, the physical and chemical characteristics of oils and fats are strongly influenced by the kind and proportion of fatty acids on triacylglycerols [13], [14]. In general, fatty acids in oils and fats extracted from plant consisted of unsaturated fatty acids in high percentage. Unsaturated fatty acids were more susceptible to oxidative damage [14], [15], [16]. Therefore, it was important to know the composition of fatty acids from oils or fats, to identify their characteristics and to determine the possibility of adulteration as well as to know the stability and physicochemical properties of oils [14], [17]. One method used to determine the composition of the fatty acids was based on chromatography analysis of their methyl ester [18]. The major components of lipids were fatty acids, and their physical, chemical, and physiological properties depend largely on its fatty acid composition. The quality and freshness of the vegetable oil were determined by the constituent fatty acid, because long chain organic acids were not highly volatile compounds [19], [20]. On the other hand, although the overall quality of the product was generally defined by the culinary benefits, the presence of inorganic compounds in the oils has very important role in terms of food safety and general product longevity. The mechanisms whereby inorganic constituents

- Fajriyati Mas'ud, is currently pursuing doctoral degree program in Food Technology Program, Hasanuddin University Makassar, and working as lecturer in Department of Chemical Engineering, State Polytechnic of Sung Pandang, Makassar, South Sulawesi, 90245, Indonesia. E-mail: fajri888@poliupg.ac.id
- Dr. Meta Mahendradatta, Professor of Food Technology, and working as lecturer in Department of Agricultural Technology, Hasanuddin University, Makassar, South Sulawesi, 90245, Indonesia
- Dr. Amran Laga, Professor of Food Technology, and working as lecturer in Department of Agricultural Technology, Hasanuddin University, Makassar, South Sulawesi, 90245, Indonesia
- Dr. Zainal Zainal, is lecturer in Department of Agricultural Technology, Hasanuddin University, Makassar, South Sulawesi, 90245, Indonesia

were incorporated into the oil include the natural uptake and accumulation of trace element by the plant and addition or adulteration at some point. The inorganic component could also reach the oil as contamination during processing [21]. Some elements might act as an accelerator auto oxidation, changing the taste and quality of products from time to time [22]. The quality of vegetable oil was directly related to the concentration of the mineral [23], the concentration of elements and substances that are significant in the end product can affect the taste, color and stability. One way of extracting neutral lipid fractions from plant sources of oil was the use of organic solvents [24]. Soxhlet extraction results in sesame revealed that polar solvent such as ethanol was good solvent compared to non-polar solvents. These results could be explained by the interaction between the unsaturated fatty acids with a polar solvent, compared with non-polar solvents [25]. On the other hand, hexane was the most commonly used, and preferred for the extraction of oil from seeds, due to its availability at a reasonable cost and its suitable functional characteristics for oil extraction. Among these characteristics are high-strength solvent to dissolve triglycerides at a fairly low temperature, non-reactivity with oil and miscella, and requires simple equipment. Related to solvents, some report studies had compared the effectiveness of hexane and ethanol in oil extraction [26]. But throughout our literature study, RBO extraction was generally carried out with one kind of solvent and repeated several times, or comparing the effectiveness of some kind of solvent with a certain ratio. Based on these, one method that was a major preoccupation in this study, for the first time, were multistage extraction of RBO by hexane followed by ethanol, so all the components of polar and non-polar of RBO and primarily long chain fatty acids and short chain contained in RBO can be extracted perfectly. As comparison, it was also presented fatty acid and mineral composition of RBO extracted directly with ethanol. In addition, analysis of mineral composition of RBO by X-ray Spectrometer and melting point of RBO by DSC was also conducted to assess its potential to be developed as raw material of oil-based food processing.

2 EXPERIMENTAL PROCEDURE

Celebes rice bran was obtained by milling rice grain in a local grinding mill Makassar-Indonesia during March to April in 2017. All chemicals and reagents used in the experiment were from Merck, Germany. Standards of Fatty Acid Methyl Ester (FAME) were from Supelco Inc., Bellefonte, PA (Supelco 37 Component FAME Mix) and other reagents from Merck, Germany. Freshly milled bran samples were directly collected from the milling system in polyethylene bags. The rice bran was screened through a 60 mesh sieve to have a uniform particle size and heated at autoclave (Hiclave HV-85 Hirayama) at 100°C for 15 min to inactivate endogenous lipase. At experimental unit 75 grams of rice bran were weighed in the reactor 1.0 L four neck flasks, RBO is extracted with 525 mL solvent using a heating mantle connected with the thermometer setting at 55°C for 5 hours, agitator on the top, speed of 200 rpm, RBO and bran residue are separated by centrifugation (refrigerated AX-521 centrifuge) at a speed of 3500 rpm, 20 min. The liquid part accommodated in the flask evaporator, solvents removed on a rotary evaporator Buchi R-215 incorporates vacuum Pump V-700, speed of 60 rpm, the heating temperature of 35°C, and the evaporation temperature 21°C. RBO obtained was packaged in a dark

glass bottle and stored in a freezer for analysis. The bran residue from hexane extraction subsequently extracted with ethanol.

2.1 Analysis component of RBO

0.05 g of RBO dissolved in methanol-chloroform (1:1) and homogenized by vortex, inserted in the vial GC-MS, the sample was injected automatically. Component of RBO was analyzed using GC-MS ultra Shimadzu QP2010, instrument uses column Rxi SH-5Sil MS (length 30 m, diameter x thickness 0.25mm ID 0.25 um df). Helium was used as a carrier gas at a pressure of 76.9 kps with a flow of 14.0 mL/min, split ratio of 1:10. The initial temperature of the oven starting at 110°C and hold for 2 min until the oven temperature to 200°C with an isothermal increase of 5°C/min, a final temperature of 280°C, hold for 9 min with an isothermal increase of 5°C/min, the total analysis time of 40 min. Scan interval 45-450 m/z. Ion temperature 200°C and detector temperature 280°C. Identification of compounds: the mass spectrum of the compound in the sample was obtained by electron ionization (EI) at 70 eV and the detector operator in scan mode from 45 to 450 m/z. Identification based on molecular weight, molecular formula, retention time and area (%).

2.2 Preparation and analysis of FAME

Fatty acid composition of RBO was verified by gas chromatography (GC-MS ultra Shimadzu QP2010). Fats were esterified by method adapted from Metcalfe [27] which consisted of lipid saponification with KOH 0.2 M in methanol solution, and catalyzed BF₃-MeOH reagent. The sample was solubilized by dichloromethane, from which 1 µL was injected for GC analysis. To separate and quantify the esterified fatty acid mixture, GC-MS QP 2010 by Shimadzu was used equipped with split/splitless injector, capillary column RTX®-1 (30 mx0.25 mmIDx0.25 µm) and flame ionization detector (FID). Helium was used as the carrier gas at flow of 1.25 mL/min. The injector and detector temperature were set to 260°C. The chromatographic conditions for separation were column initial temperature of 50°C, raising to 200°C at a flow rate of 6°C/min, holding during 4 min, the second step consisted in increasing at a heating rate of 2°C/min to 240°C, and held for 10 min. FAME peaks were identified by comparing their retention time and equivalent chain length with respect to standard FAME.

2.3 Mineral and differential scanning calorimeter (DSC) analysis

Mineral analysis by S2 Ranger X-ray Spectrometer. Analysis according the user manual XRF (2012) by Bruker AXS GmbH, Ostliche Rheinbruckenstr, 49.76187 Karlsruhe, Germany. DSC analysis according to the user manual of Shimadzu DSC-60 Plus Series.

3 RESULTS AND DISCUSSION

Rice bran oil has been extracted from Celebes rice by multistage extraction with hexane (first stage) and ethanol (second stage). Analysis by GC-MS showed that some component obtained from first stage were: undecane; octane,2,3,7-trimethyl; tridecane; tetradecane; 9-Oxononanoic acid; hexadecane; tetradecanoic acid; pentadecanone,6,10,14-trimethyl; pentadecanoic acid; hexadecanoic acid, methyl ester; n-hexadecanoic acid;

hexadecanoic acid, 1-methylethyl ester; l-(-)-ascorbic acid 2,6-hexadecanoate; n-hexadecanoic acid; heptadecanoic acid; pregn-5-ene-3,11-dione, 17,20:20,22-bis[methylenebis(oxy)]-, cyclic 3-(1,2-ethanediyol acetal); 9,12-octadecadienoic acid (Z,Z)-, methyl ester; 9-octadecenoic acid, methyl ester; 9-octadecenoic acid (Z,Z); octadecanoic acid, methyl ester; l-(-)-ascorbic acid 2,6-dihexadecanoate; 9,12-Octadecadienoic acid, ethyl ester; ethyl oleate; cis-9-hexadecenal; 4,8,12,16-Tetramethylheptadecan-4-ol; hexadecanal; hexadecanal 2-methyl; tetratriacontane; hexadecanoic acid, 2-hydroxy-1-(hydroxymethyl) ethyl ester; 1,2-benzenedicarboxylic acid; 9,12-Octadecadienoic acid (Z,Z)-, 2,3-dihydroxypropyl ester; 9,12-octadecadienoic acid (Z,Z)-, trimethylsilyl ester; 2-[1,4,6-dioxiranyl]dodecylloxirane; oleoyl chloride; tetracosanoic acid, methyl ester; 9,12-octadecadienoic acid (Z,Z)-, 2,3-dihydroxypropyl ester; cedrane, 8-propoxy; 2,6,10,14,18,22-tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-E); 17-pentatriacontene; tetratetracontane; oryzanol, vitamin E; and 2H-1-benzopyran-6-Ool, 3,4-dihydro-2,7,8-trimethyl-2-(4,8,12,16,20,24,28,32-octamethyl. In second stage obtained some component of RBO were: tetradecane; phenol, 2,4-bis(1,1-dimethylethyl); dodecanoic acid; hexadecane; tetradecanoic acid; butanamine, n-(1-methylpropylidene); octadecane; 2-pentadecanone, 6,10,14-trimethyl; pentadecanoic acid; heneicosane; hexadecanoic acid, methyl ester; n-hexadecanoic acid; 4-ethenylhexahydro-4,7a-dimethyl-17-H)-benzofuranone, [3ara-(3a.alpha.,4.beta.,7a.beta)]; 9,12-octadecadienoic acid (Z,Z)-, methyl ester; 9-octadecenoic acid, methyl ester, (E); octadecanoic acid, methyl ester; 9-octadecenoic acid, (E); heneicosane; 9,12-octadecadienoic acid (Z,Z)-, methyl ester; hexatriacontane; pregn-20-YN-17-ol; 1,2-benzenedicarboxylic acid, oryzanol, ditridecyl ester; 9,12-octadecadienoic acid (Z,Z)-, 2,3-dihydroxypropyl ester; hexadecanoic acid, tetradecyl ester; and hexatriacontane. Table 1 showed that the group of SFA was predominantly of Celebes RBO, although the composition was slightly. Oleic, linoleic, and palmitic acids were found in large numbers. In RBO extracted with hexane, all of fatty acids could be well extracted though the result of second stage extraction with ethanol was still found. RBO extracted with hexane had 18.6% of SFA and 81.4% of unsaturated fatty acids with ratio of SFA : MUFA : PUFA of approximately 1 : 2.3 : 1.3. It contained about 56.7% of MUFA, 24.7% of PUFA, and 18.6% of SFA. Oleic acid which found to be a predominant MUFA in the majority of samples was 56.39%. Concentration of palmitic acid as SFA and linoleic acid as PUFA were 15.51% and 24.74%, respectively.

9

Table 1: Fatty acid composition of Celebes RBO obtained by multistage extraction

Fatty acid	Lipid Numbers	Group	Multistage extraction	
			Stage 1 (Hexane) (mg/L)	Stage 2 (Ethanol) (mg/L)
Palmitic acid	C12:0	SFA	0.002	4.78
Myristic acid	C14:0	SFA	0.002	2.22
Palmitic acid	C16:0	SFA	1021.89	6.34
Pentadecyclic acid	C15:0	SFA	1.73	0.03
Palmitoleic	C16:1	MUFA	17.03	0.03
Stearic acid	C18:0	SFA	125.81	1.81
Oleic acid	C18:1	MUFA	3716.56	0.55
Linoleic acid	C18:2	PUFA	1630.78	3.52
Arachidic acid	C20:0	SFA	47.74	0.03
Behenic acid	C22:0	SFA	17.5	0.05

Tricosylic acid	C23:0	SFA	1.43	0.02
Lignoceric acid	C24:0	SFA	9.97	0.01

9

Table 2: Fatty acid composition of Celebes RBO obtained by directly extraction using ethanol

Fatty acid	Lipid Numbers	Group	Single stage extraction (Ethanol) (mg/L)
Lauric acid	C12:0	SFA	4.78
Myristic acid	C14:0	SFA	2.02
Palmitic acid	C16:0	SFA	0.01
Pentadecyclic acid	C15:0	SFA	0.01
Palmitoleic	C16:1	MUFA	0.08
Stearic acid	C18:0	SFA	2.88
Oleic acid	C18:1	MUFA	0.08
Linoleic acid	C18:2	PUFA	28.85
Arachidic acid	C20:0	SFA	0.04
Behenic acid	C22:0	SFA	0.04
Tricosylic acid	C23:0	SFA	0.01
Lignoceric acid	C24:0	SFA	0.01

In the second stage extracted with ethanol produced the highest levels of linoleic acid (PUFA) amounted to 84.44%. As for the results of direct extraction with ethanol obtained linoleic (28.85 mg/L), stearic (2.88 mg/L) and myristic (2.02 mg/L) (Table 2). Extraction with ethanol could produce the highest levels of PUFA [28]. Ethanol resulted the food-grade extract which was rich in polar lipid and PUFA [29] and food grade hexane extract which rich in neutral lipids [30]. For absorption in the human body and their biological functions, differences in lipid structure were considered as very important [31]. Figure 1 shows the GC-MS chromatogram of fatty acid on RBO extracted with hexane. It can be seen that oleic acid (9-octadecenoic acid) is a fatty acid mostly in RBO (peak number: 7). Oleic peak appearing at retention time of 21.215 with 36.05 %area, area/high (A/H) of 4.99, the area of 157103345. The area indicates the amount of compound concentration, the larger the area, the greater the concentration of these compounds. Can be seen on the enlarged scale that concentration of oleic of 3716.56 mg/L. The next dominant fatty acids were linoleic and palmitic seen in peaks numbered 6 and 4 with retention times of 21.075 and 18.291, respectively.




Fig. 1: GC-MS chromatogram of fatty acid on Celebes RBO

Rice bran contained specific lipids in distinct ratios [32]. RBO has an optimal ratio of SFA, MUFA and PUFA of about 1 : 2.2 : 1.5, which was very close to the recommendation of WHO, RBO contained about 47% of MUFA, 37% PUFA and 20% SFA [33]. Major fatty acid of RBO was 84% linoleic, 33.6% oleic and 20.2% of palmitic acid [34]. Unsaturated fatty acids were about 75% (w/w) in oil; the major fatty acid compositions of RBO were oleic, linoleic and palmitic acids. It

mainly contained oleic (36-38%), linoleic (35-38%) and α -linolenic (from 1.8 to 2.4%) as unsaturated fatty acids. Palmitic (21-44%) and stearic (2.7 to 3.0%) acids were SFA [35], [36], [37]. Recent studies had clearly shown a significant impact of PUFA in human health in disease prevention, particularly cardiovascular disease, coronary heart disease and cancer, inflammatory, thrombotic and autoimmune diseases, hypertension, type two diabetes, kidney disease, rheumatoid arthritis, ulcerative colitis, and Crohn's disease [38], [39]. RBO might assist reducing LDL cholesterol due to its fatty acid profiles and high quantity of unsaponifiable compounds [40]. Relative concentration and distribution of fatty acids in dietary fats had been reported to lower the risk of cardiovascular disease [41]. Diet with increasing the intake of linoleic and linolenic acids increased HDL-cholesterol and LDL-cholesterol lowering, while the intake of higher oleic acid could decrease LDL-cholesterol, but did not affect the HDL cholesterol level [42]. High content PUFA in oil indicate the high nutritional quality of oil. Owing to this high PUFA content, it exhibits properties prevention of thrombosis, inhibition of cardiovascular diseases, reduction of cholesterol in serum, dilation of blood vessels, reduction cancer and regulation of autonomic nerves [43], [44]. Reports the results of the study show the potential of RBO as functional food. Short chain fatty acids generally had lower melting point and more water soluble. Long chain fatty acids had higher melting points. Unsaturated fatty acids had lower melting point compared to SFA of similar chain length. The rate of unsaturated fatty acid to SFA in edible oils and fats was very important for human nutrition. Therefore, high levels of SFA desirable on oil were beneficial to improve the stability. On the other hand, the presence of SFA in the oil was undesirable, because high levels of SFA might increase the concentration of low density lipoprotein (LDL), thus affecting the ratio of LDL to HDL (high density lipoprotein), promoting clothing and vascular smooth muscle proliferation [45]. Edible oils which rich in unsaturated fatty acids, especially MUFA had ability to lower the total and LDL cholesterol, triacylglycerol levels and increased HDL cholesterol that helps control blood pressure. In addition, they had anti thrombotic properties and also improved insulin sensitivity [46]. In rice bran lipids were concentrated approximately 19.4 to 25.5% [7], it was about 15-25%, which contained 95% of triacylglycerols and 4% of unsaponifiable lipids [47]. RBO main lipid fraction composed of neutral lipids, was the unsaponifiable fraction which included antioxidant, but 3 to 5% (w/w) [19], [48]. However, the component and composition of fatty acid in the oil varied depending on the source of oils, and technology process used for oil extraction, but each vegetable oils had a distribution of certain fatty acids depending on the plant source, so the impact on human health could be assessed in accordance with the fatty acid content, because the fatty acids had different effects on human health and the risk of disease that could be caused [49]. In neem oil extraction, ethanol was good substitute for hexane, and able to provide the oil yield 41.11% compared to 42.29% at 50°C when hexane was used at the same temperature. A study proved the solubility of oil in higher ethanol at high temperatures, for example in the extraction of soybean oil. Hence, there was a possibility that more oil could still be extracted from the neem seed using ethanol at temperatures higher than 50°C, and possibly more oil than that extracted using hexane at 50°C, which was about the possible temperature limit of hexane due to its high risk of flammability

[50]. Long chain fatty acid was generally well extracted by hexane, but it also depend on their solubility limit in hexane, so that the short chain fatty acid was few to be obtained from extraction with hexane. The long chain fatty acids were rather non-polar, the polarity would decrease with the increasing of chain. In contrast, short chain fatty acids were more extractable in ethanol, and solubility in ethanol depend on the length of the chain. This might explain why the long chain fatty acids were troublesome extractable by ethanol. The longer the carbon chain, the more difficult to dissolve oils and fats. Unsaturated oils and fats were more soluble in organic solvents than in SFA with the equal length of carbon chain. The fatty acids which had a high level of unsaturated were more soluble than fatty acids with low level of unsaturated [51]. The polarity of molecules depend on: (1) differences in solubility, the greater the difference, the more polar the molecule, and (2) the molecular structure, if the molecule was bound to organic substances, the molecule was more sensitive and the unpolarity increased. On the other hand, the determination of minerals in oils and fats has been the attention of the researchers, as mineral causes deleterious effects on the quality and stability of oils and fats. Minerals such as copper (Cu) and iron (Fe) had been known for fats and oils oxidative activity at low concentrations of 0.005 and 0.05 ppm, respectively. Cu was the strongest pro-oxidant in oil. To the best stability of the oil, Cu content should be below 0.02 ppm [52]. For the best stability of the oil, Fe levels recommended by the Codex STAN 19 [53] should be below 51 ppm and 0.10 - 0.40 ppm for Cu, this would increase stability and edibility of oil and extended the shelf life. Results were shown in Table 2 that the RBO which extracted with ethanol contained Cu over the recommended limit, whereas Fe was detected in the RBO extracted with hexane, the numbers also over the provisions recommended. Mineral generally a central mineral ions in organic molecules. The planes that symmetry which can be done by the molecule molecular symmetry. A molecule will be polar if the ability to separate the charge (ion positive and negative) is greater. Mineral solubility in the solvent depends on the nature of the compound, if these minerals are associated with the non-polar compounds such minerals will dissolve both the non-polar compounds, and vice versa. For example, potassium (K) in Table 3 was not detected in hexane solvent, but was detected in ethanol (Table 4), because K is generally a polar compound in salt form.

Table 3: Mineral composition of the Celebes RBO

Mineral	Multistage extraction	
	Stage 1 (Hexane) (mg/L)	Stage 2 (Ethanol) (mg/L)
K	Nd	96
Cl	Nd	23
P	955	18
Si	13	907
S	Nd	566
Ba	26	35
Fe	17	Nd
Sb	Nd	13
Al	Nd	Nd
Sn	Nd	10
Cu	Nd	20
Zn	Nd	11


Nd, not detected

Table 4: Mineral composition of the Celebes RBO extracted by ethanol

Mineral	Single stage extraction (Ethanol) (mg/L)
K	245
Cl	101
P	25
Si	14
S	826
Ba	35
Sb	14
Al	353
Sn	11
Cu	26
Zn	15

Nd, not detected

In relation to the potential of RBO to be further processed as an oil-based processed feedstock, it is helpful to find data on the melting point and heat capacity of RBO analyzed using DSC. The data shown in Figure 2 RBO melts first at the temperature of 31.08°C, peak melting point of 31.90°C, at this point RBO requires the greatest energy to be able to melt at the perfect temperature of 32.89°C. The data were in accordance with the results shown in Table 1 that the dominant fatty acids constituting RBO were oleic, linoleic and palmitic. Oleic, linoleic and palmitic have a melting point of 13°C, -5°C and 63°C, respectively, so the mixture of them has a melting point around 31-32°C. The data also informs that heat capacity RBO of 0.03 J/g.

**Fig. 2:** DSC analysis of Celebes RBO

4 CONCLUSIONS

The results of this study show that the fatty acid profile of Celebes RBO contains a high percentage of oleic and linoleic acid indicates that this oil is excellent for consumption as one of the best sources of fatty acids in food, and potentially can be used as a feedstock in the production of functional foods. RBO extraction using hexane is better than using ethanol associated with the concentration of fatty acids that can be obtained. Celebes RBO also contains some minerals which its needs special attention related to its effect on the quality and stability of the oil. DSC analysis has shown that RBO has a melting point of 31.90°C with a heat capacity value of 0.03 J/g.

5 ACKNOWLEDGMENTS

We do appreciate and thank to the Department of Chemical Engineering, State Polytechnic of Ujung Pandang for

supporting the laboratory works by providing GC-MS, X-ray Spectrometer instrument and DSC.

REFERENCES

- [1] Sanches, C.E.R.Amante and M.Anto, "Prevention of hydrolytic rancidity in rice bran," vol. 75, (2009), 487-491, DOI:10.1016/j.jfoodeng.2009.06.026.
- [2] V.R.Pestana, R.C.Zambiasi, C.R.B.Mendonça, M.H.Bruscatto, M.J.Lerma-García and G.Ramis-Ramos, "Quality Changes and Tocopherols and γ -Orizanol Concentrations in Rice Bran Oil During the Refining Process," J. Am. Oil Chem. Soc., 85(11), (2008), 1013-1019.
- [3] R.Chervanky, "Bioactives in rice bran and rice bran oil," in Phytochemicals as Bioactive Agents, W.R.Bidlack, S.T.Omaye, M.S.Meskin, D.K.W.Tophan, Eds. Florida, CRC Press, (2000), 213-240.
- [4] A.C.Rustan and C.A.Drevon, "Fatty Acids: Structures and Properties", Encyclopedia of Life Sciences, (2005), 1-7, DOI:10.1038/npg.els.0003894.
- [5] M.Sugano and F.Hirahara, "Polyunsaturated fatty acids in the food chain in Japan", The American Journal of Clinical Nutrition, 71(1 Suppl), (2000), 189S-193S.
- [6] FAO/WHO, "Report of the joint FAO/WHO expert consultation on the risks and benefits of fish consumption", FAO Fisheries and Aquaculture report, (2010).
- [7] L.Liu, D.L.E.Waters, T.J.Rose, "Phospholipids in rice: Significance in grain quality and health benefits: A review", Food Chemistry, 139(1-4), (2013), 1133-1145, DOI:10.1016/j.foodchem.2012.12.046.
- [8] G.C.Burdge and S.A.Wootton, "Conversion of alpha-linolenic acid to eicosapentaenoic, docosapentaenoic and docosahexaenoic acids in young women" The British Journal of Nutrition, 88(4), (2002), 411-20, DOI:10.1079/BJN2002689.
- [9] J.T.Brenna, N.Salem, A.J.Sinclair, " α -Linolenic acid supplementation and conversion to n-3 long-chain polyunsaturated fatty acids in humans, Prostaglandins Leukotrienes and Essential Fatty Acids, 80(2-3), (2009), 85-91, DOI:10.1016/j.plefa.2009.01.004.
- [10] F.Proust, M.Lucas and E.Dewailly, "Fatty acid profiles among the Inuit of Nunavik: Current status and temporal change", Prostaglandins Leukotrienes and Essential Fatty Acids, 90(5), (2014), 159-167, DOI: 10.1016/j.plefa.2014.02.001.
- [11] F.Sahena, I.S.M.Zaidul, S.Jinap, "Application of supercritical CO₂ in lipid extraction - A review", Journal of Food Engineering, Elsevier Ltd, 95(2), (2009), 240-253.
- [12] B.M.C.Soaes, F.M.C.Gamarra, L.C.Paviani, "Solubility of 43 α -glycerols in supercritical carbon dioxide", Journal of Supercritical Fluids, 43(1), (2007), 25-31,

- DOI:10.1016/j.supflu.2007.03.013.
- [13] S.P.J.N.Senayake and F.Shahidi, "Structured lipids: acydolysis of gamma-linolenic acid rich-oils with n-3 polyunsaturated fatty acids", *Journal of food lipids*, 4, (2002), 309-323, DOI:10.1111/j.1745-4522.2002.tb00228.x.
- [14] J.R.Vercellotti, A.J.Stangelo, A.M.Spanier, "Lipid Oxidation in Foods - an Overview", *Acs Symposium Series*, (1992), DOI:10.1021/bk-1992-0500.ch001.
- [15] D.G.Bradley and B.M.David, "Singlet oxygen oxidation of foods", *Critical Review Food Science and Nutrition*, 31 (3) (1992) 211-36.
- [16] R.Z.Zambiasi and M.W.Zambiasi, "Vegetable oil oxidation: effects of endogenous components", *Revista da Sociedade Brasileira de Ciencia e Tecnologia de Alimentos*, Camphinas, 34(1), (2000), 22-32.
- [17] R.Z.Zambiasi, "Oxidation reactions of vegetable oils and fats", *Revista da Sociedade Brasileira de Ciencia e Tecnologia de Alimentos*, Camphinas, 33 (1), (1999), 1-7.
- [18] J.S.Casas, C.Gordillo, M.De, "Characteristics of virgin olive oils from the Olive Zone of Extremadura (Spain), and an approximation to their varietal origin", *J. Am. Oil Chem. Soc.*, 86, (2009), 933-940.
- [19] K.Ichihara and Y.Fukubayashi, "Preparation of fatty acid methyl esters for gas-liquid chromatography". *The Journal of Lipid Research*, 51(3), (2010), 635-640, DOI: 10.1194/jlr.D001065.
- [20] J.Gromadzka and W.Wardencki, "Trends in Edible Vegetable Oils Analysis", Part A. Determination of Different Components of Edible Oils-a Review, 61(1), (2011), 33-43.
- [21] M.G.Megahed, E.H.A.Nashy, E.A.Al-Ashkar, "Evaluation of fried edible oil and determination of trace elements content by FAAS", *Agriculture and Biology Journal of North America*, 2(4), (2011), 687-692, DOI:10.5251/abjna.2011.2.4.687.692.
- [22] E.Choe and D.B.Min, "Comprehensive Reviews in Food Science and Food Safety Mechanisms and Factors for Edible Oil Oxidation". *Comprehensive Reviews in Food Science and Food Safety*, 5, (2006), 169-186, DOI:10.1111/j.1541-4337.2006.00009.x.
- [23] R.M.De Souza, B.M.Mathias, C.L.P.Da Silveira, "Inductively coupled plasma optical emission spectrometry for trace multi-element determination in vegetable oils, margarine and butter after stabilization with propan-1-ol and water", *Spectrochimica Acta - Part B Atomic Spectroscopy*, 60(5), (2005), 711-715, DOI:10.1016/j.sab.2005.02.025.
- [24] B.Lu, Y.Zhang, X.Wu, "Separation and determination of diversiform phytosterols in food materials using supercritical carbon dioxide extraction and ultraperformance liquid chromatography-atmospheric pressure chemical ionization-mass spectrometry", *Analytica Chimica Acta*, 588(1), (2007), 50-63, DOI:10.1016/j.aca.2007.01.067.
- [25] V.F.Péres, J.Saffi, M.I.S.Melecchi, "Comparison of soxhlet, ultrasound-assisted and pressurized liquid extraction of terpenes, fatty acids and Vitamin E from *Piper gaudichaudianum* Kunth", *Journal of Chromatography A*, 1105 (1-2 SPEC. ISS), (2006), 115-118, DOI:10.1016/j.chroma.2005.07.113.
- [26] D.K.Saxena, S.K.Sharma and S.S.Sambi, "Comparative Extraction of Cottonseed Oil by n-Hexane and Ethanol", *ARNP Journal of Engineering and Applied Sciences*, 6(1), (2011), 84-89.
- [27] Metcalfe, "The rapid preparation of fatty acid esters for gas chromatographic analysis", *Analytical chemistry*, 33(3), (1961), 363-364, DOI:10.1021/ac60171a016.
- [28] M.Schmid, F.Guihéneuf, and D.B.Stengel, "Evaluation of food grade solvents for lipid extraction and impact of storage temperature on fatty acid composition of edible seaweeds *Laminaria digitata* (Phaeophyceae) and *Palmaria palmata* (Rhodophyta)", *J. Food Chemistry*, 208, (2016), 161-168, DOI:10.1016/j.foodchem.2016.03.123.
- [29] Y.Li, N.F.Ghasemi and S.Garg, "A comparative study: the impact of different lipid extraction methods on current microalgal lipid research", *Microbial cell factories*, 13(14), (2014), 1-9, DOI:10.1186/1475-2859-13-14.
- [30] J.V.Ambrozova, "Influence of extractive solvents on lipid and fatty acids content of edible freshwater algal and seaweed products, the green microalga *Chlorella kessleri* and the cyanobacterium *Spirulina platensis*", *Molecules*, 19(2) (2014), 2344-2360, DOI: 10.3390/molecules19022344.
- [31] M.C.Michalski, C.Genot, C.Gayet, "Multiscale structures of lipids in foods as parameters affecting fatty acid bioavailability and lipid metabolism", *Progress in Lipid Research*, Elsevier Ltd, 52(4), (2013), 354-373, DOI:10.1016/j.plipres.2013.04.004.
- [32] C.V.Rao, Y.Hirose, C.Indranie, "Modulation of Experimental Colon Tumorigenesis by Types and Amounts of Dietary Fatty Acids 1", *Cancer Research*, 61(5), (2001), 1927-1933.
- [33] Y.H.Hui, "Bailey's industrial oil and fat products - edible oil and fat products", Vol. 2, A Wiley- Interscience Publication, New York, (1999), 603-75.
- [34] K.C.Massarolo, "Influence of cultivation *Rhizopus oryzae* on rice bran oil lipid fraction: Fatty acids and phospholipids". *Biocatalysis and Agricultural Biotechnology*, 8, (2016), 204-208, DOI:10.1016/j.cbab.2016.10.002.
- [35] I.S.Kahlon, E.I.Chow, M.M.Chiu, "Cholesterol-lowering by

- rice bran ³⁴ rice bran oil unsaponifiable matter in hamsters". *Cereal Chemistry*, 73(1), (1996), 69-74.
- [36] A.G.G.Krishna, S.Khatoon and R.Babylatha, "Frying performance of processed rice bran oil" *Journal of Food Lipids*, 12(1), (2005), 1-11, DOI:10.1111/j.1745-4522.2005.00001.x.
- [37] V.S.Rubalya, P.A.Sahayaraj and A.A.Prema, "Antioxidant stability in palm and rice bran oil using simple parameters", *Rasayan Journal of Chemistry*, 3(1), (2010), 44-55.
- [38] R.D.Caterina, J.K.Liao and P.Libby, "Fatty acid modulation of endothelial activation 1-3", *American Journal of Clinical Nutrition*, 71, (2000), 213S-223S.
- [39] E.Abedi and M.A.Sahari, "Long-chain polyunsaturated fatty acid sources and evaluation of their nutritional and functional properties", *Food Science and Nutrition*, 2(5): (2014) 443-463, DOI:10.1002/fsn3.121.
- [40] I.A.Wilson, L.M.Ausman and C.W.Lawton, "Comparative cholesterol lowering properties of vegetable oils: beyond fatty acids", *Journal of the American College of Nutrition*, 19(5), (2000), 601-607, DOI:10.1080/07315724.2000.10718957.
- [41] L.Cordain, S.B.Eaton and J.B.Miller, "The paradoxical nature of hunter-gatherer diets: Meat-based, yet non-¹⁰⁰rogenic", *Eur. J. Clin. Nutr.*, 56, (2002), 542-552, DOI:10.1038/sj=ejcn=1601353.
- [42] C.L.Lawton, H.J.Delargy, J.Brockman, "The degree of saturation of fatty acids influences post-ingestive satiety", *Br J Nutr*, 83(5), (2000), 473-482, DOI:10.1017/S000711450000060X.
- [43] J.M.Luque-Rodríguez, M.D.L.De Castro, P.Pérez-Juan, "Extraction of fatty acids from grape seed by superheated ⁶³ane", *Talanta*, 68(1) (2005), 126-130, DOI:10.1016/j.talanta.2005.04.054.
- [44] C.Yi, J.Shi, J.Kramer, "Fatty acid composition and phenolic antioxidants of winemaking pomace ⁵⁷nder", *Food Chemistry*. Elsevier Ltd, 114(2), (2009), 570-576, DOI:10.1016/j.foodchem.2008.09.103.
- [45] D.Dzisiak, "New oil ¹⁰²duced saturated and trans fats in processed foods", *Cereal Foods World*, 49 (6), (2004), 331-333.
- [46] V.S.Rubalya and P.Neelamegam, "Antioxidant potential in vegetable oil", *Research Journal of Chemistry and Environment*, 16(2), (2012), 87-94.
- [47] X.Wang, J.Lu, H.Liu, "Improved deacidification of high-acid rice bran ⁹⁸ oil by enzymatic esterification with phytos ⁷⁹pl", *Process Biochemistry*. 51(10), (2016), 1496-1502, DOI:10.1016/j.procbio.2016.08.013.
- [48] M.M.Strieder, C.P.Pinheiro, V.S.Borba, "Bleaching optimization and winterization step evaluation in the refinement of rice bran oil", *Separation and Purification Technology*, 175, (2017), 72-78, DOI:10.1016/j.seppur.2016.11.026.
- [49] J.Orsavova, L.Misurcova, J.V.Ambrozova, "Fatty Acids Composition of Vegetable Oils and Its Contribution to Dietary Energy Intake and Dependence of Cardiovascular Mortality on Dietary Intake of Fatty Acids", *International Journal of Molecular Sciences*, 16(6), (2015), 12871-12890, DOI:10.3390/ijms160612871.
- [50] M.Y.Liauw, F.A.Natan and P.Widiyanti, "Extraction of neem oil (*Azadirachta indica* A. Juss) using n-hexane and ethanol: Studies of oil quality, kinetic and thermodynamic", *Journal of Engineering and Applied Sciences*, 3(3), (2008), 49-54.
- [51] S.Ketaren, "Edible oils and fats", 1st Ed. Jakarta: Universitas Indonesia press. ISBN: 979-8034-05-8, (1986).
- [52] F.Anwar, T.G.Kazi and R.Saleem, "Rapid determination of ¹⁶ne trace metals in several oils and fats", *Grasas Aceites*, 55(2), (2004), 160-168, DOI: 10.3989/gya.2004.v55.i2.162.
- [53] Codex Alimentarius Commission, "Fats, oils and related products", Second Edition. Revised 2001, Joint FAO/WHO Food Standard Programme, (1981).

Component, Fatty Acid And Mineral Composition Of Rice Bran Oil Extracted By Multistage With Hexane And Ethanol

ORIGINALITY REPORT

%29
SIMILARITY INDEX

%21
INTERNET SOURCES

%23
PUBLICATIONS

%11
STUDENT PAPERS

PRIMARY SOURCES

1 www.mdpi.com Internet Source **%2**

2 Kelly Cristina Massarolo, Taiana Denardi de Souza, Anelise Christ Ribeiro, Eliana Badiale Furlong et al. "Influence of cultivation Rhizopus oryzae on rice bran on lipid fraction: Fatty acids and phospholipids", Biocatalysis and Agricultural Biotechnology, 2016
Publication **%2**

3 ejournalofsciences.org Internet Source **%1**

4 journal-of-agroalimentary.ro Internet Source **%1**

5 www.researchgate.net Internet Source **%1**

6 www.pgia.ac.lk Internet Source **%1**

7 Submitted to Federal University of Technology

% 1

8

fenbilder.sdu.edu.tr

Internet Source

% 1

9

Schmid, Matthias, Freddy Guihéneuf, and Dagmar B. Stengel. "Evaluation of food grade solvents for lipid extraction and impact of storage temperature on fatty acid composition of edible seaweeds *Laminaria digitata* (Phaeophyceae) and *Palmaria palmata* (Rhodophyta)", *Food Chemistry*, 2016.

Publication

% 1

10

Krzysztof Różyło, Sylwia Andruszczak, Ewa Kwiecińska-Poppe, Renata Różyło, Piotr Kraska. "Effect of Three Years' Application of Biogas Digestate and Mineral Waste to Soil on Phytochemical Quality of Rapeseed", *Polish Journal of Environmental Studies*, 2018

Publication

% 1

11

ajpp.in

Internet Source

<% 1

12

krishikosh.egranth.ac.in

Internet Source

<% 1

13

revistas.unal.edu.co

Internet Source

<% 1

14

Asmiaty Sahur, Ambo Ala, Baharuddin Patandjengi, Elkawakib Syam'un. "Effect of Seed Inoculation with Actinomycetes and Rhizobium Isolated from Indigenous Soybean and Rhizosphere on Nitrogen Fixation, Growth, and Yield of Soybean", International Journal of Agronomy, 2018

Publication

<% 1

15

Bligh, E. G., and W. J. Dyer. "A RAPID METHOD OF TOTAL LIPID EXTRACTION AND PURIFICATION", Canadian Journal of Biochemistry and Physiology, 1959.

Publication

<% 1

16

Ni, Zhanglin, Fubin Tang, Yihua Liu, Danyu Shen, and Runhong Mo. "Multielemental Analysis of Camellia Oil by Microwave Dry Ashing and Inductively Coupled Plasma Mass Spectrometry", Analytical Letters, 2015.

Publication

<% 1

17

Piccirillo, A.. "Comparison between off-line and on-line derivatisation methods in the characterisation of siccative oils in paint media", Journal of Analytical and Applied Pyrolysis, 200508

Publication

<% 1

18

Submitted to Universiti Teknologi MARA

Student Paper

<% 1

19

Chunyang Li, Hongying Xia, Libo Zhang, Shixing Wang, Jinhui Peng, Song Cheng, Jianhua Shu, Xin Jiang, Qi Zhang. "Analysis of dielectric characterization and microwave adsorbing properties in organism-contained spent carbon: An efficient regeneration method via microwave-assisted ultrasound", Chemical Engineering and Processing - Process Intensification, 2018

Publication

<% 1

20

scholarbank.nus.edu.sg

Internet Source

<% 1

21

M Díaz. "Physico-chemical characteristics of carcass and meat Manchego-breed suckling lambs slaughtered at different weights", Meat Science, 2003

Publication

<% 1

22

Submitted to North West University

Student Paper

<% 1

23

Jimenez-Cordero, Diana, Francisco Heras, Noelia Alonso-Morales, Miguel A. Gilarranz, and Juan J. Rodriguez. "Porous structure and morphology of granular chars from flash and conventional pyrolysis of grape seeds", Biomass and Bioenergy, 2013.

Publication

<% 1

Dan Wang, Wen Xue Chen, Yue Ying Hu, Mei

- 24 Fang Peng. "Optimization, Modeling and Antibacterial Activity of Extraction from Fructus Alpinia Oxyphylla", Advanced Materials Research, 2013
Publication <% 1
-
- 25 Sharan, D.. "The effect of cetylated fatty esters and physical therapy on myofascial pain syndrome of the neck", Journal of Bodywork & Movement Therapies, 201107
Publication <% 1
-
- 26 Lemus, Christelle, Apostolis Angelis, Maria Halabalaki, and Alexios Leandros Skaltsounis. "γ-Oryzanol", Wheat and Rice in Disease Prevention and Health, 2014.
Publication <% 1
-
- 27 fedorabg.bg.ac.rs
Internet Source <% 1
-
- 28 www.frontiersin.org
Internet Source <% 1
-
- 29 Thanonkaew, A.. "Effect of stabilization of rice bran by domestic heating on mechanical extraction yield, quality, and antioxidant properties of cold-pressed rice bran oil (Oryza sativa L.)", LWT - Food Science and Technology, 201210
Publication <% 1
-

30

Internet Source

<% 1

31

Pohndorf, R.S., T.R.S. Cadaval, and L.A.A. Pinto. "Kinetics and thermodynamics adsorption of carotenoids and chlorophylls in rice bran oil bleaching", Journal of Food Engineering, 2016.

Publication

<% 1

32

César A. Can-Cauich, Enrique Sauri-Duch, David Betancur-Ancona, Luis Chel-Guerrero et al. "Tropical fruit peel powders as functional ingredients: Evaluation of their bioactive compounds and antioxidant activity", Journal of Functional Foods, 2017

Publication

<% 1

33

Laurence S. Lim, Gus Gazzard, Yen-Ling Low, Robin Choo, Donald T.H. Tan, Louis Tong, Tien Yin Wong, Seang-Mei Saw. "Dietary Factors, Myopia, and Axial Dimensions in Children", Ophthalmology, 2010

Publication

<% 1

34

etd.lsu.edu

Internet Source

<% 1

35

www.tandfonline.com

Internet Source

<% 1

36

Inakã S. Barreto, Stefani I.E. Andrade, Francisco A.S. Cunha, Marcelo B. Lima, Mario

<% 1

Cesar U. Araujo, Luciano F. Almeida. "A robotic magnetic nanoparticle solid phase extraction system coupled to flow-batch analyzer and GFAAS for determination of trace cadmium in edible oils without external pretreatment", *Talanta*, 2018

Publication

37

ilamph.ir

Internet Source

<% 1

38

SeungOk Yang, JaeMin Lee, JaeChul Lee, JaeHwan Lee. "Effects of riboflavin-photosensitization on the formation of volatiles in linoleic acid model systems with sodium azide or D2O", *Food Chemistry*, 2007

Publication

39

Shiddiq Yunus, A. M., A. Abu-Siada, and M. A. S. Masoum. "Application of SMES Unit to Improve DFIG Power Dispatch and Dynamic Performance During Intermittent Misfire and Fire-Through Faults", *IEEE Transactions on Applied Superconductivity*, 2013.

Publication

40

www.gottswaal.nl

Internet Source

<% 1

41

Submitted to University of Newcastle upon Tyne

Student Paper

<% 1

42 doaj.org Internet Source <% 1

43 Submitted to South Dakota Board of Regents Student Paper <% 1

44 Orsavova, Jana, Ladislava Misurcova, Jarmila Ambrozova, Robert Vicha, and Jiri Mlcek. "Fatty Acids Composition of Vegetable Oils and Its Contribution to Dietary Energy Intake and Dependence of Cardiovascular Mortality on Dietary Intake of Fatty Acids", International Journal of Molecular Sciences, 2015. Publication <% 1

45 www.stanceequine.com.au Internet Source <% 1

46 www.chemie.uni-hamburg.de Internet Source <% 1

47 Submitted to Taylor's Education Group Student Paper <% 1

48 www.venuslabs.org Internet Source <% 1

49 rasayanjournal.co.in Internet Source <% 1

50 www.iasj.net Internet Source <% 1

repositorio.ufpa.br

51

Internet Source

<% 1

52

www.academicjournals.org

Internet Source

<% 1

53

www.ncbi.nlm.nih.gov

Internet Source

<% 1

54

www.cdpr.ca.gov

Internet Source

<% 1

55

opus.uni-hohenheim.de

Internet Source

<% 1

56

Soares, B.M.C.. "Solubility of triacylglycerols in supercritical carbon dioxide", *The Journal of Supercritical Fluids*, 200711

Publication

<% 1

57

file.scirp.org

Internet Source

<% 1

58

Freitas, Lisiane dos Santos, Claudio Dariva, Rosângela Assis Jacques, and Elina Bastos Caramão. "Effect of experimental parameters in the pressurized liquid extraction of brazilian grape seed oil", *Separation and Purification Technology*, 2013.

Publication

<% 1

59

Submitted to University of Brighton

Student Paper

<% 1

60

A.K. Azad, M.G. Rasul, M.M.K. Khan, S.C. Sharma. "Biodiesel From Queensland Bush Nut (*Macadamia integrifolia*)", Elsevier BV, 2017

Publication

<% 1

61

Carvalho, R. H. R., E. L. Galvão, J. A., C. Barros, M. M. Conceição, and E. M. B. D. Sousa. "Extraction, fatty acid profile and antioxidant activity of sesame extract (*Sesamum Indicum L.*)", Brazilian Journal of Chemical Engineering, 2012.

Publication

<% 1

62

Pradosh Prasad Chakrabarti, Ram Chandra Reddy Jala. "Processing Technology of Rice Bran Oil", Elsevier BV, 2019

Publication

<% 1

63

Luque-Rodriguez, J.M.. "Extraction of fatty acids from grape seed by superheated hexane", Talanta, 20051115

Publication

<% 1

64

atvb.ahajournals.org

Internet Source

<% 1

65

Shipra Shukla, Satisha Hegde, Anil Kumar, Gaurav Chaudhary, Shri Krishna Tewari, Dalip Kumar Upreti, Mahesh Pal. "Fatty acid composition and antibacterial potential of *Cassia tora* (leaves and stem) collected from different geographic areas of India", Journal of

<% 1

Food and Drug Analysis, 2018

Publication

66

A.G. GOPALA KRISHNA. "FRYING PERFORMANCE OF PROCESSED RICE BRAN OILS", Journal of Food Lipids, 3/2005

Publication

<% 1

67

Kiunga, G.A.. "Elevated insulin receptor protein expression in experimentally induced colonic tumors", Cancer Letters, 20040810

Publication

<% 1

68

Laurie-Eve Rioux, Lucie Beaulieu, Sylvie L. Turgeon. "Seaweeds: A traditional ingredients for new gastronomic sensation", Food Hydrocolloids, 2017

Publication

<% 1

69

brage.bibsys.no

Internet Source

<% 1

70

Park, Heung Soon, Sungyeul Kim, Youngkyun Lee, Myung Sun Choi, and Myung-Un Choi. "Alteration of lipid composition of rat thymus during thymic atrophy by whole-body X-irradiation", International Journal of Radiation Biology, 2006.

Publication

<% 1

71

proj3.sinica.edu.tw

Internet Source

<% 1

72

"CONCENTRATES : SCIENCE TECHNOLOGY",
Chemical & Engineering News, 1982.

Publication

<% 1

73

www.rjpbcs.com

Internet Source

<% 1

74

Sarin, R.. "Terminalia belerica Roxb. seed oil: A
potential biodiesel resource", Bioresource
Technology, 201002

Publication

<% 1

75

aip.scitation.org

Internet Source

<% 1

76

Karimi, Mehdi, Shayessteh Dadfarnia, Ali
Mohammad Haji Shabani, Fatemeh Tamaddon,
and Davood Azadi. "Deep eutectic liquid
organic salt as a new solvent for liquid-phase
microextraction and its application in ligandless
extraction and preconcentration of lead and
cadmium in edible oils", Talanta, 2015.

Publication

<% 1

77

Marc WÄltermann, Alexander SteinbÄ¼chel. "
In vitro effects of sterculic acid on lipid
biosynthesis in strain PD630 and isolation of
mutants defective in fatty acid desaturation ",
FEMS Microbiology Letters, 2000

Publication

<% 1

78

Rohman, Abdul. "Rice Bran Oil's Role in Health

and Cooking", Wheat and Rice in Disease Prevention and Health, 2014.

Publication

<% 1

79

Xiaosan Wang, Jiyuan Lu, Hong Liu, Qingzhe Jin, Xingguo Wang. "Improved deacidification of high-acid rice bran oil by enzymatic esterification with phytosterol", Process Biochemistry, 2016

Publication

<% 1

80

Cristina Perez-Tenero, Maria Alvarez de Sotomayor, Maria Dolores Herrera.

"Contribution of ferulic acid, γ -oryzanol and tocotrienols to the cardiometabolic protective effects of rice bran", Journal of Functional Foods, 2017

Publication

<% 1

81

Vercellotti, J. R., Allen J. St. Angelo, and Arthur M. Spanier. "Lipid Oxidation in Foods", ACS Symposium Series, 1992.

Publication

<% 1

82

www.ijcrr.com

Internet Source

<% 1

83

scholarspace.manoa.hawaii.edu

Internet Source

<% 1

84

Monique M. Strieder, Cláudio P. Pinheiro, Verônica S. Borba, Ricardo S. Pohndorf, Tito R.S. Cadaval, Luiz A.A. Pinto. "Bleaching

<% 1

optimization and winterization step evaluation
in the refinement of rice bran oil", Separation
and Purification Technology, 2017

Publication

85 www.ajol.info <% 1
Internet Source

86 paperzz.com <% 1
Internet Source

87 www.i-scholar.in <% 1
Internet Source

88 ricebranoilthailand.blogspot.com <% 1
Internet Source

89 www.stu.edu.vn <% 1
Internet Source

90 digitalcommons.brockport.edu <% 1
Internet Source

91 Lu, Weidong, Zhongming Wang, and Zhenhong Yuan. "Characteristics of lipid extraction from Chlorella sp. cultivated in outdoor raceway ponds with mixture of ethyl acetate and ethanol for biodiesel production", Bioresource Technology, 2015.
Publication

92 Submitted to CSU, Pomona <% 1
Student Paper

93

Marina S. Nogueira, Bianca Scolaro, Ginger L. Milne, Inar A. Castro. "Oxidation products from omega-3 and omega-6 fatty acids during a simulated shelf life of edible oils", LWT, 2019

Publication

<% 1

94

fr.readkong.com

Internet Source

<% 1

95

Karel Ventura. "Comparison of Modern Extraction Techniques in Analysis of Soil Contaminated with Fuel and Crude Oil Spills", Journal of Liquid Chromatography & Related Technologies, 2003

Publication

<% 1

96

Suzana Ferreira-Dias, Natália M. Osório, Joana Rodrigues, Carla Tecelão. "Structured Lipids for Foods", Elsevier BV, 2018

Publication

<% 1

97

Vânia Maria Borges Cunha, Marcilene Paiva da Silva, Sérgio Henrique Brabo de Sousa, Priscila do Nascimento Bezerra et al. "Bacaba-de-leque (Oenocarpus distichus Mart.) oil extraction using supercritical CO₂ and bioactive compounds determination in the residual pulp", The Journal of Supercritical Fluids, 2019

Publication

<% 1

98

Yuanrong Jiang. "Bioprocessing Technology of Rice Bran Oil", Elsevier BV, 2019

<% 1

99 Sun Xiaoyang, Tian Shaojun, Zhang Lifen, Xie Jianchun. " Effect of Phospholipase A - Catalyzed Degumming on Oryzanol, Tocopherols, and Tocotrienols of Dewaxed Rice Bran Oil ", Journal of Chemistry, 2019 <% 1
Publication

100 Marinangeli, Christopher P. F., and Peter J. H. Jones. "Deconstructing the Paleolithic Diet: Components that Reduce Cardiovascular Disease Risk", Current Nutrition Reports, 2014. <% 1
Publication

101 H. Hadiyanto, Nais Pinta Adetya. "Response surface optimization of lipid and protein extractions from Spirulina platensis using ultrasound assisted osmotic shock method", Food Science and Biotechnology, 2018 <% 1
Publication

102 documents.mx <% 1
Internet Source
