

# 43\_RJPTJournalScopusQ3\_15\_1\_ 2022.pdf

*by* Rahmawati Minhajat

---

**Submission date:** 18-Jul-2022 03:55PM (UTC+0700)

**Submission ID:** 1872084108

**File name:** 43\_RJPTJournalScopusQ3\_15\_1\_2022.pdf (256.32K)

**Word count:** 5594

**Character count:** 31500

9

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/358137733>

# Qualitative Phytochemical Screening and Effectiveness Analysis of *Batissa violacea celebensis* Martens 1897 Crude extract against Antioxidant and Cytotoxic Activity

Article in *Research Journal of Pharmacy and Technology* · January 2022

DOI: 10.52711/0974-360X.2022.00043

CITATIONS

0

READS

89

6 authors, including:



Sri Anggarini Rasyid

Universitas Mandala Waluya Kendari

16 PUBLICATIONS 8 CITATIONS

[SEE PROFILE](#)



Rahmawati Minhajat

Universitas Hasanuddin

38 PUBLICATIONS 184 CITATIONS

[SEE PROFILE](#)



Ridwan Adi Surya

Universitas Haluoleo

31 PUBLICATIONS 16 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



15

Anti-angiogenic targeting therapy of colorectal cancer: focus on target non-VEGF antiangiogenesis [View project](#)



23

Community service [View project](#)

**RESEARCH ARTICLE**

## Qualitative Phytochemical Screening and Effectiveness Analysis of *Batissa violacea celebensis* Martens 1897 Crude extract against Antioxidant and Cytotoxic Activity

Sri Anggarini Rasyid<sup>1,2\*</sup>, Maria Bintang<sup>3</sup>, Suryani As'ad<sup>4</sup>, Upik Miskad<sup>5</sup>, Rahmawati Minhajat<sup>6</sup>,  
Ridwan Adi Sya<sup>7</sup>

<sup>1</sup>Doctoral Program, Graduate School of Medicine, Faculty of Medicine,  
Hasanuddin University, Makassar, Indonesia.

<sup>2</sup>D-IV Medical Laboratory Technology Study Program Mandala Waluya Health Science Coll<sup>21</sup>

<sup>3</sup>Department of Biochemistry, Faculty of Mathematics and Natural Sciences, IPB University, Bogor, Indonesia.

<sup>4</sup>Department of Nutrition, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia.

<sup>5</sup>Department of Pathology Anatomy, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia.

<sup>6</sup>Department of Internal Medicine, Division of Hematology Medical Oncology,  
Faculty of Medicine, Hasanuddin University, Makassar, Indonesia.

<sup>7</sup>Department of Environmental Science, Faculty of Forestry and Environmental Science Halu Oleo University  
Kendari South Sulawesi, Indonesia.

\*Corresponding Author E-mail: [anggarini.09@gmail.com](mailto:anggarini.09@gmail.com)

### ABSTRACT:

The aim of this study was to conduct phytochemical tests aimed at identifying the bioactive components present in the powder crude extract of pokea shell can be applied in analyzing antioxidant activity and testing cytotoxic activity. This study consists of phytochemical tests, including alkaloids, steroids, triterpenoids, saponins, and flavonoids. The antioxidant activity measured by the DPPH method from the crude extract of the pokea shell is still relatively weak but the crude extract of methanol of the boiled and dried pokea shells in toxic category so potentially as a cytotoxic compound. The result of phytochemical test of crude extract of pokea shell meat, is extract of fresh and boiled pokea shell contain flavonoid, saponin, steroid and triterpenoid while extract of dried pokea shell only contain saponin, steroid and triterpenoid. The result of the antioxidant activity test of crude extract of pokea shell shows that the crude extract of dried pokea shell has the highest value of antioxidant activity. The result of the toxicity test of crude extract pokea shell boiled and dried pokea shell have an LC50 value of less than 1000 ppm. These values indicate that the extract of the boiled and dried Pokea shell is showed potential for development as an alternative anticancer agents in the future.

**KEYWORDS:** Phytochemical, Antioxidant, Cytotoxicity, Pokea shell, Crude extract.

### INTRODUCTION:

The pokea (*Batissa violacea celebensis* Martens 1897) is a endemic bivalve species of the Corbiculidae Family found in the Pohara River of Konawe Regency, Southeast Sulawesi<sup>1</sup>. *Batissa violacea* is a freshwater mollusk spread in Southeast Asia and Northern Australia. In Indonesia, the animals are spread in Sumatra, Java, West Papua, and Sulawesi. Particularly in Southeast Sulawesi, the pokea shell *B. violacea celebensis* martens 1897 is found in the Pohara River at a depth of about 1-9 m at the edge and middle of the river with sand sediments and gravel. Pohara River is a

watershed of Konaweha which is the source of raw water belonging to PDAM (Local water company) of Kendari City Southeast Sulawesi Province Indonesia. Pokea shells are found in strong current conditions with sandy substrate textures, pokea are commonly found with the clustered spreading type<sup>2</sup>.

*Batissa violacea celebensis* is one type of clam that lives in freshwater. In general, bivalves or better known as shellfish is a group of animals with no vertebrae. Pokea shell is classified by Kingdom: Animalia, Phylum: Mollusca, Class: Bivalvia, Order: Eulamellibranchia, Family: Corbiculidae, Genus: *Batissa*, Species: *Batissa violacea celebensis* (Martens, 1897), and area name: Pokea Shells. Pokea shell is a type of bivalves that lives on the bottom of the water and has two shells that can open and close. The shell on the dorsal part is thick, and the ventral part is thin. The top of the eggplant is called the umbo and is the oldest part of the shell. A circular line around the umbo shows the growth of the shell. The mantle in the pelecypoda is shaped as a thin, wide tissue, covering the whole body located under the shell<sup>3</sup>.

Pokea shells are empirically believed to treat various diseases such as jaundice, malaria, asthma, lowering blood pressure and fever. The disease occurs due to infection by foreign materials and microorganisms<sup>1</sup>.

When the body is infected with microorganisms, the body responds with macrophage and neutrophil activity mechanisms. In this case, the oxidase and oxygenase enzymes will form various free radical compounds and reactive oxygen compounds, including hypochlorous acid (HOCl) that will attack and destroy viruses and bacteria. Radical compounds are also very dangerous because of the potential to attack the body's cells. If this is not controlled, it will trigger the emergence of various chronic diseases<sup>1</sup>.

Natural products isolated from bivalves and gastropods have been used, among others, as antioxidants, antitumors, antiviral, antibacterial, antifungal, anticancer, cytotoxic, and enzyme inhibitors<sup>4,5,6</sup>. Some secondary metabolites of aquatic organisms exhibit pharmacological activity<sup>7</sup>. Marine organisms are potential in terms of their ability to produce secondary metabolites which can be utilized as lead component in drug discovery. The search for marine derived products started 50 years ago and over the past few decades, about 3000 new compounds from various marine sources have been tested and determined so far. Metabolites from ascidians, sponges, mollusks, and sea hare, have been described as potent anti-cancer agent. Among the marine organisms, Ascidians rank second with most promising source of drugs. Ascidians are being utilized as food in various countries like Japan,

China. Many anticancer compounds entered clinical trail from ascidian (ET-743, Apilidine, Kahalalide F). It is possible to isolate new molecules from marine organisms that could contribute to a better health, serve as food ingredient and also a new source of new medicine<sup>8</sup>. Research on bioactive components in mollusks, especially bivalves and gastropods that have the potential to be nutraceutical or pharmaceutical has been widely practiced.

In modern life we are constantly exposed free radicals, which can damage any cell in the body with which they come into contact and cause hazardous effects. The free radicals cause a state of increased oxidative stress which is mainly responsible for all the damage done to the body by free radicals. Though the body has its own mechanism to combat oxidative stress, when exposure is more than body's capacity, the problem aggravates. The harmful effects of free radicals can be prevented by use of substances called antioxidants. Antioxidants are a group of compounds that neutralise or prevent free radicals or reactive species and avoid cell or tissue damage caused by oxidative potential of free radicals. Many natural substances have been explored for their antioxidant activities. There is still need to evaluate new substances for their antioxidant potential to find better alternative sources for treatment of conditions related to oxidative stress caused by free radicals. DPPH and Hydrogen Peroxide free radical scavenging methods were used for evaluating antioxidant activities. Ascorbic acid was used as standard antioxidant substance for comparative evaluation<sup>9</sup>.

The antioxidant compounds in the diet plays an important role as a health protective factor<sup>10</sup>. Scientific evidence suggests that antioxidants reduce the risk of chronic diseases including cancer and heart disease. The main feature of antioxidants is the ability to trap free radicals. Free radicals are molecules having an unpaired electron in the outer orbit. They are not stable molecules and ready to react. Free radicals play an important role in living systems. The beneficial effects of free radicals occur at low or moderate concentrations and those levels are balanced by an efficient antioxidant network in the body<sup>11</sup>.

Many synthetic compounds like butylated hydroxytoluene (BHT) and butylated hydroxy anisole (BHA) are commercially obtainable as antioxidants. However, they are not suggested for usage due to the toxicity associated with them. Therefore, many research groups have driven efforts to evaluate the antioxidant properties of natural products<sup>12</sup>. The same was reported that Antioxidant compounds may function as free radical scavengers, complexes of pro-oxidant metals, reducing agents and quenchers of singlet oxygen formation.

4 Synthetic antioxidants, such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA), have restricted use in foods as they are suspected to be carcinogenic. Therefore, the importance of search for natural antioxidants has greatly increased in the recent years and researchers focus on natural antioxidants<sup>13</sup>. Antioxidant compounds, phenolic acids, polyphenols and flavonoids capture free radicals that inhibit oxidative mechanisms that are the cause of degenerative diseases.

## MATERIAL AND METHODS:

### Sampling and sample preparation:

This research phase starts from sampling and sample preparation with materials and tools preparation for nutrient content testing and active compound extraction. The extraction method that used in this study is the stratified extraction or maceration method<sup>14</sup> was performed using the methods previously described<sup>1</sup>

### Mineral analysis (SNI 01-2891-1992):

Mineral analysis was carried out using the Atomic Absorption Spectrophotometer (SAA) to determine the levels of minerals present in food, was performed using the methods previously described<sup>1</sup>.

### Phytochemical analysis<sup>15</sup>

Phytochemical analysis was performed to determine the bioactive components found on pokea shells crude extract powder.

### Alkaloid:

5 Several examples were dissolved in a few drops of 2 N sulfuric acid then tested with three alkaloid reagents, that is dragendrof, meyer, and wagner reagents. The test results are positive if the meyer reagent formed yellowish white sediment, with a wagner reagent forming a brown sediment, and with a dragendrof reagent forming a red to orange sediment. The meyer reagent was prepared by adding 1.36HgCl with 0.5g of potassium iodide then dissolved and diluted with distilled water to 100mL in a measure's flask. This reagent is colorless. The wagner reagent was prepared using 10mL distilled aquades and then added 2.5g of iodine and 2g of potassium iodide, then dissolved and diluted with aquades to 200mL in a flask. This reagent is brown. Dragendrof reagents are prepared by 0.8g of bismuth subnitrate added with 10mL acetic acid and 40 mL of water. This solution is mixed with a solution made from 8g of potassium iodide in 20mL water. Prior to use, 1mL of this mixed volume was diluted with 2.3 mixed volumes of 20mL glacial acetic acid and 100mL of water. The reagent is orange.

### Steroid<sup>5</sup>terpenoid:

Several samples were dissolved in 2ml of chloroform in the test tube. 10 drops Acetic anhydride and 3 drops concentrated sulfuric acid were added to the mixture. The positive test results of samples containing steroid and triterpenoid are the formation of red, blue, and green solutions.

### Saponin: <sup>16</sup>

Saponins can be detected by foam test in hot water. A stable foam for 30 minutes and not lost on the addition of 1 drop of HCl 2 N shows a sample containing saponins.

### Flavonoid:

22 Several samples plus 0.1mg magnesium powder and 0.4 mL of amyl alcohol (37% hydrochloric acid mixture and 95% ethanol with equal volume) and 4mL of alcohol. then mixed were shaken. The result of 25 sample test positive contains flavonoids, which is the formation of red, yellow or orange on amyl alcohol layer.

### Analysis of antioxidant activity<sup>16</sup>

Extract of Fresh Pokea Shell, extract of boiled Pokea shell and extract of dried Pokea shell, extraction using methanol solvent p.a. (polar) is dissolved in methanol p.a. with concentrations of 0, 250, 500, 1000, 2000 and 4000 ppm. BHT Synthetic antioxidants used as positive control, made by BHT dissolved in methanol solvent p.a. with concentrations of 0, 0,3125, 0,625, 1,25, 2,5, 5, 10 and 20ppm. The DPPH solution to be used was prepared by dissolving the DPPH crystals in a methanol solvent with a concentration of 1mM. The process of making a 1mM DPPH solution is carried out under low temperature conditions and protected from sunlight. The extract solution and the BHT comparative antioxidant solution were each taken 4.5mL and treated with 500µL of 1mM DPPH solution in different and labeled reaction tubes. The mixture was then incubated at 37°C for 30 minutes and measured its absorbance using a spectrophotometer at 517nm wavelength. The absorbance of the blank solution was also measured to perform percent inhibition calculations. The blank solution was prepared by reacting 4.5mL of a methanol solvent with 500µL of 1mM DPPH solution in the test tube. After that, the antioxidant activity of each sample and antioxidant comparative BHT/Vitamin C.

### Cytotoxicity assay<sup>17</sup>

The BSLT method is usually performed in a preliminary test for pharmacological screening activity on natural products. This test used *Artemia salina naupli* (Golden West Supreme Plus Great Salt Lake, USA) as an animal test. At first *A. salina* eggs dripped in seawater under 40 watt TL lamps for 48 hours.

A total of 10 larvae *A. salina* were inserted into the test tube, then incorporated an extract of the sample solution with varying concentrations and added artificial seawater to the 5 mL volume. Artificial seawater without gill extract (0 ppm) is used as a control. All test tubes were incubated at room temperature for 24 hours under a 40 watt TL lamp. Observations were made after 24 hours by counting the amount of *A. salina* that died at each concentration. The pricing of LC<sub>50</sub> (ppm) is performed using probit analysis and regression equation.

### RESULTS AND DISCUSSION:

The results of the analysis of mineral content in fresh, boiled, and dried Pokea clam meat can be seen in Table 1.

**Table 1. Mineral content (ppm dry weight) fresh, boiled, and dried Pokea clam meat**

Parameter	Boil	Fresh	dried
Mineral Makro :			
Fosfor (P)	-	2.336,6289	3.217,5233
Kalsium (Ca)	2.321,1247	-	1.456,5214
Magnesium (Mg)	688,256	-	661,8654
Kalium (K)	910,5868	-	1.712,5860
Mineral Mikro :			
Besi (Fe)	-	1.703,1455	4.622,6855
Seng (Zn)	-	62,521	142,1247
Selenium (Se)	-	-	-

Macro minerals in fresh, boiled, and dried Pokea shells are dominated by the minerals phosphorus, calcium, and potassium. Micro minerals are dominated by iron minerals. The content of heavy metals in dried pokea shell meat has been reported<sup>18</sup>. The concentration of heavy metal in dried pokea shell was still below the limit of AAS detection device used was Pb <0.01 ppm; Cd <0.001ppm; and Hg<0.0002 ppm. According to BPOM RI (2009) and SNI 7387 (2009), that maximum Pb in food is 1.5ppm; Cd = 1.0ppm<sup>1</sup>.

Phytochemical tests were performed to detect the bioactive components found in pokea shell extract from different solvents of polarity Tests with this phytochemical method can detect bioactive components not limited to secondary metabolites itself, but also to primary metabolites that provide functional biological activity, namely proteins and peptides<sup>19</sup>. The result of phytochemical test of crude extract of pokea shell meat can be seen in Table 2.

Various biological effects of saponins are antibacterial, antiprotozoal and anticancer activities. Saponins are secondary metabolites of glycosidic nature widely distributed in higher plants. Saponins occur constitutively in a great many plants species, in both wild plants and cultivated crops. In cultivated crops the triterpenoid saponins are generally predominant, while steroid saponins are common in plants used as herbs or for their health-promoting properties<sup>20</sup>.

**Table 2. Phytochemical test results of crude extract of pokea shell meat**

Phytochemical Test	Extract of Fresh Pokea Shell	Extract of Boiled Pokea Shell	Extract of Dried Pokea Shell
Flavanoid	+	+	-
Alkaloid	-	-	-
Tanin	-	-	-
Saponin	+	+	+
Quinon	-	-	-
Steroid	+	+	+
Triterpenoid	+	+	+

Legume contain 5-13% triterpenoid saponins. Many researchers have shown the relationship between legume consumption and health benefits, such as protection from cardiovascular disease, breast cancer, colon cancer, other cancers, diabetes, anti-inflammatory, antiarthritis, anti-oxidant and laxatives effect<sup>20</sup>.

The different types of bioactive components extracted are, presumably due to differences in polarity of each solvent, materials and chemical compounds will dissolve readily on relatively<sup>21</sup>.The bioactive components present in a crude extract, such as alkaloids, steroids, saponins, phenols, free amino acids, and carbohydrates are thought to act as antioxidants<sup>22</sup>.

Flavonoid compounds are antioxidants and can inhibit the action of xanthine oxidase enzymes and superoxide reactions<sup>23</sup>. The extracts which have higher content of phenol also indicated that they have better antioxidant activities. So, antioxidative activity may be correlated with total phenolic contents of the extracts. Flavonoids are one of the most important natural phenols. They also possess several biological and chemical activities and are having free radical scavenging properties too. Its antioxidant property depends primarily on their hydroxyl group position in the molecule and their ability as electron donor to a free radical<sup>24</sup>.

Flavonoid compounds from *Stereospermum personatum* in addition to antioxidants, these compounds can also inhibit the action of xanthine oxidase enzyme.

Flavonoids and triterpenoids are known to have important roles in cancer chemoprevention and chemotherapy<sup>25</sup>. These compounds have antioxidant activity, with the capacity to capture reactive oxygen species (ROS), inhibit enzymes involved in the formation of ROS, and block the oxidation of cellular and extracellular compounds<sup>26,27,28</sup>. In addition, flavonoids play a role in the modulation pathway of cancer proliferation, cell cycle, inducing apoptosis, and inhibiting angiogenesis<sup>29</sup>.

Several flavonoid compounds which have bonds with sugar groups so that flavonoids are easily dissolved in water or polar-containing solvents such as ethanol, methanol, butanol, acetone, dimethylsulfoxide, dimethylformamide, and air<sup>30</sup>.

Flavonoids also have anticancer potential with the mechanism of inhibition of the abnormal tyrosine kinase enzyme. The tyrosine kinase enzyme is an enzyme that plays a role in cell transduction signals, namely by regulating the cell's continued cycle, transcription regulation, cell transformation, proliferation, differentiation, and apoptosis. Oncogenes will alter protein kinase activity and result in the formation of leukemia tumor types<sup>31,32</sup>. The same was reported that Antioxidant compounds include vitamins, carotenoids, flavonoids and phenolics. Among them, phenolics and flavonoids are the most important and exhibit substantial antioxidant activity<sup>33</sup>.

Triterpenoids are carbon skeletons composed of 6 isoprene units and made biosynthetic from squalene (C<sub>30</sub> acyclic hydrocarbons). Steroids are triterpene components found in the form of glycosides. Natural triterpenoids have antitumor activity. Terpenoids have antioxidant capacity functions but are not followed by cancer cell antiproliferation functions<sup>34</sup>.

Steroid is a class of triterpenoid compounds that can be classified into steroids with no more than 21 carbon atoms namely sterols, sapogenins, cardiac glycosides, and vitamin D. Natural steroids are derived from various chemical transformations of two triterpenes namely lanosterol and cycloartenol. Steroid compounds can be used as the basic ingredients for the manufacture of drugs<sup>15</sup>. There are several steroids such as fukosterol that are isolated from non toxic marine biological resources and have the characteristic of lowering cholesterol in the blood and promoting antidiabetic activity<sup>35</sup>.

Based on the standard curve of absorbance spectrum, DPPH solution showed maximum absorption at 517 nm wavelengths using UV-VIS spectrophotometer Hitachi U-2800, so that sample absorbance measurements are performed at those wavelengths. The antioxidant activity of crude extracts from pokea shells can be seen in Table 3. Antioxidant are the molecules that can effectively neutralize the free radicals. They safely interact with free radicals and terminate the chain reaction before vital molecules get damaged. Phenolics, flavonoids, vitamins (E and C), numerous minerals (Cu, Mn, Zn, Se and Fe), glutathione are some of the most widely recognised antioxidants. Antioxidants in blood, cells, and tissue fluids play an important role in neutralizing

the normal level of oxidative damage caused by these free radicals<sup>36</sup>.

**Table 3. Result of antioxidant activity test of crude extract of pokea shell**

Parameter	Extract of Fresh Pokea Shell	Extract of Boiled Pokea Shell	Extract of Dried Pokea Shell
Antioxidant IC <sub>50</sub>	2483.74 ppm	3338.61 ppm	>4000 ppm

Table 3 shows that the crude extract of dried pokea shell has the highest value of antioxidant activity. The IC<sub>50</sub> value of the crude extract of the pokea shell is still higher than the IC<sub>50</sub> value of vitamin C used as the standard of 6.02ppm. IC<sub>50</sub> value is one of the commonly used parameters to interpret the results of DPPH testing. This IC<sub>50</sub> value can be defined as a substrate concentration which can lead to a 50% reduction in DPPH activity. The smaller IC<sub>50</sub> value means the higher antioxidant activity<sup>16</sup>. The antioxidant activity of crude extract of pokea shells measured by DPPH method is still very weak. This is presumably because the DPPH method only measures antioxidant compounds with mechanisms capable of donating their hydrogen atom to radical compounds. DPPH is a constant free radical at room temperature and accepts an electron or hydrogen radical to develop into a stable diamagnetic molecule. The decrease in absorbance of DPPH radical caused by antioxidants, because of the reaction between antioxidant molecules and radical, progresses, which consequences in the scavenging of the radical by hydrogen donation<sup>37</sup>. Lipid peroxidation has gained more importance nowadays because of its involvement in the pathogenesis of many diseases like atherosclerosis, cancer, diabetes mellitus, myocardial infarction, immunological incompetence, neurodegenerative disorders and also in aging<sup>37</sup>.

The antioxidant mechanism is not limited to the ability of a compound to provide a hydrogen atom, but also in its ability to inhibit the formation of reactive oxygen compounds by metal chelating, bypassing the chain-oxidation reaction of free radicals or by capturing them<sup>38</sup>.

Antioxidant activity measurements can be performed with several methods and all methods provide a picture of different mechanisms, so to obtain an adequate assessment of the antioxidant capability in a biological system a sample is required the use of more than one method<sup>39</sup>.

Toxicity test by Brine Shrimp Lethality Test (BSLT) method is pre screening against active compounds contained in a sample. Extract is considered highly toxic when having an LC<sub>50</sub> value below 30 ppm, considered

toxic when it has a LC<sub>50</sub> value of 30-1000 ppm and is considered non-toxic when the LC<sub>50</sub> value is above 1000 ppm<sup>40</sup>.

Table 4. Result of toxicity test of crude extract of pokea shell

Parameter	Extract of Fresh Pokea Shell	Extract of Boiled Pokea Shell	Extract of Dried Pokea Shell
Toxicity LC <sub>50</sub>	>1000 ppm	962.93 ppm	780.81 ppm

The LC<sub>50</sub> value is the concentration that can cause death of 50% of population *A. salina* used. The value of LC<sub>50</sub> can be seen from the table above shows that extract of boiled and dried pokea shell have LC<sub>50</sub> value less than 1000 ppm. These values indicate that the extract of the boiled and dried Pokea shell is included in the toxic category. This is suspected because in the crude extract type of methanol solvent there is a cytotoxic bioactive component. This Extract dried Pokea showed the potential for development as an alternative anticancer agent.

The types of cytotoxic substances commonly present in crude extracts include tannins, flavonoids, triterpenoids and coumarins. This is consistent with the phytochemical results of the crude extract of ethyl acetate pokea shell containing flavonoids and steroids/triterpenoids. The presence of flavonoid and steroid compounds in rough extract of pokea from all solvents is allegedly different, both the number and the type of cytotoxic compounds so as to have a different impact on shrimp larvae<sup>41</sup>.

Fraction is potentially studied as a candidate compound with cytotoxic activity if it has a relatively small LC<sub>50</sub> value or less than 1000ppm. The LC<sub>50</sub> value of ethyl acetate extract of pokea shell is smaller than the macroalgae extract of *Turbunaria decurrens* type<sup>42</sup>.

The difference in the level of secondary metabolites in extracts is proportional to the level of cytotoxicity. The study supports the results of this study that the cytotoxic activity of extract of pokea depends on their phytochemical content<sup>43</sup>.

The death of shrimp larvae was ascertained due to administration of extracts caused by the absence of shrimp larvae death in the negative control group. Ethyl acetate fraction and water fraction have LC<sub>50</sub> values greater than 1000µg/ml. Air extract, ethanol extract, and n-hexane fraction have LC<sub>50</sub> values smaller than 1000µg/ml<sup>44</sup>.

## CONCLUSION:

This study shows the results of phytochemical tests which include steroids, triterpenoids, saponins, and flavonoids. The antioxidant activity of the crude pokea

shell extract is still relatively weak, and the crude methanol extract of boiled and dry pokea shell is in the toxic category so that it can be used as a cytotoxic compound. Extract purification studies should be carried out. In addition, further work is needed to resolve the specific agent anticancer activity of cancer cell lines and immunostimulant.

## ACKNOWLEDGEMENT:

The authors are grateful to the Ministry of Research, Technology and Higher Education, and we thank our colleagues from IPB University who provided insight and expertise that greatly assisted in this research. We are grateful to all of those with whom we have had the pleasure to work during this and other related projects.

## CONFLICT OF INTEREST:

The authors declare no conflict of interest.

## REFERENCES:

1. Rasyid, Sri Anggarini, Maria Bintang, Bambang P. Priosoeryanto, Ratna Umi Nurlila, dan Ridwan Adi Surya. 2018. Analysis chemical compound of pokea (*Batissa violacea celebensis* martens 1897) the origin of konawe regency Southeast Sulawesi. *Indian Journal of Public Health Research and Development*. <https://doi.org/10.5958/0976-5506.2018.00576.4>.
2. Bahtiar., 2005 Kajian Populasi Pokea (*Batissa violacea celebensis* Martens, 1897) di Sungai Pohara Kendari Sulawesi Tenggara [tesis]. Bogor: Program Pascasarjana, Institut Pertanian Bogor
3. Bahtiar, Hamzah, dan Hari, H. 2015, Studi Struktur dan Pertumbuhan Populasi Kerang Pokea (*Batissa violacea var. celebensis*, von Martens 1897) di Sungai Pohara Sulawesi Tenggara, *Jurnal Biologi Tropis*, Juli 2015: Volume 15 (2): 110-120 ISSN: 1411-9587.
4. Tadesse M, Gulliksen B, Strim MB, Styrvoid OB, Haug T., 2008., Screening for antibacterial and antifungal activities in marine benthic invertebrates from northern Norway. *J Invertebr Pathol* 99:286-293
5. Defer, D., N. Bourgougnon, and Y. Fleury. 2009. Screening for antibacterial and antiviral activities in three bivalve and two gastropod marine molluscs. *J. Aquaculture*, 293 : 1-7
6. Zhou, D.Y., B.W.Zhu, L. Qiao, H.T. Wu, D.M. Li, J.F. Yang, and Y. Murata. 2011. In vitro antioxidant activity of enzymatic hydrolysates prepared from abalone (*Haliotis discus hannai* ino) viscera. *Food and Bioproducts Processing*, in press
7. Pringgienies, D., 2010., Karakteristik senyawa bioaktif bakteri simbiosis moluska dengan GC-MS. *J. Ilmu dan Teknologi Kelautan Tropis*, 2(2): 34-40
8. Natarajan K., Ramanathan S., Sathish R.,C. Amudhadevi. In vitro Cytotoxic Study on crude extract of Marine Invertebrata Animal *Polyclinum madrasensis* Sebastian. *Asian J. Pharm. Tech.* 1(4): Oct – dec 2011; Page 152-154
9. Mrunali Potbhare, Deepak Khobragade. In Vitro Evaluation of Antioxidant Potential of Ayurvedic Preparations Lauha Bhasma and Mandura Bhasma. *Asian J. Pharm. Res.* 2017; 7(2): 63-66. doi: 10.5958/2231-5691.2017.00011.9
10. Prakash A., Rigelhof F., Miller E., Antioxidant Activity. *Medallion Laboratories Analytical Progress*. <http://www.medallionlabs.com>
11. Krishna Kondragunta. V, Karuppuraj. V, Perumal. K. Antioxidant activity and Folic acid content in indigenous isolates of *Ganoderma lucidum*. *Asian J. Pharm. Ana.* 2016;6(4): 213-215
12. Priyanga Rangasamy, Vadakkenchery Salimudheen Hansiya, Palanisamy Uma Maheswari, Thamburaj Suman, Natesan Geetha. *Phytochemical Analysis and Evaluation of in vitro Antioxidant*

- and anti-urolithiatic Potential of Various Fractions of *Clitoria ternatea* L. Blue Flowered Leaves. *Asian J. Pharm. Ana.* 2019;9(2):67-76. Doi : 10.5958/2231-5675.2019.00014.0
13. Jiby Elias, Rajesh M.G., Anish N.P., Manu M.S., Iwin C. Varkey. *Terminalia chebula* Retz. Stem Bark Extract: A Potent Natural Antioxidant. *Asian J. Research Chem.* 4(3): March 2011; Page 445-449.
  14. Rasyid S A, Sugireng, Ridwan Adi Surya, Sanatang, Rosdarni, Wa Ode Rejeki Natalia. The antibacterial activity of Tembelekan leaf (*Lantana camara* L.) and Kopasanda leaf (*Chromolaena odorata* L.) extracts against *Staphylococcus aureus*. *Infectious Disease Reports* 2020; 12(s1):8734 doi:10.4081/idr.2020.8734
  15. Harborne, J.B. 1987. *Metode Fitokimia*. Edisi kedua. Padmawinata K, Soediro L, penerjemah. Bandung. ITB. Terjemahan dari: *Phytochemical Methods*.
  16. Molyneux P., 2004., The use of the stable free radicals diphenylpicrylhydrazyl (DPPH) for estimating antioxidant activity. *Songklanakarin J Sci Technol* 26:211-219
  17. Carballo JL, Hernandez-Inda ZL, Perez P, Garcia-Gravalos MD., 2002. A comparison between two brine shrimp assay to detect in vitro cytotoxicity in marine natural product (methodology article). *Bioorganic Mar Chem Biotech* 2(1):1-5
  18. Yenni. 2012. Pengaruh Perebusan Terhadap Kandungan Gizi Kerang Pokea (*Batissa Violacea Celebensis* Martens 1897) Dan Aktivitas Antioksidannya. [tesis]. Bogor: Program Studi Teknologi Hasil Perairan, Fakultas Pertanian, Institut Pertanian Bogor.
  19. Kannan M., and Singh R., 2009., *Phytochemistry and immunopharmacological investigation of Rubia cordifolia* Linn (Rubiaceae), *Pharmacology online*, 3: 653-662
  20. Anjali Soni, Patel Femida, Preeti Sharma. In vitro Cytotoxic Activity of Plant Saponin Extracts on Breast Cancer Cell Line. *Res. J. Pharmacognosy and Phytochem.* 2017;9(1): 17-22. Doi ; 10.5958/0975-4385.2017.00003.6
  21. Khopkar SM., 2003. *Basic concept analytical Chemistry*. Jakarta: UI-Press
  22. Nurjanah., Laili Izzati dan Asadatul Abdullah., 2011. Aktivitas Antioksidan dan Komponen Bioaktif Kerang Pisau (*Solen spp*). *J. Ilmu Kelautan. Vol. 16 (3) 119-124*
  23. Cos P, Ying L, Calomme M, Hu JP, Cimanga K, Poel BV, Pieters L, Vlietinck AJ, Berghe DV., 1998. Structure activity relationship and classification of flavonoids as inhibitors of xanthine oxidase and superoxide scavengers. *J Nat Prod* 61:71-76
  24. Amit Roy, Pushpa Prasad. Study of Physicochemical, Preliminary Phytochemical and Antioxidant activity of *Calocybe indica*. *Research J. Pharm. and Tech.* 9(1): Jan., 2016; Page 53-59. doi: 10.5958/0974-360X.2016.00010.X
  25. Kumar US, Tiwari AK, Reddy SV, Apama P, Rao RJ, Ali AZ, Rao JM., 2006., Free radical scavenging and xanthine oksidase inhibitory constituents from *Stereospermum personatum*. *J Nat Prod* 68:1611-1621
  26. Hertzog DI, Tica OS. 2012. Molecular mechanisms underlying the anti- cancerous action of flavonoids. *Current Health Sci J.* 38(4), pp. 145-149.
  27. Kumar PS, Febriyanti RM, Sofyan FF, Luftimas DE, Abdulah R. 2014. Anticancer potential of *Syzygium aromaticum* L. in MCF-7 human breast cancer cell lines. *Pharmacognosy Res.* 6: 350-354.
  28. Han N, Bakovic M. 2015. Biologically active triterpenoids and their cardioprotective and anti-inflammatory effects. *J Bioanalysis Biomed.* vol. S12, article 005, pp. 1-11.
  29. Katyal P, Bhardwaj N, Khajuria R. 2014. Flavonoids and their therapeutic potential as anticancer agents; biosynthesis, metabolism and regulation. *World J Pharmacy Pharm Sci.* 3(6), pp. 2188-2216
  30. Mierziak, J. Kostyn, K., and Kulma, A. 2014. Flavanoid as important Molecules of Plant Interactions with the Environment. *Mol. Basel Switz.* 19, 16240-16265.
  31. Hui LQ. 2009. Total Phenolic Content and Total Flavonoid of *H. polyrhizus* Waste Extract by Using Ultrasonic Solvents Extraction. Faculty of Chemical Engineering and Natural Resources. Project Paper. Malaysia: Universiti Malaysia Pahang.
  32. Genin M, Clement F, Fattaccioli A, Raes M, Michiels C. 2015. M1 and M2 macrophages derived from THP-1 cells differentially modulate the response of cancer cells to etoposide. *BMC Cancer.* 15: 577.
  33. Preeti Tiwari. Phenolics and Flavanoids and Antioxidant Potential of Balaristha Prepared by Traditional and Modern Methods. *Asian J. Pharm. Ana.* 4(1): Jan-Mar. 2014; Page 05-10
  34. Firdiyani F, Agustini TW, Ma'ruf WF. 2015. Extraction of bioactive compounds as natural antioxidants from fresh spirulina platensis using different solvents. *J Pengolahan Hasil Perikanan Indonesia.* 18(1): 23-37
  35. Bhakuni DS and Awat DS., 2005 *Bioactive marine natural Product*. New Delhi: Anamaya Publisher 315
  36. Abhishek Jain, Varsha Ojha, Gaurav Kumar, Loganathan Karthik, Kokati Venkata, Bhaskara Rao. *Phytochemical Composition and Antioxidant Activity of Methanolic Extract of Ficus benjamina (Moraceae) Leaves.* *Research J. Pharm. And Tech.* 6(11): November 2013; Page 1184-1189
  37. Yogesh Shivhare, Priya Singh, K.K. Pate, Prashant Soni. Preliminary Phytochemical Screening and Antioxidant Activity of Petroleum Ether Extract of *Trichosanthes dioica* Roxb (Fruits). *Research J. Pharmacognosy and Phytochemistry* 2010; 2(2):155-158
  38. Winarsi H., 2007., *Antioksidan Alami dan Radikal Bebas*. Yogyakarta: Kanisius
  39. Bohm V., 2010. Determination of hydrophilic and lipophilic antioxidant capacity- comments and results [abstrak]. On the: Workshop Antioxidant Measurement Assay Methods; Istanbul, Turkey 21 Apr 2010. hlm 14. Abstr no PL03
  40. Meyer BN, Ferrigni NR, Putnam JE, Jacobsen LB, Nicholas DE, McLaughlin JL., 1982., Brine shrimp: a convenient general bioassay for active plant constituents. *Planta Medica* 45(3):31-34
  41. Peteros NP, and Uy MM., 2010., Antioxidant and cytotoxic activities and phytochemical screening of four Philippine medicinal plants. *Journal of Medicinal Plants Research* 4(5):407-414
  42. Pisutthanan, S., Plianbangchang, P., Pisutthanan, N., Ruanruay, S., Muanrit, O., 2004., Brine shrimp lethality activity of Thai medicinal plants in the family Meliaceae. *Naresuan Uni J.* 12, 13-18.
  43. Handayani D, Rasyid W, Rustini, Zainudin EN, Hertiani T. 2018. Cytotoxic activity screening of fungal extracts derived from the west sumatran marine sponge *Haliclona fascigera* to several human cell lines: Hela, WiDr, T47D and Vero. *J App Pharm Sci.* 8 (01): 055-058
  44. Puspita R, Bintang M, Priosoeryanto BP. Possible antiproliferative activity of longan (*Dimocarpus longan* Lour.) leaf extracts. *J Appl Pharm Sci.* 2019; 9(05):102-106.

ORIGINALITY REPORT

---

**23%**  
SIMILARITY INDEX

**20%**  
INTERNET SOURCES

**17%**  
PUBLICATIONS

**12%**  
STUDENT PAPERS

---

PRIMARY SOURCES

---

**1** [storage.googleapis.com](https://storage.googleapis.com) Internet Source **1%**

---

**2** [www.sciencepub.net](http://www.sciencepub.net) Internet Source **1%**

---

**3** [text-id.123dok.com](http://text-id.123dok.com) Internet Source **1%**

---

**4** [www.ijbio.com](http://www.ijbio.com) Internet Source **1%**

---

**5** [files.aiscience.org](http://files.aiscience.org) Internet Source **1%**

---

**6** [www.acarindex.com](http://www.acarindex.com) Internet Source **1%**

---

**7** Gaurav Kumar, Loganathan Karthik, Kokati Vankata Bhaskara Rao. "Phytochemical composition and in vitro antioxidant activity of aqueous extract of *Aerva lanata* (L.) Juss. ex Schult. Stem (Amaranthaceae)", Asian Pacific Journal of Tropical Medicine, 2013  
Publication **1%**

---

8	ijrap.net Internet Source	1 %
9	repository.politeknikmfh.ac.id Internet Source	1 %
10	academicjournals.org Internet Source	1 %
11	www.ijptonline.com Internet Source	1 %
12	www.pagepress.org Internet Source	1 %
13	Submitted to University of Portsmouth Student Paper	1 %
14	Submitted to iGroup Student Paper	1 %
15	www.researchgate.net Internet Source	1 %
16	Submitted to International Islamic University Malaysia Student Paper	1 %
17	Mirwa Adiprahara Anggarani, Erlix Rakhmad Purnama, Jihan Shofwatul Islam Dalilah Aziz. "Antioxidant Potential of the Shell of Razor Clams (Solen spp) in Antidiabetic Mellitus Type II Therapy", Journal of Physics: Conference Series, 2021	1 %

---

18	<a href="mailto:scialert.net">mail.scialert.net</a> Internet Source	1 %
19	<a href="http://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet Source	1 %
20	<a href="http://repository.unhas.ac.id">repository.unhas.ac.id</a> Internet Source	1 %
21	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet Source	1 %
22	Submitted to Universitas Brawijaya Student Paper	<1 %
23	<a href="http://www.coursehero.com">www.coursehero.com</a> Internet Source	<1 %
24	<a href="http://www.ijppsjournal.com">www.ijppsjournal.com</a> Internet Source	<1 %
25	Abdul Hapid, Mery Napitupulu, Muhammad Sulaiman Zubair. "Ethnopharmacology and Antioxidant Activity Studies of Woody Liana Original Wallacea", International Journal of Design & Nature and Ecodynamics, 2021 Publication	<1 %
26	<a href="http://www.mongoliajol.info">www.mongoliajol.info</a> Internet Source	<1 %
27	<a href="http://ijsrm.humanjournals.com">ijsrm.humanjournals.com</a> Internet Source	<1 %

---

28

[ayujournal.org](http://ayujournal.org)

Internet Source

&lt;1 %

29

[www.open.edu](http://www.open.edu)

Internet Source

&lt;1 %

30

Laurentius Hartanto Nugroho, Anna Estyaniyana. "The potency of gadung (*Dioscorea hispida* Dennst.) tuber as a functional food: Toxicity, phytochemical content and starch characters", AIP Publishing, 2018

Publication

&lt;1 %

31

[Repository.Unej.Ac.Id](http://Repository.Unej.Ac.Id)

Internet Source

&lt;1 %

32

[repositori.uin-alauddin.ac.id](http://repositori.uin-alauddin.ac.id)

Internet Source

&lt;1 %

33

Nyi Mekar Saptarini, Irma Erika Herawati. "Comparative Antioxidant Activity on The *Ficus benjamina* and *Annona reticulata* Leaves", International Journal of Public Health Science (IJPHS), 2015

Publication

&lt;1 %

34

[devileye.net](http://devileye.net)

Internet Source

&lt;1 %

35

[faculty.uobasrah.edu.iq](http://faculty.uobasrah.edu.iq)

Internet Source

&lt;1 %

[www.mdpi.com](http://www.mdpi.com)

36

Internet Source

&lt;1 %

37

Ajay Kumar Mahapatra, Sweta Smita Pani, Atish Kumar Sahoo. " Free radical-scavenging activities of species – An endangered medicinal plant of Eastern Ghats of India ", Natural Product Research, 2014

Publication

&lt;1 %

38

[www.i-scholar.in](http://www.i-scholar.in)

Internet Source

&lt;1 %

39

[doczz.net](http://doczz.net)

Internet Source

&lt;1 %

40

[www.neliti.com](http://www.neliti.com)

Internet Source

&lt;1 %

41

"Abstracts of the Asian Congress of Nutrition 2019", Annals of Nutrition and Metabolism, 2019

Publication

&lt;1 %

42

"Marine Niche: Applications in Pharmaceutical Sciences", Springer Science and Business Media LLC, 2020

Publication

&lt;1 %

43

Wilis A. Setyati, Rini Pramesti, Muhamad Zainuddin, Maya Puspita, Person P Renta. " Cytotoxicity and Phytochemical Profiling of Extract As Anti-Mdr Bacteria ", IOP

&lt;1 %

# Conference Series: Earth and Environmental Science, 2018

Publication

---

44	<a href="http://ppm-poltekkeskemenkesbanjarmasin.com">ppm-poltekkeskemenkesbanjarmasin.com</a> Internet Source	<1 %
45	<a href="http://repositori.usu.ac.id">repositori.usu.ac.id</a> Internet Source	<1 %
46	<a href="http://www.phytojournal.com">www.phytojournal.com</a> Internet Source	<1 %
47	<a href="http://www.gssrr.org">www.gssrr.org</a> Internet Source	<1 %

---

Exclude quotes  On

Exclude matches  < 5 words

Exclude bibliography  On