

Bladder stones associated with vesicovaginal fistula-TURNITIN

by Asykar Palinrungi

Submission date: 14-Jan-2022 12:41AM (UTC+0000)

Submission ID: 1741401442

File name: stones_associated_with_vesicovaginal_fistula-September_2020.pdf (1.07M)

Word count: 2529

Character count: 13752



ELSEVIER

Contents lists available at ScienceDirect

International Journal of Surgery Case Reports

journal homepage: www.casereports.com

Bladder stones associated with vesicovaginal fistula: A case report

Francisca^a, Khoirul Kholis^b, Muhammad Asykar Palinrungi^{b,*}, Syakri Syahrir^b, Syarif^b, Muhammad Faruk^a^a Department of Surgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia^b Division of Urology, Department of Surgery, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

ARTICLE INFO

Article history:

Received 11 July 2020

Received in revised form 13 August 2020

Accepted 5 September 2020

Available online 11 September 2020

Keywords:

Bladder stone

Vesicovaginal fistula

Vesicovaginal fistula repair

Cystolithotomy

Case report

ABSTRACT

INTRODUCTION: The presentation of bladder stones is the most common manifestation of a lower urinary tract stone. In severe cases, it can cause pressure necrosis and lead to vesico-vaginal fistula (VVF). This case reports a female patient diagnosed with a bladder stone accompanied by VVF.

PRESENTATION OF CASE: A 56-year-old female came to the hospital with complaints of dysuria accompanied by continual urinary incontinence symptoms dating back about a month. During vaginal examination, we found a fistula and a stone. An ultrasound (US) showed a hyperechoic shadow measuring 1.27 × 1.36 cm in the vesica urinary and plain kidney, ureter, and bladder (KUB) x-ray showed the presence of a radiopaque shadow of about 7.5 × 5 cm in the pelvic area. This stone was removed through an open cystolithotomy. Intraoperatively, a fistula of about 1 cm in diameter was found in the trigonum close to the internal meatus. A fistula repair was performed immediately without complications, and the follow-up results were satisfactory.

CONCLUSION: The presentation of a bladder stone is an uncommon condition that can cause VVF. Therefore, it is important to pay attention to its clinical signs.

© 2020 The Author(s). Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Bladder stones are the most common manifestation of lower urinary tract stones, accounting for 5% of urinary stones and nearly 1.5% of all patients in the urology hospitalizations [1]. A vesico-vaginal fistula (VVF) is an abnormal communication between the bladder and the vagina that results in the unknown leakage of urine [2]. This condition is the most common type of urogenital fistula [1].

VVF resulting from giant bladder stones is uncommon; the bladder stone may put pressure on the bladder wall, causing tissue necrosis and eventually VVF [3]. Thus, this study reports the first case of bladder stone causing VVF in an elderly female in Indonesia. We reported the case in accordance with the SCARE 2018 guidelines [4].

2. Case presentation

A 56-year-old female came to hospital with a complaint of pain during urination over the past month. There was also a complaint of continuous urinary incontinence, which had resulted in frequently wearing diapers over the last six months. She was embarrassed to spend time with others and thus had been isolating herself because she felt she smelled of urine. Since developing incontinence, the patient stated that she had not had sexual intercourse.

Meanwhile, the patient has a history of non-radiating low back pain. The patient claimed there was no blood or stones in her urine. Twenty years ago, her second pregnancy ended with a caesarean delivery, but a history of radiation was denied. Her menstruation was normal until her menopause began eight years ago. The patient had no significant record of previous medical history nor drug, cigarette or alcohol abuse.

On physical examination, the patient's general condition was good and vital signs were within normal limits with a body mass index (BMI) of 17.3 kg/m² (underweight). For the urological status, there were no abnormalities in the right or left costovertebral regions. Furthermore, there was no bladder distension. However, the vaginal examination revealed large stones protruding in the VVF area. Additionally, there was urine dribbling continually during a vaginal speculum examination. Based on laboratory results (routine blood), leucocytosis was found, and a leukocyte was discovered

* Corresponding author at: Division of Urology, Department of Surgery, Faculty of Medicine, Hasanuddin University, Jalan Perintis Kemerdekaan KM 11, Makassar, 90245, South Sulawesi, Indonesia.

E-mail addresses: ikaduma987@gmail.com (Francisca), khoirulkholis@yahoo.com (K. Kholis), apalinrungi@yahoo.com (M.A. Palinrungi), syahrirsyakri@gmail.com (S. Syahrir), syarifbakri@hotmail.com (Syarif), farox8283@gmail.com (M. Faruk).

<https://doi.org/10.1016/j.ijscr.2020.09.028>

2210-2612/© 2020 The Author(s). Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. The results of ultrasonography showing a hyperechoic shadow in the bladder measuring 1.27×1.36 cm (yellow arrow) accompanied by posterior acoustic shadow (white arrow). The bladder wall irregular and thickened (blue arrow).



Fig. 2. A kidney, ureter, and bladder (KUB) x-ray showing a radiopaque shadow in the pelvic area measuring about 7.5 cm.

on urinalysis tests. Also, a urological ultrasound (USG) examination (Fig. 1) showed a hyperechoic shadow measuring 1.27×1.36 cm in the vesica urinary. The vesica urinary walls were thickened and had an irregular surface. The size of both kidneys were normal with no stones observed. A kidney, ureter, and bladder (KUB) x-ray showed the presence of a radiopaque shadow of about 7.5×5 cm in the pelvic area (Fig. 2).

Based on the patient's history and the physical and supporting examinations, we diagnosed this patient with bladder stones accompanied by VVF. We (an urologic surgeon and a general surgery resident) performed an open cystolithotomy and fistula repair under general anesthesia. The stone, which was approximately $8 \times 5 \times 5$ cm, was successfully removed (Fig. 3) and a fistula hole of about 1 cm in diameter was found in the trigonum close to the internal meatus and subsequently repaired (Fig. 4).

Postoperatively, the patient was catheterized for fourteen days without the installation of peri-vesical drainage. The patient was advised to abstain from sexual intercourse for three months. At the patient follow-up at one, three, and six months postoperative, the results were satisfactory. She was voiding normally without incontinence and sexual function was significantly improved.

3. Discussion

The risk factors of bladder stone formation are continuous stagnation of infected urine [5,6], urinary tract infections [7,8], foreign bodies [9,10], diet [11], and the debris desquamation of vaginal epithelium into the bladder [6]. In this case, primary bladder stones were accompanied by VVF. Its exact relationship with VVF is not known. In India, Mukerji et al. reported three cases of bladder stones, the first case with VVF and the rest with rectourethral fis-




Fig. 3. There was oval-shaped bladder stones sized approximately $8 \times 5 \times 5$ cm.

tula [3]. Meanwhile, Lavery et al., found that 1 of 160 case of VVF accompanied by bladder stone due to ulceration and erosion [12].

VVF has many etiologies, which may be acquired or congenital, and the major causes in developed countries are radiotherapy, pelvic region malignancy, and surgical procedures. Meanwhile, the most common cause developing countries is obstructed labor [3,13]. Foreign bodies are the most common cause, which is



Fig. 4. Vaginal examination revealed the existed of yellowish-white stones (marked with yellow arrow) and vesicovaginal fistulas (blue arrow).

reported in about 10% of females with urinary disorders [14]. Meanwhile, other sources stated that VVF can be caused by pressure necrosis on the bladder wall [3]. In this study, the primary cause of the condition was pressure necrosis.

Clinical symptoms of bladder stones with VVF can include recurrent urinary tract infections, pyuria, suprapubic pain, severe swelling of the perineum, hematuria, or urinary incontinence. The severity of leakage can be reduced due to the “ball valve” effect caused by vesicle stones [13]. The patients in this study presented with the main symptoms of dysuria and continuous incontinence due to VVF.

Abdominal plain x-ray, ultrasonography (US), intravenous urography (IVU) and Multi-slice computed tomography (MSCT) are the most important tools for bladder stone evaluation and diagnosis. Most clinicians use KUB and US for initial examination [15].

Stone management depends on the size. It can be treated through cystolitholapaxy when it is less than 3 cm whereas open cystolithotomy is recommended when it is more than 3 cm. Existing literature suggested that handling bladder stones with VVF should follow a two-stage operation (Delayed Repair). The first stage is the removal of the stone with an endoscopic technique or open cystolithotomy, followed by VVF management for three months postoperative [5,6,16]. The rationale for the two-stage surgery is that the urine may contain infections or stones that can lead to inflammatory reactions in the bladder mucosa (edema) that will inhibit fistula healing [17].

The management of VVF operations is usually transvaginal or abdominal through a trans vesical approach. This treatment depends on the size and location of the fistula(e), the surgeon's experience, and the need for concurrent action [5,6]. Dalela et al. always performs the surgery by creating an omentum or flap Martius to strengthen the repair, and they succeeded in 84% of the cases [6].

In this study's patient, a one-stage operation (immediate repair) was performed because there were no signs of infection or edema. However, there was a possibility of fistula(e) repair failure in our

patient, but at the postoperative follow-up, the results were satisfactory.

4. Conclusion

Bladder stone is an uncommon condition that can cause VVF. Therefore, it is important to pay attention to its clinical signs.

Declaration of Competing Interest

Nothing to declare.

Funding

No funding or sponsorship.

Ethical approval

The study is exempt from ethical approval in our institution.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Author contribution

Francisca, Muhammad Asykar Palinrungi and Muhammad Faruk: study concept, surgical therapy for this patient. **Syahrir:** Data collection, Writing - Original draft preparation. **Khoirul Kholis:** senior author and the manuscript reviewer. **Syarif:** reviewed the manuscript. **Muhammad Faruk:** Editing, Writing. **All authors** read and approved the final manuscript.

Registration of research studies

Not applicable – single case report.

Guarantor

Muhammad Asykar Palinrungi.

Provenance and peer review

Not commissioned, externally peer-reviewed.

References

- [1] A. Cardoso, A. Paes, T. Teixeira, Giant bladder stone in a patient with vesicovaginal fistula caused by bladder wall ischemia due to unassisted prolonged labor in the amazon jungle, *Int. J. Med. Rev. Case Rep.* 2 (2018) 82, <http://dx.doi.org/10.5455/IJMRCR.giant-bladder-stone-vesicovaginal-fistula>.
- [2] I. Sunday-Adeoye, B. Daniyan, K. Ekwedigwe, D. Dantani, S. Uguru, *A Review of the Management of Vesico-Vaginal Fistula With Co-Existing Bladder Calculi in South-East Nigeria*, 2020.
- [3] A.C. Mukerji, B.C. Ghosh, M.M. Sharma, O.P. Taneja, Spontaneous urinary fistula caused by vesical calculus, *Br. J. Urol.* 42 (1970) 208–212, <http://dx.doi.org/10.1111/j.1464-410X.1970.tb10024.x>.
- [4] R.A. Agha, M.R. Borrelli, R. Farwana, K. Koshy, A.J. Fowler, D.P. Orgill, H. Zhu, A. Alsawadi, A. Noureldin, A. Rao, A. Enam, A. Thoma, M. Bashashati, B. Vasudevan, A. Beamish, B. Challacombe, R.L. De Wilde, D. Machado-Aranda, D. Laskin, D. Muzumdar, A. D'cruz, T. Manning, D. Healy, D. Pagano, P. Goel, P. Ranganathan, P.S. Pai, S. Raja, M.H. Ather, H. Kadioazlu, I. Nixon, I. Mukherjee, J. Gómez Rivas, K. Raveendran, L. Derbyshire, M. Valmasoni, M. Chalkoo, N. Raison, O. Muensterer, P. Bradley, C. Roberto, R. Afifi, D. Rosin, R. Klappenbach, R. Wynn, S. Giordano, S. Basu, S. Surani, P. Surman, M. Thorat, V. Kasi, The SCARE 2018 statement: updating consensus Surgical CAse REport (SCARE)

- guidelines, *Int. J. Surg.* 60 (2018) 132–136, <http://dx.doi.org/10.1016/j.ijss.2018.10.028>.
- [5] S. Rajaian, N.S. Kekre, Vesicovaginal fistula with large bladder calculus, *ANZ J. Surg.* 82 (2012) 278–279, <http://dx.doi.org/10.1111/j.1445-2197.2012.06014.x>.
- [6] D. Dalela, G. Apul, S.N. Shakhwar, K.M. Singh, Vesical calculi with unrepaired vesicovaginal fistula: a clinical appraisal of an uncommon association, *J. Urol.* 170 (2003) 2206–2208, <http://dx.doi.org/10.1097/01.ju.0000095503.76155.30>.
- [7] N. Segawa, Y. Katsuoka, K. Kaneda, [Vesico-vaginal fistula with a giant vesico-vaginal stone: a case report], *Hinyokika Kyo* 44 (1998) 517–520.
- [8] L. Yongzhi, Y. Shi, L. Jia, L. Yili, Z. Xingwang, G. Xue, Risk factors for urinary tract infection in patients with urolithiasis-primary report of a single center cohort, *BMC Urol.* 18 (2018) 45, <http://dx.doi.org/10.1186/s12894-018-0359-y>.
- [9] F. Kamal, A.T.D. Clark, L.T. Lavallée, M. Roberts, J. Watterson, Intravesical foreign body-induced bladder calculi resulting in obstructive renal failure, *Can. Urol. Assoc. J.* 2 (2008) 546–548, <http://dx.doi.org/10.5489/cuaj.927>.
- [10] M.A. Palinrungi, S. Syahrir, K. Kholis, Syarif, M. Faruk, Giant bladder stone formed around sewing-needle in childhood: a case report and literature review, *Urol. Case Rep.* 29 (2020), 101101, <http://dx.doi.org/10.1016/j.eucr.2019.101101>.
- [11] İ.N. Tahtalı, T. Karataş, Giant bladder stone: a case report and review of the literature, *Turk. J. Urol.* 40 (2014) 189–191, <http://dx.doi.org/10.5152/tud.2014.02603>.
- [12] D.W. Lavery, Vesico-vaginal fistulae: a report on the vaginal repair of 160 cases, *J. Obstet. Gynaecol. Br. Emp.* 62 (1955) 530–539, <http://dx.doi.org/10.1111/j.1471-0528.1955.tb14785.x>.
- [13] M.-H. Cheng, H.-T. Chao, H.-C. Horng, M.-J. Hung, B.-S. Huang, P.-H. Wang, Vesical calculus associated with vesicovaginal fistula, *Gynecol. Minim. Invasive Ther.* 3 (2014) 23–25, <http://dx.doi.org/10.1016/j.gmit.2013.07.009>.
- [14] N. Radnia, T. Eftekhari, S. Sohbati, T. Mohajery, Vesicovaginal fistula associated with a vaginal foreign body in a 16-year-old girl, *Nephrourol. Mon.* 10 (2018), e82387, <http://dx.doi.org/10.5812/numonthly.82387>.
- [15] K.-S. Chow, C.-Y. Chou, A boy with a large bladder stone, *Pediatr. Neonatol.* 49 (2008) 150–153, [http://dx.doi.org/10.1016/S1875-9572\(08\)60031-5](http://dx.doi.org/10.1016/S1875-9572(08)60031-5).
- [16] T.P. Mahapatra, M.S. Rao, K. Rao, S.K. Sharma, S. Vaidyanathan, Vesical calculi associated with vesicovaginal fistulas: management considerations, *J. Urol.* 136 (1986) 94–95, [http://dx.doi.org/10.1016/s0022-5347\(17\)44742-2](http://dx.doi.org/10.1016/s0022-5347(17)44742-2).
- [17] S.N. Shephard, S.J. Lengman, C.V. Kirschner, Bladder stones in vesicovaginal fistula: is concurrent repair an option? Experience with 87 patients, *Int. Urogynecol. J.* 28 (2017) 569–574, <http://dx.doi.org/10.1007/s00192-016-3142-1>.

Open Access

This article is published Open Access at scimedirect.com. It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.

Bladder stones associated with vesicovaginal fistula-TURNITIN

ORIGINALITY REPORT

3%

SIMILARITY INDEX

2%

INTERNET SOURCES

2%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.e-yujm.org Internet Source	1%
2	"Acknowledgement to Reviewers", International Journal of Surgery Case Reports, 2020 Publication	1%
3	mafiadoc.com Internet Source	1%
4	Shanmugasundaram Rajaian. "Vesicovaginal fistula with large bladder calculus : Images for surgeons", ANZ Journal of Surgery, 04/2012 Publication	1%
5	Rony A. Adam. "Diagnosis and Treatment of Fistulas", Clinical Gynecology, 2006 Publication	1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Bladder stones associated with vesicovaginal fistula-TURNITIN

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
