

Hepatitis C Virus Genotype in Blood Donors and Associated Liver Disease in Indonesia

by Upik Miskad

Submission date: 25-Jul-2021 09:15AM (UTC+0700)

Submission ID: 1623596331

File name: utama2008.pdf (208.21K)

Word count: 4994

Character count: 23278

Hepatitis C Virus Genotype in Blood Donors and Associated Liver Disease in Indonesia

Andi Utama^a Bugi Ratno Budiarto^a Dewi Monasari^a Theresia Imelda Octavia^a
Ivan Stevanus Chandra^a Rino Alvani Gani^b Irsan Hasan^b Andri Sanityoso^b
Upik Anderiani Miskad^c Irawan Yusuf^c Laurentius Adrianus Lesmana^b
Ali Sulaiman^b Susan Tai^a

^aMolecular Epidemiology Division, Mochtar Riady Institute for Nanotechnology, Lippo Karawaci, Tangerang,

^bDivision of Hepatology, Department of Internal Medicine, Faculty of Medicine, University of Indonesia, Jakarta, and ^cFaculty of Medicine, Hasanuddin University, Makassar, Indonesia

Key Words

Hepatitis C virus · Genotype · Chronic hepatitis C · Liver cirrhosis · Hepatocellular carcinoma · Indonesia

Abstract

Objective: The aim of this study was to investigate the distribution of hepatitis C virus (HCV) genotype and the possible association between genotype and HCV-associated liver disease in Indonesia. **Methods:** 32 anti-HCV-positive asymptomatic carriers (AC), 55 chronic hepatitis (CH), 41 liver cirrhosis (LC), and 35 hepatocellular carcinoma (HCC) patients were included in this study. HCV genotyping was performed by phylogenetic analysis of the NS5B and 5'-UTR regions. **Results:** The HCV subtype 1b (36.5%), based on NS5B region, was the most prevalent, followed by subtypes 3k (15.4%), 2a (14.4%), 1a (12.5%) and 1c (12.5%), and 2e (4.8%). Subtypes 2f, 3a, 3b, and 4a were also found in some of the samples. HCV subtypes 3k (40.0%) and 1a (35.0%) were the two major subtypes in AC. HCV subtype 1b was not found in AC, but it was common in CH (31.3%), LC (50.0%), and HCC (57.1%). **Conclusion:** HCV subtype 1b was prevalent in samples of HCV-as-

sociated liver disease patients, including CH, LC and HCC. The percentage of subtype 1b was increased with the disease severity (AC < CH < LC < HCC).

Copyright © 2009 S. Karger AG, Basel

Introduction

Hepatitis C virus (HCV) infection is known to be a major contribution to chronic liver disease such as chronic hepatitis (CH) and liver cirrhosis (LC), often leading to hepatocellular carcinoma (HCC). Worldwide, more than 170 million people are infected with HCV [1]. HCV is an enveloped virus with a single-stranded, positive-sense, non-segmented RNA genome of approximately 9,500 nucleotides, encoding a polyprotein precursor consisting of about 3,000 amino acids [2]. The polyprotein is cleaved by the host signal peptidase and two intrinsic viral encoded proteases to generate at least 10 viral proteins, including the core protein (C), the envelope glycoprotein (E1) and two types of envelope glycoprotein (E2), and 6 nonstructural proteins (NS2, NS3, NS4A, NS4B, NS5A, and NS5B) [3, 4].

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2009 S. Karger AG, Basel
0300-5526/08/0516-0410\$24.50/0

Accessible online at:
www.karger.com/int

Andi Utama, PhD
Molecular Epidemiology Division, Mochtar Riady Institute for Nanotechnology
Jalan Boulevard Jend. Sudirman 1688, Lippo Karawaci
Tangerang 15810, Banten (Indonesia)
Tel. +62 21 542 10123, Fax +62 21 542 10110, E-Mail autama@mrinstitute.org

55 Analysis of the HCV genome has revealed high heterogeneity of the virus. 44 Relatively well-conserved regions of the genome (5'-UTR, C, E1, and NS5B) have been extensively studied and used as the basis for the classification of at least six different genotypes in human [5]. The varied genotypes differ in distribution both by geographical region and by mode of transmission. Genotype 17-3 occur globally, while HCV genotype 4 is common in the Middle East and Africa [6-8], genotypes 5 and 6 are found 13 South Africa [9] and Southeast Asia [10], respectively. Subtypes 1a and 3a are associated with intravenous drug use, whereas subtype 1b is common in blood donors 58 Europe [11-13]. In Indonesia, the genotypes 1-3 are found in both blood donors and patients diagnosed with CH 15 C, and HCC [14-16].

HCV infection becomes chronic in about 85% of individuals as adjudged by the persistence of HCV RNA in serum [17]. CH may progress to worsening stages of fibrosis and cirrhosis which can ultimately lead to the development of HCC [18]. Several case-control and cohort studies of HCC and LC patients in Europe and Asia found a weak, albeit consistently, increased relative risk of HCV subtype 1b with the development of severe and advanced liver disease including cirrhosis and HCC [19-23]. However, 27 the studies did not confirm these findings [24, 25]. Thus, the association of HCV genotype and pathogenesis of liver disease including HCC remains 27 controversial and there is still limited information about the association 51 of HCV genotype and HCC in Indonesian patients. In this study, we investigated the distribution of HCV genotype in blood donor carriers and patients of CH, LC, and HCC, analyzing the possible association between HCV genotype and HCC development.

59 Materials and Methods

Samples

Serum samples were obtained from 163 blood donors and HCV-associated liver disease patients. There were 32 asymptomatic carriers (AC), 55 patients with CH, 41 patients having LC, and 35 HCC patients. AC sera were collected from blood donors in the Blood Transfusion Unit, Red Cross Makassar, South Sulawesi, between January 2007 and January 2008. Sera of CH, LC, and HCC patients were collected from Cipto Mangunkusumo Hospital and Klinik Hati 45arta, from the period of May 2005 until January 2008. All sera were positive for anti-HCV antibody as determined by using a third-generation HCV enzyme immunoassay (AXSYM, Abbott Laboratories, Chicago, Ill., USA). Blood samples were collected from each patient at the time of their clinical evaluation, then separated in 29 sera and stored at -70° until use for viral RNA extraction. The study was approved by the Institutional Ethic Committee and informed consent was obtained from each patient.

30 Viral RNA Extraction

Serum samples that had been stored at -70° were retrieved for analysis. HCV RNA was extracted from 200 µl serum using High Pure Viral RNA kit (Roche Diagnostics, Mannheim, Germany) according 47 the manufacturer's protocol. The RNA was finally eluted in 50 µl of ribonuclease-free water and stored at -70° until further analysis.

Reverse Transcriptase-Nested PCR NS5B Region

The viral RNA was used for genome amplification of partial NS5B region 28 a nested reverse transcriptase-PCR (RT-PCR). First-round RT-PCR was performed using the Access RT-PCR System (Promega, 34 Madison, Wisc., USA) in 25-µl aliquots containing 5 µl RNA, 0.2 mM of each dNTP, 1× AMV buffer, 2 mM MgSO₄, 2.5 units of AMV reverse transcriptase, 2.5 units of Tfl polymerase, and 1 µM of NS5B-1 and NS5B-2 primers (table 1). The following cycling parameters were used for the first-round RT-PCR: cDNA synthesis at 45° (45 min), enzyme inactivation at 95° (5 min), DNA amplification: denaturation at 95° (30 s), annealing at 80-55° (30 s) and elongation at 72° (1 min). For the first 6 cycles, the annealing temperature was reduced by 5° per cycle (touchdown PCR), and 55° for the rest of 23 cycles. Two microliters of PCR product from first-round PCR was used as a template for second-round PCR. The second-round PCR was performed using the Go Taq PCR Core System (Promega) in 25-µl aliquots containing 2 mM MgCl₂, 1× Green Go Taq Flexi Buffer, 0.625 units of Go Taq DNA polymerase, and 0.5 µM of NS5B-3 and NS5B-4 primers (table 1). The following cycling parameters were used 38 for 30 cycles of second-round PCR: denaturation at 94° (1 min), annealing at 50° (2 min) and elongation at 72° (2 min).

5'-UTR Region

A nested 48-PCR was devised to amplify the 5'-UTR region. First-round RT-PCR was performed using Access 1 34 PCR System (Promega) in 25-µl aliquots containing 5 µl RNA, 0.2 mM of each dNTP, 1× AMV buffer, 2 mM MgSO₄, 2.5 units of AMV reverse transcriptase, 2.5 units of Tfl polymerase, and 1 µM of 5'-UTR-1 and 5'-UTR-2 primers (table 1). The following cycling parameters were used for the first-round RT-PCR: cDNA synthesis at 45° (60 min), enzyme inactivation at 95° (5 min), 35 cycles of DNA amplification: denaturation at 95° (1 min), annealing 21 7.7° (30 s) and elongation at 72° (1 min). Two microliters of PCR product from the first-round PCR was taken as a template for the second-round PCR with Go Taq PCR Core System (Promega) using 0.5 µM of 5'-UTR-3 and 5'-UTR-4 primers (table 1). The following cycling parameters were used for 30 cycles in the second-round PCR: denaturation at 95° (1 min), annealing at 53.7° (30 s) and elongation at 72° (1 min).

Sequencing

PCR product of 16 NS5B and 5'-UTR regions were purified from agarose gel using Wizard SV Gel and PCR Clean Up system kit (Promega), according to the manufacturer's protocol. Purified D 19 fragments were directly sequenced employing an ABI 3130 xl Genetic Analyzer (Applied Biosystems, Inc., Foster City, Calif., USA) with the Big Dye Terminator V3.1 Cycle Sequencing kit (Applied Biosystems, Inc.).

36

Table 1. Primers used in this study

Primer	Nucleotide sequence (5' → 3')	Polarity	Reference
NS5B			
NS5B-1	TATGAYACCCGYTGCTTTGAC	forward	Cantaloube et al., 2005
NS5B-2	GAGGAGCAAGATGTTATCAGCTC	reverse	
NS5B-3	GATACCCGCTGCTTTGACTC	forward	
NS5B-4	GAATACCTGGTCATAGCCTCCG	reverse	
5'-UTR			
5'-UTR1	CCCTGTGAGGAACTWCTGTCTTCACGC	forward	Stuyver et al., 1996
5'-UTR2	TCTAGCCATGGCGTTAGTAYGAGTGT	reverse	
5'-UTR3	CACTCGCAAGCACCTATCAGGCAGT	forward	
5'-UTR4	GCTCATGRTGCACGGTCTACGAGACCT	reverse	

63

Table 2. Positivity of RT-PCR and sequencing of the NS5B region of samples from different clinical diagnoses

Clinical diagnosis	n	PCR-positive n (%)	Sequenced n (%)
AC	32	20 (62.5)	20 (100)
CH	55	32 (58.2)	32 (100)
LC	41	25 (61.0)	24 (96.0)
HCC	35	28 (80.0)	28 (100)
Total	163	105 (64.4)	104 (99.0)

Phylogenetic Analysis

The nucleotide sequences of the HCV and a panel of sequences retrieved from the GenBank were aligned using BioEdit Sequence Alignment Editor. Phylogenetic analysis of NS5B (367 bp) and 5'-UTR (236 bp) regions was performed with ClustalW version 1.83.

Statistical Analysis

Statistical analyses were performed using SPSS 15.0 (SPSS, Inc., Chicago, Ill., USA). χ^2 test, unpaired *t* test, and ANOVA were used to assess the statistical significance of the difference between groups. *p* values <0.05 were considered statistically significant.

Results

HCV Genotype Distribution

The NS5B region was used for HCV genotyping and 105 out of the 163 samples (64.4%) were positive by RT-PCR (table 2). The number of samples that could be amplified was higher in HCC (80.0%) compared with other

groups; 62.5% in AC, 58.2% in CH, and 61.0% in LC patients. Of the 105 RT-PCR positive samples, 104 samples (99.0%) could be sequenced and classified not only into genotype, but also subtype. By phylogenetic analysis of partial NS5B region (367 bp), the HCV strains from 104 samples were classified into the genotype 1 (with subtypes 1a, 1b, and 1c), 2 (with subtypes 2a, 2e, and 2f), 3 (with subtypes 3a, 3b, and 3k), and 4 (subtype 4a) (table 3). The subtype 1b was the most prevalent, accounting for 36.5% of the total samples, followed by subtypes 3k (15.4%), 2a (14.4%), 1a (12.5%) and 1c (12.5%), and 2e (4.8%) (table 3). One of each subtype 2f, 3a, 3b, and 4a was also found in the samples.

HCV Genotype and Clinical Diagnosis

When the genotype distribution was compared with clinical diagnosis, it was found that subtype 3k (40.0%) and 1a (35.0%) were dominantly detected in the AC group (table 3). The HCV subtype 1b was prevalent in CH (31.3%), LC (50.0%), and HCC (57.1%) groups. In the CH group, HCV genotype 1 was common (62.5%), which composed of subtypes 1a (9.4%), 1b (31.3%), and 1c (21.9%), in addition the genotypes 2 (subtypes 2a and 2e) and 3 (subtypes 3a and 3k) were also detected. Similarly, genotype 1 was also the major genotype in LC (70.8%), which composed of subtypes 1a (8.3%), 1b (50.0%), and 1c (12.5%). A high percentage of genotype 1 (57.1%) was found in HCC. Particular to subtype 1b, the prevalence of this subtype was increased from AC to HCC (0.0% in AC, 31.3% in CH, 50.0% in LC, and 57.1% in HCC) (table 3). Statistical analysis revealed a significant difference of HCV subtype 1b among all clinical stages ($p < 0.001$); however, it was not significant across all clinical stages. For instance, there was no significant difference between

Table 3. HCV genotype prevalence in different clinical diagnosis based on NS5B region

Genotype	AC (n = 20)	CH (n = 32)	LC (n = 24)	HCC (n = 28)	Total
Genotype 1	8 (40.0)	20 (62.5)	17 (70.8)	19 (67.9)	64 (61.5)
1a	7 (35.0)	3 (9.4)	2 (8.3)	1 (3.6)	13 (12.5)
1b	0 (0.0)	10 (31.3)	12 (50.0)	16 (57.1)	38 (36.5)
1c	1 (5.0)	7 (21.9)	3 (12.5)	2 (7.1)	13 (12.5)
Genotype 2	2 (10.0)	6 (18.8)	5 (20.8)	8 (28.6)	21 (20.2)
2a	2 (10.0)	5 (15.6)	3 (12.5)	5 (17.9)	15 (14.4)
2e	0 (0.0)	1 (3.1)	1 (4.2)	3 (10.7)	5 (4.8)
2f	0 (0.0)	0 (0.0)	1 (4.2)	0 (0.0)	1 (1.0)
Genotype 3	9 (45.0)	6 (18.8)	2 (8.3)	1 (3.6)	18 (17.3)
3a	0 (0.0)	1 (3.1)	0 (0.0)	0 (0.0)	1 (1.0)
3b	1 (5.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.0)
3k	8 (40.0)	5 (15.6)	2 (8.3)	1 (3.6)	16 (15.4)
Genotype 4	1 (5.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.0)
4a	1 (5.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.0)
Grand total	20 (100.0)	32 (100.0)	24 (100.0)	28 (100.0)	104 (100.0)

Table 4. HCV genotype prevalence in different clinical diagnoses based on the 5'-UTR region

Genotype	AC (n = 20)	CH (n = 31)	LC (n = 24)	HCC (n = 28)	Total
Genotype 1	9 (45.0)	18 (58.0)	17 (70.8)	20 (71.4)	64 (62.1)
Genotype 2	1 (5.0)	6 (19.4)	5 (20.8)	6 (21.4)	18 (17.5)
Genotype 3	9 (45.0)	7 (22.6)	2 (8.3)	2 (7.1)	20 (19.4)
3a	0 (0.0)	1 (3.2)	0 (0.0)	0 (0.0)	1 (1.0)
3b	1 (5.0)	1 (3.2)	0 (0.0)	0 (0.0)	2 (2.0)
3k	8 (40.0)	5 (16.1)	2 (8.3)	2 (7.1)	17 (16.5)
Genotype 4	1 (5.0)	0 (0.0)	0 (0.0)	0 (0.0)	1 (1.0)
Grand total	20 (100.0)	31 (100.0)	24 (100.0)	28 (100.0)	103 (100.0)

Table 5. Samples which showed different genotypes from an analysis based on 5'-UTR and NS5B regions

No.	Report No.	Clinical diagnosis	HCV genotype based on	
			5'-UTR	NS5B
1	07.22.133	AC	1	2a
2	P.X01.27	LC	1	2a
3	06.10.009	HCC	3k	2a

43

CH and LC ($p = 0.121$), and between LC and HCC ($p = 0.565$), but there was significant difference between AC and CH ($p = 0.015$). Moreover, the prevalence of subtype 1₅₇ was significantly higher in HCC compared to both CH ($p < 0.001$) and AC ($p < 0.001$).

Comparison of NS5B and 5'-UTR Regions

In samples where the NS5B region could be amplified, the 5'-UTR region was explored to confirm the genotype. Phylogenetic analysis of the 5'-UTR region showed that it could identify the genotype, but not the subtype of HCV, particularly for genotypes 1 and 2 as compared to




Fig. 1. Unrooted neighbor-joining phylogenetic tree from (a) partial 5'-UTR and (b) NS5B. Samples which are different genotypes in 5'-UTR and NS5B regions are highlighted.

analysis based on NS5B region (fig. 1); however, it could distinguish the subtype of genotype 3 (fig. 1a). By phylogenetic analysis of the 5'-UTR region (236 bp), the HCV strains from 103 samples were classified into genotypes 1 (62.1%), 2 (17.5%), 3 (19.4%) and 4 (1.0%) (table 4), and were identical to genotyping based on NS5B region (table 3). However, the genotyping results based on 5'-UTR and NS5B regions were different in three samples (07.22.133, P.X01.27, and 06.10.009 from the AC, LC, and

HCC groups, respectively) (table 5). These three samples were identified as subtype 2a based on NS5B region, but 07.22.133 and P.X01.27 were classified as HCV genotype 1 and 06.10.009 was identified as subtype 3k based on the 5'-UTR region (table 5, fig. 1). These findings indicate that the HCV isolates in those samples might be recombinant viruses resulted from recombination of HCV with different genotypes.

Discussion

Phylogenetic analysis of partial NS5B region demonstrated that subtype 1b was the most prevalent in our samples, although other subtypes (3k, 2a, 2e, 1a, and 1c) were also detected. Generally, our findings are similar with previous data of HCV genotype in Indonesia. Hotta et al. [14] identified HCV subtypes 1b, 1d, and 2a from CH and LC patients and subtype 1b was found in 45.0 and 34.0% samples of CH and LC, respectively. In a study of blood donors, hemodialysis and HCC patients, subtypes 1a, 1b, 1d, 2a, and genotype 4 were found [15]. The subtype 2a was dominant in blood donors (51.9%), whilst subtype 1b was more common in hemodialysis and HCC patients (31.3 and 57.1%, respectively). HCV genotyping of 64 blood donor samples found that the predominant genotype is 1b (57.8%); this followed by 2a (17.2%), and 3b (10.9%) [16]. In contrast with this finding, our results did not identify genotype 1b from the blood donor (AC) group. In addition, we also found one subtype 4a, the genotype that is common in the Middle East and Africa, in AC samples.

When genotype distribution was analyzed based on clinical diagnosis, it was determined that subtype 1b was not found in AC, but was prevalent in CH and increased in LC and HCC. Statistical analysis did not show a significant difference between CH and LC, and between LC and HCC. It is suggested that there is an association between HCV subtype 1b and CH development, but no association with development of HCC from LC or CH. A weak but consistently increased relative risk of HCC in patients with subtype 1b has been found in several case-control and cohort studies of HCC and LC patients in Europe and Asia [19–23], which is different to our findings. However, further studies have not confirmed these findings [24, 25]. Therefore, the association of HCV genotype and pathogenesis of liver disease, particularly HCC, is still debatable, and needs prospective cohort studies to obtain more conclusive data.

References

- 1 World Health Organization: Global surveillance and control of hepatitis C. *J Viral Hepat* 1999;6:35–47.
- 2 Choo QL, Richman KH, Han JH, Berger K, Lee C, Dong C, Gallegos C, Coit D, Medina J, Barr PJ, Weiner A, Bradley DW, Kuo G, Houghton M: Genetic organization and diversity of the hepatitis C virus. *Proc Natl Acad Sci USA* 1991;88:2451–2455.
- 3 Akouli A, McCourt DW, Wychowski C, Feinstone SM, Rice CM: Characterization of the hepatitis C virus-encoded serine protease: determination of proteinase-dependent polyprotein cleavage sites. *J Virol* 1993;67:2832–2843.
- 4 Mizushima H, Hijikata M, Asabe S, Hirota M, Kimura K, Shimotohno K: Two hepatitis C virus glycoproteins E2 products with different C termini. *J Virol* 1994;68:6215–6222.

Comparison of phylogenetic analysis in both NS5B and 5'-UTR regions showed differences of HCV genotypes in three samples, implying the possible recombination between different genotypes. Analysis of those samples was confirmed by repeating the sequencing experiment and data analysis. It is hypothesized that recombination might have occurred between genotype 1 and 2a, which results in 1/2a (for 07.22.133 and P.X01.27) or between subtype 3k and 2a, which results in 3k/2a (for 06.10.009). Further sequence analysis of full-length genome is needed to prove that a recombination event has occurred. Nevertheless, prior investigators reported the natural recombination of HCV from samples in St. Petersburg (2k/1b), Peru (1b/1a), the Philippines (2b/1b), Vietnam (2/6), Ireland (2k/1b), and France (2/5) [26–31]. Since there is no prevalence of recombination in certain stage of liver disease, it is suggested that virus recombination event could naturally occur during virus replication. The HCV recombination is a cause of genetic diversity of the virus, which may have an important implication for the pathogenesis, laboratory diagnosis, and treatment of HCV infection. However, so far there is no direct evidence on the clinical implication of the recombination, including response to certain therapy. In summary, our study demonstrated that HCV subtype 1b was prevalent in HCV-associated liver disease in the order of CH < LC < HCC. By comparing NS5B and 5'-UTR regions, putative recombinant of HCV was found in three samples.

Acknowledgments

The authors thank Mardiana Radjuni, Tities Anggraeni Indra, Marcell Harlan for sample collection. The authors also thank Drs. John D. Groopman (Johns Hopkins Bloomberg School of Public Health) and Ivori Aripranata (Mochtar Riady Institute for Nanotechnology) for critical reading of the manuscript. This work was supported by MRIN Funding (Budget No. cc042/2007).

- ▶ 5 Simmonds P, Bukh J, Combet C, Deleage G, Enomoto N, Feinstone S, Halfon P, Inchauspe G, Kuiken C, Maertens G, Mizokami M, Murphy DG, Okamoto H, Pawlotsky JM, Penin F, Sablon E, Shin-i T, Stuyver LJ, Thiel HJ, Viazov S, Weiner AJ, Widell A: Consensus proposals for a unified system of nomenclature of hepatitis C virus genotypes. *Hepatology* 2005;42:962-973.
- ▶ 6 Bukh J, Purcell RH, Miller RH: At least 12 genotypes of hepatitis C virus predicted by sequence analysis of the putative E1 gene of isolates collected worldwide. *Proc Natl Acad Sci USA* 1993;90:8234-8238.
- ▶ 7 Ndjimou J, Pybus OG, Matz B: Phylogenetic analysis of hepatitis C virus isolates indicates a unique pattern of endemic infection in Cameroon. *J Gen Virol* 2003;84:2333-2341.
- ▶ 8 Abdel-Hamid M, El-Daly M, Molnegren V, El-Kafrawy S, Abdel-Latif S, Esmat G, Strickland G, Loffredo C, Albert J, Widell A: Genetic diversity in hepatitis C virus in Egypt and possible association with hepatocellular carcinoma. *J Gen Virol* 2007;88:1526-1531.
- ▶ 9 Smuts HE, Kannemeyer J: Genotyping of hepatitis C virus in South Africa. *J Clin Microbiol* 1995;33:1679-1681.
- ▶ 10 Lu L, Li C, Fu Y, Thaikruea L, Thongsawat S, Maneekarn N, Apichartpiyakul C, Hotta H, Okamoto H, Netski D, Pybus OG, Murphy D, Hahn CH, Nelson E: Complete genome for hepatitis C virus subtypes 6f, 6i, 6j and 6m: viral genetic diversity among Thai blood donors and infected spouses. *J Gen Virol* 2007;88:1505-1518.
- ▶ 11 Haushofer AC, Koptcy C, Hauer R, Brunner H, Halbmayer WM: HCV genotypes and age distribution in patients of Vienna and surrounding areas. *J Clin Virol* 2001;20:41-47.
- ▶ 12 Bourliere M, Barberin JM, Rotily M, Gualdiardo V, Portal I, Lecomte L, Benali S, Rustiere C, Perrier H, Jullien M, Lombot G, Loyer R, LeBars O, Daniel R, Khiri H, Halfon P: Epidemiological changes in hepatitis C virus genotypes in France: evidence in intravenous drug users. *J Viral Hepat* 2002;9:62-70.
- ▶ 13 Van Asten L, Verhaest I, Lamzira S, Hernandez-Aguado I, Zangerle R, Boufassa F, Rezza G, Broers B, Robertson JR, Brettle RP, McMenamin J, Prins M, Cochrane A, Simmonds P, Coutinho RA, Bruisten S: Spread of hepatitis C virus among European injection drug users infected with HIV: a phylogenetic analysis. *J Infect Dis* 2004;189:292-302.
- ▶ 14 Hatta H, Handajani R, Lusida MI, Soemarto W, Doi H, Miyajima H, Homma M: Subtype analysis of hepatitis C virus in Indonesia on the basis of NS5B region sequences. *J Clin Microbiol* 1994;32:3049-3051.
- ▶ 15 Soetjipto, Handajani R, Lusida MI, Darmadi S, Adi P, Soemarto, Ishido S, Katayama Y, Hatta H: Differential prevalence of hepatitis C virus subtypes in healthy blood donors, patients on maintenance hemodialysis, and patients with hepatocellular carcinoma in Surabaya, Indonesia. *J Clin Microbiol* 1996;34:2875-2880.
- ▶ 16 Inoue Y, Sulaiman HA, Matsubayashi K, Julitasari, Iinuma K, Ansari A, Laras K, Corwin AL: Genotypic analysis of hepatitis C virus in blood donors in Indonesia. *Am J Trop Med Hyg* 2000;62:92-98.
- ▶ 17 Di Bisceglie AM: Hepatitis C. *Lancet* 1998;351:351-355.
- ▶ 18 Marcellin P: Hepatitis C: the clinical spectrum of the disease. *J Hepatol* 1999;31:9-16.
- ▶ 19 Silini E, Bottelli R, Asti M, Bruno S, Canino ME, Brambilla S, Bono F, Iamoni G, Tinelli C, Mondelli MU, Ideo G: Hepatitis C virus genotypes and risk of hepatocellular carcinoma in cirrhosis: a case-control study. *Gastroenterology* 1996;111:199-205.
- ▶ 20 Hatzakis A, Katsoulidou A, Kaklamani E, Touloumi G, Koumantaki Y, Tassopoulos NC, Karvountzis G, Gioustozi A, Hadziyan G, Trichopoulos D: Hepatitis C virus 1b is the dominant genotype in HCV-related carcinogenesis: a case-control study. *Int J Cancer* 1996;68:51-53.
- ▶ 21 Ikeda K, Kobayashi M, Someya T, Saitoh S, Tsubota A, Akuta N, Suzuki F, Suzuki Y, Iwase Y, Kumada H: Influence of hepatitis C virus subtype on hepatocellular carcinogenesis: a multivariate analysis of a retrospective cohort of 593 patients with cirrhosis. *Intervirology* 2002;45:71-78.
- ▶ 22 Chen CM, Hung CH, Lu SN, Wang JH, Tung HD, Huang WS, Chen CL, Chen WJ, Changchien CS: Viral etiology of hepatocellular carcinoma and HCV genotypes in Taiwan. *Intervirology* 2006;49:76-81.
- ▶ 23 Bruno S, Silini E, Grosignani A, Borzio F, Leandro G, Bono F, Asti M, Rossi S, Larghi A, Cerino A, Podda M, Mondelli MU: Hepatitis C virus genotypes and risk of hepatocellular carcinoma: a prospective study. *Hepatology* 1997;25:754-758.
- ▶ 24 Bruno S, Yokosuka O, Imazeki F, Tagawa M, Omata M: Incidence of hepatocellular carcinoma in chronic hepatitis B and C: a prospective study of 251 patients. *Hepatology* 1995;21:650-655.
- ▶ 25 Benvegno L, Pontisso P, Cavalletto D, Nascimben F, Chemello L, Alberti A: Lack of correlation between hepatitis C virus genotypes and clinical course of hepatitis C virus-related cirrhosis. *Hepatology* 1997;25:211-215.
- ▶ 26 Kalinina O, Norder H, Mukomolov S, Magnius LO: A natural intergenotypic of hepatitis C virus identified in St. Petersburg. *J Virol* 2002;76:4034-4043.
- ▶ 27 Colina R, Casane D, Vasquez S, Garcia-Irre L, Chunga A, Romero H, Khan B, Cristina J: Evidence of intratypic recombination in natural populations of hepatitis C virus. *J Gen Virol* 2004;85:31-37.
- ▶ 28 Kageyama S, Agdamag DM, Alesna ET, Leano PS, Heredia AML, Abellanosa-Tac-An IP, Jereza T, Tanimoto T, Yamamura J, Ichimura H: A natural inter-genotypic (2b/1b) recombinant of hepatitis C virus in the Philippines. *J Med Virol* 2006;78:1423-1428.
- ▶ 29 Noppornpanth S, Lien TX, Poovorawan Y, Smits SL, Osterhaus AD, Haagmans BL: Identification of a naturally occurring recombinant genotype 2/6 hepatitis C virus. *J Virol* 2006;80:7569-7577.
- ▶ 30 Moreau I, Hegarty S, Levis J, Sheehy P, Crosbie O, Kenny-Walsh E, Fanning LJ: Serendipitous identification of natural intergenotypic recombinants of hepatitis C virus in Ireland. *Virology* 2006;3:95.
- ▶ 31 Grand-Abravanel F, Claudinon J, Nicot F, Dubois M, Chapuy-Regaud S, Sandres-Saune K, Pasquier C, Izopet J: New natural intergenotypic (2/5) recombinant of hepatitis C virus. *J Virol* 2007;81:4357-4362.

Hepatitis C Virus Genotype in Blood Donors and Associated Liver Disease in Indonesia

ORIGINALITY REPORT

23%

SIMILARITY INDEX

17%

INTERNET SOURCES

21%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1	www.karger.jp Internet Source	1%
2	www.jidc.org Internet Source	1%
3	João Manuel Antunes Bras. "Effects of Peripheral Axotomy on Cholecystinin Neurotransmission in the Rat Spinal Cord", <i>Journal of Neurochemistry</i> , 06/13/2002 Publication	1%
4	www.nlm.nih.gov Internet Source	1%
5	hepatmon.com Internet Source	1%
6	Chuang, C.K.. "Experimental evidence that RNA recombination occurs in the Japanese encephalitis virus", <i>Virology</i> , 20091125 Publication	1%
7	app.trdizin.gov.tr Internet Source	1%

8

www.ints.fr

Internet Source

1 %

9

GERRY C MACQUILLAN. "Does sequencing the PKRBD of hepatitis C virus NS5A predict therapeutic response to combination therapy in an Australian population?", *Journal of Gastroenterology and Hepatology*, 5/2004

Publication

1 %

10

Fujita, T.. "Suppression of Actinomycin D-Induced Apoptosis by the NS3 Protein of Hepatitis C Virus", *Biochemical and Biophysical Research Communications*, 19961224

Publication

<1 %

11

Ting-Liang Wang, Hong-Lei Li, Wen-Ye Tjong, Qian-Su Chen et al. "Genetic factors contribute to patient-specific warfarin dose for Han Chinese", *Clinica Chimica Acta*, 2008

Publication

<1 %

12

Indra Bachtiar, Julian Mulya Santoso, Benny Atmanegara, Rino Alvani Gani et al. "Combination of alpha-1-acid glycoprotein and alpha-fetoprotein as an improved diagnostic tool for hepatocellular carcinoma", *Clinica Chimica Acta*, 2009

Publication

<1 %

13

O. Kalinina. "Shift in predominating subtype of HCV from 1b to 3a in St. Petersburg mediated by increase in injecting drug use", *Journal of Medical Virology*, 11/01/2001

Publication

<1 %

14

R. Trowbridge, E. J. Gowans. "Molecular cloning of an Australian isolate of hepatitis C virus", *Archives of Virology*, 2014

Publication

<1 %

15

Adrian M Di Bisceglie. "Hepatitis C", *The Lancet*, 1998

Publication

<1 %

16

www.theses.fr

Internet Source

<1 %

17

Aline G Vigani. "Comparative study of patients with chronic hepatitis C virus infection due to genotypes 1 and 3 referred for treatment in southeast Brazil", *BMC Infectious Diseases*, 2008

Publication

<1 %

18

Rinonce, Hanggoro Tri, Yoshihiko Yano, Takako Utsumi, Didik Setyo Heriyanto, Nungki Anggorowati, Dewiyani Indah Widasari, Maria Inge Lusida, Soetjipto, Heru Prasanto, Hak Hotta, and Yoshitake Hayashi. "Hepatitis B and C virus infection among hemodialysis patients in yogyakarta, Indonesia: Prevalence

<1 %

and molecular evidence for nosocomial transmission : Hepatitis B and C in Indonesian Hemodialysis Patients", Journal of Medical Virology, 2013.

Publication

19

Submitted to Universidad Nacional de Colombia

Student Paper

<1 %

20

Yuki Ishida, Tsunefusa Hayashida, Masaya Sugiyama, Haruka Uemura et al. "Full - genome analysis of hepatitis C virus in HIV - coinfecting hemophiliac Japanese patients", Hepatology Research, 2020

Publication

<1 %

21

Submitted to University of Hyderabad, Hyderabad

Student Paper

<1 %

22

ddd.uab.cat

Internet Source

<1 %

23

open.library.ubc.ca

Internet Source

<1 %

24

Giovanni Matera, Angelo Lamberti, Angela Quirino, Domenico Focà et al. "Changes in the prevalence of hepatitis C virus (HCV) genotype 4 in Calabria, Southern Italy", Diagnostic Microbiology and Infectious Disease, 2002

Publication

<1 %

25 Seiji Kageyama. "A natural inter-genotypic (2b/1b) recombinant of hepatitis C virus in the Philippines", *Journal of Medical Virology*, 11/2006
Publication <1 %

26 www.em-consulte.com
Internet Source <1 %

27 Francesc X. López-Labrador, Sergi Ampurdanés, Xavier Forn, Antoni Castells et al. "Hepatitis C virus (HCV) genotypes in Spanish patients with HCV infection: relationship between HCV genotype 1b, cirrhosis and hepatocellular carcinoma", *Journal of Hepatology*, 1997
Publication <1 %

28 Jose de la Fuente. "Gene expression profiling of human promyelocytic cells in response to infection with *Anaplasma phagocytophilum*", *Cellular Microbiology*, 4/2005
Publication <1 %

29 Takao SHIBAYAMA. "Risk Factors of Hepatocellular Carcinoma in Chronic Hepatitis C and Cirrhosis: Special Reference to Laparoscopic Findings", *Digestive Endoscopy*, 1/1999
Publication <1 %

30 Submitted to University of Hong Kong
Student Paper

<1 %

31

www.annalsgastro.gr

Internet Source

<1 %

32

Avó, Ana Patrícia, Ivone Água-Doce, Ana Andrade, and Elizabeth Pádua. "Hepatitis C virus subtyping based on sequencing of the C/E1 and NS5B genomic regions in comparison to a commercially available line probe assay", *Journal of Medical Virology*, 2013.

Publication

<1 %

33

M. S. Choi. "Clinical significance of pre-S mutations in patients with genotype C hepatitis B virus infection", *Journal of Viral Hepatitis*, 3/2007

Publication

<1 %

34

Victoria Naranjo, Ursula Häßfle, Joaquín Vicente, Ma Paz Martín et al. " Genes differentially expressed in oropharyngeal tonsils and mandibular lymph nodes of tuberculous and nontuberculous European wild boars naturally exposed to ", *FEMS Immunology & Medical Microbiology*, 2006

Publication

<1 %

35

ueg.eu

Internet Source

<1 %

36

www.j3.jstage.jst.go.jp

Internet Source

<1 %

37

www.lifesciencesite.com

Internet Source

<1 %

38

Liu, Z.. "Phosphorylation of Canine Distemper Virus P Protein by Protein Kinase C- α and Casein Kinase II", *Virology*, 19970526

Publication

<1 %

39

bmccancer.biomedcentral.com

Internet Source

<1 %

40

cpb-us-w2.wpmucdn.com

Internet Source

<1 %

41

eprints.whiterose.ac.uk

Internet Source

<1 %

42

G.A.S. Pereira. "Human immunodeficiency virus type 1 and hepatitis C virus Co-infection and viral subtypes at an HIV testing center in Brazil", *Journal of Medical Virology*, 06/2006

Publication

<1 %

43

M. Bortolami. "Fas / FasL system, IL-1 β expression and apoptosis in chronic HBV and HCV liver disease", *Journal of Viral Hepatitis*, 3/6/2008

Publication

<1 %

44

M. Kew. "Prevention of hepatitis C virus infection*", *Journal of Viral Hepatitis*, 5/2004

<1 %

45 bmcgastroenterol.biomedcentral.com <1 %
Internet Source

46 dev.endocrine.org <1 %
Internet Source

47 www.agriculturejournals.cz <1 %
Internet Source

48 www.plantcell.org <1 %
Internet Source

49 www.suodangzhi.com <1 %
Internet Source

50 www.theses.ulaval.ca <1 %
Internet Source

51 Dong, Zhi Xia, Hui Juan Zhou, Jian Hua Wang, Xiao Gang Xiang, Yan Zhuang, Si Min Guo, Hong Lian Gui, Gang De Zhao, Wei Liang Tang, Hui Wang, and Qing Xie. "Distribution of hepatitis C virus genotypes in Chinese patients with chronic hepatitis C: Correlation with patients's characteristics and clinical parameters : HCV genotypes in Chinese patients", Journal of Digestive Diseases, 2012.
Publication

52 Haushofer, A.C.. "HCV genotypes and age distribution in patients of Vienna and <1 %

surrounding areas", Journal of Clinical
Virology, 200101

Publication

53

Indra Bachtiar. "Alpha-1-acid glycoprotein as potential biomarker for alpha-fetoprotein-low hepatocellular carcinoma", BMC Research Notes, 2010

Publication

<1 %

54

www.fma.org.tw

Internet Source

<1 %

55

" Hepatitis ", Wiley, 2006

Publication

<1 %

56

Cesar A. B. Duarte, Leonardo Foti, Sueli M. Nakatani, Irina N. Riediger, Celina O. Poersch, Daniela P. Pavoni, Marco A. Krieger. "A Novel Hepatitis C Virus Genotyping Method Based on Liquid Microarray", PLoS ONE, 2010

Publication

<1 %

57

Osamu Nunobiki. "Significance of hormone receptor status and tumor vessels in normal, hyperplastic and neoplastic endometrium", Pathology International, 12/2003

Publication

<1 %

58

R María. "Identification of hepatitis C virus (HCV) genotypes in infected patients from the west of Mexico", Hepatology Research, 1998

Publication

<1 %

59 Teeraporn Chinchai, Suwanna Noppornpanth, Kavita Bedi, Apiradee Theamboonlers, Yong Poovorawan. "222 Base Pairs in NS5B Region and the Determination of Hepatitis C Virus Genotype 6", Intervirology, 2006
Publication

60 [documents.mx](#)
Internet Source

61 [journalijar.com](#)
Internet Source

62 [openaccess.leidenuniv.nl](#)
Internet Source

63 [osshhm.org](#)
Internet Source

64 [webhome.weizmann.ac.il](#)
Internet Source

65 Arief Suriawinata, Swan N. Thung. "Hepatitis C virus and malignancy", Hepatology Research, 2007
Publication

66 Denise P.C. Chan, Shui Shan Lee, Krystal C.K. Lee. "The effects of widespread methadone treatment on the molecular epidemiology of hepatitis C virus infection among injection drug users in Hong Kong", Journal of Medical Virology, 2011

67 Sukowati, Caecilia HC. "Significance of hepatitis virus infection in the oncogenic initiation of hepatocellular carcinoma", *World Journal of Gastroenterology*, 2016. <1 %
Publication

68 vir.sgmjournals.org <1 %
Internet Source

69 www.kjim.org <1 %
Internet Source

70 bmcbioinformatics.biomedcentral.com <1 %
Internet Source

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On