

03_PP_YT_IJAST1.pdf

by

FILE	03_PP_YT_IJAST1.PDF (271.89K)	WORD COUNT	3284
TIME SUBMITTED	13-JAN-2021 01:12PM (UTC+0700)	CHARACTER COUNT	17713
SUBMISSION ID	1486735501		

Analysis of Factors Affecting Musculoskeletal Disorders to Female Seaweed Workers

Adhinda Putri Pratiwi¹, Yahya Thamrin¹, Suriah², Masyitha Muis¹, Syamsiar S. Russeng¹, Fridawaty Rivai³

¹Department of Occupational Safety and Health, Faculty of Public Health, Hasanuddin University, Indonesia

²Department of Health Promotion and Behavioral Science, Faculty of Public Health, Hasanuddin University, Indonesia

³Department of Hospital Management and Administration, Faculty of Public Health, Hasanuddin University, Indonesia

*Corresponding author E-mail: adhinda.putripratiwi@gmail.com

Abstract

The purpose of this study was to examine the factors affecting the occurrence of musculoskeletal disorders in female seaweed workers in Mangarabombang District, Takalar Regency. The design used in this study used a mix method with a sample size of 103 quantitative respondents and 10 qualitative informants. Quantitative data obtained through direct observation by filling out the questionnaire, then the data is processed using SPSS, then presented in tabular form through its explanation, while qualitative data is focus group discussion and obtained through interviews, then interpreted and presented in narrative form. The results showed that the factors that influence musculoskeletal disorders are age (0,0001), anthropometry of the upper arm (0,008), anthropometry of the forearm, wrist and back (0,0001), BMI (0,007), length of work (0,001), workload (0,0001), work posture (0,0001), then it was discovered that workload was the most influential variable on musculoskeletal disorders (OR = 58,098), then the informants supported that age, anthropometry, length of work, workload and work posture were the causes of musculoskeletal disorders among female seaweed workers in Mangarabombang District, Takalar Regency. There is a significant relationship between age, anthropometry of the upper arm, anthropometry of the forearm, wrist and back, BMI, length of work, workload, work posture, then workload is the variable that most influential variable on musculoskeletal disorders.

Keywords: MSDs, BMI, length of work, workload, work posture.

1. Introduction

Work-related accidents and illnesses can have a profound impact on individuals and their families. Potential hazards in the workplace can pose risks that affect the health of workers (ILO, 2013). One of the hazards in the workplace is causing health problems, in which ergonomic factors are one of the causes (Rachmawati, 2019). Work processes that are not ergonomic can cause musculoskeletal disorders (Tarwaka, 2015).

Work-related musculoskeletal disorders (MSDs) refer to injuries or disorders of the muscles, nerves, tendons, joints, cartilage, and spinal discs associated with exposure to risk factors in the workplace (Maimaiti et al., 2019). MSDs are a significant problem for workers (Prawira et al., 2017).

These disruptions are associated with social and economic costs, ultimately impacting the quality of life (Liao et al., 2016). Work-related musculoskeletal disorders affect the back, neck, shoulders, and upper and lower limbs. They cover up damage or abnormalities in joints or other tissues (European Agency for Safety and Health at Work, 2020).

In 2018 the World Health Organization (WHO) stated that musculoskeletal disorders were the second highest cause in the world (World Health Organization, 2018). The prevalence of musculoskeletal diseases in Indonesia based on data that has been diagnosed by health personnel is 11.9% and based on diagnosis or a symptom is 24.7%. The highest prevalence of musculoskeletal diseases is among workers such as farmers, fishermen, and laborers, which are 31.2% (Risksedas, 2013). The informal sector business is one business that has a very high health risk. The workforce in 2000 was 95,650,691 people, of which 70–80% was in the informal business sector. Informal sector workers are the population that does not get health services, especially occupational health (Icsal et al., 2016).

One of the workers in the informal sector is a seaweed farmer. Currently, the seaweed industry is one of the commodities that attract many people to enter the world of work. Seaweed cultivation has recently become the main occupation of farmers in many coastal areas including Takalar District, South Sulawesi. The district produced 996,550 tonnes in 2017 from a harvested area of around 17,448 hectares (Thamrin et al., 2020). The work process of seaweed cultivation for seaweed workers in Takalar Regency is still manual handling; in this case, it includes seeding, planting, and harvesting (drying). Traditionally, farmers rely on their feet and hands when doing their work (Faujijah, 2020).

Many factors can influence the occurrence of musculoskeletal disorders as has been done in previous studies, such as individual factors, occupational factors, environmental factors, and psychosocial factors (Tarwaka, 2015; Kuswana, 2014). Research conducted by Wijnhvon, et al (2020) shows that the prevalence rate of musculoskeletal problems for all occupational groups of women is greater than men. This is due to physiological factors of muscle strength in women, which are around 2/3 the muscle strength of men. Work posture can trigger muscle complaints in workers (Tarwaka, 2015). Things that are often done by humans in doing work include standing, sitting, squatting, bending, walking, and others (Nurmianto, 2004). Work postures that are not ergonomic when done at work can often cause problems for workers such as complaints of the musculoskeletal muscles felt by workers (Tarwaka, 2015).

Based on the background description above, the author wants to examine more deeply about ergonomic problems, in this case, to find out what affects the occurrence of musculoskeletal disorders in female seaweed workers in Takalar Regency so that it can be a prevention for seaweed workers from experiencing musculoskeletal disorders.

2. Methods

The method used in this research is mixed methods. Collecting quantitative data using this type of analytic observational research with a cross-sectional approach, while for qualitative data collection using the FGD method and obtained through interviews. The purpose of this study was to examine the factors affecting the occurrence of musculoskeletal disorders in female seaweed workers in Mangarabombang District, Takalar Regency.

The population in this study was 138 people. The sample in this study were 103 quantitative respondents and 10 qualitative informants. Quantitative data were obtained through direct observation by filling out questionnaires, then the data were processed using SPSS, then presented in tabular form through explanations, while qualitative data was in the form of focus group discussions and obtained through interviews, then interpreted and presented in narrative form.

3. Results and Discussion

The results of this study provided information on the characteristics of female seaweed workers who had musculoskeletal disorders

Tabel 1. Results of Univariate Test Analysis Based on the Variables Researched on Female Seaweed Workers

Variable	Frequency (n)	Percent (%)
Age		
Young	25	24,3
Old	78	75,7
Anthropometry (upper arm)		
Appropriated	38	36,9
Unappropriated	65	63,1
Anthropometry (forearm)		
Appropriated	35	34,0
Unappropriated	68	66,0
Anthropometry (Wrist)		
Appropriated	31	30,1
Unappropriated	72	69,9
Anthropometry (Back)		
Appropriated	30	29,1
Unappropriated	73	70,9
BMI		
Normal	51	49,5
Fat	52	50,5
Length of worth		
Qualified	64	62,1

Unqualified	39	37,9
Workload		
Mild	47	45,6
Heavy	56	54,4
Work Posture		
Ergonomic	39	37,9
Not Ergonomic	64	62,1
Musculoskeletal Disorders		
Mild	29	28,2
Heavy	74	71,8

Source: Primary Data, 2020

Table 1 above shows that from 103 samples, female seaweed workers who are older, are 78 people (75.7%). Anthropometric measurements on the upper arm were more in the unsuitable category as many as 65 people (63.1%), for the forearm the most were in the unsuitable category of 68 people (66.0%), the wrist part was the most in the unsuitable category. as many as 72 people (69.9%), and on the measurement of the back there were 73 people (70.9%) who were at most in the unsuitable category. BMI of female seaweed workers is more in the fat category, namely as many as 52 people (50.5%). The length of work done by the farmers is more, that meets the requirements, namely 64 people (62.1%). The most moderate workload felt by female seaweed workers was 56 people (54.4%). Work posture is more in the non-ergonomic category of 64 people (62.1%) and MSDs was mostly in the heavy category of 74 people (71,8%).

Tabel 2. The Results of the Bivariate Analysis

Variabel	Musculoskeletal Disorders				Total		p-value
	Mild		Heavy		n	%	
	n	%	N	%			
Age							
Young	22	88,0	3	12,0	25	100	0,0001
Old	7	9,0	71	91,0	78	100	
Anthropometry (upper arm)							
Appropriated	17	44,7	21	55,3	38	100	0,008
Unappropriated	12	18,5	53	81,5	65	100	
Anthropometry (forearm)							
Appropriated	18	51,4	17	48,6	35	100	0,0001
Unappropriated	11	17,5	57	83,8	68	100	
Anthropometry (wrist)							
Appropriated	17	54,8	14	45,2	31	100	0,0001
Unappropriated	12	16,7	60	83,3	72	100	
Anthropometry (Back)							

Appropriated	19	63,3	11	36,7	30	100	0,0001
Unappropriated	10	13,7	63	86,3	73	100	
BMI							
Normal	21	41,2	30	58,8	51	100	0,007
Fat	8	15,4	44	84,6	52	100	
Length of worth							
Qualified	26	40,6	38	77,2	64	100	0,001
Unqualified	3	7,7	36	84,8	39	100	
Workload							
Mild	25	53,2	22	46,8	47	100	0,0001
Heavy	4	7,1	52	92,9	56	100	
Work Posture							
Ergonomics	19	48,7	20	51,3	39	100	0,0001
Not Ergonomics	1	1,6	63	98,4	64	100	

Source: Primary Data, 2020

Table 2 shows the chi-square test results show that there is a significant influence between musculoskeletal disorders and age, upper arm anthropometry, forearm anthropometry, wrist and back, BMI, length of work, workload, and work posture.

Table 3. Multivariate Analysis

Variables	B	S.E.	Wald	Df	Sig.	Exp(B)
KAT_age	3.951	1.378	8.219	1	.004	52.011
KAT_workposture	2.246	1.084	4.296	1	.038	9.452
KAT_workload	4.062	1.420	8.179	1	.004	58.098
Constant	-4.630	1.444	10.27	1	.001	.010
			9			

Source: Primary Data, 2020

Table 3 show, it is known that workload is the most influential variable on musculoskeletal disorders with an OR value of 58.098, which means that the risk workload variable is 58.098 times greater and has an effect on musculoskeletal disorders in female seaweed workers in Mangarabombang District, Takalar Regency.

Musculoskeletal disorders are one of the most common and costly occupational (Saberipour et al., 2019). Many different factors contribute to the risk of MSD (Panel on Musculoskeletal Disorders the Workplace, & Institute of Medicine (2001). Musculoskeletal disorders system will usually be felt in workers aged 35 years and the level of complaints felt can continue to increase with age (Tarwaka, 2015). This is the same as what was obtained from informants during the FGD:

“Because I’m old, my body hurts sometimes ...”

“Yes, it usually hurts my back and calves when I work...”

Symptoms are generally experienced at age 35 years and continue to increase. This is because, during the middle age phase, muscle strength and endurance tend to decrease (Arifin et al., 2020). The

body mass index of workers also has the risk of developing pain in their body parts if a person is overweight because the load on the joints for weight will be increased, thus allowing pain in the body, especially at the waist. This is in line with research conducted by Sumardiyono et al (2018) on written batik workers who found that there was a significant relationship between IMT and musculoskeletal disorders.

Anthropometry of workers' bodies can also affect the occurrence of musculoskeletal disorders, such as research conducted by Susana (2016) on fish drying workers which found that anthropometry is not in accordance with the work aids used by drying workers causes musculoskeletal disorders. In-depth interviews, the informant also revealed that:

"When I do the nursery I have to sit on a short stool, so it keeps my legs bent for hours..."

Occupational diseases arise because of work relationships or those caused by work and work attitudes (Suryadi & Rachmawati, 2020). The working posture done by the female seaweed workers is bending over and arms pointing upwards and resting on her legs. The position of the head is down and the back is slightly bent. This kind of work posture is carried out for hours by female seaweed workers in the nursery. This is consistent with the results of in-depth interviews and FGDs, where the informant said:

"All of us who work on the nursery process are short stools for long hours..."

"In the nursery process, usually the position of the arms is bent and then rotated while doing the seeding, and this position is carried out continuously..."

Women seaweed workers usually do their work for more than 8 hours. This is because the wages received will be more when workers can maximize the yield from tying seaweed seeds. The addition of working hours can reduce productivity, worker efficiency, cause fatigue, and can lead to illness and accidents (Suma'mur, 2009). These results were obtained from informants during the FGD:

"In a day I used to do my work from morning to evening..."

The physical load that is justified is that it does not exceed 30-40% of the maximum work capacity of the workforce in 8 hours a day by observing the prevailing working hours regulations. More heavy the load, the shorter the work time⁽²²⁾. The results of the FGD showed that female seaweed workers have a heavy workload because in addition to working as a seaweed workers she is also a housewife who has to take care of her home and family. As the informant said:

"Before this job, I had to clean my house first, prepare clothes for my children who wanted to go to school, I also had to cook for my family first. After everything was finished I continued to go to work as a seaweed farmer..."

This is in line with the research conducted by Tjahjuningtyas (2019) on informal workers who found that there is a significant relationship between workload and musculoskeletal disorders. MSDs can be reduced in various ways, for example, by paying attention to work posture while working, increasing physical fitness, adjusting diet, exercise, and adequate rest (Hanif, 2020).

4. Conclusion

13
There is a significant relationship between age, anthropometry of the upper arm, anthropometry of the forearm, wrist, and back, BMI, length of work, workload and work posture, then it was discovered that workload was the most influential variable on musculoskeletal disorders. Then the informants supported that age, anthropometry, length of work, workload, and work posture were the causes of musculoskeletal disorders among female seaweed workers in Mangarabombang District, Takalar Regency. It is recommended for seaweed workers to notice to work posture and rest time while working.

References

1. ILO. (2013). Occupational Safety and Health: Means for Productivity. Available from: https://www.ilo.org/wcmsp5/groups/public/---ed_protect/---protrav/---safework/documents/publication/wcms_301214.pdf
2. Rachmawati, D. (2019). Musculoskeletal Disorders and Its Related Factors among Workers in Circulator Loom Unit. *Indonesian Journal Occupation Safe Health*, 2019, 8(3):265.
3. Tarwaka, H. (2015). Ergonomi Industri Dasar-Dasar Pengetahuan Ergonomi dan Aplikasi di Tempat Kerja Edisi II. Surakarta: Harapan Press.
4. Maimaiti, N., Wang, J., Jin, X., Wang, S., Qin, D., He, L., & Yang, Q. (2019). Cervical musculoskeletal disorders and their relationships with personal and work-related factors among electronic assembly workers. *Journal of safety research*, 71, 79-85.
5. Prawira, M. A., Yanti, N. P. N., Kurniawan, E., & Artha, L. P. W. (2017). Factors related musculoskeletal disorders on students of Udayana University on 2016. *Journal of Industrial Hygiene and Occupational Health*, 1(2), 101-118.
6. Liao, J. C., Ho, C. H., Chiu, H. Y., Wang, Y. L., Kuo, L. C., Liu, C., & Kuo, J. R. (2016). Physiotherapists working in clinics have increased risk for new-onset spine disorders: a 12-year population-based study. *Medicine*, 95(32).
7. European Agency for Safety and Health at Work. (2020). *Musculoskeletal Disorders*. Available from: <https://osha.europa.eu/en/themes/musculoskeletal-disorders>.
8. World Health Organization. (2018). *Musculoskeletal Disorders*. Geneva: WHO.
9. Riskesdas. (2013). Laporan Hasil Riset Kesehatan Dasar Nasional. Badan Penelitian dan Pengembangan Kesehatan. Available from: <https://www.litbang.kemkes.go.id/laporan-riset-kesehatan-dasar-riskesdas/>
10. Icsal, M., Sabilu, Y., & Pratiwi, A. D. (2016). Faktor yang berhubungan dengan keluhan musculoskeletal disorders (msds) pada penjahit wilayah pasar panjang kota kendari tahun 2016. (*Jurnal Ilmiah Mahasiswa Kesehatan Masyarakat*), 1(2).
11. Thamrin, Y., Wahyu, A., Russeng, S. S., Wahyuni, A., & Hardianti, A. (2020). Ergonomics and musculoskeletal disorders among seaweed workers in Takalar Regency: A mixed method approach. *Medicina Clínica Práctica*, 3, 100110.
12. Faujiyah, F. (2020). Studi Prevalensi Keluhan Muskuloskeletal Pada Petani Indonesia. *Jurnal TEDC*, 14(2), 113-119.
13. Nurmiyanto, E. (2004). *Ergonomi konsep dasar dan aplikasinya edisi kedua*. Surabaya: Guna Widya.
14. Kuswana, W. R. (2014). *Ergonomi dan Keselamatan dan Kesehatan Kerja*. Jakarta: Rusda.
15. Wijnhoven, H. A., De Vet, H. C., & Picavet, H. S. J. (2006). Prevalence of musculoskeletal disorders is systematically higher in women than in men. *The Clinical journal of pain*, 22(8), 717-724.
16. Saberipour, B., Ghanbari, S., Zarea, K., Gheibizadeh, M., & Zahedian, M. (2019). Investigating prevalence of musculoskeletal disorders among Iranian nurses: a systematic review and meta-analysis. *Clinical Epidemiology and Global Health*, 7(3), 513-518.
17. Panel on Musculoskeletal Disorders, A., the Workplace, & Institute of Medicine. (2001). *Musculoskeletal disorders and the workplace: Low back and upper extremities*. National Academy Press.
18. Arifin, A. N., Rahayu, S. R., & Rustiana, E. R. (2020). Analysis of Postures Towards

14671

- Musculoskeletal Disorders Experienced By Batik Artisans of Sukamaju Giriloyo Yogyakarta. *Public Health Perspective Journal*, 5(1).
19. Sumardiyono, S., Wijayanti, R., Probandari, A., Larasati, G., Dewi, A. K., & Ardiani, R. F. (2018). Faktor Risiko Kesehatan Kerja Pada Pekerja Pembatik TuliS. *Prosiding SNST Fakultas Teknik*, 1(1).
 20. Susana, I. G. B. (2016). Rancangan ruang pengering berbasis ergonomi menurunkan keluhan muskuloskeletal perajin ikan. *Dinamika Teknik Mesin*, 6(1).
 21. Suryadi, I., & Rachmawati, S. (2020). Work Posture Relations With Low Back Pain Complaint on Partners Part of Pt'x'manufacture Of Tobacco Products. *Journal of Vocational Health Studies*, 3(3), 126-130.
 22. Suma'mur, P. K. (2009). Higiene perusahaan dan kesehatan kerja (HIPERKES). *Jakarta: Sagung Seto*, 116-32.
 23. Vithalbhai, V. S. . (2020). Financial Performance of Banks in India: A Study of Selected Private Sector Banks .*Journal of Advanced Research in Economics and Administrative Sciences*, 1(1), 45-52.
 24. Tjahayuningtyas, A. (2019). Faktor yang Mempengaruhi Keluhan Musculoskeletal Disorders (MSDs) pada Pekerja Informal. *The Indonesian Journal of Occupational Safety and Health*, 8(1), 1-10.
 25. Hanif, A. (2020). Hubungan Antara Umur Dan Kebiasaan Merokok Dengan Keluhan Musculoskeletal Disorders (Msd) Pada Pekerja Angkat Angkut Ud Maju Makmur Kota Surabaya. *Medical Technology and Public Health Journal*, 4(1), 7-15.
 26. Borela, V. T. ., & Apolinar , D. A. D. . (2020). Banana Peel Cellulose Nanofibers (CNFs) as Retrofitting Material to Soy-Protein in Manufacturing Biodegradable Food Packaging. *Journal of Scientific Research in Medical and Biological Sciences*, 1(1), 10-29.

ORIGINALITY REPORT

%**9**

SIMILARITY INDEX

%**8**

INTERNET SOURCES

%**5**

PUBLICATIONS

%**0**

STUDENT PAPERS

PRIMARY SOURCES

1

event.ners.unair.ac.id

Internet Source

%**2**

2

www.hindawi.com

Internet Source

%**1**

3

e-journal.unair.ac.id

Internet Source

%**1**

4

onlinelibrary.wiley.com

Internet Source

%**1**

5

"IUNS. 21st International Congress of Nutrition. Buenos Aires, Argentina, October 15-20, 2017: Abstracts", *Annals of Nutrition and Metabolism*, 2017

Publication

%**1**

6

Fatemeh Rahimi, Khadijeh Kazemi, Shahla Zahednejad, Daniel López-López, César Calvo-Lobo. "Prevalence of Work-Related Musculoskeletal Disorders in Iranian Physical Therapists: A Cross-sectional Study", *Journal of Manipulative and Physiological Therapeutics*,

%**1**

2018

Publication

7	www.utccmbaonline.com Internet Source	% 1
8	www.researchgate.net Internet Source	<% 1
9	www.osteopathic-services.co.uk Internet Source	<% 1
10	www.tandfonline.com Internet Source	<% 1
11	jcdr.in Internet Source	<% 1
12	link.springer.com Internet Source	<% 1
13	Uly Agustine, Maria Endang, Era Kale, Emilia Akoit. "Risk Factors of Malaria Events among Pregnant Women in East Sumba Regency, Indonesia", Research Square, 2020 Publication	<% 1
14	roderic.uv.es Internet Source	<% 1
15	Ralalicia Limato, Rukhsana Ahmed, Amelia Magdalena, Sudirman Nasir, Fiona Kotvojs. "Use of most significant change (MSC) technique to evaluate health promotion training	<% 1

of maternal community health workers in Cianjur
district, Indonesia", Evaluation and Program
Planning, 2018

Publication

EXCLUDE QUOTES ON

EXCLUDE ON

BIBLIOGRAPHY

EXCLUDE MATCHES

< 5

WORDS