

Progesterone Receptor Expression and Score Differences in Determining Grade and Subtype of Meningioma

by Upik Miskad

FILE	E_DIFFERENCES_IN_DETERMINING_GRADE_AND_SUBTYPE_OF_MENINGIOMA.PDF (918.23K)	WORD COUNT	3714
TIME SUBMITTED	27-DEC-2020 09:27AM (UTC+0700)	CHARACTER COUNT	18852
SUBMISSION ID	1481407505		



Progesterone Receptor Expression and Score Differences in Determining Grade and Subtype of Meningioma

Jeni Poniman¹ Muh Husni Cangara¹ Cahyono Kaelan¹ Upik A Miskad¹ Gunawan Arsyadi¹
Mahmud Ghaznawe² Dasril Daud³ Andi Ihwan⁴ Rohadi Muhammad Rosyidi⁵

21

¹Department of Anatomical Pathology, Faculty of Medicine, Hasanudin University, Makassar, Indonesia

²Faculty of Medicine, Muhammadiyah University, Makassar, Indonesia

³Department of Pediatrics, Faculty of Medicine, Hasanuddin University, Makassar, Indonesia

⁴Department of Neurosurgery, Faculty of Medicine, Hasanuddin University Makassar, Indonesia

⁵Department of Neurosurgery, Faculty of Medicine, Mataram University, West Nusa Tenggara Province General Hospital, Mataram, Indonesia

11

Address for correspondence Rohadi Muhammad Rosyidi, Department of Neurosurgery Medical Faculty of Mataram University, West Nusa Tenggara Province General Hospital, Mataram, 83127, Indonesia (e-mail: rha.ns2010@gmail.com).

J Neurosci Rural Pract

Abstract

6

Background Meningioma is the most common brain tumor in which therapy and monitoring depend on the histopathological grade (World Health Organization [WHO] Grade). Progesterone receptor (PR) expression was reported positive in meningeothelial cells and meningiomas with various degrees of positivity. We evaluated PR expression to determine its correlation with WHO Grade and each subtype of meningioma.

Materials and Methods This study used 10 samples of paraffin block that were diagnosed as meningioma and classified into WHO Grade I, II, and III. The paraffin blocks were sectioned in 5 µm thickness and immunohistochemically stained with the anti-PR antibody.

Results PR expression was found positive in WHO Grade I and II groups, but negative in WHO Grade III group with the score of +2 found in clear cell and atypical subtype. These results were statistically significant with p -value < 0.05.

Conclusion PR can be used as an additional marker to determine WHO Grade and subtype of meningioma.

Keywords

- ▶ progesterone receptor
- ▶ meningioma
- ▶ WHO Grade

Introduction

Meningioma is the most frequent brain tumor originating from meningeothelial cells in the arachnoid layer with an incidence reported in the USA found in 36% in adults and 2.8% in children with a higher incidence in women. Meningioma is classified in grade I, II, and III based on World Health Organization (WHO) 2016 criteria. The incidence of each grade is 90%, 20 to 25%, and 1 to 6%, respectively.^{1,2} Grade I meningioma generally could be treated with surgery followed by routine observations. However, Grade II meningioma needs a more complex approach such as subtotal

resection, and radiotherapy must be done for the treatment of grade III meningioma. Besides therapy, monitoring after therapy also depends on the extent of tumor resection and grading.^{3,4} The WHO Grade is also the most useful predictor of recurrence, where Grade I, II, and III have recurrence 7 to 25%, 29 to 52%, and 50 to 94% in 20 years, respectively.²

It is important to determine the appropriate grade of meningioma based on WHO 2016 criteria to determine appropriate therapy and monitoring and to determine the patient's prognosis. However, there are some borderline cases that cannot meet the minimum criteria for a particular grade, so the additional marker is needed.

9

DOI <https://doi.org/10.1055/s-0040-1714043>
ISSN 0976-3147.

©2020 Association for Helping Neurosurgical Sick People

License terms



2 Progesterone is an essential regulator of female reproductive function in the uterus, ovary, mammary gland, and brain, which also has an important role in nonreproductive tissues such as the cardiovascular system, bone, and central nervous system.⁵ Progesterone receptor (PR) expression is reported to be found positive in 50 to 80% meningioma³ with various degrees of expression, generally inversely related to the grade of meningioma, Grade I has a positive expression,⁶⁻⁹ while Grade II and III have negative expressions or only show focal positive expressions.³ So, PR can be useful in borderline cases.^{3,10}

Based on the importance of clinical application and role⁸ in biological concepts, a study was conducted to evaluate PR expression in Meningioma WHO Grade I, II, and III.

18 Materials and Methods

This study was an analytic-observational study with a cross-sectional design, using paraffin blocks of central nervous system tumor that were diagnosed as meningioma in Anatomical Pathology Laboratories of DR. Wahidin Sudirohusodo Hospital, and Sentra Diagnostik Patologia, Makassar. Samples were grouped as Meningioma WHO Grade I, II, and III based on WHO 2016 grading¹² system. This study was ethically approved by the Health Research Ethics Committee of Faculty of Medicine, Hasanuddin University, (457/H4.8.4.5.31/PP36-KOMETIK/2018).

Immunohistochemical Staining²⁶ PR

Paraffin blocks were sectioned in 5 μ m thickness by microtome and mounted on Poly-L-lysine¹⁶ object-glass. These tissue sections were deparaffinized¹⁶ with xylene and rehydrated with gradient alcohol. The endogenous peroxidase activity of tissue was blocked by 3% hydrogen peroxide for 10 minutes. The tissue sections were treated with retrieval antigen solution in the microwave for 20 minutes and protein block for 10 minutes, incubated with primary antibody anti-PR (DB-Biotech) for 30 to 60 minutes that was detected with secondary antibody for 10 minutes and conjugated with horseradish peroxidase for 15 minutes. The enzymes were²⁵ labeled with 3,3'-diaminobenzidine for 10 minutes. The tissue sections were then counterstained with Mayer's hematoxylin, and covered with Entellan and deck glass.

Immunohistochemical Staining Interpretation

PR expression is an accumulation of PR protein at the nucleus that is stained brown as a positive expression by immunohistochemical staining. The expression was scored semiquantitatively based on the staining¹³ intensity, a score of stained nuclei of the tumor cells. (0 = no staining, 1+ = staining in \leq 10% nuclei, 2+ = staining in 11–50% nuclei, 3+ = staining in 51–80% nuclei and, 4+ = staining in³³ 81–100% nuclei). PR expression was considered negative if scored as 0 and 1+ (\rightarrow Fig. 1), and positive if scored as 2+, 3+, and 4+ (\rightarrow Fig. 2)^{11,12}.




Fig. 1 Negative progesterone receptor (PR) expression. (A) PR score 0, (B) PR score 1+ (Anaplastic Meningioma, World Health Organization [WHO] Grade III). (400X)




Fig. 2 Positive progesterone receptor (PR) expression. (A) PR score 2+ (Atypical Meningioma, World Health Organization [WHO] Grade II), (B) PR score 3+ (Psammomatous Meningioma, WHO Grade I), (C) PR score 4+ (Meningothelial Meningioma, WHO Grade I). (400X)

14

Statistical Analysis

The results of this study were analyzed with SPSS (Statistical Package for Social Sciences) version 17 (IBM Company). The correlation of PR expression with Meningioma WHO Grade I, II, and III and its subtypes was analyzed with Mann-Whitney U and chi-squared tests.

Results

In this study, 70 samples were collected from the Anatomical Pathology Laboratories of DR. Wahidin Sudirohusodo Hospital and Sentra Diagnostik Patologia Makassar from January 2012 to June 2018 that were diagnosed as Meningioma WHO Grade I, II, and III as many as 27, 23, and 20 samples, respectively. The subtypes of each grade were Meningothelial, Fibrous, Transitional, Psammomatous and Microcystic; Chordoid, Clear Cell and Atypical; and Papillary and Anaplastic (Table 1).

The samples in this study consisted of 24 men and 45 women with a mean age of 49.26 (from the range of 17–98) years old (Table 1). The mean age of each Meningioma WHO Grade I, II, and III was 49.15, 47.3, and 51.65 years old, respectively, and the mean age with positive and negative PR expression was 48.3 and 51.65 years old, respectively. The distribution of sex and age is not significantly related to WHO Grade and PR expression (chi-squared test and one-way analysis of variance; $p > 0.05$). (Table 2)

PR expression scores and PR expression in each Meningioma WHO Grade I, II, and III were 4+ and 3+ (positive 100%), 4+ and 2+ (positive 100%), and 1+ and 0 (negative 100%), respectively. These data indicates that there were differences in PR expression scores between Meningioma WHO Grade I and II, between WHO Grade I and III, and between WHO Grade II and III (Mann-Whitney U test; $p = 0.000$ [$p < 0.01$]), and there were different PR expression in Meningioma WHO Grade I, II, and III (chi-squared test; $p = 0.000$ [$p < 0.05$]). PR expression scores and PR expression based on subtypes of each Meningioma WHO Grade I, II, and III were Meningothelial, Fibrous, Transitional, and Microcystic

4+ (positive), and Psammomatous 4+ and 3+ (positive); Clear Cell and Atypical 2+ (positive), and Chordoid 4+ (positive); and Papillary 1+ and 0 (negative), and Anaplastic 1+ and 0 (negative). This data also showed that there were differences in PR expression scores and PR expression between each subtype of Meningioma WHO Grade I, II, and III (chi-squared test; $p = 0.000$ [$p < 0.05$]) (Table 3).

20

Table 1 General characteristics

Characteristics	n	%
Gender	Male	25 (35.7)
	Female	45 (64.3)
WHO Grade	I	27 (38.6)
	II	23 (32.9)
	III	20 (28.6)
Subtypes	Meningothelial	6 (8.6)
	Fibrous	6 (8.6)
	Mixed	7 (10)
	Psammomatous	5 (7.1)
	Microcystic	3 (4.3)
	Chordoid	7 (10)
	Clear cell	2 (2.9)
	Atypical	14 (20)
	Papillary	2 (2.9)
PR Score	0	13 (18.6)
	1+	7 (10.0)
	2+	16 (22.9)
	3+	2 (2.9)
	4+	32 (45.7)
PR Expression	Positive	50 (71.4)
	Negative	20 (28.6)

Abbreviations: PR, progesterone receptor; WHO, World Health Organization.

Table 2 Comparison between WHO Grade of Meningioma and PR expression to gender and age characteristics

	WHO Grade			p-Value	PR Expression		p-Value
	I (%)	II (%)	III (%)		Positive (%)	Negative (%)	
Gender				0.174 ^a			0.229 ^a
Male	6 (22.22)	10 (43.48)	9 (45)		16 (32)	9 (45)	
Female	21 (77.78)	13 (56.52)	11 (55)		34 (68)	11 (55)	
Age				0.597 ^b			0.365 ^b
Mean	49.15	47.30	51.65		48.30	51.65	
Median	49	48	51.50		48.5	51.50	
Minimum	26	26	17		26	17	
Maximum	89	68	98		89	98	

Abbreviations: PR, progesterone receptor; WHO, World Health Organization.

^aChi-squared test.

^bOne-way analysis of variance test.

Table 3 Comparison between PR score and PR expression to WHO Grade of Meningioma and its subtypes characteristics

	PR score					p-Value	PR expression		p-Value
	4+ (%)	3+ (%)	2+ (%)	1+ (%)	0 (%)		Positive (%)	Negative (%)	
WHO Grade						0.000 ^a			0.000 ^b
I	25 (92.59)	2 (7.41)	0	0	0		27 (100)	0	
II	7 (30.43)	0	16 (69.57)	0	0		23 (100)	0	
III	0	0	0	6 (30)	14 (70)		0	20 (100)	
Subtypes						0.000 ^b			0.000 ^b
Meningothelial	6 (100)	0	0	0	0		6 (100)	0	
Fibrous	6 (100)	0	0	0	0		6 (100)	0	
Mixed	7 (100)	0	0	0	0		7 (100)	0	
Psammomatous	3 (60)	2 (40)	0	0	0		5 (100)	0	
Microcystic	3 (100)	0	0	0	0		3 (100)	0	
Chordoid	7 (100)	0	0	0	0		7 (100)	0	
Clear Cell	0	0	2 (100)	0	0		2 (100)	0	
Atypical	0	0	14 (100)	0	0		14 (100)	0	
Papillary	0	0	0	1 (50)	1 (50)		0	2 (100)	
Anaplastic	0	0	0	5 (27.78)	13 (72.22)		0	18 (100)	

Abbreviations: PR, progesterone receptor; WHO, World Health Organization.

^aMann-Whitney U test.

^bChi-square test.

Discussion

The risk of having meningioma increases with age. The tendency of women to have Grade I meningioma is higher than men, while for Grade II and III meningioma the tendency of men is higher than women. Because of the differences in gender, one of the other proposed etiology of meningioma is hormonal.² But based on the data of age and gender in this study, there were no significant differences related to the distribution in WHO Grade I, II, and III meningioma and also no significant differences with PR expression. So that it supports the general etiologies of meningioma that are idiopathic and sporadic rather than the proposed one which is hormonal.^{2,13} This is in line with the research conducted by Mukhopadhyay et al, which reported there was no relationship between age and WHO Grade⁹ and also another research conducted by Roser et al, which reported there was no relationship between sex and age with the PR expression.¹⁰

The data characteristics of this study based on WHO Grade I, II, and III were 27 (38.6%), 23 (32.9%), and 20 (28.6%) samples, respectively, with each subtype consisting 6 Meningothelial (8.6%), 6 Fibrous (8.6%), 7 Transitional (10%), 5 Psammomatous (7.1%), and 3 Microcystic (4.3%) samples (Grade I); 7 Chordoid (10%), 2 Clear Cell (2.9%), and 14 Atypical (20%) samples (Grade II); and 2 Papillary (2.9%) and 18 Anaplastic (25.7%) samples (Grade III). This is in accordance with the nature of meningioma that is a slow-growing neoplasm and generally benign so that the incidence of Meningioma WHO Grade I is more common than WHO Grade II and III, with the subtypes commonly are Meningothelial and Transitional.^{2,14}

30

Progesterone is an essential regulator of reproductive function and also plays an important role in nonreproductive tissues such as the cardiovascular system, bone, and central nervous system.⁵ In the central nervous system, progesterone is a neurosteroid and has a variety of important roles. Besides that, progesterone is also an indirect precursor of cortisol. Cortisol itself has a role in the response to stress and has an anti-inflammatory effect.¹⁵ Progesterone is normally expressed in non-neoplastic meningothelial cells in small amounts, which then increases when there is cell proliferation, such as in meningioma, and decreases when there are changes in cell differentiation.¹⁶ However, this progesterone expression is not under the influence of the estrogen as in reproductive organs.^{9,12,17} Meningothelial cells primarily line the edges of arachnoid granulation so-called arachnoid cap cells,¹⁸ and this arachnoid granulation plays a role in the absorption of cerebrospinal fluid into dural sinuses.^{17,19} The role of PR in the regulation of the fluid transportation is not yet clear, but there are some indications that steroid hormones can play a role in which glucocorticoid dexamethasone is reported to increase the absorption of cerebrospinal fluid into the sagittal sinus. In addition, metabolites from progesterone and mineralocorticoid deoxycorticosterone have been reported to modulate the function of the receptor-operated chloride channel in the nervous system, which is associated in fluid and ion transportation.¹⁷

PR expression in meningioma is inversely related to WHO Grade, which in WHO Grade I there is an increase of 50 to 29% PR expression; so it shows positive PR expression while in WHO Grade II and III there is a loss of 60 to 80% and 80 to 90% PR expression, respectively; so it shows focal or negative

PR expression.¹³ Almost all WHO Grade III shows negative expression.² In this study, obtained PR expression scores 4+ and 3+ were found in WHO Grade I, 4+ and 22 were found in WHO Grade II, and 1+ and 0 were found in WHO Grade III, with 100% positive PR expression in WHO Grade I and II, and 100% negative in WHO Grade III. These data showed significant difference in PR expression scores and PR expression in Meningioma WHO Grade I, II, and III. Based on each subtype, a score of 4+ was obtained 100% in Meningeal, Fibrous, Transitional, and Microcystic (WHO Grade I), 60% in Psammomatous (WHO Grade I) and 100% in Chordoid (WHO Grade II); 3+ was obtained 40% in Psammomatous; 2+ was obtained 100% in Clear Cell and Atypical (WHO Grade II); 1+ was obtained 50% in Papillary and 27.78% in Anaplastic (WHO Grade III); and 0 was obtained 50% in Papillary and 72.22% in Anaplastic (WHO Grade III). This result is in line with the other studies, which reported that positive PR expression was found more in WHO Grade I than II and negative PR expression was found in WHO Grade III.^{6-8,10,16} In one study, Ikeri et al reported that PR expression scores decrease along with the increasing of grade in which PR expression scores of WHO Grade I were higher than II and III, and WHO Grade II was higher than III.¹⁶

Based on these data from this study, PR expression scores and PR expression can be used to help diagnose Meningioma WHO Grade I, II, and III, especially for Atypical and Anaplastic subtypes that are sometimes difficult to diagnose using 2016 WHO criteria, because there are some cases that cannot meet the minimum diagnostic criteria (borderline cases), while the other subtypes were diagnosed based on each its histopathological characteristics.

Other studies also recommended PR expression to be used as a routine marker to help assess the nature of meningioma, which can distinguish the degrees of differentiation with the value inversely related to WHO Grade, although there was the same value between WHO Grade I and II, both had a positive expression.^{7,8,10,16} Roser et al added that PR expression should be combined with other markers to help assess the nature of meningioma, and also to predict recurrence and to determine more effective follow-up strategies.¹⁰

Conclusion

We can conclude that if the PR expression is negative, then it is grouped into WHO Grade III, and if the PR expression is positive, then it is grouped into WHO Grade I and II. If the PR expression score is +2, then it is grouped in WHO grade II with Clear Cell or Atypical subtypes. Based on this, PR expression can be used as an additional marker to help determine WHO grade and several subtypes especially in borderline cases.

Highlights

- Meningioma is the most common brain tumor which therapy and monitoring depend on the histopathological grade (WHO Grade).
- Progesterone is not only an essential regulator of reproduction but also an important role in the nonreproductive organ as in the central nervous system.

- PR expression was reported positive in meningothelial cells and meningiomas with various degree of positivity.
- There was a significant correlation between PR expression to WHO Grade of meningioma and its subtypes.

Conflict of Interest

None declared.

References

- Kleinschmidt-DeMasters B, Rodríguez FJ, Tihan T, Neoplastic: meninges. In: Kleinschmidt-DeMasters B, Rodríguez FJ, Tihan T, eds. *Diagnostic Pathology: Neuropathology*. 2nd edition. Philadelphia: Elsevier; 2016:450-467
- Perry A, Louis DN, Budka H, et al. Meningioma. In: Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, eds. *WHO Classification of Tumours of the Central Nervous System*. Revised 4th edition. Lyon: International Agency for Research on Cancer (IARC); 2016:232-237
- Perry A, Meningiomas. In: Perry A, Brat DJ, eds. *Practical Surgical Neuropathology: A Diagnostic Approach*. 1st edition. Philadelphia: Churchill Livingstone an affiliate of Elsevier Inc; 2010:185-217
- Goldbrunner R, Minniti G, Preusser M, et al. EANO guidelines for the diagnosis and treatment of meningiomas. *Lancet Oncol* 2016;17(9):e383-e39110.1016/S1470-2045(16)30321-7[Internet]
- Scarpin KM, Graham JD, Mote PA, Clarke CL. Progesterone action in human tissues: regulation by progesterone receptor (PR) isoform expression, nuclear positioning and coregulator expression. *Nucl Recept Signal* 2009;7:e009
- Wolfsberger S, Doostkam S, Boecher-Schwarz HG, et al. Progesterone-receptor index in meningiomas: correlation with clinico-pathological parameters and review of the literature. *Neurosurg Rev* 2004;27(4):238-245
- Shayanfar N, Mashayekh M, Mohammadpour M. Expression of progesterone receptor and proliferative marker Ki 67 in various grades of meningioma. *Acta Med Iran* 2010;48(3):142-147
- Mukherjee S, Ghosh SN, Chatterjee U, Chatterjee S. Detection of progesterone receptor and the correlation with Ki-67 labeling index in meningiomas. *Neurol India* 2011;59(6):817-822
- Mukhopadhyay M, Das C, Kumari M, Sen A, Mukhopadhyay B, Mukhopadhyay B. Spectrum of meningioma with special reference to prognostic utility of ER, PR and Ki67 expression. *J Lab Physicians* 2017;9(4):308-313
- Roser F, Nakamura M, Bellinzona M, Rosahl SK, Ostertag H, Samii M. The prognostic value of progesterone receptor status in meningiomas. *J Clin Pathol* 2004;57(10):1033-1037
- El-Badawy NM, Farid RM, Nagib LN, Ibrahim RA. Role of progesterone receptor expression and proliferative activity in predicting the recurrence of meningioma. *Egypt J Pathol* 2013;76-81
- Shanthi V, Grandhi B, Mohan Rao N, Sundar Rao BS, Muram Reddy VL. Assessing the prognostic importance of ER, PR expression in meningiomas by comparing with proliferative rate using Ki67. *Indian J Pathol Res Pract*. 2017;6(2):431-434
- Perry A, Meningioma. In: Perry A, Brat DJ, eds. *Practical Surgical Neuropathology: A Diagnostic Approach*. 2nd edition. Philadelphia: Elsevier; 2018:259-298
- Perry A, Louis DN, Budka H, et al. Meningioma variants. In: Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, eds. *WHO Classification of Tumours of the Central Nervous System*. Revised 4th edition. Lyon: International Agency for Research on Cancer (IARC); 2016: 237-245
- Carisieri M, Niyonkuru C, McCullough EH, et al. Cerebrospinal fluid cortisol and progesterone profiles and

- 15 outcomes prognostication after severe traumatic brain injury. *J Neurotrauma* 2014;31(8):699–712
- 16 Ikeri NZ, Anunobi CC, Bankole OB. Progesterone receptor expression and Ki-67 labelling index of meningiomas in the Lagos university teaching hospital. *Niger Postgrad Med J* 2018;25(1):17–20 www.npmj.org [Internet]
- 17 Verhagen A, Go KG, Visser GM, Blankenstein MA, Vaalburg W. The presence of progesterone receptors in arachnoid granulations and in the lining of arachnoid cysts: its relevance to expression of progesterone receptors in meningiomas. *Br J Neurosurg* 1995;9(1):47–50
- 18 Brat DJ, Normal brain histopathology. In: Perry A, Brat DJ, eds. *Practical Surgical Neuropathology: A Diagnostic Approach*. Philadelphia: Churchill Livingstone an affiliate of Elsevier Inc; 2010 15–33
- 19 Lopes MBS, Meninges: embryology. In: Lee JH, ed. *Meningioma*. London: Springer; 2009:25–29

Progesterone Receptor Expression and Score Differences in Determining Grade and Subtype of Meningioma

ORIGINALITY REPORT

% **19**
SIMILARITY INDEX

% **10**
INTERNET SOURCES

% **15**
PUBLICATIONS

% **4**
STUDENT PAPERS

PRIMARY SOURCES

1 "Meningiomas", Springer Science and Business Media LLC, 2020 % **2**
Publication

2 Isabella dos Santos Guimares, Renata Dalmaschio, Alice Laschuk Herlinger, Klesia Pirola et al. "Chapter 1 Conventional Cancer Treatment", IntechOpen, 2013 % **2**
Publication

3 www.gssrr.org % **1**
Internet Source

4 Serge Weis, Michael Sonnberger, Andreas Dunzinger, Eva Voglmayr, Martin Aichholzer, Raimund Kleiser, Peter Strasser. "Imaging Brain Diseases", Springer Science and Business Media LLC, 2019 % **1**
Publication

5 Submitted to State Islamic University of Alauddin Makassar % **1**
Student Paper

6	journal.uin-alauddin.ac.id Internet Source	% 1
7	"ESP Abstracts 2014", Virchows Archiv, 2014 Publication	% 1
8	Madhumita Mukhopadhyay, Chhanda Das, Madhu Kumari, Ankita Sen, Bedabrata Mukhopadhyay, Biswanath Mukhopadhyay. "Spectrum of meningioma with special reference to prognostic utility of ER,PR and Ki67 expression", Journal of Laboratory Physicians, 2020 Publication	% 1
9	Submitted to Parkland College Student Paper	% 1
10	jimdc.org.pk Internet Source	% 1
11	doaj.org Internet Source	% 1
12	Submitted to Universitas Hasanuddin Student Paper	% 1
13	Shengqi Wang, Neng Wang, Yifeng Zheng, Bowen Yang, Pengxi Liu, Fengxue Zhang, Min Li, Juxian Song, Xu Chang, Zhiyu Wang. "Caveolin-1 inhibits breast cancer stem cells via c-Myc-mediated metabolic reprogramming", Cell	% 1

Death & Disease, 2020

Publication

14

microbiologyjournal.org

Internet Source

<% 1

15

bmcneurosci.biomedcentral.com

Internet Source

<% 1

16

scholarbank.nus.edu.sg

Internet Source

<% 1

17

Santarsieri, Martina, Christian Niyonkuru, Emily H. McCullough, Julie A. Dobos, C. Edward Dixon, Sarah L. Berga, and Amy K. Wagner. "Cerebrospinal Fluid Cortisol and Progesterone Profiles and Outcomes Prognostication after Severe Traumatic Brain Injury", *Journal of Neurotrauma*, 2014.

Publication

<% 1

18

repository.unhas.ac.id

Internet Source

<% 1

19

www.ruralneuropractice.com

Internet Source

<% 1

20

ocean.kisti.re.kr

Internet Source

<% 1

21

Piyapharom Intarawichian, Supattra Kumdinpitak, Sakda Waraasawapati, Chat Sumanont, Supinda Koonmee. "Extraskeletal

<% 1

myxoid chondrosarcoma: a case report high grade tumor in breast", Pathology, 2014

Publication

22

Ashraf Fakhrijou, Ali Meshkini, Sepideh Shadravan. "Status of Ki-67, Estrogen and Progesterone Receptors in Various Subtypes of Intracranial Meningiomas", Pakistan Journal of Biological Sciences, 2012

Publication

<% 1

23

apsjournals.apsnet.org

Internet Source

<% 1

24

med.uth.edu

Internet Source

<% 1

25

s3-eu-west-1.amazonaws.com

Internet Source

<% 1

26

Jian-Fa Wang, Shou-Peng Fu, Su-Nan Li, Zhan-Qing Yang, Wen-Jing Xue, Zhi-Qiang Li, Wei Wang, Ju-Xiong Liu. "Establishment and characterization of dairy cow growth hormone secreting anterior pituitary cell model", In Vitro Cellular & Developmental Biology - Animal, 2013

Publication

<% 1

27

neuro.wustl.edu

Internet Source

<% 1

28

Liong Boy Kurniawan, Endy Adnan, Budi

Mulyono. "Insulin resistance and testosterone level in Indonesian young adult males",
Romanian Journal of Internal Medicine, 2020

Publication

<% 1

29

Nancy Wang, Matthias Osswald. "Meningiomas: Overview and New Directions in Therapy",
Seminars in Neurology, 2018

Publication

<% 1

30

John C Stevenson, Serge Rozenberg, Silvia Maffei, Christian Egarter, Petra Stute, Thomas Römer. "Progestogens as a component of menopausal hormone therapy: the right molecule makes the difference",
Drugs in Context, 2020

Publication

<% 1

31

www.birpublications.org

Internet Source

<% 1

32

SUHEYLA UYAR BOZKURT.
"Immunohistochemical expression of SPARC is correlated with recurrence, survival and malignant potential in meningiomas",
Apmis, 09/2009

Publication

<% 1

33

Smitha Makkat, Robert Luybaert, Tadeusz Stadnik, Claire Bourgain et al. "Deconvolution-based Dynamic Contrast-enhanced MR Imaging of Breast Tumors: Correlation of Tumor Blood

<% 1

Flow with Human Epidermal Growth Factor Receptor 2 Status and Clinicopathologic Findings—Preliminary Results", Radiology, 2008

Publication

34

www.bmlsearch.com

Internet Source

<% 1

35

"Breast", Modern Pathology, 02/2011

Publication

<% 1

36

mafiadoc.com

Internet Source

<% 1

37

test.dovepress.com

Internet Source

<% 1

38

"Brain Tumors in Children", Springer Science and Business Media LLC, 2018

Publication

<% 1

39

Maryam Ahmad Sharifuddin, Sharifah Emilia Tuan Sharif, Hasnan Jaafar. "Prognostic Role of Matrix Metalloproteinase Expression in Meningioma: A Cross-Sectional Study.", Research Square, 2020

Publication

<% 1

EXCLUDE QUOTES ON

EXCLUDE ON

EXCLUDE MATCHES

< 5 WORDS

BIBLIOGRAPHY